

# Ectopic overexpression of the aluminum-induced protein gene from *Panax ginseng* enhances heavy metal tolerance in transgenic *Arabidopsis*

Moon-Gi Jang · Yu-Jin Kim · Gyu-Hwan Jang ·  
Johan Sukweenadhi · Woo-Saeng Kwon ·  
Deok-Chun Yang

Received: 6 February 2014 / Accepted: 17 May 2014 / Published online: 1 June 2014  
© Springer Science+Business Media Dordrecht 2014

**Abstract** Aluminum (Al), one of the most abundant metals in agricultural soils, significantly limits crop growth and productivity in acidic soil by inhibiting root elongation. Al ions, especially  $\text{Al}^{3+}$ , have a toxic effect on both plant and animal cells under low-pH conditions. We first isolated and characterized *aluminum-induced protein (AIP)* cDNA from a 4-year-old root of *Panax ginseng* Meyer. This cDNA encodes an open reading frame of 711 bp with a deduced amino acid sequence of 236 residues. The calculated molecular mass of the mature protein is approximately 58.9 kDa with a predicted isoelectric point of 5.13. The *Panax ginseng AIP (PgAIP)* contains a domain also present in *wheat aluminum-induced protein 7 (Wali7)* and shares homology with the AIPs of other species, including *Codonopsis* and *Arabidopsis*. The *PgAIP* gene was abundantly expressed in the plant's leaves and was up-regulated by Al exposure. The functional role of *PgAIP* in Al tolerance was further validated through its overexpression in *Arabidopsis*. Transgenic *Arabidopsis* plants overexpressing the *PgAIP* gene showed enhanced Al tolerance in terms of root growth when compared to wild-type plants, suggesting *PgAIP* is important in plant defense against Al toxicity. Confocal analysis of CFP-tagging *PgAIP* in *Arabidopsis* showed subcellular localization in the plasma membrane. Our results suggest that *PgAIP* in the plasma

membrane plays an important role in the protection of plant cells against heavy metal exposure.

**Keywords** Aluminum-induced protein · Gene expression · Heavy metal · *Panax ginseng* · *Wali7*

## Abbreviations

AIP	Aluminum-induced protein
Al	Aluminum
cDNA	Complementary DNA
EST	Expressed sequence tag
ORF	Open reading frame
qRT-PCR	Quantitative reverse transcription–polymerase chain reaction
HM	Heavy metal
Wali7	Wheat aluminum-induced protein 7

## Introduction

In many parts of the world, agricultural soils are contaminated with heavy metals that pose a serious health hazard to humans, animals, plants, and soil microorganisms (Ghnaya et al. 2010). Aluminum (Al) is the most abundant metal (Goodwin and Sutter 2009), composing approximately 7.5 % of the elements in the earth's crust (Haug and Foy 1984). Although Al is an important mineral in plant growth and development (Kim et al. 2004), it has a toxic effect under low-pH conditions (Ezaki et al. 2004). Present in over 50 % of the world's arable lands (von Uexküll and Mutert 1995), Al toxicity in acidic soils is a major factor limiting crop growth and productivity by inhibiting root elongation (Kochian 1995; Goodwin and Sutter 2009). The inhibition of root elongation impairs the uptake of water

**Electronic supplementary material** The online version of this article (doi:10.1007/s11240-014-0516-2) contains supplementary material, which is available to authorized users.

Moon-Gi Jang and Yu-Jin Kim have equally contributed to this work.

M.-G. Jang · Y.-J. Kim · G.-H. Jang · J. Sukweenadhi ·  
W.-S. Kwon · D.-C. Yang (✉)  
Department of Oriental Medicinal Materials and Processing,  
College of Life Science, Kyung Hee University,  
Suwon 449-701, Korea  
e-mail: dcyang@khu.ac.kr