


ISSN 1411-9420

Accepted by IIRCI
No. 13/2010/10/2010

Indonesian Journal of Chemistry

Vol. 12, No. 2, July, 2012



ISSN 1411-9420

Accredited by DIKTI
No. 51/DIKTI/Kep/2010

Indonesian Journal of Chemistry

Vol. 12, No. 2, July, 2012

MENCEKAM

Salinan/fotocopy sesuai dengan aslinya

Surabaya, _____


UNIVERSITAS SURABAYA

FAKULTAS TEKNIK

Dekan



Dr. Amelia Santoso





SJR

Scimago Journal & Country Rank

Enter Journal Title, ISSN or Publisher Name

[Home](#)[Journal Rankings](#)[Country Rankings](#)[Viz Tools](#)[Help](#)[About Us](#)

Indonesian Journal of Chemistry


Country [Indonesia](#) -  [SJR Ranking of Indonesia](#)**Subject Area and Category** [Chemistry](#)
[Chemistry \(miscellaneous\)](#)**Publisher** [Gadjah Mada University](#)**Publication type** Journals**ISSN** 14119420**Coverage** 2012-ongoing [Join the conversation about this journal](#)

8


H Index

Quartiles

Chemistry (miscellaneous)



SJR







Citations per document

Total Cites

Self-Cites





Indonesian Journal of Chemistry

Q3
Chemistry (miscellaneous)
best quartile

SJR 2018
0.22

powered by scimagojr.com

← Show this widget in your own website

Just copy the code below and paste within your html code:

```
<a href="https://www.scimagojr.com" style="color: #0070C0; text-decoration: none;">

```

M **Melanie Cornelia** 7 months ago

selamat pagi, saya dosen di Teknologi Pangan UPH....saya mau kirim article penelitian ke journal ini, rata-rata berapa lama review nya? terimakasih infonya

reply

M **Manton Rais** 12 months ago

Source details

Indonesian Journal of Chemistry

Open Access ⓘ

Scopus coverage years: from 2012 to Present

Publisher: Gadjah Mada University

ISSN: 1411-9420

Subject area: Chemistry: General Chemistry[View all documents >](#)[Set document alert](#)[Journal Homepage](#)

CiteScore 2018

0.95 ⓘ[Add CiteScore to your site](#)

SJR 2018

0.215 ⓘ

SNIP 2018

1.044 ⓘ[CiteScore](#) [CiteScore rank & trend](#) [CiteScore presets](#) [Scopus content coverage](#)CiteScore 2018 ▾Calculated using data from **30 April, 2019**

CiteScore rank ⓘ

$$0.95 = \frac{\text{Citation Count 2018}}{\text{Documents 2015 - 2017}^*} = \frac{153 \text{ Citations } >}{161 \text{ Documents } >}$$

*CiteScore includes all available document types

[View CiteScore methodology >](#)[CiteScore FAQ >](#)

Category	Rank	Percentile
Chemistry		
General Chemistry	#222/371	40th

[View CiteScore trends >](#)

CiteScoreTracker 2019 ⓘ

Last updated on *12 August, 2019*
Updated monthly

$$0.55 = \frac{\text{Citation Count 2019}}{\text{Documents 2016 - 2018}} = \frac{126 \text{ Citations to date } >}{229 \text{ Documents to date } >}$$

Metrics displaying this icon are compiled according to Snowball Metrics ↗, a collaboration between industry and academia.

About Scopus

[What is Scopus](#)
[Content coverage](#)
[Scopus blog](#)
[Scopus API](#)
[Privacy matters](#)

Language

[日本語に切り替える](#)
[切换到简体中文](#)
[切换到繁體中文](#)
[Русский язык](#)

Customer Service

[Help](#)
[Contact us](#)

ELSEVIER

[Terms and conditions ↗](#) [Privacy policy ↗](#)

Copyright © Elsevier B.V. ↗. All rights reserved. Scopus® is a registered trademark of Elsevier B.V.

We use cookies to help provide and enhance our service and tailor content. By continuing, you agree to the use of cookies.

RELX

Editor in Chief

Prof. Dr. Harno Dwi Pranowo, M.Si Email: harnodp@ugm.ac.id or harnoprano@yahoo.com
Department of Chemistry, Universitas Gadjah Mada
Sekip Utara, Yogyakarta Indonesia 55281, Tel/Fax (0062-274)-545188.
Website : <http://www.ijc.chemistry.ugm.ac.id>
Email : ijcugm@yahoo.com

Vice Editor in Chief

Prof. Dr. Mudasir, M.Eng Email: mudasir@ugm.ac.id or m_mudasir@hotmail.com
Drs. Iqmal Tahir, M.Si. Email: iqmal@ugm.ac.id or iqmal.tahir@yahoo.com

Editorial Board

Prof. Dr. Kama Wijaya, M.Eng. (*Physical/Material Chemistry*)
Prof. Dr. Jumina (*Organic Chemistry*)
Prof. Dr. Nuryono, MS (*Inorganic Chemistry*)
Dr. Ria Armunanto, M.Si. (*Computational/Physical Chemistry*)
Dr. Tri Joko Raharjo, M.Si. (*Biochemistry/Bioanalysis*)
Dr. Nurul Hidayat Aprilita (*Analytical/Environmental Chemistry*)

Advisory Editorial Board

Dr. Brian Williams (Adelaide University Australia)	Prof. Dr. Muhammad Idris Saleh (University Sains Malaysia)
Prof. Dr. Dr. Hc. Bernd M Rode (University of Innsbruck, Austria)	Dr. Hery Haerudin (Pertamina, Indonesia)
Dr. Dirk Bax (Utrecht University, Netherlands)	Prof. Dr. Iip Izul Falah (Universitas Gadjah Mada, Indonesia)
Prof. Dr. M. Gross (Louis Pasteur University, France)	Prof. Sri Juari Santosa, M.Eng, Ph.D. (Universitas Gadjah Mada, Indonesia)
Prof. Dr. Hardjono Sastroamidjojo (Universitas Gadjah Mada, Indonesia)	Prof. Dr. Endang Tri Wahyuni, MS (Universitas Gadjah Mada, Indonesia)
Prof. Dr. David. St. C. Black (University of New South Wales, Australia)	Prof. Dr. Bambang Rusdiarso, DEA (Universitas Gadjah Mada, Indonesia)
Prof. Dr. Max Lu (University of Queensland, Australia)	Prof. Dr. Wega Trisunaryanti, M.S., Ph.D. Eng (Universitas Gadjah Mada, Indonesia)
Prof. Dr. Buchori (Bandung Institute of Technology, Indonesia)	Prof. Dr. Sabirin Matsjeh (Universitas Gadjah Mada, Indonesia)
Prof. Dr. Abdul Ra'uf Pathong (Hasanudin University, Indonesia)	Dr. Winarto Haryadi, M.Si. (Universitas Gadjah Mada, Indonesia)
Prof. Dr. Naoki Yoshioka (Keio University, Japan)	
Assoc. Prof. Dr. Wan Ahmad Kamil Mahmood (University Sains Malaysia)	

Administrator

Dr. Akhmad Syoufian
Warakustarti Listyariwangi, A.Md
Djoko Prihandono

Robby Noor Cahyono, S.Si., M.Sc
Emie Kurniawati, SH

Aims and Scope

Indonesian Journal of Chemistry is an international journal covering all aspects of Chemistry, including *Chemical Education and Chemical Engineering*. The journal publishes original research papers, short communications, and review articles. The paper published in this journal implies that the work described has not been, and will not be published elsewhere, except in abstract, as part of a lecture, review or academic thesis.

Indonesian Journal of Chemistry (ISSN 1411-9420) is published by the Department of Chemistry, Faculty of Mathematics and Natural Sciences, Universitas Gadjah Mada, Sekip Utara, Yogyakarta, Indonesia. All orders accompanied by payment should be sent directly to The Department of Chemistry, Universitas Gadjah Mada. Annual subscription rate is IDR 150,000.00 (Java-Bali), IDR 200,000.00 (outside Java-Bali) (shipping included). Reprint order price is IDR 40,000.00 per article (shipping not included). Customers may make payments by transfer on **BNI 46 Universitas Gadjah Mada Filial (Emie Kurniawati, SH cq IJC, Account Nr. 0150173640, SWIFT code: BNINIDJAXXX)**.

CONTENTS

Transport of Phenol Through Inclusion Polymer Membrane (PIM) Using Copoly(eugenol-DVB) as Membrane Carriers by Agung Abadi Kiswandono, Dwi Siswanta, Nurul Hidayat Aprilita, and Sri Juari Santosa	105-112
Synthesis of Poly(acrylamide-co-acrylic Acid)-Starch based Superabsorbent Hydrogels by Gamma Radiation: Study Its Swelling Behavior by Erizal	113-118
Effect of Pyrolysis Temperatures on Composition and Electrical Conductivity of Carbosil Prepared from Rice Husk by Wasinton Simanjuntak, Simon Sembiring, and Kerista Sebayang	119-125
Kinetics Study on Nitration of Methyl Ricinoleate by Abdullah, Triyono, Wega Trisunaryanti, and Winarto Haryadi	126-130
Kinetic Investigation of Carbon Dioxide, Acetic Acid, Acrylic Acid Formation on Diluted and Leached MoVTeNb Catalyst by Restu Kartiko Widi	131-134
Theoretical Study on 15-crown-5 Complex with Some Metal Cations by Yahmin, Harno Dwi Pranowo, and Ria Armunanto	135-140
Construction and Optimization of Structure-Based Virtual Screening Protocols to Identify Cyclooxygenase-1 Inhibitors using Open Babel, Spores and Plants by Enade Perdana Istyastono	141-145
Synthesis, Characterization and Antioxidant Activity of 7-hydroxy-3',4'-dimethoxyflavone by Elfi Susanti VH, Sabirin Matsjeh, Tutik Dwi Wahyuningsih, Mustofa, and Tri Redjeki	146-151
Synthesis and Antiplasmodial Activity Testing of (1)-N-alkyl- and (1)-N-benzyl-6-nitro-1,10-phenanthroline Salts as New Potential Antimalarial Agents by Ruslin Hadanu, Sabirin Mastjeh, Jumina, Mustofa, Eti Nurwening Sholikhah, and Mahardika Agus Wijayanti	152-162
Rapid Solvent-Free Microwave Assisted Synthesis of Some N'-benzylidene Salicylic Acid Hydrazides by Tutuk Budiati, Stephanie D.A., and Elisabeth C. Widjajakusuma	163-166
Isolation Anthocyanin from Roselle Petals (<i>Hibiscus sabdariffa</i> L) and the Effect of Light on The Stability by Siti Nuryanti, Sabirin Matsjeh, Chairil Anwar, and Tri Joko Raharjo	167-171
In Vitro Activity and Comparative Studies of Some Organotin(IV) Benzoate Derivatives Against Leukemia Cancer Cell, L-1210 by Sutopo Hadi, Mita Rilyanti, and Suharso	172-177
Synthesis of 2-methoxy-4,6-di(prop-1-enyl) Phenol from Eugenol and Its Activity as an Antioxidant by Hernawan, Bambang Purwono, and Tutik Dwi Wahyuningsih	178-183
An Ester of 4-methoxy Cinnamic Acid Isolated from <i>Xylocarpus moluccensis</i> (Lamk) M. Roem (Meliaceae) by Tukiran, Septiani Setyo Cahyasari, and Kuniyoshi Shimizu	184-188
Synthesis of Zinc-difattyalkyldithiocarbamates and Their Antioxidant Activities by Komar Sutriah, Zainal Alim Mas'ud, and Tun Tedja Irawadi	189-194
Short Communication: Secondary Metabolite from Endophytic Fungi <i>Aspergillus niger</i> of the Stem Bark of Kandis Gajah (<i>Garcinia griffithii</i>) by Elfita, Muharni, Munawar, and Septa Aryani	195-200
Note: Determination of Inulin from Multivitamin Syrup Product by High Performance Liquid Chromatography with RI Detector by Yuni Retnaningtyas	201-205

KINETIC INVESTIGATION OF CARBON DIOXIDE, ACETIC ACID, ACRYLIC ACID FORMATION ON DILUTED AND LEACHED MoVTeNb CATALYST

Restu Kartiko Widi

Department of Chemical Engineering, Faculty of Engineering, University of Surabaya (UBAYA),
TG Building 6th floor, Jln. Raya Kalirungktu Tenggilis, Surabaya, 60293, Indonesia

Received January 24, 2012; Accepted February 8, 2012

ABSTRACT

Reaction kinetics for the oxidation of propane over diluted-leached MoVTeNb is described. This paper is focused on the study of determination of the orders of carbon dioxide, acetic acid and acrylic acid formation. Deep oxidation of propane to carbon dioxide is first order with respect to hydrocarbon, and partial order (0.31) with respect to oxygen. The selective oxidation of propane to acrylic acid is partial order (0.49) with respect to hydrocarbon and partial order (0.09) with respect to oxygen.

Keywords: propane oxidation; catalyst; selective oxidation; kinetic; acrylic acid

ABSTRAK

Telah dipaparkan kinetika reaksi oksidasi propana pada katalis MoVTeNb dengan pengembangan dan pembilasan. Artikel ini fokus pada studi penentuan orde reaksi pembentukan karbon dioksida, asam asetat dan asam akrilat. Oksidasi lanjut propana menjadi karbon dioksida mengikuti reaksi orde pertama terhadap hidrokarbon dan orde parsial (0,31) terhadap oksigen. Sedangkan oksidasi propana menjadi asam akrilat mengikuti orde parsial baik terhadap hidrokarbon maupun oksigen, masing-masing sebesar 0,49 dan 0,09.

Kata Kunci: oksidasi propana; katalis; oksidasi selektif; kinetika; asam akrilat

INTRODUCTION

The selective oxidation of light alkanes (C_1 - C_5), especially propane, to form the corresponding olefins, alcohols, aldehydes, anhydrides or acids has been largely studied in the past 20 years for two main reasons. The first one is economical since alkanes are much cheaper and more abundant than the corresponding olefins. The second one is fundamental since the way by which the alkane is activated and functionalized into an oxygenate remains unclear and challenging [1].

Many factors play a role in selective oxidation reactions on metal oxide catalysts and the complexity of such catalysts is widely recognized [2]. A number of assumptions on how the MoVTeNb catalyst works suggested [3-5]

In the previous article, it has been described the reaction kinetics and mechanism of propane partial oxidation to acrylic acid on diluted MoVTeNbOx mixed-oxide catalysts using high-throughput system (nanoflow reactor) [6]. This article more focus and deals with the similar reaction oxidation over diluted and leached MoVTeNbOx mixed-oxide catalysts. This paper is the second part and focused on the study of determination

of the orders of carbon dioxide, acetic acid and acrylic acid formation divided into two parts. Part one which is focused on the profile of products selectivity, the rate of propane disappearance and the rate of propene formation has been reported [7].

EXPERIMENTAL SECTION

Materials

Steam (demineralized water), Nitrogen (99.9%, MOX-Malaysian Oxygen Bhd), Propane (98.9%, MOX-Malaysian Oxygen Bhd), Oxygen (99.9%, MOX-Malaysian Oxygen Bhd).

Instrumentation

Nanoflow catalytic reactor on-line with gas chromatography (HP6890-Agilent Technology).


Procedure

The preparation of Diluted $Mo_1V_{0.30}Te_{0.23}Nb_{0.125}O_x$ and its characterization have been described in more detail before [8-9]. In addition, in this article the catalyst

* Corresponding author. Tel/Fax : +62-312981398/312981387
Email address : restu@ubaya.ac.id

Table 1. Apparent activation energy

Process	Activation energy (kJ. mol ⁻¹)
Propane disappearance	27.3
Propene formation	40.9
CO ₂ formation	177.3
Acetic acid formation	91.6
Acrylic acid formation	30.3

**Fig 1.** Rate of carbon dioxide formation with respect to propane concentration at 653, 663, 673, and 683 K

preparation is followed by leaching process using water for 1 h, and then it is dried and calcined according to the method described in the previous literature. The kinetic studies were conducted by analysis of the products at low propane and oxygen concentrations (lower than 10%), which is required for operating the reactor in a differential mode. This was achieved by introducing small amount of catalyst (0.09 g) diluted in SiO₂ to a total volume of 0.5 mL and using high total flow rates 15 mLN/min and Gas Hourly Space Velocity (GHSV) of 1800 h⁻¹. Gas hourly space velocity (GHSV) is defined as the ratio of the volumetric flow rate of reactants at standard conditions to the total catalyst volume yielding a quantity with units of inverse time [13]. By this definition, it means, the higher GHSV the higher volumetric flow rate is. As a consequence, the contact time between gas feed and catalyst surface (active site) will be less by increasing the GHSV value.

Only one of the reactants was varied at the time and the rest were chosen to be in excess and outside of the explosive limits for the mixture of propane and oxygen. The feed composition was propane, oxygen, and steam (water) balance in nitrogen. The study of the kinetic dependence on propane was performed by varying the propane concentration and maintaining the others. For the oxygen dependence, oxygen concentration was varied and concentrations of the others were maintained. Finally, for the water dependence, water concentration was varied and concentrations of the others were maintained. The effect of reactant concentration was studied at four different temperatures, i.e., 653, 663, 673, and 683 K. Particle size of catalytic material was 212-425 μm or less [6,9].

For all experimental results, the reaction rate and the concentration of reactants are calculated using Equations shown below:

$$\text{Reaction rate (mol/h.g}_{\text{cat}}) = \frac{\% \text{vol} \times \text{total gas flow} \times d \times 60}{M_r \times \text{catalyst weight}} \quad (1)$$

$$\text{Concentration (mol/L)} = (\% \text{vol} \times d \times 1000) / M_r \quad (2)$$

where, %vol is concentration of product in % (v/v); total gas flow is the total flow of product stream from the reactor in mLN/min; M_r and d are molecular weight and density of the product; and catalyst weight is the weight of catalyst in gram.

The activation energy is determined experimentally by carrying out the reaction at several different temperatures. After taking the natural logarithm of equation

$$\ln k = \ln A - \frac{E_a}{R} \left(\frac{1}{T} \right) \quad (3)$$

RESULT AND DISCUSSION

The structure and surface texturing of MoVTeNb oxide catalyst particles composed of two phases referred to in the literature as M1 and M2. The M1 phase is the cluster that active as catalyst for the conversion of propane to acrylic acid [10-11]. Chemical composition of the catalyst surface is modified by treatment in water to obtain a significant increment in yield of acrylic acid. The process is called as leaching. It means the leaching process using water for 1 h support to release the M2 phase from the catalyst. It can be expected that the catalyst more active since only M1 phase stay in the catalyst. This phenomenon has been reported clearly by Jacob and coworkers [8]. The paper showed the SEM images of diluted catalyst before and after leaching and it showed that the distribution of catalyst was improved by leaching. The paper also showed high-resolution images of the termination of M1. It was clearly observed such structural rearrangement to exist at the surface of the M1 and the M2 phase [8]. The reproducibility of catalytic activity test has been carried out by repeated the experiments under the same condition and showed very good reproducibility in terms of both the propane conversion and acrylic acid selectivity [9].

In part one, it has been described the product selectivity profiles for propane oxidation over diluted and leached MoVTeNb oxide catalyst under reactor differential mode. According to the data, the reaction can be proposed by the following reaction $C_3H_8 + O_2 \rightarrow$ products, where the products are propene, acrylic acid, acetic acid, and carbon dioxide [7].

Table 1 depicts the activation energy which is needed in the total reaction, i.e. oxidation of propane to acrylic acid.




Fig 2. Rate of carbon dioxide formation with respect to oxygen concentration at 653, 663, 673, and 683 K




Fig 3. Rate of acetic acid formation with respect to propane concentration at 653, 663, 673, and 683 K

Rate of CO₂ and acetic acid formation

Fig. 1 shows that the rate of carbon dioxide formation is first order with respect to propane concentration at all reaction temperatures. It is strong indicated that this product is formed during the gas reactant (propane) adsorbed on the catalyst surface (not in the gas phase), it means that the carbon dioxide is formed by catalytic surface reaction. Fig. 2 shows that the rate of partial oxidation of propane to carbon dioxide is partial order (0.31) with respect to oxygen concentration. It is believed that this product is produced by direct combustion of propane. But it is also possible that the product is produced from further oxidation of propane or intermediate product (acetic acid), since the order differ compare to that of propane disappearances. But, the high activation energy (177.3 kJ/mol) which is required for the formation of carbon dioxide over this catalyst strongly indicated that carbon dioxide originates from a direct combustion of propane and not from over-oxidation of acetic acid. The total oxidation of the desired product acrylic acid is a less likely contribution as its




Fig 4. Rate of acetic acid formation with respect to oxygen concentration at 653, 663, 673, and 683 K




Fig 5. Rate of acrylic acid formation with respect to propane concentration at 653, 663, 673, and 683 K

formation and desorption requires smaller activation energies.

This description is supported by Fig. 3 and 4. The figures show that the acetic acid formation from oxidation of propane is not a linear function with respect to propane concentration and oxygen concentration at all temperatures. These indicate that the production of acetic acid is hindered by desorption from catalyst. During the hinder process, propane can be further oxidizing to produce carbon dioxide. The high activation energy describe that the acetic acid production is quite difficult. This is also support that acetic acid production is strongly hindered by desorption from the catalyst surface. In addition, extrapolation to zero propane concentration indicates that the curve does not pass through the origin of the graph; their rate of appearance is zero at a propane concentration is about 0.15 mol/L. The reasons for this at low propane concentration (0.15 mol/L), an intermediate product or propane does not convert to acetic acid but it convert to other products, i.e. over-oxidation to produce carbon dioxide.




Fig 6. Rate of acrylic acid formation with respect to oxygen concentration at 653, 663, 673, and 683 K

Rate of acrylic acid (AA) formation

Fig. 5 and 6 depict that the acrylic acid formation from oxidation of propane is not a linear function with respect to propane concentration and oxygen concentration at all temperatures. In addition, extrapolation to zero propane concentration indicates that the curve does not pass through the origin of the graph; their rate of appearance is zero at a propane concentration of about 0.05 – 0.075 mol/L. The reasons for this is that the experimentally determined rate for a product (i.e. AA) is the difference between the true rate of formation and the rate of intermediate disappearance in deep oxidation of this intermediate. This indicates that a kinetically intermediate must exist before the formation of AA. It is believed that the intermediate is acrolein [5,10,12]. Because of acrolein is very easy to convert to AA, acrolein can not be detected during the reaction. This description is supported by the activation energy of AA formation which is quite small, indicate that AA quite easy to produce from acrolein. This is also describe that the formation of AA is easier than formation of acetic acid (require very high activation energy). As the activation energy for total combustion is so high it is assumed that the possibility of a fast consecutive total oxidation seems less likely than the existence of a difficult-to-form intermediate before the formation of AA.

CONCLUSION

The result supported the argument that the structure and surface texturing of MoVTenb oxide catalyst particles composed of two phases referred to in

the literature as M1 and M2 [8-9,12]. The M1 phase is the cluster that active as catalyst for the conversion of propane to acrylic acid. The leaching process using water for 1 h support to release the M2 phase from the catalyst. The very high activation energy of carbon dioxide formation indicates that the M2 which is responsible for total oxidation release during the leaching process. The result also shows that AA quite easy produce from intermediate since the activation energy quite low.

ACKNOWLEDGEMENT

The research has been carried out in the COMBICAT Laboratory, University of Malaya, Kuala Lumpur, Malaysia.

REFERENCES

1. Boizumault-Moriceau, P., Pennequin, A., Grzybowska, B., and Barbaux, Y., 2003, *Appl. Catal. A*, 245, 55–67.
2. Grzybowska Swierkosz, B., 2000, *Top. Catal.*, 11-12, 23–42.
3. Grzybowska-Swierkosz, B., 2002, *Top. Catal.*, 21, 175
4. Novakova, E.K., Vedrine, J.C., and Derouane, E.G., 2002, *J. Catal.*, 211, 226–234.
5. Vedrine, J.C., Novakokva, E.K., and Derouane, E.G., 2003, *Catal. Today*, 81, 2, 247–262.
6. Widi, R.K., Hamid, S.B.A., and Schlogl, R., 2009, *React. Kinet. Catal. Lett.*, 98, 2, 273–286.
7. Widi, R.K., 2010, *Indo. J. Chem.*, 10, 2, 172–176.
8. Jacob, B.W., Olaf, T., Fazliana, A.H., Annette, T., Ute, W., Dang, S.S., Restu, K.W., Sharifah, B.A.H., and Robert, S., 2006, *Top Catal.*, 39, 1-2, 51–61.
9. Widi, R.K., Hamid, S.B.A., and Schlogl, R., 2008, *J. Nat. Gas Chem.*, 17, 2, 130–134.
10. Lin, M.M., 2001, *Appl. Catal. A*, 207, 1–7.
11. Botela, P., Solsona, B., Martinez-Arias, A., and Lopez-Nieto, J.M., 2001, *Catal. Lett.*, 74, 3-4, 149–155.
12. Lin, M., Desai, T.B., Kaiser, F.W., and Klugherz, P.D., 2000, *Catal. Today*, 61, 223–229.
13. Fogler, H.S., 1999, *Elements of Chemical Reaction Engineering*, 3rd ed., Prentice-Hall Inc., New Jersey, 126.