

ISSN: 2088-5334

Vol 8, No 3 (2018)

International Journal on Advanced Science Engineering Information Technology

Available online at
<http://ijaseit.insightsociety.org>

Advanced Science Image processing image retrieval microprocessor
Back Projection fuzzy Engineering alumina Gait analysis Numeric
Mobile Agent Interpolation Soft Tissue hydrothermal stability Outlier line
Sampling Theorem Magnetic field porous structure acidity Nitro
Parsimony Database Server Robot Operation Life Cycle Cost
Graphical User Interface Virus Compensatory Strategy Energy Saving
Current shaping Phylogenetic Analysis Sensor Networks Image enhancement
Metamorphic Virus neural network Predictive Tools Energy Optimization
parallel port Obfuscation Techniques Admixture Router Image matching
Power factor catalyst Construction Delay Predicted Strength Regression Analysis
Boost regulator breast phantom Effects of Delay Support Vector Machine
Information Technology soybean Current harmonics Surveillance








SJR

Scimago Journal & Country Rank

Enter Journal Title, ISSN or Publisher Name

[Home](#)[Journal Rankings](#)[Country Rankings](#)[Viz Tools](#)[Help](#)[About Us](#)

International Journal on Advanced Science, Engineering and Information Technology

Country	Indonesia -  SJR Ranking of Indonesia	11 H Index
Subject Area and Category	Agricultural and Biological Sciences Agricultural and Biological Sciences (miscellaneous) Computer Science Computer Science (miscellaneous) Engineering Engineering (miscellaneous)	
Publisher	INSIGHT - Indonesian Society for Knowledge and Human Development	
Publication type	Journals	
ISSN	20885334, 24606952	
Coverage	2015-ongoing	
Scope	International Journal on Advanced Science, Engineering and Information Technology (IJASEIT) is an international peer-reviewed journal dedicated to interchange for the results of high quality research in all aspect of science, engineering and information technology. The journal publishes state-of-art papers in fundamental theory, experiments and simulation, as well as applications, with a systematic proposed method, sufficient review on previous works, expanded discussion and concise conclusion. As our commitment to the advancement of science and technology, the IJASEIT follows the open access policy that allows the published articles freely available online without any subscription	
	Homepage How to publish in this journal Contact  Join the conversation about this journal	


Quartiles




Agricultural and Biological Sciences (miscellaneous)

Computer Science (miscellaneous)


SJR




Citations per document




Total Cites Self-Cites




External Cites per Doc Cites per Doc




% International Collaboration



Citable documents Non-citable documents



Cited documents Uncited documents



Show this widget in your own website

Just copy the code below and paste within your html code:

Source details

International Journal on Advanced Science, Engineering and Information Technology

Open Access ⓘ

Scopus coverage years: from 2015 to Present

Publisher: INSIGHT - Indonesian Society for Knowledge and Human Development

ISSN: 2088-5334 E-ISSN: 2460-6952

Subject area: Agricultural and Biological Sciences: General Agricultural and Biological Sciences
Engineering: General Engineering Computer Science: General Computer Science

[View all documents >](#) [Set document alert](#) [Journal Homepage](#)

CiteScore 2018
1.07 ⓘ
[Add CiteScore to your site](#)

SJR 2018
0.230 ⓘ

SNIP 2018
0.802 ⓘ

[CiteScore](#) [CiteScore rank & trend](#) [CiteScore presets](#) [Scopus content coverage](#)

CiteScore 2018 ▾

Calculated using data from 30 April, 2019

1.07 = $\frac{\text{Citation Count 2018}}{\text{Documents 2015 - 2017}^*}$

678 Citations >

634 Documents >

*CiteScore includes all available document types

[View CiteScore methodology >](#) [CiteScore FAQ >](#)

CiteScoreTracker 2019 ⓘ

Last updated on 12 August, 2019
Updated monthly

0.51 = $\frac{\text{Citation Count 2019}}{\text{Documents 2016 - 2018}}$

447 Citations to date >

877 Documents to date >

Metrics displaying this icon are compiled according to Snowball Metrics ↗ , a collaboration between industry and academia.

CiteScore rank ⓘ

Category	Rank	Percentile
Agricultural and Biological Sciences	#70/185	62nd
General Agricultural and Biological Sciences		
Engineering		

[View CiteScore trends >](#)

About Scopus

- What is Scopus
- Content coverage
- Scopus blog
- Scopus API
- Privacy matters

Language

- 日本語に切り替える
- 切换到简体中文
- 切换到繁體中文
- Русский язык

Customer Service

- Help
- Contact us

ELSEVIER

Terms and conditions ↗

Privacy policy ↗

Copyright © Elsevier B.V. All rights reserved. Scopus® is a registered trademark of Elsevier B.V.

We use cookies to help provide and enhance our service and tailor content. By continuing, you agree to the use of cookies.

RELX

Editorial Team

Editor in Chief :

Sri Atmaja Putra, (Scopus ID: 26321207400), Muhammadiyah Yogyakarta University, INDONESIA

Associate Editors :

Taufik, (Scopus ID: 23670809800), California Polytechnic State University, USA
Moustafa M Eissa, (Scopus ID: 35581693900), Helwan University, EGYPT
Wan Mohtar Wan Yusoff, (Scopus ID: 15019967700), Univ. Kebangsaan Malaysia, MALAYSIA
Halimah Badioze Zaman, (Scopus ID: 25825801600), Univ. Kebangsaan Malaysia, MALAYSIA
Son Radu, (Scopus ID: 7005251005), Universiti Putra Malaysia, Malaysia, MALAYSIA
Mohd Razi Ismail, (Scopus ID: 25957691400), Universiti Putra Malaysia, MALAYSIA
Takashi Oku, (Scopus ID: 56275094900), Prefectural University of Hiroshima, JAPAN
Kohei Nakano, (Scopus ID: 7402011766), Gifu University, JAPAN
Nurul Huda, (Scopus ID: 6701695514), Universiti Sultan Zainal Abidin, MALAYSIA
Yandra Arkeman, (Scopus ID: 55946558300), Bogor Agriculture University, INDONESIA
Sate Sampattagul, (Scopus ID: 7801640861), Chiangmai University, THAILAND
Peeyush Soni, (Scopus ID: 9248907800), Asian Institute of Technology, THAILAND
Yolanda Lechon Perez, (Scopus ID: 6602826000), Ciemat, Madrid, SPAIN
Gabriele Arcidiacono (Scopus ID: 56656284600), G. Marconi University, ITALY
Alessandra Pieroni (Scopus ID: 25929524500). , Marconi International University, Florida - USA
Nguyen Hay , (Scopus ID: 15834645900) Nong Lam University. VIETNAM
Rita Muhamad Awang, (Scopus ID: 55957782400), Universiti Putra Malaysia, MALAYSIA
Anton S Prabuwno, (Scopus ID: 18134309800), King Abdulaziz Univ, SAUDI ARABIA
P Mangala C S De Silva, (Scopus ID: 7006461145), University of Ruhuna, SRI LANKA
Bich Huy Nguyen, (Scopus ID: 36191086100), Nong Lam University, VIETNAM
Paul Kristiansen, (Scopus ID: 23097563600), University of New England, AUSTRALIA
Amitava Basu, (Scopus ID: 21833738300), Bidhan Chandra Krishi Vidyalaya, INDIA
Haitham Alali, (Scopus ID: 49963007000), Amman Arab University, JORDAN
Shahrul Azman Mohd Noah, (Scopus ID: 35087633200), Universiti Kebangsaan Malaysia, MALAYSIA
Luca Di Nunzio (Scopus ID: 57195199010), University of Rome Tor Vergata, ITALY

Managing Editor :

Rahmat Hidayat, (Scopus ID: 57191265401) Politeknik Negeri Padang, INDONESIA

Editors :

Nurhamidah, (Scopus ID: 57191636504), Andalas University, INDONESIA
Ario Betha Juansilfero, (Scopus ID: 57189369470), Kobe University, JAPAN
Ifa Puspasari, (Scopus ID: 55180322800), National University of Malaysia - MALAYSIA
Zairi Ismael Rizman, (Scopus ID: 36959761800), Universiti Teknologi MARA (UiTM) (Terengganu) MALAYSIA
Shahreen Kasim, (Scopus ID: 36155431900), Universiti Tun Hussein Onn - MALAYSIA
Chi-Hua Chen, (Scopus ID: 35799698800), National Chiao Tung University, TAIWAN
Gusri Yaldi, (Scopus ID: 55359203100), Polytechnics State of Padang, INDONESIA
Abrar Ismardi, (Scopus ID: 26633102900), Telkom University - INDONESIA

Vol. 8 (2018) No. 3

Articles

Methods to Characterize Fly Ash Quality in the Field

- Triwulan, Khorin Agus Priadana, Januarti Jaya Ekaputri
pages: 646-651 [Full text](#) [DOI:10.18517/ijaseit.8.3.4312](https://doi.org/10.18517/ijaseit.8.3.4312)

Finite Element Analysis of Osteosynthesis Miniplate for the Reconstruction of Parasymphyseal Compound Fracture

Muslim Mahardika, Romario Aldrian Wicaksono, Maria Goreti Widiastuti, Budi Arifvianto, - Suyitno
pages: 652-656 [Full text](#) [DOI:10.18517/ijaseit.8.3.4465](https://doi.org/10.18517/ijaseit.8.3.4465)

Neutronic Analysis of a Thorium-Fueled Reduced Moderation Boiling Water Reactor

Dayu Fajrul Falaakh, Alexander Agung, Andang Widi Harto
pages: 657-663 [Full text](#) [DOI:10.18517/ijaseit.8.3.2834](https://doi.org/10.18517/ijaseit.8.3.2834)

Tobermorite Microstructure Resulted from Reaction of Low Content of Silica of Rice Husk Ash and Quicklime Mixture

Bakri Bakri
pages: 664-670 [Full text](#) [DOI:10.18517/ijaseit.8.3.1892](https://doi.org/10.18517/ijaseit.8.3.1892)

Identification of Slope Stability Analysis

Ika Juliantina, Yulindasari Sutejo, Ratna Dewi, Bimo Brata Adhitya, Reffanda Kurniawan Rustam
pages: 671-677 [Full text](#) [DOI:10.18517/ijaseit.8.3.2685](https://doi.org/10.18517/ijaseit.8.3.2685)

Preliminary Investigations for Policy Framework to Regulate the Utilisation of Residual Strength of Demolition Waste Aggregate in Cement Concrete Mix

M. R. Raja Shekhara, Reshma. E. K.
pages: 678-684 [Full text](#) [DOI:10.18517/ijaseit.8.3.4175](https://doi.org/10.18517/ijaseit.8.3.4175)

Capillary Shock Phenomenon of Groundwater at the Beginning of Rainy Season

Darwis Panguriseng, Abd. Rakhim Nanda
pages: 685-693 [Full text](#) [DOI:10.18517/ijaseit.8.3.3818](https://doi.org/10.18517/ijaseit.8.3.3818)

Conversion of Polypropylene Plastic Waste Into Liquid Fuel with Catalytic Cracking Process Using Al₂O₃ as Catalyst

Devi Rachmadena, Muhammad Faizal, Muhammad Said
pages: 694-700 [Full text](#) [DOI:10.18517/ijaseit.8.3.2586](https://doi.org/10.18517/ijaseit.8.3.2586)

Farm Machinery Development and Utilization System Policies for Small-Scale Rice Farming

Ujang Paman, Saipul Bahri, - Asrol, - Khairizal, Hajry Arief Wahyudy
pages: 701-707 [Full text](#) [DOI:10.18517/ijaseit.8.3.1758](https://doi.org/10.18517/ijaseit.8.3.1758)

Influence of Moisture Content to the Physical Properties of Unhusk Rice Grain

Renny Eka Putri, Santosa Santosa, Muhammad Makky
pages: 708-713 [Full text](#) [DOI:10.18517/ijaseit.8.3.3455](https://doi.org/10.18517/ijaseit.8.3.3455)

Parasitizations Levels and Temperature Tolerance of Rice Bug (*Leptocorisa oratorius* Fabricius) Egg Parasitoids: Mass Rearing for Biological Control

Fri Maulina, Novri Nelly, Hidrayani Hidrayani, Hasmiandy Hamid
pages: 714-719 [Full text](#) [DOI:10.18517/ijaseit.8.3.1241](https://doi.org/10.18517/ijaseit.8.3.1241)

Oil-Palm Plantation Identification from Satellite Images Using Google Earth Engine

Supattra Puttinaovarat, Paramate Horkaew
pages: 720-726 [Full text](#) [DOI:10.18517/ijaseit.8.3.2415](https://doi.org/10.18517/ijaseit.8.3.2415)

Soil Mounding Practices Towards Yield Performances of Oil Palm on Peat Soil

Shampazuraini binti Samsuri, Husein bin Abdul Gani, Siti Aminah Binti Wahab, Nurul Farhana Hazira H
pages: 727-731 [Full text](#) [DOI:10.18517/ijaseit.8.3.2751](https://doi.org/10.18517/ijaseit.8.3.2751)

An Immunological-Based Simulation: A Case Study of Risk Concentration for Mobile Spam Context Assessment

Kamahazira Zainal, Mohd Zalisham Jali
pages: 732-742 [Full text](#) [DOI:10.18517/ijaseit.8.3.2719](https://doi.org/10.18517/ijaseit.8.3.2719)

Implementation of Otsu's Method in Vein Locator Devices

Gwo-Jia Jong, Hendrick Hendrick, Zhi-Hao Wang, Dedi Kurniadi, Aripriharta Aripriharta, Gwo-Jiun Horng
pages: 743-748 [Full text](#) [DOI:10.18517/ijaseit.8.3.4414](https://doi.org/10.18517/ijaseit.8.3.4414)

A Novel Packet Scheduling with Channel-Aware Algorithm for Multi-Service Flow in the LTE Network

Ida Nurcahyani, Sisdarmanto Adinandra
pages: 749-755 [Full text](#) [DOI:10.18517/ijaseit.8.3.5579](https://doi.org/10.18517/ijaseit.8.3.5579)

Component-connected Feature for Signature Identification

Naeli Umniati, Achmad Benny Mutiara, Tubagus Maulana Kusuma, Suryarini Widodo
pages: 756-761 [Full text](#) [DOI:10.18517/ijaseit.8.3.2880](https://doi.org/10.18517/ijaseit.8.3.2880)

Different Applications of the Genetic Mutation Operator for Symetric Travelling Salesman Problem

Velin Kralev
pages: 762-770 [Full text](#) [DOI:10.18517/ijaseit.8.3.4867](https://doi.org/10.18517/ijaseit.8.3.4867)

Continuous Innovation: A Literature Review and Future Perspective

Benny Lianto, M. Dachyar, Tresna Priyana Soemardi
pages: 771-779 [Full text](#) [DOI:10.18517/ijaseit.8.3.4359](https://doi.org/10.18517/ijaseit.8.3.4359)

Synthesis and Optimization of Acrylic-N-Acryloxysuccinimide Copolymer Microspheres

Alizar Ulianas, Nurlily Nurlily, Lee Yook Heng, Tan Ling Ling
pages: 780-784 [Full text](#) [DOI:10.18517/ijaseit.8.3.3336](https://doi.org/10.18517/ijaseit.8.3.3336)

Tree Vegetation Structure at the Realolo Village Bantimurung Bulusaraung National Park Maros District

Muhammad Wiharto, Siti Fatmah Hiola, Syamsinar S, Muhammad Wijaya, Hamka L
pages: 785-791 [Full text](#) [DOI:10.18517/ijaseit.8.3.4725](https://doi.org/10.18517/ijaseit.8.3.4725)

Compression Behavior of Fibrous Peat Stabilized with Admixtures of Lime CaCO₃+Rice Husk Ash and Lime CaCO₃+Fly Ash

Noor Endah Mochtar, Faisal Estu Yulianto
pages: 792-798 [Full text](#) [DOI:10.18517/ijaseit.8.3.4317](https://doi.org/10.18517/ijaseit.8.3.4317)

Comparison of WF & Tubular Links on the Eccentrically Brace Frame System due to Cyclic Loads

Budi Suswanto, Aniendhita Rizki Amalia, Endah Wahyuni, Najibullah Al Farisy
pages: 799-804 [Full text](#) [DOI:10.18517/ijaseit.8.3.4337](https://doi.org/10.18517/ijaseit.8.3.4337)

Development of Thermal Insulation Material Using Coconut Fiber to Reuse Agricultural Industrial Waste

Ryushi Kimura, Masato Ohsumi, Lusi Susanti
pages: 805-810 [Full text](#) [DOI:10.18517/ijaseit.8.3.4610](https://doi.org/10.18517/ijaseit.8.3.4610)

Disinfection of Escherichia coli Bacteria Using Combination of Ozonation and Hydrodynamic Cavitation Method with Venturi Injector

Eva Fathul Karamah, Fitri Amalia, Fitri Amalia, Setijo Bismo, Rioneli Ghaudenson, Setijo Bismo, Rioneli Ghaudenson

pages: 811-817 [Full text](#) [DOI:10.18517/ijaseit.8.3.3922](https://doi.org/10.18517/ijaseit.8.3.3922)

Adsorption of Phosphate in Aqueous Solutions Using Manganese Dioxide

Yunus Fransiscus, Restu K. Widi, Gracia O. Aprilasti, Marta D. Yuharma

pages: 818-824 [Full text](#) [DOI:10.18517/ijaseit.8.3.3866](https://doi.org/10.18517/ijaseit.8.3.3866)

The Effect of TiO₂ Particles Addition on The Characteristics of Polysulfone Membrane

Nurul Nadiyah Said, Fazlena Hamzah, Nur Azrini Ramlee, Nur Najwa Yunus

pages: 825-831 [Full text](#) [DOI:10.18517/ijaseit.8.3.3901](https://doi.org/10.18517/ijaseit.8.3.3901)

Physicochemical Properties of Duck Feet Collagen with Different Soaking Time and Its Application in Surimi

Chong Hui Theng, Nurul Huda, Nik Aisyah Nik Muhammad, Chatarina Wariyah, Haslaniza Hashim

pages: 832-841 [Full text](#) [DOI:10.18517/ijaseit.8.3.2670](https://doi.org/10.18517/ijaseit.8.3.2670)

Utilization of Empty Fruit Bunch Fiber of Palm Oil Industry for Bio-Hydrogen Production

Eka Sari, Mohammad Effendy, Nufus Kanani, - Wardalia, - Rusdi

pages: 842-848 [Full text](#) [DOI:10.18517/ijaseit.8.3.3985](https://doi.org/10.18517/ijaseit.8.3.3985)

Responses Growth and Yield of Three Shallot Cultivars in Sandy Coastal Land with PGPR (Plant Growth Promoting Rhizobacteria)

Sumiyati Tuhuteru, Endang Sulistyaningsih, Arif Wibowo

pages: 849-855 [Full text](#) [DOI:10.18517/ijaseit.8.3.1093](https://doi.org/10.18517/ijaseit.8.3.1093)

Geological Mapping and Assessment for Measurement the Electric Grid Transmission Lines in West Sumatera Area, Indonesia

Husnul Kausarian, Batara Batara, Dewandra Bagus Eka Putra, Adi Suryadi, Muhammad Zainuddin Lubis

pages: 856-862 [Full text](#) [DOI:10.18517/ijaseit.8.3.4069](https://doi.org/10.18517/ijaseit.8.3.4069)

Tectonic Geomorphology of Upper Cimanuk Drainage Basin, West Java, Indonesia

Emi Sukiyah, Edy Sunardi, Nana Sulaksana, P. P. Raditya Rendra

pages: 863-869 [Full text](#) [DOI:10.18517/ijaseit.8.3.5441](https://doi.org/10.18517/ijaseit.8.3.5441)

Geotechnical Engineering Aspect Related to Pidie Jaya–Aceh Earthquake Disaster and Mitigation

- Munirwansyah, Reza Pahlevi Munirwan, Halida Yunita

pages: 870-875 [Full text](#) [DOI:10.18517/ijaseit.8.3.4189](https://doi.org/10.18517/ijaseit.8.3.4189)

Evaluation of Seismic Analysis Procedures for Concrete Moment-Resistant Frames with Horizontal Re-entrant Corners Irregularity

Anis S Shatnawi, Mazen Musmar, Laith I Gharaibeh

pages: 876-881 [Full text](#) [DOI:10.18517/ijaseit.8.3.5851](https://doi.org/10.18517/ijaseit.8.3.5851)

An Experimental Study of Bending Behaviour of Double Channel and Hollow Sections of Light Gauge Steel

Sabril Haris, Andrey Prasetyo, Rendy Thamrin, Hazmal Herman

pages: 882-888 [Full text](#) [DOI:10.18517/ijaseit.8.3.4192](https://doi.org/10.18517/ijaseit.8.3.4192)

Methodology on Empirical Study in Determining Critical Energy Efficiency Factors for Building Structural Components

Saeed Balubaid, Rosli Mohammed Zin, Shaik Hussein Mydin

pages: 889-896 [Full text](#) [DOI:10.18517/ijaseit.8.3.4191](https://doi.org/10.18517/ijaseit.8.3.4191)

Experimental Study on High Strength Steel-Fiber Concrete

Sisi Nova Rizkiani, James Saputra, Johannes Adhijoso Tjondro

pages: 897-903 [Full text](#) [DOI:10.18517/ijaseit.8.3.3962](https://doi.org/10.18517/ijaseit.8.3.3962)

Synthesis of Tobermorite Structure with Non-traditional Silica Components

Vit Cerny, Jan Fleischhacker, Magdalena Kocianova, Rostislav Drochytka

pages: 904-910 [Full text](#) [DOI:10.18517/ijaseit.8.3.5853](https://doi.org/10.18517/ijaseit.8.3.5853)

Assessment of Time and Costs of Two Formwork Methodologies in the Philippines using BIM Simulation

Jason Maximino C. Ongpeng

pages: 911-917 [Full text](#) [DOI:10.18517/ijaseit.8.3.5838](https://doi.org/10.18517/ijaseit.8.3.5838)

Airline Service Quality Analysis Using Integration of Fuzzy Servqual, PGCV Index, and TRIZ Methods in Indonesian Full-Service Carrier Airlines

Diana Puspita Sari, Dyah Ika Rinawati, - Midiawati

pages: 918-923 [Full text](#) [DOI:10.18517/ijaseit.8.3.3889](https://doi.org/10.18517/ijaseit.8.3.3889)

A Simple Transistors Width Adjustment Method on CMOS Transmission Gate Switch to Reduce Hold Error of S/H Circuit

Agung Setiabudi, Hiroki Tamura, Koichi Tanno

pages: 924-929 [Full text](#) [DOI:10.18517/ijaseit.8.3.4329](https://doi.org/10.18517/ijaseit.8.3.4329)

CDM Based Servo State Feedback Controller with Feedback Linearization for Magnetic Levitation Ball System

Alfian Ma'arif, Adha imam Cahyadi, Oyas Wahyunggoro
pages: 930-937 [Full text](#) [DOI:10.18517/ijaseit.8.3.1218](https://doi.org/10.18517/ijaseit.8.3.1218)

Delay Tolerant Energy Efficient protocol for Inter-BAN Communication in Mobile Body Area Networks

Aarti Sangwan, Partha Pratim Bhattacharya
pages: 938-948 [Full text](#) [DOI:10.18517/ijaseit.8.3.4502](https://doi.org/10.18517/ijaseit.8.3.4502)

Mobile Forensic Tools Evaluation for Digital Crime Investigation

Rusydi Umar, Imam Riadi, Guntur Maulana Zamroni
pages: 949-955 [Full text](#) [DOI:10.18517/ijaseit.8.3.3591](https://doi.org/10.18517/ijaseit.8.3.3591)

A Study of Delay and Data Traffic of IEEE 802.15.4 ZigBee-Based WSN in a Smart Home

Naseem Kadhim Baqer, Ameen Mohammed-Taqy Al-Modaffer, Esam A. AlKaldy
pages: 956-962 [Full text](#) [DOI:10.18517/ijaseit.8.3.5483](https://doi.org/10.18517/ijaseit.8.3.5483)

Developing an Artificial Neural Network Algorithm for Generalized Singular Value Decomposition-based Linear Discriminant Analysis

Rolysent K Paredes, Ariel M Sison, Ruji P Medina
pages: 963-969 [Full text](#) [DOI:10.18517/ijaseit.8.3.5677](https://doi.org/10.18517/ijaseit.8.3.5677)

Efficient Handwritten Digit Classification using User-defined Classification Algorithm

R. Vijaya Kumar Reddy, U. Ravi Babu
pages: 970-979 [Full text](#) [DOI:10.18517/ijaseit.8.3.5397](https://doi.org/10.18517/ijaseit.8.3.5397)

Data Mining for Detecting E-learning Courses Anomalies: An Application of Decision Tree Algorithm

Fatiha Elghibari, Rachid Elouahbi, Fatima El Khoukhi
pages: 980-987 [Full text](#) [DOI:10.18517/ijaseit.8.3.2756](https://doi.org/10.18517/ijaseit.8.3.2756)

Study of Convective Cloud Lifetime and Movement Using Radar Image and ECMWF Model

Mochammad Donny Anggoro, Bagus Pramujio
pages: 988-997 [Full text](#) [DOI:10.18517/ijaseit.8.3.4212](https://doi.org/10.18517/ijaseit.8.3.4212)

Optimization of Fingerprint Indoor Localization System for Multiple Object Tracking Based on Iterated Weighting Constant - KNN Method

Asti Putri Rahmadini, Prima Kristalina, Amang Sudarsono
pages: 998-1007 [Full text](#) [DOI:10.18517/ijaseit.8.3.6086](https://doi.org/10.18517/ijaseit.8.3.6086)

Adsorption of Phosphate in Aqueous Solutions Using Manganese Dioxide

Yunus Fransiscus^{1,2}, Restu K. Widi², Gracia O. Aprilasti², Marta D. Yuharma²

¹ Centre for Environmental Studies, The University of Surabaya, Raya Kalirungkut, Surabaya, 60293, Indonesia
E-mail:yunus@staff.ubaya.ac.id

² Chemical Engineering Department, The University of Surabaya, Raya Kalirungkut, Surabaya, 60293, Indonesia

Abstract. Effort to remove phosphate from aquatic ecosystem is of great interest, not only for preventing algae bloom problem but also for recovering phosphate, as this is an essential material. Adsorption is considered as an effective method especially because the nutrient loaded adsorbent can be directly used for fertilizer. Therefore, this study investigated the potential use of manganese dioxide in natural form (pyrolusite) to adsorb phosphate in aqueous solutions. A series of batch experiments were done to elaborate the adsorption process of phosphate onto manganese dioxide. Several environmental conditions such as pH, temperature and ionic strength were applied in order to get a better understanding of the process mechanism. The results indicated that pH was obviously affected the adsorption process, meanwhile ionic strength did not play significant role. The adsorption of phosphate was higher at a lower pH and getting reduced as the pH increased. Similar to that, the percentage removal of phosphate was declined significantly in higher ionic strength, indicated that the interaction between phosphate and manganese dioxide was mainly controlled by electrostatic force. The adsorption isotherm data correlated better with Langmuir model rather than Freundlich model. The maximum adsorption capacity of manganese dioxide was 11.40 mg P/g. The kinetic data was very well fitted to pseudo second order equation suggested that chemical reaction involved in the adsorption process. Moreover, thermodynamic data confirmed that phosphate adsorption onto manganese dioxide was an endothermic process.

Keywords— adsorption; phosphate; manganese dioxide; isotherm; kinetic; thermodynamic.

I. INTRODUCTION

Phosphate is an essential material for many applications in industrial sector such as paint and coating, a cleaning agent (detergent), pharmaceutical, water treatment and agriculture. However, besides its positive function, the accumulation of phosphate in the aquatic system creates serious problems. Phosphate may naturally occur in water body, but anthropogenic sources from domestic wastewater, industrial wastewater and runoff water from agriculture/feedlot area increase the concentration significantly [1]. It has been reported that 1.3 Mt of phosphate every year discharged into aquatic systems all over the world [2]. This oversupply condition will stimulate eutrophication condition. More than 38% of water bodies in many regions of the world were considered to have eutrophication problem [3]. Eutrophication will lead to algae bloom, abnormal growth of aquatic plants and algae. It is reported that once the concentration of phosphate in lakes or sea is over 0.03 mg/L, algae bloom will occur. The further effect will then arise since algae bloom potentially shifts the water system into anaerobic condition result in aesthetical

(e.g., odor, turbidity) and health problem by producing cyanotoxins. Considering the tremendous effect of phosphate in the water system, some countries applied strict regulation. Environmental Protection Agency (EPA) stated that the maximum permissible concentration and the stringent discharge limit of phosphate ions are 0.1 mg/L and less than 0.05 mg/L respectively [4]. In Australia, the total phosphorus contaminant level in rivers and streams is controlled in the range of 0.01 – 0.1 mg/L, while the requirement for lakes and reservoirs is more stringent as 0.005 – 0.05 mg/L [5].

Based on that condition, strategic effort to eliminate phosphate has been continually developed. Several techniques have been proposed to remove phosphates such as chemical precipitation, biological process, ion exchange, membrane technology and adsorption. Chemical precipitation (by adding ferric or aluminum salts) is a relatively easy method, but the excessive sludge as by-product requires post-treatment. The biological process is an effective method to remove phosphate from water. However stability is another issue since the performance of microorganism is strongly depend on nutrients (e.g., carbon,

nitrogen, iron) availability and operational condition (e.g., pH, temperature). Meanwhile, ion exchange and membrane technology only work well for phosphate elimination if there is no other contaminant exist [6]. Therefore, among those methods, adsorption is considered the most economical, effective and reliable method [7]. Adsorption is a simple technology that requires a relatively low-cost facility, yet has been proved to be able to reduce contaminant in a high percentage. Regards to phosphate, adsorption can be applied not only for removal but also to recover it from the water system. This is important since the demand for phosphate still growing, especially for fertilizer production and detergent manufacturing, but at the same time, the phosphate reserves are gradually diminishing. Another interesting feature from this technique is that nutrient loaded filter can be used directly as phosphate fertilizer and soil conditioner [8], [9].

Various materials have been tested to be an effective adsorbent for phosphate; it was including natural minerals, industrial by-products (steel slag, fly ash, and red mud) and synthetic compound. Focus on the usage of natural minerals such as iron (hydr)oxides including amorphous hydrous ferric oxide, poorly crystalline hydrous ferric oxide goethite and akaganeite has increased as these materials have high affinity to phosphate, low cost and environmentally friendly [10]. Manganese oxides are another natural mineral with poor crystalline oxides, like a ferric oxide, which generally found in manganese-rich coatings. This mineral is considered as a potent scavenger of trace metals in soil, sediment, and rocks [11]. There was much research has been done to test the ability of manganese oxides as an adsorbent for various metals in aqueous solutions. However, there was a limited study to investigate the usage of manganese oxides for phosphate removal.

Therefore, a set of experiments was done to get further information on phosphate removal by using manganese oxides. In this work manganese oxides, ore was used as an adsorbent for phosphate in aqueous solutions. The experiments were done in a batch system with several environmental conditions, such as pH, ionic strength and temperature to elucidate the processing mechanism. The kinetic data, as well as adsorption isotherm, were also determined.

II. MATERIAL AND METHOD

A. Material

All chemicals including KH_2PO_4 , KMnO_4 , H_2SO_4 , Sodium oxalate, HNO_3 , NaOH , HCl were purchased in analytical grade and used directly without any further treatment. A private company in Surabaya supplied pyrolusite as the manganese oxide ore for this research. This material was sieved to get a size of 100 mesh and washed with demineralized water before the usage. To determine the manganese oxides content in the minerals, a simple titration was performed by weighing out 1 g of pyrolusite and put it into a conical flask with 40 mL of 0.5N Sodium oxalate. 50 mL of diluted H_2SO_4 was added before heating process. After black colored of particles disappeared, the solution was titrated against 0.3N KMnO_4 until the colorless turn into pink. The result indicated that the content of manganese oxides (MnO_2) was 84.76% (w/w). Also, the characterization

of surface functional groups of pyrolusite was done by Fourier Transform Infra Red (FTIR, Shimadzu).




Fig.1. FTIR spectrum of MnO_2 (Pyrolusite)

Refers to the Figure 1, the peak at about 590 cm^{-1} is a result of stretching vibration of the Mn-O and Mn-O-Mn bonds. The peak at 1085 cm^{-1} indicates a vibration of hydroxyl groups that are joined to Mn atoms. As for peak at 1519 cm^{-1} arises from bending vibration of O-H and H_2O , which implies that water molecule exists in nanostructures. Meanwhile, peak at about 3464 cm^{-1} describes the stretching vibration on O-H bond and bending vibration of the absorbed water molecule in the lattice [12].

B. Method

1) *Effect of pH and Ionic Strength:* Adsorption batch experiments to investigate the effect of pH and ionic strength was done by preparing 200 mL of phosphate solution with an initial concentration of 10 mg/L into a conical flask. 2.5 g/L of MnO_2 (pyrolusite) was added into the flask before mixing process on thermoshaker at 160 rpm. A variation on pH was set at 2, 4, 6 and 8. pH adjustment was done by using either 0.1 M HCl or NaOH . Sampling was done in a certain interval until the phosphate concentration reached an equilibrium condition. Solution samples were filtered using membrane filter $0.45\text{ }\mu\text{m}$ (Agilent) before phosphate analysis with UV-Spectrometer (Agilent). Ionic strength variation was arranged by using NaCl at 0.01 M, 0.1 M and 1 M. All tests were conducted at temperature 25.5°C .

2) *Adsorption Isotherms:* Experiments for isotherms study were performed with initial concentrations of 10 – 100 mg/L. All processes were the same with the previous explanation, except for sampling time, which was set 24 hours to make sure equilibrium condition was reached. The collected data of adsorbed phosphate onto MnO_2 at equilibrium condition were then analyzed with two adsorption models, Langmuir and Freundlich.

Langmuir model :

$$q_e = \frac{b q_{\max} C_e}{1 + b \cdot C_e} \quad (1)$$

Freundlich model :

$$q_e = k_f C_e^n \quad (2)$$

where C_e is the phosphate concentration in the solution at equilibrium (mgP/L), b is Langmuir constant related to the affinity of binding sites (L/mg), q_{max} is the maximum adsorption capacity (mg/g), k_f is a Freundlich constant related to roughly adsorption capacity (mgP/L), n is the constant related to adsorption density.

3) *Adsorption Kinetics*: Adsorption kinetics was defined from an experiment with one initial concentration of phosphate (10 mg/L) at 25.5°C. Phosphate concentration was analysed until an equilibrium condition achieved. Two equations were applied to define whether the process fit to pseudo-first order or pseudo-second order.

Pseudo first order:

$$\log(q_e - q_t) = \frac{K_1 t}{2.303} + \log q_e \quad (3)$$

Pseudo-second order :

$$\frac{t}{q_t} = \frac{1}{2Kq_e^2} + \frac{1}{q_e} t \quad (4)$$

Where q_t is phosphate concentration over the sampling time (mgP/L), q_e is phosphate concentration at equilibrium condition (mgP/L), K_1 is pseudo first order constant (L/min) and K is a pseudo second order constant ($g/mg.min$)

4) *Thermodynamic Data*: Thermodynamic data were collected from experiments with different temperature applied (25.5°C, 35°C and 45°C). T , e values of standard enthalpy change (ΔH°), standard entropy change (ΔS°) moreover, Gibb's free energy (ΔG°) were determined to ascertain the nature of adsorption process under different temperature condition. Those three parameters can be calculated by applying the following equations :

$$K_d = \frac{C_s}{C_e} \quad (5)$$

$$\ln K_d = \frac{\Delta S^\circ}{R} - \frac{\Delta H^\circ}{RT} \quad (6)$$

where C_s and C_e are the removed and remaining concentrations of phosphate, respectively. R (8.314 Jmol⁻¹K⁻¹) is the ideal gas constant, T is the temperature in Kelvin. While free energy changes (ΔG°) is defined from :

$$\Delta G^\circ = \Delta H^\circ - T\Delta S^\circ \quad (7)$$

The values of ΔH° and ΔS° were determined from the slope and y-intercept of the plot between $\ln K_d$ versus $1/T$ [2].

III. RESULTS AND DISCUSSION

Results of adsorption experiment in different conditions are presented below.

A. Effect of pH and Ionic Strength




Fig.2. Phosphate adsorbed in different pH




Fig.3. % Phosphate removal in different ionic strength ("0" mean there was no NaCl addition)

In all pH condition, the amount of adsorbed phosphate onto manganese dioxide was high at the beginning, indicating a spontaneous reaction, and gradually decreased until an equilibrium stage (after 2.5 hours). Fig. 2 portrays that pH is a significant factor in determining sorption process of phosphate onto manganese dioxide. The higher the pH value, the amount of absorbed phosphate reduced. In this experiment, the most suitable pH for the adsorption of phosphate onto manganese dioxide was 2. The effect of pH can be explained, related to the pH_{pzc} (point of zero charges) of manganese dioxide. pH_{pzc} is a pH in which the net charge of chemical substances in the aqueous solution is zero. At $pH < pH_{pzc}$ the adsorbent's surface will positively charge, while at $pH > pH_{pzc}$ the surface charge will be negative. The manganese dioxide has a low pH_{pzc} ($1.4 < pH_{pzc} < 4.5$) [8], later on, it was reconfirmed that the pH_{pzc} is about 4.2 [11], [12]. Thus, if the pH solution above 4.2 the surface site of adsorbent will have more negative charges because of deprotonation, in which more hydrogen ions leave the sites. Since the domination species of phosphate are negatively charged ($H_2PO_4^-$, HPO_4^{2-}), the interaction with adsorbent will be unlikely. More favorable interaction exists at pH below 4.2 due to protonation at the surface site of manganese dioxide results in more positively charged. In addition to that, at lower pH value the presence of phosphate hydrate ions is ampler than its anhydrate. Thus interaction with the active sites of adsorbent will be much more comfortable. A pH-dependent process indicates that the adsorption of phosphate onto manganese dioxide is driven

by different charges interaction or so-called surface complexation. Phosphate adsorption onto metal oxides has been reported as a dependent process. The amount of adsorbed phosphate tended to increase by decreasing solution pH [5].

Application of ionic strength in different level was intended to explain the sorption mechanism. The increase of ionic strength will reduce the adsorption capacity if electrostatic interaction applies between adsorbate and adsorbent. In contrary, the adsorption capacity will increase along with the increase of ionic strength if different mechanism pertains. The result, as can be seen in Fig. 3., shows a higher level of ionic strength (indicated by the higher concentration of NaCl) the removal of phosphate decreased significantly (about six times lower). The additional concentration of NaCl produces more ions speciation that inhibits the interaction between phosphate and manganese dioxide. This is in another way, indicates that the sorption mechanism of phosphate onto manganese dioxide is driven by electrostatic interaction or surface complexation rather than ligand exchange.

B. Adsorption Isotherms

Adsorption isotherm was used to explain the distribution of adsorbate in liquid and solid phase. Two models, Langmuir and Freundlich, were applied to identify the adsorption isotherm. Data analysis for adsorption isotherms (as can be seen in Fig.4a and 4b) suggested that the adsorbent performance fit better to Langmuir model than Freundlich model, indicated by the higher value of R^2 (0.9955 compared to 0.9397) assuming:

- Manganese dioxide has homogeneous active sites
- A monolayer adsorption process between manganese dioxide and phosphate
- There is no further interaction among adsorbed phosphate

Thus, based on the preferred model, maximum adsorption capacity of manganese dioxide was 11.40 mgP/g (Table I).




Fig. 4(a). The linear plot of Freundlich isotherm




Fig. 4(b). The linear plot of Langmuir isotherm

TABLE I
LANGMUIR AND FREUNDLICH ISOTHERM PARAMETERS

Langmuir	b	0.02 (L/mg)
	q_{\max}	11.40 (mg/g)
Freundlich	K_f	22.67 (mg/g)
	n	2.06

Comparison of the performance of manganese dioxide for adsorbing phosphate in aqueous solution to other natural materials or industrial by-products as reported from previous works can be seen in Table II. The maximum adsorption capacity of manganese dioxide is similar to magnetite, about 1.4 times of iron oxide tailings, almost two times higher than vesuvianite and fly ash and 2.4 times of dolomite. This result is promising for further development.

TABLE II
COMPARISON OF ADSORBED PHOSPHATE IN SEVERAL NATURAL MATERIALS

Adsorbent	Adsorption Capacity (mgP/g)	Reference
La-doped vesuvianite	6.70	[7]
Magnetite	11.20	[13]
Iron oxide tailings	8.20	[14]
Class F fly ash	6.60	[15]
Natural palygorskite	4.00	[16]
Dolomite	4.76	[17]
Manganese dioxide	11.40	This study

C. Adsorption Kinetics

Adsorption kinetics was applied as one of the most important factors to identify the adsorption efficiency and working mechanism of adsorbent. The removal of phosphate from aqueous solutions was occurred relatively fast, in 2.5 hours the adsorption process reached an equilibrium condition. A rapid adsorption rate could be ascribed to the high driving force provided by the concentration gradient of phosphate in aqueous solutions and the large number availability of active sites on the surface of manganese dioxide [4]. Measurement on phosphate concentration at 24 and 48 hours was done, and there was no indication of

further reaction (no additional removal) or desorption confirming that effective process was found at the initial phase (Fig.5.)




Fig. 5. % Removal of Phosphate as a Function of Time, pH = 2, T = 25.5°C, initial concentration = 10 mg/L

To further evaluate the adsorption mechanism of phosphate onto manganese dioxide, two model equations, pseudo first order, and pseudo second order were employed. Experimental data, the concentration of phosphate over the time periods of monitoring (q_t) and concentration of phosphate at equilibrium condition (q_e) were fitted to both models (Fig. 6a and 6b).

Based on regression coefficients, adsorption process of phosphate onto manganese dioxide was much better represented by pseudo-second-order model ($R^2 = 0.9999$) compared to the pseudo-first-order model ($R^2 = 0.8341$). This indicates that chemisorption rather than physisorption drives adsorption mechanism and only induces a monolayer system. This finding inline with isotherm model that follow Langmuir equation, which is also, assume a monolayer adsorption process. A similar finding was reported by previous work, suggesting that the adsorption process of phosphate onto manganese dioxide involves valency forces through sharing or exchanging electrons between adsorbate and adsorbent [2], [9], [17].




Fig. 6(a). pseudo-first-order model for phosphate adsorption




Fig. 6(b). pseudo-second-order model for phosphate adsorption

D. Thermodynamic Data

Temperature is a significant factor in determining the adsorption process. The result of adsorption experiment in different temperature conditions showed that the removal of phosphate from aqueous solutions slightly increased with the increase of temperature. The highest removal, 62.63%, was achieved at the highest temperature applied, 45°C (Fig.7.).




Fig.7. % removal of Phosphate in Different Temperature Conditions

This phenomenon can be explained as the increase in temperature there will be dehydration on phosphate ions, results in unstable binding between the ions. The more unstable the binding will make the adsorption of phosphate onto manganese dioxide sites easier. It seems more energy needed to enhance the adsorption process. By increasing temperature, more energy will be provided thus the interaction between phosphate ions and active sites of manganese dioxide will be more intense, results in more adsorbed phosphate.

To evaluate the adsorption mechanism of phosphate onto manganese dioxide, the thermodynamic parameters (ΔG^0 , ΔH^0 , ΔS^0) were determined in various temperatures (298.65 ; 308.15 and 318.15 K). A plot of $\ln K_d$ versus $1/T$ was found to be linier, as can be seen in Fig.8. From that plot, the values of ΔH^0 and ΔS^0 were ascertained. Then, from both parameters the value of ΔG^0 could be calculated.




Fig.8. Van't Hoff plot of adsorption equilibrium constant, K_d

TABLE III
Thermodynamic Parameters of Phosphate Adsorption

T (K)	C ₀ (ppm)	ΔS^0 (J/mol.K)	ΔH^0 (J/mol)	ΔG^0 (J/mol)
298.65	106.80	10.81	2710.63	-517.26
308.15	106.40	10.81	2710.63	-619.93
318.15	106.55	10.81	2710.63	-728.02

The analysis of thermodynamic data revealed several essential pieces of information (Tabel III) that explain the adsorption mechanism. The decrease in ΔG^0 value with the increase of temperature suggests that the adsorption process is more favour at high than at low temperature. This is an indication that the nature of phosphate adsorption by manganese dioxide was not spontaneous. Or, it can be said that the spontaneity of adsorption is driven by the temperature. The negative value of ΔG^0 confirms that energy is needed to enhance the reaction, the energy requirement is bigger at lower temperature. The positive value of ΔH^0 denotes that the adsorption process is endothermic and irreversible, probably due to non polar interactions [2]. However, the low value of ΔH^0 the point to the adsorption of phosphate onto manganese dioxide is weak. Meanwhile, the positive value of ΔS^0 indicates that randomness at the solid-liquid interface increases during the adsorption process of phosphate onto manganese dioxide. It also reflects the affinity of the adsorbent for phosphate and suggests some structural changes in adsorbate and adsorbent [3].

Many scientific reports have mentioned that endothermic reaction drives the adsorption process of phosphate for most adsorbents, for example, iron oxide tailings, bentonite, red mud, [3], [14], [18]. Referring to the low values of standard Gibbs free energy that have been obtained in this study, the use of manganese dioxide as an adsorbent for phosphate will save energy consumption.

IV. CONCLUSIONS

Manganese dioxide in natural form (Pyrolusite) can be used as an adsorbent to remove phosphate from aqueous solutions. The adsorption process was pH dependent. In this study, the most effective process was at pH = 2. Adsorption isotherm was fitted to Langmuir better compared to

Freundlich model. The maximum adsorption capacity of manganese dioxide following the Langmuir is 11.40 mg Phosphate/g. Pseudo-second order model best represented the adsorption kinetics characteristic. Indicating that adsorption process is mainly driven by chemical reaction (chemisorption). Thermodynamic data demonstrated that the adsorption process of phosphate onto manganese dioxide is endothermic and at high affinity. Compared to several natural materials, manganese dioxide (Pyrolusite) presented a considerably higher adsorption capacity for phosphate removal. However, modification to enlarge its capacity needs to be investigated to find the optimum condition of the adsorbent utilization.

ACKNOWLEDGMENT

The authors would like to thank ECOMEC for supplying Pyrolusite and some preliminary technical data used in this study. Special thanks to Laboratory of Bioprocess Technology and Environmental Process for providing all necessary instruments for analysis

REFERENCES

- [1] M. Dimitris, P. Maria, B. Joannis, I.J. Vassilis, K. Nikolaos, T. Nikolaos, P. Dimitrios, K. Efstratios, O. Georgios, M. Giorgos, "Removal of phosphate from aqueous solutions by adsorption onto Ca(OH)₂ treated natural clinoptilolite", *Chemical Engineering Journal*, vol. 320, pp. 510 – 522, 2017.
- [2] W. Xiong, J. Tong, Z. Yang, G. Zeng, Y. Zhou, D. Wang, P. Song, R. Xu, C. Zhang, M. Cheng, "Adsorption of phosphate from aqueous solution using iron-zirconium modified activated carbon nanofiber: performance and mechanism," *J. Colloid Interf. Sci.* vol. 493, pp. 17 – 23, 2017
- [3] E.B. Mohamed, A.M. Alaa, "Adsorption of phosphate ions from aqueous solutions by modified bentonite with magnesium hydroxide Mg(OH)₂", *Applied Clay Science*, vol. 140, pp.157 – 164, 2017
- [4] X. Liu, L. Zhang, "Removal of phosphate anions using the modified chitosan beads: adsorption kinetic, isotherm and mechanism studies," *Powder Technology*, vol. 277, pp 112 – 119, 2015
- [5] W. Huang, Y. Zhang, D. Li, "Adsorptive removal of phosphate from water using mesoporous materials: a review," *Journal of Environmental Management*, vol. 193, pp. 470 – 482, 2017
- [6] Y. Yu, J.P. Chen, "Key factors for optimum performance in phosphate removal from contaminated water by a Fe-Mg-La tri-metal composite sorbent," *J. Colloid Interf. Sci.* vol. 445, pp. 303 – 311, 2015
- [7] H.Li, J. Ru, W. Yin, X. Liu, J. Wang, W. Zhang, "Removal of phosphate from polluted water by lanthanum doped vesuvianite," *J. Hazard. Mater.* vol.168, pp. 326-330, 2009
- [8] LD. Hylander, A. Kietlinska, G. Renman, G. Siman, "Phosphorus retention in filter materials for wastewater treatment and its subsequent suitability for plant production," *Bioresource Technol.*, vol. 97, pp. 914-921, 2006.
- [9] J. Lalley, C. Han, X. Li, D.D. Dionysiou, M.N. Nadagouda, "Phosphate adsorption using modified iron oxide-based sorbents in lake water: kinetics, equilibrium, and column tests," *Chemical Engineering Journal*, vol 284), pp. 1386 – 1396, 2016
- [10] G. Li, S.Gao, G. Zheng, X. Zhang, "Enhanced adsorption of phosphate from aqueous solution by nanostructured iron(III)-copper(II) binary oxides," *Chemical Engineering Journal*, vol. 235, pp.124-131, 2014.
- [11] Zhao, X.Yang, H. Zhang, C. Chen, X. Wang, "Effect of environmental conditions on Pb(II) adsorption on β -MnO₂", *Chemical Engineering Journal*, vol.164, pp. 49-55, 2010
- [12] Q. Qin, Q. Wang, D. Fu, J. Ma, "An efficient approach for Pb(II) and Cd(II)removal using manganese dioxide formed in situ," *Chemical Engineering Journal*, vol. 172, pp. 68-74, 2011
- [13] L. Lai, Q. Xie, L.Chi, W. Gu, D. Wu, "Adsorption of phosphate from water by easily separable Fe₃O₄@SiO₂ core/shell magnetic nanoparticles functionalized with hydrous lanthanum oxide", *J. Colloid Interf. Sci.* vol. 465, pp. 76-82, 2016

- [14] L. Zeng, X.M. Li, J.D. Liu, "Adsorptive removal of phosphate from aqueous solutions using iron oxide tailings." *Water Research*, vol. 38, pp. 1318-1326, 2004
- [15] N.M. Agyei, C.A. Strydom, J.H. Potgieter. "The removal of phosphate ions from aqueous solution by fly ash, slag, ordinary Portland cement and related blends," *Chem. Conc. Res.* Vol. 30, pp. 823-826, 2002
- [16] H.P. Ye, F.Z. Chen, Y.Q. Sheng, G.Y. Sheng, J.M. Fu, "Adsorption of phosphate from aqueous solution onto modified palygorskite," *Sep. Purif. Technol.*, vol. 50, pp. 283 – 290, 2006
- [17] X. Yuan, W. Xia, J. An, J. Yin, X. Zhou, W. Yang, "Kinetic and thermodynamic studies on the phosphate adsorption removal by dolomite mineral," *Journal of Chemistry*, vol. 2015, pp. 1 – 8, 2015
- [18] W. Huang, S. Wang, Z. Zhu, L. Li, X. Yao, V. Rudolph, F. Haghseresht, "Phosphate removal from wastewater using red mud," *J. Hazardous Materials*, vol. 158, pp. 35 – 42, 2008