

PAPER • OPEN ACCESS

## Scientific & Editorial Boards the 2<sup>nd</sup> NRLS-2018

To cite this article: 2019 *IOP Conf. Ser.: Earth Environ. Sci.* **293** 011004

View the [article online](#) for updates and enhancements.



**IOP | ebooks™**

Bringing you innovative digital publishing with leading voices  
to create your essential collection of books in STEM research.

Start exploring the **collection** - **download the first chapter of  
every title for free.**



## SCIENTIFIC & EDITORIAL BOARDS the 2<sup>nd</sup> NRLS-2018

- Andoniana Rakoto Malala, Centre de Formation et d'Application du Machinisme Agricole (CFAMA), Antsirabe, MDG
- Christina Avanti, Faculty of Pharmacy, University of Surabaya, IDN
- Fauna Herawati, Faculty of Pharmacy, University of Surabaya, IDN
- Finna Setiawan, Faculty of Pharmacy, University of Surabaya, IDN
- Ida Bagus Made Artadana, Faculty of Biotechnology, University of Surabaya, IDN
- Johan Sukweenadhi, Faculty of Biotechnology, University of Surabaya, IDN
- Juris Burlakovs, Faculty of Health and Life Sciences, Linnaeus University, SWD
- Kartini, Faculty of Pharmacy, University of Surabaya, IDN
- Lieke Riadi, Faculty of Chemical Engineering, University of Surabaya, IDN
- Lim Kok Kuan, Nippon Biodiesel Fuel, Odawara-city, Kanagawa, JPN
- Maizirwan Mel, Department of Biotechnology Engineering, Faculty of Engineering International Islamic University Malaysia. MYS
- Mariana Wahjudi, Faculty of Biotechnology, University of Surabaya, IDN
- Maria Goretti Marianti Purwanto, Fac.of Biotechnology, University of Surabaya, IDN
- Oeke Yunita, Faculty of Biotechnology, University of Surabaya, IDN
- Olga Anne, Faculty of Marine Technology and Natural Sciences of Klaipeda University, LTU
- Peeyush Soni, Department of Agricultural and Food Engineering, Indian Institute of Technology Kharagpur, IND
- Popy Hartatie Hardjo, Faculty of Biotechnology, University of Surabaya, IDN
- Praptiningsih Gamawati Adinurani, Department of Agrotechnology, Faculty of Agriculture, Merdeka University of Madiun, IDN
- Rangga Kala Mahaswa, Graduate Program, Faculty of Philosophy, Universitas Gadjah Mada, Yogyakarta, IDN
- Rika Yulia, Faculty of Pharmacy, University of Surabaya, IDN
- Risma Ikawaty, Faculty of Medical, University of Surabaya, IDN
- Roy Hendroko Setyobudi, Postgraduate of Agriculture Science, Waste Laboratory University of Muhammadiyah Malang, IDN
- Sulistyono Emantoko, Faculty of Biotechnology, University of Surabaya, IDN
- Tjie Kok, Faculty of Biotechnology, University of Surabaya, IDN
- Tsitsino Turkadze, Department of Chemical and Environmental Technologies, Akaki Tsereteli State University, Kutaisi, GEO
- Yahya Jani, Departement of Biology and Environmental Science, Linnaeus University, Kalmar, SWD
- Yalun Arifin, Faculty of Food Business and Technology, Prasetya Mulya University, Jakarta, IDN
- Zahrah Nurfadhilah, Faculty of Biology, Universitas Gadjah Mada, Yogyakarta, IDN
- Zane Vincēviča-Gaile, Department of Environmental Science, University of Latvia, Riga – LVA





This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our Privacy and Cookies policy.



# Table of contents

Volume 293

2019

◀ Previous issue

**The 2nd International Conference on Natural Resources and Life Sciences (NRLS-2018)**  
**23–24 August 2018, Ibis Styles Hotel, Surabaya, Indonesia**

[View all abstracts](#)

**Accepted papers received: 17 May 2019**

**Published online: 27 June 2019**

## Preface

---

**OPEN ACCESS** 011001

The 2nd International Conference on Natural Resources and Life Sciences (NRLS-2018)

[+ View abstract](#) [View article](#) [PDF](#)

---

**OPEN ACCESS** 011002

Conference Photos

[+ View abstract](#) [View article](#) [PDF](#)

---

**OPEN ACCESS** 011003

Conference Sponsors

[+ View abstract](#) [View article](#) [PDF](#)

---

**OPEN ACCESS** 011004

Scientific & Editorial Boards the 2<sup>nd</sup> NRLS-2018

[+ View abstract](#) [View article](#) [PDF](#)

---

**OPEN ACCESS** 011005

## Peer review statement

[+ View abstract](#)[View article](#)[PDF](#)

---

**Papers****OPEN ACCESS**

012001

## Android based rice pest detection system using learning vector quantization method

A Budiman, P Utomo and S Rahayu

[+ View abstract](#)[View article](#)[PDF](#)

---

**OPEN ACCESS**

012002

## Community behavior and single-use plastic bottle consumption

A Khoironi, S Anggoro and S Sudarno

[+ View abstract](#)[View article](#)[PDF](#)

---

**OPEN ACCESS**

012003

Genetic variability of soybean (*Glycine max* L. Merrill) genotypes for pod shattering resistance

A Krisnawati and M M Adie

[+ View abstract](#)[View article](#)[PDF](#)

---

**OPEN ACCESS**

012004

## Review on biogas from palm oil mill effluent (POME): Challenges and opportunities in Indonesia

A Rajani, Kusnadi, A Santosa, A Saepudin, S Gobikrishnan and D Andriani

[+ View abstract](#)[View article](#)[PDF](#)

---

**OPEN ACCESS**

012005

## Early detection of somaclonal variation in oil palm callus culture through cytological and SDS-PAGE protein analysis

A Sahara, Reflini, C Utomo and T Liwang

[+ View abstract](#)[View article](#)[PDF](#)

---

**OPEN ACCESS**

012006

Genetic diversity of Indonesian soybean (*Glycine max* L. Merrill) germplasm based on morphological and microsatellite markers

A Sulisty, F C Indriani, M J Mejaya, A N Sugiharto and J Agranoff

[+ View abstract](#)[View article](#)[PDF](#)

---

**OPEN ACCESS**

Fermentation quality of *Pennisetum purpureum* cv. Mott ensiled with *Lactobacillus plantarum* and sugarcane molasses in tropic

012007

A Wahyudi, L Hendraningsih, Sutawi, R H Setyobudi and M Mel

[+ View abstract](#)

[View article](#)

[PDF](#)

**OPEN ACCESS**

012008

The evaluation of estimated breeding value and the most probable producing ability for the basis selection of Ettawa crossbred goat (*Capra hircus* sp.) at Malang, East Java, Indonesia

A Winaya, Suyatno, P Coy and N Fauzi

[+ View abstract](#)

[View article](#)

[PDF](#)

**OPEN ACCESS**

012009

Menthol from the stem and leaf *in-vitro* *Mentha piperita* Linn.

B K Wijaya, P H Hardjo and S Emantoko

[+ View abstract](#)

[View article](#)

[PDF](#)

**OPEN ACCESS**

012010

A review: Biomass-based fuel pellet usage in biomass gasifier-an optimization on non-wood material

D Andriani, T D Atmaja, M Arifin, A Rajani and Kusradi

[+ View abstract](#)

[View article](#)

[PDF](#)

**OPEN ACCESS**

012011

Effectiveness of the use of organic waste as fertilizer and physical scarification of seeds on growth of seeds nila plants (*Indigofera* sp.)

D Roeswitawati, M Huda, D Indratmi and M Mel

[+ View abstract](#)

[View article](#)

[PDF](#)

**OPEN ACCESS**

012012

Physiology and genotyping of adaptive and sensitive oil palm progenies under unwatered stress condition

D Yono, E Purwanti, A Sahara, Y A Nugroho, Z A Tanjung, R Aditama, C Dewi, A E Sihotang, C Utomo and T Liwang

[+ View abstract](#)

[View article](#)

[PDF](#)

**OPEN ACCESS**

012013

Effect of phosphoric acid pretreatment on characterization of gelatin from broiler chicken (*Gallus gallus domesticus* L.) bones

D Yuliani, A Maunatin, A Jannah and H H Fauziyyah

[+ View abstract](#)

[View article](#)

[PDF](#)

**OPEN ACCESS**

012014

Selection for drought-resistant rice (*Oryza sativa* L.) using polyethylene glycol

E D Purbajanti, F Kusmiyati, E Fuskhah, R Rosyida, P G Adinurani and Z Vincēviča-Gaile

[+ View abstract](#)[View article](#)[PDF](#)**OPEN ACCESS**

012015

Effectiveness the source of nitrogen from NO<sub>3</sub> and NH<sub>4</sub> for *Panicum maximum* Jacq. growth in saline soil

E D Purbajanti, P G Adinurani, T Turkadze, Z Vincēviča-Gaile and R H Setyobudi

[+ View abstract](#)[View article](#)[PDF](#)**OPEN ACCESS**

012016

Anticancer potential from ethanol extract of *Zanthoxylum acanthopodium* DC. seed to against MCF-7 cell line

E V Arsita, D E Saragih and K Aldrin

[+ View abstract](#)[View article](#)[PDF](#)**OPEN ACCESS**

012017

## Long-term Proton Pump Inhibitors induces recurrent Urinary Tract Infections: A case study

F Cokro and S T Arrang

[+ View abstract](#)[View article](#)[PDF](#)**OPEN ACCESS**

012018

## Molecular docking, drug-likeness, and ADMET study of 1-benzyl-3-benzoylurea and its analogs against VEGFR-2

F Suhud, D H Tjahjono, T A Yuniarta, G S Putra and J Setiawan

[+ View abstract](#)[View article](#)[PDF](#)**OPEN ACCESS**

012019

Microcontroller application for pH and temperature control system in liquid sugar liquefaction process made from cassava (*Manihot esculenta* Crantz.)

H Santosa, Yuliati, I J Mulyana, D S Sirait and S D Novitasari

[+ View abstract](#)[View article](#)[PDF](#)**OPEN ACCESS**

012020

The effect of fermentation process on physical and chemical characteristics of pitaya (*Hylocereus polyrhizus* [F.A.C. Weber] Britton & Rose) stem flour

H Soedjatmiko, R Chrisnasari and P H Hardjo

[+ View abstract](#)[View article](#)[PDF](#)

**OPEN ACCESS**

012021

**The performance of three local rice (*Oryza sativa* L.) cultivar from East Kalimantan - Indonesia under drought stress at early seedling stage**

I B M Artadana, I T Dewi and J Sukweenadhi

[+ View abstract](#) [View article](#) [PDF](#)**OPEN ACCESS**

012022

**Coffee reduced the production of neutrophil superoxide radical *in vitro***

I D A Susilawati, A Safaatin and J Burlakovs

[+ View abstract](#) [View article](#) [PDF](#)**OPEN ACCESS**

012023

**The potential of extract of *Zingiber zerumbet* (L.) Smith as a feed additive to improve the production performances and meat nutritional composition of broiler chickens**

I D Rahayu, W Widodo, I Prihartini, A Winaya, L Zalizar, T Untari and M Mel

[+ View abstract](#) [View article](#) [PDF](#)**OPEN ACCESS**

012024

**Callus induction and *in vitro* mass culture of adventitious roots from leaf segment explants of *Dendropanax morbifer* Lev.**

J Sukweenadhi, J Y Choi, Y J Kim, L Kaliraj, S Abid, J C Ahn and D C Yang

[+ View abstract](#) [View article](#) [PDF](#)**OPEN ACCESS**

012025

**Comparison of energy production, net energy balance, net energy ratio, and renewable index for biodiesel production from oil palm (*Elaeis guineensis* Jacq.) and jatropha (*Jatropha curcas* L.) based on life cycle assessment**

K Siregar, A H Tambunan, Sholihati, S S Wirawan and T Araki

[+ View abstract](#) [View article](#) [PDF](#)**OPEN ACCESS**

012026

**Potency of *Fibraurea tinctora* Lour. extract as anti-bacterial agents towards pathogenic bacteria**

L Zalizar, I D Rahayu, Sujono and Y A Nor

[+ View abstract](#) [View article](#) [PDF](#)**OPEN ACCESS**

012027

**Characterization of adaptive and productive soybean (*G. max* L.) genotypes in dry land of Kalimantan, Indonesia**

M M Adie, A Krishawati and D Suryati



[+ View abstract](#)[View article](#)[PDF](#)**OPEN ACCESS**

012028

### Inhibition of *Burkholderia cenocepacia* H111 quorum sensing system by environmental bacterial isolates

M Wahjudi, W D Kurniawan, M T Gultom, I B M Artadana and N I M Puad

[+ View abstract](#)[View article](#)[PDF](#)**OPEN ACCESS**

012029

### The drought stress tolerance of physic nut (*Jatropha curcas* Linn.) genotypes

Maftuchah, I Z Fahmi, A Zainudin, A Ikhwan, Djumali and L K Kuan

[+ View abstract](#)[View article](#)[PDF](#)**OPEN ACCESS**

012030

### Characterization of thermostable chitinase from *Bacillus licheniformis* B2

N Jayanthi, M G M Purwanto, R Chrisnasari, T Pantjajani, A Wahjudi and M Sugiarto

[+ View abstract](#)[View article](#)[PDF](#)**OPEN ACCESS**

012031

### Antibacterial activity roll on deodorant with *Pluchea indica* (L.) leaf extract against *Staphylococcus epidermidis* (Evans 1916) *in vitro*

O Komala, I Y Wiendarlina and N Rizqiyana

[+ View abstract](#)[View article](#)[PDF](#)**OPEN ACCESS**

012032

### Production potensial of sweet corn (*Zea mays* Linn. var. *Saccharata* Sturt) 'Bonanza' to different planting pattern and phosphorus sources

P G Adinurani, S Rahayu, L S Budi, S Pambudi and P Soni

[+ View abstract](#)[View article](#)[PDF](#)**OPEN ACCESS**

012033

### Genome-wide identification of oil palm (*Elaeis guineensis* Jacq.) chitinases and their response to *Ganoderma boninense* Pat. infection

R Aditama, R Tryono, Z A Tanjung, C Utomo and T Liwang

[+ View abstract](#)[View article](#)[PDF](#)**OPEN ACCESS**

012034

### Effect of new NPK fertilizer on lowland rice (*Oryza sativa* L.) growth

R Budiono, P G Adinurani and P Soni

[+ View abstract](#)[View article](#)[PDF](#)

**OPEN ACCESS**

012035

**Prospect of Fe non-heme on coffee flour made from solid coffee waste: Mini review**

R H Setyobudi, L Zalizar, S K Wahono, W Widodo, A Wahyudi, M Mel, B Prabowo, Y Jani, Y A Nugroho, T Liwang and A Zaebudin

[+ View abstract](#)[View article](#)[PDF](#)**OPEN ACCESS**

012036

**Biomass to methanol plant based on gasification of palm empty fruit bunch**

R Heryadi, A S Uyun, E Yandri, S M Nur, K Abdullah and O Anne

[+ View abstract](#)[View article](#)[PDF](#)**OPEN ACCESS**

012037

**Review article: Myopia - Genetically inherited or environmental influences**

S E E Tjoa and S E D Putra

[+ View abstract](#)[View article](#)[PDF](#)**OPEN ACCESS**

012038

**Rapid reversed-phase high performance liquid chromatography assay of Tert-butylhydroquinone content in food products**

S Sutanto and M G M Purwanto

[+ View abstract](#)[View article](#)[PDF](#)**OPEN ACCESS**

012039

**Developmental effect of cashew nut shell extract against nymphal instar of Silver leaf Whitefly (*Bemisia tabaci* Genn.)**

W R Andayanie and N Ermawati

[+ View abstract](#)[View article](#)[PDF](#)**OPEN ACCESS**

012040

**Identification of a local variety of 'uwi' (*Dioscorea alata* Linn.) in four agro-climate regions of East-West Java - Indonesia based on tuber character**

Wuryantoro, I R Puspitawati, R I Fitriyani and P Soni

[+ View abstract](#)[View article](#)[PDF](#)**OPEN ACCESS**

012041

**Genome-wide SNP-discovery and analysis of genetic diversity in oil palm using double digest restriction site associated DNA sequencing**

Y A Nugroho, Z A Tanjung, D Yono, A S Mulyana, H M Simbolon, A S Ardi, Y Y Yong, C Utomo and T Liwang

[+ View abstract](#)[View article](#)[PDF](#)

**OPEN ACCESS**

012042

**Application of blood donor routine detector using K-Nearest neighbors**

Y Nurdiansyah, P Pandunata, N D Prasetyo, A Trihartono, F G Putrianti and F Wijayanto

[+ View abstract](#)[View article](#)[PDF](#)**OPEN ACCESS**

012043

**Identification of the bacterium isolate from Mackerel Fish (*Rastrelliger* sp.) using 16S rRNA Gene**

Y S Ismail, Febriani, C Yulvizar and R Ramadhani

[+ View abstract](#)[View article](#)[PDF](#)**JOURNAL LINKS**[Journal home](#)[Information for organizers](#)[Information for authors](#)[Search for published proceedings](#)[Contact us](#)[Reprint services from Curran Associates](#)

PAPER • OPEN ACCESS

## Menthol from the stem and leaf *in-vitro* *Mentha piperita* Linn.

To cite this article: B K Wijaya *et al* 2019 *IOP Conf. Ser.: Earth Environ. Sci.* **293** 012009

View the [article online](#) for updates and enhancements.



**IOP | ebooks™**

Bringing you innovative digital publishing with leading voices  
to create your essential collection of books in STEM research.

Start exploring the collection - download the first chapter of  
every title for free.

## Menthol from the stem and leaf *in-vitro* *Mentha piperita* Linn.

B K Wijaya, P H Hardjo\* and S Emantoko

Faculty of Biotechnology, University of Surabaya, Raya Kalirungkut Road, Surabaya 60292, East Java, Indonesia

\*Corresponding author: poppyhardjo@gmail.com

**Abstract.** The need for menthol in Indonesia is increasing annually, but it is not followed by increases the availability in the country, so the number of imports is increasing each year. Therefore, in this research conducted a study using plant tissue culture techniques to produce secondary metabolite especially menthol using shoot multiplication. A single node of *Mentha piperita* L. was inoculated on three kinds of growth medium, which are MS + 0.1 mg L<sup>-1</sup> NAA + 0.1 mg L<sup>-1</sup> BA, MS + 0.5 mg L<sup>-1</sup> BA, and MS + 2 mg L<sup>-1</sup> BA. Based on fresh weight and morphological observation, MS + 0.1 mg L<sup>-1</sup> NAA + 0.1 mg L<sup>-1</sup> BA was the best growth medium. Based on the previous experiment, the study was aiming to continue observing the concentration of menthol found in *M. piperita* shoot. Stems and leaves of *M. piperita* were dried in room temperature, the dried stems and leaves were extracted using steam distillation method with ratio 0.3 g of dried stems and leaves in 25 mL aquadest. The crude extract was analyzed by gas chromatography (HP 6890) using INNOWAX 19095N-123 column. The results showed that 2 wk was the best amount of time to obtain the highest concentration of menthol specifically (1 218.5 ± 47.1) mg L<sup>-1</sup> (yield = 9.748 %).

**Keywords:** Gas chromatography, menthol, shoot multiplication, steam distillation

### 1. Introduction

*Mentha* spp. belongs to Labiatae family, there are three main species which have high economic value like *Mentha piperita* L. The industrial demands of *M. piperita* product are high, but Indonesia cannot fulfill the demands without import. In 2006 Indonesia imported *Mentha* spp. product worth the USD  $3.78 \times 10^6$  [1]. *M. piperita* is a first sterile hybrid generation between *M. spicata* and *M. aquatica* [2]. The main compound from *Mentha* essential oil is menthol; this compound is produced in peltate glandular trichome which located at leaf [3].

The purpose of this research is to increase the production of menthol using plant tissue culture technique. Cytokinin is phytohormone that induced shoot proliferation and cell division [4]. Plant tissue culture is known to produce plants that grow faster [5]. In addition, the concentration of secondary metabolites, especially monoterpenes in *Lavandula pedunculata in vitro*, is known to be greater than plants grown in their natural [6]. The growth medium used for multiplication of shoots with the aim of increasing menthol concentration is Murashige & Skoog media (MS) + 6-Benzylaminopurine (BAP) + Naphthaleneacetic acid (NAA) [7]. The advantage of *in vitro* shoot multiplication techniques is that it is able to produce target compounds under controlled conditions from changes in weather and soil conditions, plants obtained free from microorganisms and insects, all types of plants can be used in this technique and reduce labor costs and increase productivity [8].



## 2. Materials and methods

### 2.1. Plant materials, treatment, and growing condition

This research was carried out at the Laboratory of Plant Biotechnology and Biopurification and Biomolecular Laboratory, Faculty of Biotechnology, University of Surabaya. The peppermint seeds (Biopot<sup>®</sup>) used in this study were pre-sterilized using bactericidal and fungicidal solution for an hour. The surface sterilization of explants was continued in Laminar Air Flow Cabinet with immersion in 70 % ethanol for 1 min, and 5.25 % NaOCl solution for 5 min followed by rinsing with sterile H<sub>2</sub>O at least three times. Furthermore, sterilization was carried out again by immersion in 1.75 % NaOCl solution for 10 min followed by rinsing with sterile water at least three times. Sterilized mint seeds are planted on MS media without hormones. The culture bottle was incubated under the white fluorescent lamps. Explants used are nodes from mint plants that are approximately 3 mo. The plant is cut to only one node. Furthermore, nodules cut into various growth media (table 1.) The culture was incubated in a culture room with a temperature of  $(25 \pm 2) ^\circ\text{C}$  with the lighting of white fluorescent lamps and observed the formed tissue.

**Table 1.** Growth media variation.

Media
MS + 0.5 mg L <sup>-1</sup> 2.4-D
MS + 1 mg L <sup>-1</sup> 2.4-D
MS + 2 mg L <sup>-1</sup> 2.4-D
MS + 4 mg L <sup>-1</sup> 2.4-D
MS + 6 mg L <sup>-1</sup> 2.4-D
MS + 0.5 mg L <sup>-1</sup> NAA
MS + 2 mg L <sup>-1</sup> NAA
MS + 4 mg L <sup>-1</sup> NAA
MS + 6 mg L <sup>-1</sup> NAA
MS + 8 mg L <sup>-1</sup> NAA
MS + 2 mg L <sup>-1</sup> NAA + 0.5 mg L <sup>-1</sup> BAP
MS + 2 mg L <sup>-1</sup> NAA + 1 mg L <sup>-1</sup> BAP
MS + 2 mg L <sup>-1</sup> NAA + 2 mg L <sup>-1</sup> BAP
MS + 1 mg L <sup>-1</sup> BAP
MS + 2 mg L <sup>-1</sup> BAP
MS + 2 mg L <sup>-1</sup> BAP + 1 mg L <sup>-1</sup> NAA
MS + 0.1 mg L <sup>-1</sup> NAA + 0.1 mg L <sup>-1</sup> BAP

### 2.2. Extraction and determination of menthol

The extraction samples used were leaves and stems of mint plants obtained from the bud multiplication process that have been dried in room temperature under air conditioning. The solvent used was 25 mL of water (distilled water) to extract 0.3 g of dried stems and *M. piperita* leaves, the sample was extracted using the steam distillation method. Then the extract obtained was analyzed for menthol content using a gas chromatography device (HP 6890) with INNOWAX 19095N-123 column [9]. For the ex vitro sample used stem and leaves of *M. piperita ex vitro* at 3 mo.

### 2.3. Data analyses

The research conducted was an experimental study in the laboratory with a Completely Randomized Design (CRD). In this study two non-parametric data were obtained because the data wasn't normally distributed, namely the best media used to grow *M. piperita* plants in vitro, and data on menthol production monitored every week. To determine the best growth media, and the best time to harvest, need to obtain the fresh weight of *M. piperita in vitro*, and menthol concentration for 7 wk. The

statistic test used in this research is the Kruskal-Wallis test and if significance found in the Kruskal-Wallis test continue using Dunn-Bonferroni as post hoc test.

### 3. Results and discussions

#### 3.1. Preliminary test

Preliminary tests were carried out to determine the three types of growth media to be tested. Further, the aspects considered were morphology of *M. piperita* grown *in vitro*. From this preliminary test selected, MS medium + 0.1 mg L<sup>-1</sup> NAA + 0.1 mg L<sup>-1</sup> BA, MS + 0.5 mg L<sup>-1</sup> NAA and MS + 2 mg L<sup>-1</sup> NAA to be used at the next research stage because it has the best morphology. The reason for choosing MS media was 0.5 mg L<sup>-1</sup> NAA and MS + 2 mg L<sup>-1</sup> NAA compared to MS + 1 mg L<sup>-1</sup> BA and MS + 2 mg L<sup>-1</sup> BA, because the hormones NAA and IAA (Indole-3-Acetic Acid) were able to increase the oil content of essential oils and menthol in *M. piperita* [9]. Then MS + 0.5 mg L<sup>-1</sup> NAA and 2 mg L<sup>-1</sup> NAA has a better morphology than MS + 2 mg L<sup>-1</sup> BA + 1 mg L<sup>-1</sup> NAA, the morphology in question is the appearance of the plant (abnormal).

**Table 2.** Preliminary test result.

Media	Tissue/organ formed
MS + 0.5 mg L <sup>-1</sup> 2,4-D	Callus
MS + 1 mg L <sup>-1</sup> 2,4-D	Callus
MS + 2 mg L <sup>-1</sup> 2,4-D	Callus
MS + 4 mg L <sup>-1</sup> 2,4-D	Browning
MS + 6 mg L <sup>-1</sup> 2,4-D	Browning
MS + 0.5 mg L <sup>-1</sup> NAA	Shoot + callus
MS + 2 mg L <sup>-1</sup> NAA	Shoot + callus
MS + 4 mg L <sup>-1</sup> NAA	Browning
MS + 6 mg L <sup>-1</sup> NAA	Browning
MS + 8 mg L <sup>-1</sup> NAA	Browning
MS + 2 mg L <sup>-1</sup> NAA + 0.5 mg L <sup>-1</sup> BA	Shoot + callus
MS + 2 mg L <sup>-1</sup> NAA + 1 mg L <sup>-1</sup> BA	Browning
MS + 2 mg L <sup>-1</sup> NAA + 2 mg L <sup>-1</sup> BA	Browning
MS + 1 mg L <sup>-1</sup> BA	Shoot + callus
MS + 2 mg L <sup>-1</sup> BA	Shoot + callus
MS + 2 mg L <sup>-1</sup> BA + 1 mg L <sup>-1</sup> NAA	Shoot + callus
MS + 0.1 mg L <sup>-1</sup> NAA + 0.1 mg L <sup>-1</sup> BA	Shoot


#### 3.2. Growth index curve

Growth Index Curve is obtained from the final weight of the stem and fresh leaves divided by fresh initial weight. The calculation of the growth index was done by weighing fresh initial weight before planting and after being cultivated for (1, 2, 3, 4, 5, 6 and 7) wk. Observation of the growth index stopped at week 7 because the index value had decreased compared to the 6th week.


**Table 3.** Fresh weight of *M. piperita* at 6 wk old.

Media	Fresh weight (g)
MS + 0.1 mg L <sup>-1</sup> NAA + 0.1 mg L <sup>-1</sup> BA	0.136 ± 0.071
MS + 0.5 mg L <sup>-1</sup> NAA	0.129 ± 0.035
MS + 2 mg L <sup>-1</sup> NAA	0.179 ± 0.121

The three growth media did not have a significant effect using the Kuskal-Wallis test at  $\alpha = 0.05$



**Figure 1.** *M. piperita* growth index curve on MS media + 0.1 mg L<sup>-1</sup> NAA + 0.1 mg L<sup>-1</sup> BA, MS + 0.5 mg L<sup>-1</sup> NAA and MS + 2 mg L<sup>-1</sup> NAA.




**Figure 2.** Morphology of *M. piperita* *in-vitro*.

Fresh weight of *M. piperita* grown on MS + 2 mg L<sup>-1</sup> NAA is the largest among other growth media. This is due to the presence of callus at the base of the plant stem. Callus also found at the base of the plant stem that grown using MS + 0.5 mg L<sup>-1</sup> NAA. *M. piperita* that grew at MS + 0.1 mg L<sup>-1</sup> NAA + 0.1 mg L<sup>-1</sup> BA obtained plants do not form callus and have a heavier mass than MS media + 0.5 mg L<sup>-1</sup> NAA. Menthol compounds are stored in the peltate glandular trichome [3]. The peltate glandular trichome is found in parts of plants exposed to air [10]. So it can be said that the larger/, the heavier plant, the more peltate glandular trichome, in other words, the more menthol that can be extracted. In addition, other factors considered by the morphology of plants planted *in vitro*. There was no monoterpene found in *Mentha spicata* callus because the glandular oil gland or peltate glandular trichome had not been formed because callus cells were undifferentiated plant cells [11]. Therefore, the MS medium was selected + 0.1 mg L<sup>-1</sup> NAA + 0.1 mg L<sup>-1</sup> BA as the best growth medium.




### 3.3. Extraction and determination of menthol

Extraction of menthol compounds was carried out using the steam distillation method because it is commonly used to isolate essential oils. The distillation process is stopped until all the water in the flask has evaporated. From steam distillation results obtained  $\pm 24$  mL of crude menthol extract from steam distillation. The crude extract obtained was analyzed using a gas chromatography device (HP 6890) with the INNOWAX 19095N-123 column.




**Figure 3.** Profile of gas chromatography on crude extract *M. piperita* grown on MS + 0.1 mg L<sup>-1</sup> NAA + 0.1 mg L<sup>-1</sup> BA medium at 2 wk old.



**Figure 4.** Profile of gas chromatography on crude extract *M. piperita* has grown on *ex-vitro*.

There is a difference between the crude extracts of *M. piperita in-vitro* and *ex-vitro*, the peaks that appeared at retention time was around 5 min in the crude extract profile of *ex-vitro* plants but were not found in crude plant extracts *in-vitro*. And the crude extracts of plants *in-vitro* found peaks at retention time about 8 min which were not found in crude extracts of *ex-vitro* plants. In other words, the *M. piperita* plant *in-vitro* is not capable of producing secondary metabolites that are exactly the same as their original plants and is able to form secondary metabolites that are completely different from the

original plants. Several factors that influence the difference in results is nutrition availability and environmental conditions. The availability of nutrients includes the concentration of sugar, nitrate, phosphate and growth regulating substances whereas environmental conditions include temperature, light intensity and pH [12]. Plant tissue culture systems are able to produce secondary metabolites that are completely different from the original plants because of very different environmental conditions, while the levels can be the same, larger or smaller [13]. *In-vitro* plant growth media have been given nutrition, sufficient growth regulating substances, and the growth environment is conditioned stable with adequate lighting, the temperature of 20 °C and pH of 5.8. All treatments given to plants *in-vitro* are common treatments to optimize growth in plant tissue culture while the *ex-vitro* plants all of these factors are not controlled at all, or in other words depending on nature.



**Figure 5.** Menthol concentration curve and *M. piperita* growth index at 0.1 mg L<sup>-1</sup> NAA and 0.1 mg L<sup>-1</sup> BA compared to incubation time

**Table 4.** Menthol concentration observed every week grown at MS + 0.1 mg L<sup>-1</sup> BA + 0.1 mg L<sup>-1</sup> NAA

Plant age (wk)	Menthol concentration (mg L <sup>-1</sup> )	Yield (% w/w)
0	231.3 <sup>ab</sup> ± 59.9	1.85
1	586.3 <sup>ab</sup> ± 126.6	4.69
2	1 218.5 <sup>a</sup> ± 47.1	9.748
3	599 <sup>ab</sup> ± 179	4.792
4	366.5 <sup>ab</sup> ± 85.4	2.932
5	405.6 <sup>ab</sup> ± 50.7	3.245
6	228.997 <sup>ab</sup> ± 0.642	1.832
7	314.93 <sup>ab</sup> ± 9.58	2.519
<i>Ex-vitro plant</i>	97.21 <sup>b</sup> ± 1.373	0.778

The different annotation followed by average value showed significance using Dunn-Bonferroni test  $\alpha = 0.05$

The highest concentration of menthol was 2 wk old *M. piperita* that have grown using MS + 0.1 mg L<sup>-1</sup> NAA + 0.1 mg L<sup>-1</sup> BA, and the least concentration of menthol was *ex-vitro M. piperita*. This phenomenon is not in accordance with the theory, that the production of secondary metabolites generally occurs at the end of the stationary period when the nutrient supply is depleted [4]. This may be caused by menthol has been further metabolized to menthyl acetate [14]. There are also indications that the terpenoids used for defense have decreased due to the absence of threats.

Another factor that might cause the highest phenomenon of menthol concentration in the second week of culture is stress caused by injury at the beginning of cutting explants. This phenomenon was also showed on another research, where at 2 wk after the plant was injured there was a significant increase in the concentration of monoterpenes compared to non-injured plants [15]. This increase in the concentration of monoterpenes may be caused by an increase in the activity of the enzyme monoterpene cyclase [16]. Monoterpene cyclase plays a role in converting Geranyl pyrophosphate (GPP) to 4s-limonene [17]. GPP is the earliest compound in menthol biosynthesis pathway or it can be said that it is a compound that initiates menthol formation so that plants will tend to produce menthol in this condition. *M. piperita* grown in vitro is able to produce more menthol than *ex vitro*. This may be due to the availability of sufficient nutrients so that in vitro plants are able to produce more menthol. Besides that phytohormones are able to increase the levels of essential oil and menthol oil in *Mentha* spp. [18].

#### 4. Conclusion

Based on this research, it can be concluded that the best growth medium for multiplying *M. piperita* shoots *in-vitro* was MS + 0.1 mg L<sup>-1</sup> NAA + 0.1 mg L<sup>-1</sup> BA. Chromatogram profile on crude extracts of *M. piperita* plant in vitro showed three peaks, namely at about two retention times: 4.5 min and 8 min. Chromatogram profile on crude extract of *M. piperita ex-vitro* showed three peaks, namely at about two retention times: 4.5 min and 5 min. The menthol concentration observed for 7 wk in the largest *M. piperita* plant *in-vitro* was at the culture time of the 2<sup>nd</sup> week which was (1 218.5 ± 47.1) mg L<sup>-1</sup>. The menthol concentration observed in the *M. piperita ex-vitro* plant was (97.21 ± 1.373) mg L<sup>-1</sup>.

#### References

- [1] Pribadi E R 2010 Peluang Pemenuhan Kebutuhan Produk *Mentha* Spp. di Indonesia [Chance to Fulfill the Need of *Mentha* Spp. Product] *Perspektif* **2**(9) 66–77  
[http://perkebunan.litbang.pertanian.go.id/wp-content/uploads/2011/03/perkebunan\\_perspektif\\_N-2-Rini-Pribadi-mentha\\_.pdf](http://perkebunan.litbang.pertanian.go.id/wp-content/uploads/2011/03/perkebunan_perspektif_N-2-Rini-Pribadi-mentha_.pdf)
- [2] Abbaszadeh B A Valadabadi S A Farahani H A and Darvishi H H 2009 Studying of essential oil variations in leaves of *Mentha* species *Afr. J. Plant Sci.* **3**(10) 217–21  
<https://academicjournals.org/journal/AJPS/article-full-text-pdf/32A65E611115>
- [3] Turner G W and Croteau R 2004 Organization of monoterpene biosynthesis in *Mentha*: immunocytochemical localizations of geranyl diphosphate synthase, limonene-6-hydroxylase, isopiperitenol dehydrogenase, and pulegone reductase *Plant. Physiol.* **136** 4215–27  
<https://www.ncbi.nlm.nih.gov/pubmed/15542490>
- [4] Davies P J 2010 The plant hormones: Their nature, occurrence, and functions ed Davies PJ *Plant Hormones* (Springer: Dordrecht) pp 1–15  
[https://link.springer.com/chapter/10.1007/978-1-4020-2686-7\\_1](https://link.springer.com/chapter/10.1007/978-1-4020-2686-7_1)
- [5] Gao S L, Zhu D N, Cai Z H, Jiang Y and Xu D R 2004 Organ culture of a precious Chinese medicinal plant—*Fritillaria unibracteata* plant cell *Tiss. Org. Cult.* **59** 197–201  
<https://www.researchgate.net/publication/225220689>
- [6] Zuzarte M R, Dinis A M, Cavaleiro C, Salgueiro L R and Canhoto J M 2010 Trichomes, essential oils and in vitro propagation of *Lavandula pedunculata* (Lamiaceae) *Ind. Crops. Prod.* **32** 580–87  
<http://www.uc.pt/fctuc/dcv/Projectos/ProfJCanhoto/Lavandula>
- [7] Karuppusamy S 2009 A Review on trends in production of secondary metabolites from higher plants by in vitro tissue, organ and cell cultures *J. Med. Plants Res.* **3**(13) 1222–39  
<http://academicjournals.org/journal/JMPR/article-abstract/EEE689215753>
- [8] Debnath M, Malik C P and Bisen P S 2006 Micropropagation: A tool for the production of high quality plant-based medicines *Curr. Pharm. Biotechnol.* **7** 33–49

- <https://www.researchgate.net/publication/7304696>
- [9] Kim J, Seo C and Shin H 2010 Simultaneous determination of (–)-menthone and (–)-menthol in menthae herba by gas chromatography and principal component analysis *Nat. Prod. Sci.* **16**(3) 180–4  
<http://www.papersearch.net/thesis/article.asp?key=2870278>
- [10] Croteau R B, Davis M E, Ringer K L and Wildung M R 2005 Menthol biosynthesis and molecular genetics *Naturwissenschaften* **92** 562–77  
<https://www.ncbi.nlm.nih.gov/pubmed/16292524>
- [11] Santoro M V, Nievas F, Zygadlo J, Giordano W and Banchio E 2013 Effects of growth regulators on biomass and the production of secondary metabolites in peppermint (*Mentha piperita*) micropropagated in vitro *Am. J. Plant. Sci.* **4** 49–55  
<http://ri.conicet.gov.ar/handle/11336/22617>
- [12] Oh M Trick H N and Rajashekar C B 2009 Secondary metabolism and antioxidants are involved in environmental adaptation and stress tolerance in lettuce *J. Plant Physiol.* **166** 180–91  
<https://www.sciencedirect.com/science/article/pii/S0176161708001107>
- [13] Murthy H N and Lee E 2014 Production of secondary metabolites from cell and organ cultures: Strategies and approaches for biomass improvement and metabolite accumulation *Plant Cell Tiss. Organ Cult.* **118** 1–16  
<https://link.springer.com/article/10.1007/s11240-014-0467-7>
- [14] Dolzhenko Y, Berteau C M, Occhipinti A, Bossi S and Maffei M E 2010 UV-B Modulates the interplay between terpenoids and flavonoids in peppermint (*Mentha piperita* L.) *J. Photochem. Photobiol. B: Biology* **100** 67–75  
<https://www.sciencedirect.com/science/article/pii/S1011134410001107>
- [15] Wang M and Lincoln D E 2004 Effects of light intensity and artificial wounding on monoterpene production in *Myrica cerifera* from two different ecological habitats *Can. J. Bot.* **82** 1501–08  
<http://www.nrcresearchpress.com/doi/abs/10.1139/b04-107>
- [16] Ringer K L, Davis E M and Croteau R 2005 Monoterpene metabolism. Cloning, expression, and Characterization of (2)-Isopiperitenol/(2)-Carveol Dehydrogenase of Peppermint and Spearmint *Plant Physiology* **137** 863–72  
<http://www.plantphysiol.org/content/137/3/863.short>
- [17] Ringer K L, Davis E M and Croteau R 2005 Monoterpene metabolism. Cloning, expression, and characterization of (2)-Isopiperitenol/(2)-Carveol dehydrogenase of peppermint and spearmint *Plant. Physiol.* **137** 863–72  
<http://www.plantphysiol.org/content/137/3/863.short>
- [18] Prins C L, Vieira I J C and Freitas S P 2010 Growth regulators and essential oil production *Braz. J. Plant Physiol.* **22**(2) 91–102  
[http://www.scielo.br/scielo.php?pid=S1677-04202010000200003&script=sci\\_arttext](http://www.scielo.br/scielo.php?pid=S1677-04202010000200003&script=sci_arttext)

Final Review The 2nd NRLS Ubaya x

mail.google.com/mail/u/1/#inbox/FMfcgxwBVgtkcDwXrXBgDrKbwZnCtZlg

Gmail Search mail

Compose

Inbox 3,957

Starred

Snoozed

Important

Sent

Poppy +

hidayah murtiyaningsih

Joule Hardjo

Final Review The 2nd NRLS Ubaya

Rangga Mahaswa <rangga.mahaswa@gmail.com> to me

Tue, Feb 19, 9:57 AM

Yth. Poppy H. Hardjo

Selamat Pagi,

Perkenalkan saya Rangga Kala Mahaswa (salah satu editor teknis untuk The 2nd NRLS Ubaya).

Berikut saya lampirkan naskah dan beberapa catatan yang tercantum di balon komentar pada manuskrip yang berjudul "Menthol from Stem and Leaf in-vitro Mentha piperita".  
Penyelarasan akhir ini akan berlangsung secara bertahap, kami harap proses perbaikan dapat diselesaikan sebelum 23 Februari 2019.

Sebagai catatan, ada dua komentar jika yang berbahasa Inggris berarti dari bagian reviewer substansi dan dapat ditanggapi atau direvisi sekaligus, sedangkan komentar berbahasa Indonesia dari saya selaku editor teknis.  
Mohon untuk segera menghubungi saya apabila terdapat kesulitan dalam proses perbaikan melalui email ini atau no. wa saya 085726387667.

Terima kasih atas perhatian dan kerjasamanya.

Tim Editor Teknis

Revisi manuskrip Menthol - pop x

mail.google.com/mail/u/1/#sent/QgrCJHsNhMptUKjczbPFgQChLXsvNMwFBI

Gmail in:sent

Compose

Inbox 3,956

Starred

Snoozed

Important

Sent

Poppy +

hidayah murtiyaningsih

Joule Hardjo

Revisi manuskrip Menthol

Poppy Hardjo <poppyhardjo@gmail.com> to Rangga

Thu, Mar 7, 6:14 PM

Dear Editor,

berikut saya kirimkan ulang hasil revisi manuskrip sesuai arahan Editor Mas Rangga (yaitu revisi no referensi yang sama, no 15 dan 16)

terima kasih bantuan dan kerjasamanya.

Salam  
Poppy

Fullpaper-2.4 [Rev ...]

# Full\_Paper-3.4.doc

*by* Bintang Wijaya

---

**Submission date:** 28-Nov-2018 11:21AM (UTC+0700)

**Submission ID:** 1046161820

**File name:** Full\_Paper-3.4.doc (527.5K)

**Word count:** 3172

**Character count:** 15441

## *Menthol from in vitro Stem and Leaf Mentha piperita*

Formatted: Font: Times New Roman, 17 pt, Bold

Formatted: subsection

BK Wijaya<sup>1</sup>, PH Hardjo<sup>1</sup> and S Emantoko<sup>1</sup>

<sup>1</sup>Faculty of Biotechnology, University of Surabaya, Raya Kalirungkut Road, Surabaya 60292, East Java, Indonesia, Phone: +6231 2981399, Fax: +62 31 2981278

poppyhardjo@gmail.com

**Abstract.** The need for menthol in Indonesia is increasing annually, but its availability isn't enough to fulfill the need. Therefore, in this research we conducted a study using plant tissue culture techniques to produce secondary metabolite especially menthol using shoot multiplication. Since nodes of *Mentha piperita* were inoculated on three kinds of growth medium, which are MS + 0.1 mg L<sup>-1</sup> NAA + 0.1 mg L<sup>-1</sup> BAP, MS + 0.5 mg L<sup>-1</sup> NAA, and MS + 2 mg L<sup>-1</sup> NAA. Based on fresh weight and morphological observation, MS + 0.1 mg L<sup>-1</sup> NAA + 0.1 mg L<sup>-1</sup> BAP was the best growth medium. Based on the previous experiment, the study was aiming to continue observing the concentration of menthol found in *Mentha piperita* shoot. Stems and leaves of *Mentha piperita* were dried in room temperature, the dried stems and leaves were extracted using steam distillation method with ratio 0.3 g of dried stems and leaves in 25 mL aquadest. The crude extract was analyzed by gas chromatography (HP 6890) using INNOWAX 19095N-123 column. The results showed that two weeks was the best amount of time to obtain highest concentration of menthol specifically 1.218.5 ± 47.1 mg L<sup>-1</sup> (yield = 9.748%).

**Keywords:** Gas chromatography, *Mentha piperita*, menthol, shoot multiplication, steam distillation

### 1. Introduction

*Mentha* spp. belongs to Labiatae family, there are three main species which have high economic value like *Mentha piperita*. The industrial demands of *Mentha piperita* product is high but Indonesia cannot fulfill the demands without import. In 2006 Indonesia imported *Mentha* spp. product worth US \$ 3.78 million [1]. *M. piperita* is first sterile hybrid generation between *M. spicata* and *M. aquatica* [2]. The main compound from *Mentha* essential oil is menthol, this compound is produced in peltate glandular trichome which located at leaf [3].

The purpose of this research is to increase the production of menthol using plant tissue culture technique. Cytokinin is phytohormone that induced shoot proliferation and cell division [4]. Plant tissue culture is known to produce plants that grow faster [5]. In addition, the concentration of secondary metabolites, especially monoterpenes in *Lavandula pedunculata* in vitro, is known to be greater than plants grown in their natural [6]. The growth medium used for multiplication of shoots with the aim of increasing menthol concentration is Murashige & Skoog media (MS) + 6-Benzylaminopurine (BAP) + Naphthaleneacetic acid (NAA) [7]. The advantage of in vitro shoot multiplication techniques is that it is able to produce target compounds under controlled conditions from changes in weather and soil conditions, plants obtained free from microorganisms and insects, all types of plants can be used in this technique, and reduce labor costs and increase productivity [8].

### 2. Materials and Methods

#### 2.1. Plant Materials, treatment and growing condition

This research was carried out at the Laboratory of Plant Biotechnology and Biopurification and Biomolecular Laboratory, Faculty of Biotechnology, University of Surabaya. The peppermint seeds

(Biopot®) used in this study were pre-sterilized using bactericidal and fungicidal solution for an hour. The surface sterilization of explants was continued in Laminar Air Flow Cabinet with immersion in 70% ethanol for 1 minute, and 5.25% NaOCl solution for 5 minutes followed by rinsing with sterile H<sub>2</sub>O at least three times. Furthermore, sterilization was carried out again by immersion in 1.75% NaOCl solution for 10 minutes followed by rinsing with a sterile water at least three times. Sterilized mint seeds are planted on MS media without hormones. The culture bottle was incubated under the white fluorescent lamps. Explants used are nodes from mint plants that are approximately 3 months old. The plant is cut to only one node. Furthermore, nodules cut into various growth media (Table 1.) The culture was incubated in a culture room with a temperature of  $25 \pm 2^\circ\text{C}$  with the lighting of white fluorescent lamps and observed the formed tissue.

**Table 1.** Growth Media Variation

Media
MS + 0.1 mg L <sup>-1</sup> 2,4-D
MS + 1 mg L <sup>-1</sup> 2,4-D
MS + 2 mg L <sup>-1</sup> 2,4-D
MS + 4 mg L <sup>-1</sup> 2,4-D
MS + 6 mg L <sup>-1</sup> 2,4-D
MS + 0.5 mg L <sup>-1</sup> NAA
MS + 2 mg L <sup>-1</sup> NAA
MS + 4 mg L <sup>-1</sup> NAA
MS + 6 mg L <sup>-1</sup> NAA
MS + 8 mg L <sup>-1</sup> NAA
MS + 2 mg L <sup>-1</sup> NAA + 0.5 mg L <sup>-1</sup> BAP
MS + 2 mg L <sup>-1</sup> NAA + 1 mg L <sup>-1</sup> BAP
MS + 2 mg L <sup>-1</sup> NAA + 2 mg L <sup>-1</sup> BAP
MS + 1 mg L <sup>-1</sup> BAP
MS + 2 mg L <sup>-1</sup> BAP
MS + 2 mg L <sup>-1</sup> BAP + 1 mg L <sup>-1</sup> NAA
MS + 0.1 mg L <sup>-1</sup> NAA + 0.1 mg L <sup>-1</sup> BAP

## 2.2. Extraction and determination of Menthol

The extraction samples used were leaves and stems of mint plants obtained from the bud multiplication process that have been dried in room temperature under air conditioning. The solvent used was 25 mL of water (distilled water) to extract 0.3 g of dried *Mentha piperita* stems and leaves. The sample was extracted using the steam distillation method. Then the extract obtained was analyzed for menthol content using a gas chromatography device (HP 6890) with INNOWAX 19095N-123 column [9]. For the ex vitro sample used stem and leaves of *Mentha piperita* ex vitro at 3 months old.

## 2.3. Data analyses

The research conducted was an experimental study in the laboratory with a Completely Randomized Design (CRD). In this study two non-parametric data were obtained because the data wasn't normally distributed, namely the best media used to grow *Mentha piperita* plants in vitro, and data on menthol production monitored every week. To determine the best growth media, and the best time to harvest, need to obtain fresh weight of *Mentha piperita* in vitro, and menthol concentration for seven weeks. The statistic test used in this research is Kruskal-Wallis test and if significance found in the Kruskal-Wallis test continue using Dunn-Bonferroni as post hoc test.



### 3. Results and Discussions

#### 3.1. Preliminary Test


Preliminary tests were carried out to determine the 3 types of growth media to be tested further, the aspects considered were morphology of *Mentha piperita* grown in vitro. From the preliminary test selected, MS medium + 0.1 mg L<sup>-1</sup> NAA + 0.1 mg L<sup>-1</sup> BAP, MS + 0.5 mg L<sup>-1</sup> NAA and MS + 2 mg L<sup>-1</sup> NAA to be used at the next research stage because it has the best morphology. The reason for choosing MS media was 0.5 mg L<sup>-1</sup> NAA and MS + 2 mg L<sup>-1</sup> NAA compared to MS + 1 mg L<sup>-1</sup> BAP and MS + 2 mg L<sup>-1</sup> BAP, because the hormones NAA and IAA (Indole-3-Acetic Acid) were able to increase the oil content of essential oils and menthol in *Mentha piperita* [10]. Then MS + 0.5 mg L<sup>-1</sup> NAA and 2 mg L<sup>-1</sup> NAA has a better morphology than MS + 2 mg L<sup>-1</sup> BAP + 1 mg L<sup>-1</sup> NAA, the morphology in question is the appearance of the plant (abnormal).

Table 2. Preliminary test result

Media	Tissue/Organ Formed
MS + 0.5 mg L <sup>-1</sup> 2,4-D	Callus
MS + 1 mg L <sup>-1</sup> 2,4-D	Callus
MS + 2 mg L <sup>-1</sup> 2,4-D	Callus
MS + 4 mg L <sup>-1</sup> 2,4-D	Browning
MS + 6 mg L <sup>-1</sup> 2,4-D	Browning
MS + 0.5 mg L <sup>-1</sup> NAA	Shoot + callus
MS + 2 mg L <sup>-1</sup> NAA	Shoot + callus
MS + 4 mg L <sup>-1</sup> NAA	Browning
MS + 6 mg L <sup>-1</sup> NAA	Browning
MS + 8 mg L <sup>-1</sup> NAA	Browning
MS + 2 mg L <sup>-1</sup> NAA + 0.5 mg L <sup>-1</sup> BAP	Shoot + callus
MS + 2 mg L <sup>-1</sup> NAA + 1 mg L <sup>-1</sup> BAP	Browning
MS + 2 mg L <sup>-1</sup> NAA + 2 mg L <sup>-1</sup> BAP	Browning
MS + 1 mg L <sup>-1</sup> BAP	Shoot + callus
MS + 2 mg L <sup>-1</sup> BAP	Shoot + callus
MS + 2 mg L <sup>-1</sup> BAP + 1 mg L <sup>-1</sup> NAA	Shoot + callus
MS + 0.1 mg L <sup>-1</sup> NAA + 0.1 mg L <sup>-1</sup> BAP	Shoot

#### 3.2. Growth Index Curve

Growth Index Curve is obtained from the final weight of the stem and fresh leaves divided by fresh initial weight. The calculation of the growth index was done by weighing fresh initial weight before planting and after being cultivated for 1, 2, 3, 4, 5, 6 and 7 weeks. Observation of the growth index stopped at week 7 because the index value had decreased compared to the 6th week.




**Figure 1.** *Mentha piperita* growth index curve on MS media + 0.1 mg L<sup>-1</sup> NAA + 0.1 mg L<sup>-1</sup> BAP, MS + 0.5 mg L<sup>-1</sup> NAA and MS + 2 mg L<sup>-1</sup> NAA.

**Table 3.** Fresh Weight of *Mentha piperita* at 6 weeks old.

Media	Fresh Weight (g)
MS + 0.1 mg L <sup>-1</sup> NAA + 0.1 mg L <sup>-1</sup> BAP	0.136 ± 0.071
MS + 0.5 mg L <sup>-1</sup> NAA	0.129 ± 0.035
MS + 2 mg L <sup>-1</sup> NAA	0.179 ± 0.121

The three growth media did not have a significant effect using the Kuskal-Wallis test at  $\alpha = 0.05$




**Figure 2.** Morphology of *Mentha piperita* in vitro.

Fresh weight of *Mentha piperita* grown on MS + 2 mg L<sup>-1</sup> NAA is the largest among other growth media. This is due to the presence of callus at the base of the plant stem. Callus also found at the base of the plant stem that grown using MS + 0.5 mg L<sup>-1</sup> NAA. *Mentha piperita* that grew at MS + 0.1 mg L<sup>-1</sup> NAA + 0.1 mg L<sup>-1</sup> BAP obtained plants do not form callus and have a heavier mass than MS media + 0.5 mg L<sup>-1</sup> NAA. Menthol compounds are stored in the peltate glandular trichome [3]. The peltate glandular trichome is found in parts of plants exposed to air [11]. So it can be said that the larger/the heavier plant, the more peltate glandular trichome, in other words, the more menthol that can be extracted. In addition, other factors considered by the morphology of plants planted in vitro. There was


no monoterpene found in *Mentha spicata* callus, because the glandular oil gland or peltate glandular trichome had not been formed because callus cells were undifferentiated plant cells [12]. Therefore, the MS medium was selected + 0.1 mg L<sup>-1</sup> NAA + 0.1 mg L<sup>-1</sup> BAP as the best growth medium.

### 3.3. Extraction and Determination of Menthol

Extraction of menthol compounds was carried out using the steam distillation method because it is commonly used to isolate essential oils. The distillation process is stopped until all the water in the flask has evaporated. From steam distillation results obtained ± 24 ml of crude menthol extract from steam distillation. The crude extract obtained was analyzed using a gas chromatography device (HP 6890) with the INNOWAX 19095N-123 column.




**Figure 3.** Chromatogram of crude extract *Mentha piperita* Grown on MS + 0.1 mg L<sup>-1</sup> NAA + 0.1 mg L<sup>-1</sup> BAP medium at 2 weeks old.



**Figure 4.** Chromatogram of crude extract *Mentha piperita* grown on ex vitro.

There is a difference between the crude extracts of *Mentha piperita* in vitro and ex vitro, the peaks that appeared at retention time were around 5 minutes in the crude extract profile of ex vitro plants but

were not found in crude plant extracts in vitro. And the crude extracts of plants in vitro found peaks at retention time about 8 minutes which were not found in crude extracts of ex vitro plants. In other words, the *Mentha piperita* plant in vitro is not capable of producing secondary metabolites that are exactly the same as their original plants and is able to form secondary metabolites that are completely different from the original plants. Several factors that influence the difference in results is nutrition availability and environmental conditions. The availability of nutrients includes the concentration of sugar, nitrate, phosphate and growth regulating substances. Whereas environmental conditions include temperature, light intensity and pH [13]. Plant tissue culture systems are able to produce secondary metabolites that are completely different from the original plants because of very different environmental conditions, while the levels can be the same, larger or smaller [14]. In vitro plant growth media have been given nutrition, sufficient growth regulating substances, and the growth environment is conditioned stable with adequate lighting, temperature of 20°C, and pH of 5.8. All treatments given to plants in vitro are common treatments to optimize growth in plant tissue culture. While the ex vitro plants all of these factors are not controlled at all, or in other words depending on nature.



**Figure 3.** Menthol concentration curve and *Mentha piperita* growth index at 0.1 mg L<sup>-1</sup> NAA and 0.1 mg L<sup>-1</sup> BAP compared to incubation time

**Table 4.** Menthol concentration observed every week grown at MS + 0.1 mg L<sup>-1</sup> BAP + 0.1 mg L<sup>-1</sup> NAA

Plant Age (Weeks)	Menthol Concentration (mg L <sup>-1</sup> )	Yield (% w/w)
0	231.3 <sup>ab</sup> ± 59.9	1.85
1	586.3 <sup>ab</sup> ± 126.6	4.69
2	1 218.5 <sup>a</sup> ± 47.1	9.748
3	599 <sup>ab</sup> ± 179	4.792
4	366.5 <sup>ab</sup> ± 85.4	2.932
5	405.6 <sup>ab</sup> ± 50.7	3.245
6	228.997 <sup>ab</sup> ± 0.642	1.832
7	314.93 <sup>ab</sup> ± 9.58	2.519
Ex vitro plant	97.21 <sup>b</sup> ± 1.373	0.778

The different annotation followed by average value showed significance using Dunn-Befferoni test  $\alpha = 0.05$

The highest concentration of menthol was 2 weeks old *Mentha piperita* that have grown using MS + 0.1 mg L<sup>-1</sup> NAA + 0.1 mg L<sup>-1</sup> BAP, and the least concentration of menthol was ex vitro *Mentha piperita*. This phenomenon is not in accordance with the theory, that the production of secondary metabolites generally occurs at the end of the stationary period when the nutrient supply is depleted [4]. This may be caused by menthol has been further metabolized to menthyl acetate [15]. There are also indications that the terpenoids used for defense have decreased due to the absence of threats. Another factor that might cause the highest phenomenon of menthol concentration in the second week of culture is stress caused by injury at the beginning of cutting explants. This phenomenon was also showed on another research, where at 2 weeks after the plant was injured there was a significant increase in the concentration of monoterpenes compared to non-injured plants [16]. This increase in the concentration of monoterpenes may be caused by an increase in the activity of the enzyme monoterpene cyclase as occurs in *Abies grandis* plants [17]. Monoterpene cyclase plays a role in converting Geranyl pyrophosphate (GPP) to 4s-limonene [18]. GPP is the earliest compound in menthol biosynthesis pathway or it can be said that it is a compound that initiates menthol formation so that plants will tend to produce menthol in this condition. *Mentha piperita* grown in vitro is able to produce more menthol than ex vitro. This may be due to the availability of sufficient nutrients so that in vitro plants are able to produce more menthol. Besides that the hormones NAA and IAA (Indole-3-Acetic Acid) are able to increase the levels of essential oil and menthol oil in *Mentha piperita* [10]. There are another research showed that increasing BAP and sugar concentration on media didn't increase the production of menthol in *Mentha piperita* [19].

#### 4. Conclusions

Based on this research, it can be concluded that the best growth medium for multiplying *Mentha piperita* shoots in vitro was MS + 0.1 mg L<sup>-1</sup> NAA + 0.1 mg L<sup>-1</sup> BAP. Chromatogram profile on crude extracts of *Mentha piperita* plant in vitro showed 3 peaks, namely at about 2 retention times: 4.5 and 8 minutes. Chromatogram profile on crude extract of *Mentha piperita* ex vitro showed 3 peaks, namely at about 2 retention times: 4.5 and 5 minutes. The menthol concentration observed for 7 weeks in the largest *Mentha piperita* plant in vitro was at the culture time of the 2<sup>nd</sup> week which was 1 218.5 ± 47.1 mg L<sup>-1</sup>. The menthol concentration observed in the *Mentha piperita* ex vitro plant was 97.21 ± 1.373 mg L<sup>-1</sup>.

#### 5. References

- [1] BPS 2007 Statistik Impor tahun 2000 – 2006 BPS Indonesia.
- [2] Hefendehl F W and Murray J M 1972 Changes in Monoterpene Composition in *Mentha aquatica* Produced by Gene Substitution *Phytochemistry* **11** 189-95.
- [3] Turner G W and Croteau R 2004 Organization of Monoterpene Biosynthesis in *Mentha*: Immunocytochemical Localizations of Geranyl Diphosphate Synthase, Limonene-6-Hydroxylase, Isopiperitenol Dehydrogenase, and Pulegone Reductase *Plant Physiology* **136** 4215–27.
- [4] Bhojwani S S and Razdan M K 1996 *Plant Tissue Culture: Theory and Practice. a Revised Edition* India Elsevier Science B. V.
- [5] Gao S L, Zhu D N, Cai Z H, Jiang Y and Xu D R 2004 Organ Culture of a Precious Chinese Medicinal Plant – *Fritillaria unibracteata* *Plant Cell Tiss. Org. Cult.* **59** 197-201.
- [6] Zuzarte M R, Dinis A M, Cavaleiro C, Salgueiro L R and Canhoto J M 2010 Trichomes, essential oils and in vitro propagation of *Lavandula pedunculata* (Lamiaceae) *Industrial Crops and Products* **32** 580-7.
- [7] Karuppusamy S 2009 A Review on Trends in Production of Secondary Metabolites from Higher Plants by In vitro Tissue, Organ and Cell Cultures *Journal of Medicinal Plants Research* **3**(13) 1222-39.
- [8] Debnath M, Malik C P and Bisen P S 2006 Micropropagation: A Tool for the Production of High Quality Plant-based Medicines *Current Pharmaceutical Biotechnology* **7** 33-49.

- [9] Iscan G, Kirimer N, Kurkcuglu M, Baser K H C, Demirci F 2002 Antimicrobial Screening of *Mentha piperita* Essential Oils *J. Agric. Food Chem.* **50** 3943–3946.
- [10] Sangwan N S, Farooqi A H A, Shabih F and Sangwan R S 2001 Regulation of Essential Oil Production in Plants *Plant Growth Regulation* **34** 3–21.
- [11] Croteau R B, Davis M E, Ringer K L and Wildung M R 2005 Menthol Biosynthesis and Molecular Genetics *Naturwissenschaften* **92** 562-77.
- [12] Suga T, Hirata T and Yamamoto T 1980 Lipid Constituents of Callus Tissues of *Mentha spicata* *Agric. Biol. Chem.* **44**(8) 1817-1820.
- [13] Rao S R and Ravinshankar G A 2002 Plant Cell Cultures : Chemical Factories of Secondary Metabolite *Biotechnology Advances* **20** 101-53.
- [14] Indrayanto G 1987 Produksi Metabolit Sekunder dengan Teknik Kultur Jaringan Tanaman *Buku Risalah Seminar Nasional Metabolit Sekunder* 32-40.
- [15] Akhila A 1986 Biosynthesis of Menthol and Related Monoterpenes in *Mentha arvensis* *J. Plant Physiol.* **126** 379-86.
- [16] Wang M and Lincoln E 2004 Effects of Light Intensity and Artificial Wounding on Monoterpene Production in *Myrica cerifera* from Two Different Ecological Habitats *Can. J. Bot.* **82** 1501-8.
- [17] Gijzen M, Lewinsohn E and Croteau R 1991 Characterization of the Constitutive and Wound-Inducible Monoterpene Cyclases of Grand Fir (*Abies grandis*) *ARCHIVES OF BIOCHEMISTRY AND BIOPHYSICS* **289**(2) 267-73.
- [18] Alonso W R, Rajaonarivony J I M, Gershenzon J and Croteau R 1992 Purification of 4s-Limonene Synthase, a Monoterpene Cyclase from the Glandular Trichomes of Peppermint (*Mentha piperita*) and Spearmint (*Mentha spicata*) *The Journal of Biological Chemistry* **267**(11) 7582-7.
- [19] Devi H P A S, Wiendi N M A 2017 Produksi Biomassa dan Kandungan Mentol Secara In vitro Melalui Proliferasi Tunas Tanaman *Mentha piperita* L. dengan Penambahan Kitosan, BA, dan Gula IPB Scientific Repository Indonesia.

ORIGINALITY REPORT

20%

SIMILARITY INDEX

18%

INTERNET SOURCES

17%

PUBLICATIONS

14%

STUDENT PAPERS

MATCH ALL SOURCES (ONLY SELECTED SOURCE PRINTED)

21%

★ "Biotechnological Approaches for Medicinal and Aromatic Plants", Springer Nature America, Inc, 2018

Publication

Exclude quotes Off

Exclude bibliography On

Exclude matches < 4 words

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8