

PAPER • OPEN ACCESS

## Molecular docking, drug-likeness, and ADMET study of 1-benzyl-3-benzoylurea and its analogs against VEGFR-2

To cite this article: F Suhud *et al* 2019 *IOP Conf. Ser.: Earth Environ. Sci.* **293** 012018

View the [article online](#) for updates and enhancements.



**IOP | ebooks™**

Bringing you innovative digital publishing with leading voices to create your essential collection of books in STEM research.

Start exploring the collection - download the first chapter of every title for free.

# Molecular docking, drug-likeness, and ADMET study of 1-benzyl-3-benzoylurea and its analogs against VEGFR-2

F Suhud<sup>1\*</sup>, D H Tjahjono<sup>2</sup>, T A Yuniarta<sup>1</sup>, G S Putra<sup>1</sup> and J Setiawan<sup>1</sup>

<sup>1</sup>Faculty of Pharmacy, University of Surabaya, Jalan Raya Kalirungkut, Surabaya, 60294 Indonesia

<sup>2</sup>School of Pharmacy, Bandung Institute of Technology, Jalan Ganesha 10, Bandung, 40132 Indonesia

\*Corresponding author: faridasuhud@yahoo.com

**Abstract.** *In silico* study was performed to predict the possibility of 1-benzyl-3-benzoylurea and 22 analogs as anticancer drug candidates, via VEGFR2 inhibition. Molecular docking studies against VEGFR2 receptor revealed that all of designed compounds have better score than the lead compound, of which three analogs (p-nitro, p-methoxy, and p-ethyl) were considered optimal among other compounds ( $< -90$  kcal mol<sup>-1</sup>). However, this result was not comparable to lenvatinib, which acts as native ligand of the receptor (-118.62 kcal mol<sup>-1</sup>). Docking poses analysis showed that 1-benzyl-3-benzoylurea analogs failed to completely occupy VEGFR2 binding site. Therefore, it is argued that this has caused the non-optimal docking score of designed compounds. Furthermore, these compounds passed five different drug-likeness criteria successfully and were predicted to be orally bioavailable in rat. Ultimately, most of the analogs were predicted to have good ADMET characteristics, notably in terms of GI absorption and the absence of P-gp interaction, and low toxicity in rat. This study can be used as a starting point to validate this model by synthesis, *in vitro* and *in vivo* assay to validate the activity of 1-benzyl-3-benzoylurea and its analogs as potential anticancer candidate.

**Keywords:** Anticancer candidate, arylurea derivatives, *in silico* study, VEGFR-2 inhibitor.

## 1. Introduction

Cancer is a terminology which is commonly used to describe the abnormality in cell growth and division [1]. In 2012, the number of population with cancer has reached  $14.1 \times 10^6$  people, with  $8.2 \times 10^6$  mortality [2]. In Indonesia itself, the prevalence of cancer is 1.4 % of total population [3]. Hydroxyurea has been used for decades and still possesses valuable therapeutic activity against several type of cancer. This hydrophilic compound is distributed evenly in the human body fluid. Under physiologic condition (pH 7.4), the majority of hydroxyurea will undergo ionization, thus reducing its ability to penetrate biological membrane. This condition will significantly diminished its biological activity [4].

Several analog of urea based compounds have been widely developed, one of which is arylurea. Numerous studies suggested that this modification showed potential anticancer activity. Furthermore, preliminary study has proven the anticancer activity of 1-benzyl-3-benzoylurea, one of the analog of benzoylurea. This compound inhibits the growth of MCF-7 cell culture in breast cancer better than hydroxyurea [5]. Based on the finding, it can be concluded that 1-benzyl-3-benzoylurea is potential to be further developed as anticancer agent. This premise is supported by the number of anticancer drugs




in the market which contain arylurea functional group, such as sorafenib and lenvatinib. These two compounds are known to act as VEGFR-2 inhibitor [6, Topliss in 7] (figure 1).

This research aimed to explore the possibility to improve the anticancer activity of 1-benzyl-3-benzoylurea analogs using *in silico* method. Several novel compounds were designed and predicted their activity using molecular docking against VEGFR-2. Furthermore, the drug-likeness and ADMET properties of those compounds were also evaluated.

## 2. Materials and methods

### 2.1. Preparation of 1-benzyl-3-benzoylurea analogs

Total of 23 novel analogs of 1-benzyl-3-benzoylurea were designed by implementing Topliss scheme [Topliss in 7], of which the steric and electronic properties of every substituent were taken into account. These compounds were drawn in 2D format (MarvinSketch 16.8.1) (<http://www.chemaxon.com>) and then transformed into 3D structure using semiempirical method of AM1 [8] available in MOPAC2016 (<http://openmopac.net/MOPAC2016.html>).



**Figure 1.** Chemical structure of 1-Benzyl-3-Benzoylurea (a) and lenvatinib (b).

### 2.2. Preparation of protein

VEGFR-2 protein was obtained from Brookhaven Protein Data Bank (PDB) (<https://www.rcsb.org/pdb/>) with PDB ID of 3WZD. This protein was chosen since it is complexed with lenvatinib, an arylurea analog [6]. Hydrogen atom was then added, followed by the assignment of Amber ff14SB partial charge [9]. Correction of sidechain residue was also performed by employing the most recent Dunbrack rotamer library [10]. The purpose of this step was to reconstruct the missing information of amino acid sidechain. Lenvatinib was also prepared by adding hydrogen atom. The whole process was performed using Chimera 1.13 [11].

### 2.3. Molecular docking

Molegro Virtual Docker 5.0 (Molegro ApS, 2013) was applied to evaluate the interaction of designed compounds with VEGFR-2.

### 2.4. ADMET and drug-likeness evaluation

Drug candidates should possess favorable ADME properties and ideally non-toxic. Therefore, the designed compounds were evaluated of their ADME profile, including drug-likeness, partition coefficient, solubility, and several other parameters using SwissADME [12] module provided in SIB (Swiss Institute of Bioinformatics) webserver (<https://www.sib.swiss>). Furthermore, the toxicity aspect of designed compound was also predicted using webserver ProTox [13] (<https://tox.charite.de>).

## 3. Results

### 3.1. Molecular docking


Molecular docking was performed to evaluate the interaction mode of designed compounds against VEGFR-2 and to assess the magnitude of interaction between them. Similar study have been performed [14] using similar receptor, albeit complexed with different arylurea analog [15]. This study

was conducted in order to verify the previous result using slightly modified method. Docking method needs to be validated by redocking the native ligand to the protein prior to the usage [16]. Based on redocking result of lenvatinib to its protein [6], RMSD result of 0.7 was obtained. This implies that the docking method used is able to predict the ligand orientation accurately. Docking method selected in this study was simplex evolution as pose generator which was evaluated using grid based MolDock score [17]. The findings showed that no designed compounds performed better than native ligand, in terms of docking score (table 1).


Docking pose analysis of three best performing analogs (Compound 11, 22, and 23) showed the dissimilarity of binding mode compared to Lenvatinib. Similarity is only observed in the urea moiety of both compound, which is with Glu 885 and Asp 1046. Lenvatinib as the newly discovered VEGFR-2 inhibitor classified as type V inhibitor, interact in a similar way with Sorafenib as the older, type II VEGFR-2 inhibitor [6, Topliss in 7]. Therefore, the pharmacophore motive for Sorafenib-like, type II VEGFR-2 inhibitor [18] could also be applicable for type V inhibitor. However, docking pose of all designed compounds showed occupation only in the RDP (Regulatory Domain Pocket) area. While type II inhibitor and Lenvatinib also occupy ATP binding domain. This made the docking score of 1-benzyl-3-benzoylurea analogs are higher than lenvatinib and sorafenib [14] (figure 2).

### 3.2. ADMET and drug-likeness evaluation

SwissADME provides detail and extensive physicochemical profile, ADME, and medicinal chemistry property of a compound. Regarding the physicochemical aspect, two parameters (partition coefficient and solubility) are considered to play important roles. Based on predicted LogP value, it is concluded that all of designed compounds lies within the range value of 1.6 to 3.6. Eventhough LogP value does not always correspond to certain ADME aspect, this parameter could depict the probability of a compound as drug candidate, where in this case all 23 compounds possess such quality to be considered as drug-like [19]. SwissADME LogP value was calculated from five different algorithm, therefore it is assumed that the value represents real condition [12]. On the other hand, solubility prediction was carried out using three different method with the output of LogS value. Generally, the designed compounds are predicted to have optimum water solubility, even if there was a slightly different result among the method. The most obvious difference was observed in SILICOS-IT (<http://silicos-it.be.s3-website-eu-west-1.amazonaws.com/software/filter-it/1.0.2/filter-it.html>) method since it employs fragment-based approach in LogS calculation, while the other methods [20, 21] based on complete molecular topology (table 2).



**Figure 2.** Docking pose superposition of one of 1-Benzyl-3-Benzoylurea analogs (compound 23) (light blue) with Lenvatinib (gray). It showed a dissimilarity in binding mode between those two compounds, notably in urea moiety (highlighted).

**Table 1.** Docking score of 1-Benzyl-3-Benzoylurea and its analogs against VEGFR-2.

Code	Compound	Substituents	Docking Score (kcal mol <sup>-1</sup> )
1	1-benzyl-3-benzoylurea	R1=H;R2=H;R3=H	-73.95
2	1-benzyl-3-(2-chloro)-benzoylurea	R1=Cl;R2=H;R3=H	-76.92
3	1-benzyl-3-(3-chloro)-benzoylurea	R1=H;R2=Cl;R3=H	-80.38
4	1-benzyl-3-(4-chloro)-benzoylurea	R1=H;R2=H;R3=Cl	-81.71
5	1-benzyl-3-(2,4-dichloro)-benzoylurea	R1=Cl;R2=H;R3=Cl	-79.14
6	1-benzyl-3-(3,4-dichloro)-benzoylurea	R1=H;R2=Cl;R3=Cl	-79.83
7	1-benzyl-3-(4-chloromethyl)-benzoylurea	R1=H;R2=H;R3=CH <sub>2</sub> Cl	-75.48
8	1-benzyl-3-(3-chloromethyl)-benzoylurea	R1=H;R2=CH <sub>2</sub> Cl;R3=H	-87.19
9	1-benzyl-3-(2-chloromethyl)-benzoylurea	R1=CH <sub>2</sub> Cl;R2=H;R3=H	-79.72
10	1-benzyl-3-(4-methyl)-benzoylurea	R1=H;R2=H;R3=CH <sub>3</sub>	-74.15
11	1-benzyl-3-(4-ethyl)-benzoylurea	R1=H;R2=H;R3=C <sub>2</sub> H <sub>5</sub>	-90.73
12	1-benzyl-3-(3-ethyl)-benzoylurea	R1=H;R2=C <sub>2</sub> H <sub>5</sub> ;R3=H	-78.99
13	1-benzyl-3-(2-ethyl)-benzoylurea	R1=C <sub>2</sub> H <sub>5</sub> ;R2=H;R3=H	-78.6
14	1-benzyl-3-(4-propyl)-benzoylurea	R1=H;R2=H;R3=C <sub>3</sub> H <sub>7</sub>	-83.51
15	1-benzyl-3-(4-t-butyl)-benzoylurea	R1=H;R2=H;R3=C(CH <sub>3</sub> ) <sub>3</sub>	-69.76
16	1-benzyl-3-(4-fluoro)-benzoylurea	R1=H;R2=H;R3=F	-82.21
17	1-benzyl-3-(2-trifluoromethyl)-benzoylurea	R1=CF <sub>3</sub> ;R2=H;R3=H	-81.13

Continue to the next page.

**Table 1.** Continued.

Code	Compound	Substituents	Docking Score (kcal mol <sup>-1</sup> )
18	1-benzyl-3-(3-trifluoromethyl)-benzoylurea	R1=H;R2=CF <sub>3</sub> ;R3=H	-87.26
19	1-benzyl-3-(4-trifluoromethyl)-benzoylurea	R1=H;R2=H;R3=CF <sub>3</sub>	-79.78
20	1-benzyl-3-(4-bromo)-benzoylurea	R1=H;R2=H;R3=Br	-76.53
21	1-benzyl-3-(4-bromomethyl)-benzoylurea	R1=H;R2=H;R3=CH <sub>2</sub> Br	-74.34
22	1-benzyl-3-(4-nitro)-benzoylurea	R1=H;R2=H;R3=NO <sub>2</sub>	-95.2
23	1-benzyl-3-(4-methoxy)-benzoylurea	R1=H;R2=H;R3=OCH <sub>3</sub>	-90.91
	Lenvatinib		-118.62

ADME prediction showed that the analog compounds possess several favorable ADME properties. Based on ratio of WlogP to tPSA [22], all of the compound was predicted to have good absorption in GIT and 22 of which could penetrate blood-brain barrier. Analog with nitro substituent possesses the lowest partition coefficient, hence too hydrophilic to penetrate blood-brain barrier. Surprisingly, there is no single compound which has a tendency to act as P-gp substrate. P-glycoprotein, a macromolecule found to be overexpressed in multi-drug resistant cancer, has an important role in transporting xenobiotics out from cell [23]. Therefore, this finding indicates that 1-benzyl-3-benzoylurea scaffold could be used as potential anticancer agent. Furthermore, SwissADME also enable one to predict the possible occurrence of CYP450-mediated biotransformation. The result showed that the majority of compound interacts with at least one of the five isoforms, where two of the analogs (4-propyl and 4-bromomethyl) would interact with four out of five isoforms. On the contrary, three of the analogs (4-fluoro, 4-nitro, and 4-methoxy) along with the parent compound possibly would not interact with any of the CYP450 isoforms. It is also concluded that 1A2 and 2C19 are the major isoform which would become the target of most of the analogs, while no interaction is predicted to happen with 3A4 (figure 3).

Drug-likeness is a key criteria in screening drug candidates at the earlier phase of drug discovery and development. This parameter can be described as a mean to correlate physicochemical aspect of a compound with its biopharmaceutical aspect in human body, especially its influence in bioavailability of per oral route [24]. Based on five commonly used drug-likeness criteria applied in the webserver [12], 1-benzyl-3-benzoylurea with its 22 analogs are considered to pass all of them, thus can be categorized as drug-like compounds. In addition, evaluation was also carried out using ABS criteria [25], where all of the compounds obtained the value of 0.55. This criteria is based on the probability value of a compound to possess optimum profile of bioavailability and permeability, where value of 0.55 implies the obedience of Lipinski rule of five [26] and 55 % probability of rat bioavailability value higher than 10 %.


In addition, detection of structural alert [27] was performed in order to identify whether the designed compound possess problematic functional group, which could lead to a toxicity, mutagenicity, or metabolic instability. It showed that five of designed compounds possess undesirable moiety (alkyl halide and nitroarene). Studies have indicated the reactivity of compounds containing such functional groups, which lead to carcinogenic, mutagenic, and hepatotoxic effect [28]. This



criteria could be used as preliminary alert to give more attention to the activity and toxicity in developing 1-benzyl-3-benzoylurea scaffold [29].

**Table 2.** Water solubility prediction of 1-Benzyl-3-Benzoylurea and its analogs.

Compounds	LogP	Water Solubility		
	(Consensus LogP)	LogS (ESOL)	LogS (Ali)	LogS (SILICOS-IT)
1	2.16	Soluble	Soluble	Moderately Soluble
2	2.79	Soluble	Moderately Soluble	Moderately Soluble
3	2.74	Soluble	Moderately Soluble	Moderately Soluble
4	2.73	Soluble	Moderately Soluble	Moderately Soluble
5	3.32	Moderately Soluble	Moderately Soluble	Poorly Soluble
6	3.26	Moderately Soluble	Moderately Soluble	Poorly Soluble
7	2.8	Soluble	Moderately Soluble	Poorly Soluble
8	2.77	Soluble	Moderately Soluble	Poorly Soluble
9	2.78	Soluble	Moderately Soluble	Poorly Soluble
10	2.59	Soluble	Moderately Soluble	Moderately Soluble
11	2.86	Soluble	Moderately Soluble	Poorly Soluble
12	2.86	Soluble	Moderately Soluble	Poorly Soluble
13	2.89	Soluble	Moderately Soluble	Poorly Soluble
14	3.25	Moderately Soluble	Moderately Soluble	Poorly Soluble
15	3.55	Moderately Soluble	Moderately Soluble	Poorly Soluble
16	2.55	Soluble	Soluble	Moderately Soluble
17	3.24	Moderately Soluble	Moderately Soluble	Poorly Soluble
18	3.29	Moderately Soluble	Moderately Soluble	Poorly Soluble
19	3.27	Moderately Soluble	Moderately Soluble	Poorly Soluble
20	2.79	Moderately Soluble	Moderately Soluble	Poorly Soluble
21	2.92	Moderately Soluble	Moderately Soluble	Poorly Soluble
22	1.62	Soluble	Moderately Soluble	Moderately Soluble
23	2.23	Soluble	Soluble	Moderately Soluble



**Figure 3.** Prediction of interaction of designed compounds with several CYP450 isoforms.

Ultimately, prediction of toxicity using ProTox [13] showed that the designed compound are predicted to have oral LD50 value ranging from 818 mg kg<sup>-1</sup> to 3 000 mg kg<sup>-1</sup> in rat model, with analogs of p-methyl and p-trifluoromethyl bearing the lowest value and four analogs (o-chloro, o-chloromethyl, m-chloromethyl, and o-trifluoromethyl) bearing the highest one. It was also observed that the substituent position in aromatic could significantly affect the predicted value of LD50 as was shown in the case of trifluoromethyl (*vide supra*).

#### 4. Conclusion

Molecular docking study showed that three of 1-benzyl-3-benzoylurea analog (p-nitro, p-methoxy, and p-ethyl) possess better docking score among the other analogs with the value around -90 kcal mol<sup>-1</sup>. However, there is no single compound which performs better than Lenvatinib (-118.62 kcal mol<sup>-1</sup>) due to the inability of all designed compound to completely occupy the VEGFR-2 binding site. This hypothesis would be used as a foundation which needs to be proven via *in vitro* assay in the future. Furthermore, it can be implied that the majority of compound have good physicochemical profile with several other ADMET properties, notably in terms of P-gp interaction. Nevertheless, the identification of several analogs with problematic moiety, should be noted in order to develop more safe and potent anticancer agent based on 1-benzyl-3-benzoylurea scaffold.

#### Acknowledgement

This research was supported by Ministry of Research, Technology, and Higher Education of the Republic of Indonesia (Grant No: 120/SP2H/LT/DRPM/2018). The authors would also like to show the author gratitude to Prof. Siswandono (Universitas Airlangga) for the possibility to use Molegro in this research.

#### References

- [1] Avendaño C and Menéndez C J 2015 *Medicinal Chemistry of Anticancer Drugs 2<sup>nd</sup> Edition* (USA: Elsevier)  
<https://www.elsevier.com/books/medicinal-chemistry-of-anticancer-drugs/avendano/978-0-444-62649-3>
- [2] WHO 2015 *World Cancer Report 2014* ed B W Steward and C P Wild (France: IARC)  
<http://publications.iarc.fr/Non-Series-Publications/World-Cancer-Reports/World-Cancer-Report-2014>
- [3] Pusat Data dan Informasi Kementerian Kesehatan RI 2015 *Situasi Penyakit Kanker* [Cancer Situation] (Jakarta, Indonesia: Ministry of Health Indonesia) [in Bahasa Indonesia]  
<http://www.depkes.go.id/resources/download/pusdatin/infodatin/infodatin-kanker.pdf>
- [4] Saban N and Bujak M 2009 Hydroxyurea and hydroxamic acid derivatives as antitumor drugs *Cancer Chemother Pharmacol.* **64**(2) 213–21  
<https://www.ncbi.nlm.nih.gov/pubmed/19350240>
- [5] Suhud F, Siswandono and Budiati T 2015 Synthesis and activity evaluation of a novel lead compound 1-benzyl-3-benzoylurea as antiproliferative agent *World J. Pharm. Sci.* **3**(2) 192–95  
[http://www.wjpsonline.org/view\\_issue.php?title=Synthesis-and-activity-evaluation-of-a-novel-lead-compound-1-benzyl-3-benzoylurea-as-antiproliferative-agent](http://www.wjpsonline.org/view_issue.php?title=Synthesis-and-activity-evaluation-of-a-novel-lead-compound-1-benzyl-3-benzoylurea-as-antiproliferative-agent)
- [6] Okamoto K, Ikemori-Kawada M, Jestel A, von König K, Funahashi Y, Matsushima T, Tsuruoka A, Inoue A and Matsui J 2014 Distinct binding mode of multikinase inhibitor lenvatinib revealed by biochemical characterization *ACS Med. Chem. Lett.* **6**(1) 89–94  
<https://www.ncbi.nlm.nih.gov/pubmed/25589937>
- [7] Richter L 2017 Topliss batchwise schemes reviewed in the era of open data reveal significant differences between enzyme and membrane receptors *J. Chem. Inf. Model* **57**(10) 2575–83  
<https://pubs.acs.org/doi/abs/10.1021/acs.jcim.7b00195>
- [8] Stewart J J 2004 Optimization of parameters for semiempirical methods IV: Extension of MINDO, AM1, and PM3 to more main group elements *J. Mol. Mod.* **10**(2) 155–64  
<https://link.springer.com/article/10.1007%2Fs00894-004-0183-z>



- [9] Maier J A, Martinez C, Kasavajhala K, Wickstrom L, Hauser K E and Simmerling C 2015 ff14SB: Improving the accuracy of protein side chain and backbone parameters from ff99SB *J. Chem. Theory Comput.* **11**(8) 3696–713  
<https://www.ncbi.nlm.nih.gov/pubmed/26574453>
- [10] Shapovalov V and Dunbrack Jr. R L 2011 A smoothed backbone-dependent rotamer library for proteins derived from adaptive kernel density estimates and regressions *Structure* **19**(6) 844–58  
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3118414/>
- [11] Pettersen E F, Goddard T D, Huang C C, Couch G S, Greenblatt D M, Meng E C and Ferrin T E 2004 UCSF chimera—A visualization system for exploratory research and analysis *J. Comput. Chem.* **25**(13) 1605–12  
<https://www.ncbi.nlm.nih.gov/pubmed/15264254>
- [12] Daina A, Michelin O and Zoete V 2017 SwissADME: a free web tool to evaluate pharmacokinetics, drug-likeness and medicinal chemistry friendliness of small molecules *Sci. Rep.* **7**(42717) 1–13  
<https://www.ncbi.nlm.nih.gov/pubmed/28256516>
- [13] Drwal MN, Banerjee P, Dunkel M, Wettig M R and Preissner R 2014 ProTox: A webserver for the in silico prediction of rodent oral toxicity *Nucleic Acid Res.* **42** W53–W58  
<https://www.ncbi.nlm.nih.gov/pubmed/24838562>
- [14] Suhud F, Effendi C and Siswandono 2017 In silico QSAR of 1-benzyl-3-benzoylurea lead and its analogue compounds as anticancer *17th Asian Conference in Clinical Pharmacy (27–30 July 2017, Yogyakarta, Indonesia)*  
<http://repository.ubaya.ac.id/31818/>
- [15] McTigue M, Murray B W, Chen J H, Deng Y L, Solowiej J and Kania R S 2012 Molecular conformations, interactions, and properties associated with drug efficiency and clinical performance among VEGFR TK inhibitors *Proc. Natl. Acad. Sci.* **109**(45) 18281–89  
<https://www.ncbi.nlm.nih.gov/pubmed/22988103>
- [16] Jain A N and Nicholls A 2008 Recommendations for evaluation of computational methods *J. Comput. Aided. Mol. Des.* **22**(3–4) 133–39  
<https://www.ncbi.nlm.nih.gov/pubmed/18338228>
- [17] Thomsen R and Christensen M H 2006 MolDock: A new technique for high-accuracy molecular docking *J. Med. Chem.* **49**(11) 3315–21  
<https://www.ncbi.nlm.nih.gov/pubmed/16722650>
- [18] Liu Y and Gray N S 2006 Rational design of inhibitors that bind to inactive kinase conformations *Nat. Chem. Biol.* **2**(7) 358–64  
<https://www.ncbi.nlm.nih.gov/pubmed/16783341>
- [19] Waring M J 2010 Lipophilicity in drug discovery *Expert Opin. Drug Discov.* **5**(3) 235–48  
<https://www.ncbi.nlm.nih.gov/pubmed/22823020>
- [20] Delaney J S 2004 ESOL: Estimating aqueous solubility directly from molecular structure *J. Chem. Inf. Comput. Sci.* **44**(3) 1000–05  
<https://pubs.acs.org/doi/abs/10.1021/ci034243x>
- [21] Ali J, Camilleri P, Brown M B, Hutt A J and Kirton S B 2012 In silico prediction of aqueous solubility using simple QSPR models: The importance of phenol and phenol-like moieties *J. Chem. Inf. Model.* **52**(11) 2950–57  
<https://www.ncbi.nlm.nih.gov/pubmed/23121381>
- [22] Daina A and Zoete V 2016 A BOILED-egg to predict gastrointestinal absorption and brain penetration of small molecules *ChemMedChem* **11**(11) 1117–21  
<https://www.ncbi.nlm.nih.gov/pubmed/27218427>
- [23] Thomas H and Coley H M 2003 Overcoming multidrug resistance in cancer: An update on the clinical strategy of inhibiting p-glycoprotein *Cancer Control* **10**(2) 159–65  
<https://www.ncbi.nlm.nih.gov/pubmed/12712010>
- [24] Bickerton G R, Paolini G V, Besnard J, Muresan S and Hopkins A L 2012 Quantifying the chemical beauty of drugs *Nat. Chem.* **4** 90–8

- <https://www.nature.com/articles/nchem.1243>
- [25] Martin Y C 2005 A bioavailability score *J. Med.Chem.* **48**(9) 3164–70  
<https://www.ncbi.nlm.nih.gov/pubmed/15857122>
- [26] Lipinski C A, Lombardo F, Dominy B W and Feeney P J 2012 Experimental and computational approaches to estimate solubility and permeability in drug discovery and development settings *Adv. Drug. Deliv. Rev.* **64** 4–17  
<https://www.sciencedirect.com/science/article/pii/S0169409X12002797>
- [27] Brenk R, Schipani A, James D, Krasowski A, Gilbert I H, Frearson J and Wyatt P G 2008 Lessons learnt from assembling screening libraries for drug discovery for neglected diseases *ChemMedChem* **3**(3) 435–44  
<https://www.ncbi.nlm.nih.gov/pubmed/18064617>
- [28] Kalgutkar A S et al. 2005, A Comprehensive listing of bioactivation pathways of organic functional groups *Curr. Drug Metab.* **6** 161–225  
<https://www.ncbi.nlm.nih.gov/pubmed/15975040>
- [29] Park B K et al. 2011 Managing the challenge of chemically reactive metabolites in drug development *Nat. Rev. Drug Discov.* **10**(4) 292–306  
<https://www.ncbi.nlm.nih.gov/pubmed/21455238>



**275**

1. **Introduction**  
 2. **Methodology**  
 3. **Results and Discussion**  
 4. **Conclusion**

**275**

1. **Introduction**  
 2. **Methodology**  
 3. **Results and Discussion**  
 4. **Conclusion**



also developed by scimago:



SCIMAGO INSTITUTIONS RANKINGS



Scimago Journal & Country Rank

Enter Journal Title, ISSN or Publisher Name



- Home
- Journal Rankings
- Country Rankings
- Viz Tools
- Help
- About Us



Ads by Google

Stop seeing this ad

Why this ad? ⓘ

# IOP Conference Series: Earth and Environmental Science

## COUNTRY

[United Kingdom](#)



Universities and research institutions in United Kingdom

## SUBJECT AREA AND CATEGORY

[Earth and Planetary Sciences](#)

[Earth and Planetary Sciences \(miscellaneous\)](#)

[Environmental Science](#)

[Environmental Science \(miscellaneous\)](#)

## PUBLISHER

[IOP Publishing Ltd.](#)

## H-INDEX

# 26

## PUBLICATION TYPE

Conferences and Proceedings

## ISSN

17551307, 17551315

## COVERAGE

2010-2020

## INFORMATION

[Homepage](#)

[How to publish in this journal](#)

[ees@ioppublishing.org](mailto:ees@ioppublishing.org)




Ads by Google


Stop seeing this ad

Why this ad?

**SCOPE**

The open access IOP Conference Series: Earth and Environmental Science (EES) provides a fast, versatile and cost-effective proceedings publication service.

 Join the conversation about this journal











### Get more leads and sales

**Boost visitors from news sites**

Guest post on news sites with high domain authority and millions of monthly visitors.

brandpush.co

OPEN



**IOP Conference Series: Earth and Environmental...**

Not yet assigned quartile

SJR 2020  
**0.18**

powered by scimagojr.com

← Show this widget in your own website

Just copy the code below and paste within your html code:

```
<a href="https://www.scimagojr.com" style="border: 1px solid #ccc; padding: 2px 5px; display: inline-block;">https://www.scimagojr.com
```

**SCImago Graphica**

Explore, visually communicate and make sense of data with our **new free tool**.

Get it





## GrainCom Event 21

October 18-20, 2021

The Game Changer Event for the Grain Industry

graincomevents.com

OPEN

Metrics based on Scopus® data as of April 2021

A **Alharia Dinata** 5 months ago

IOP Conference Series: Earth and Environmental Science - Volume 708 is not available in Scopus.

reply



**Melanie Ortiz** 5 months ago

SCImago Team

Dear Alharia,  
thank you very much for your comment, unfortunately we cannot help you with your request. We suggest you contact Scopus support: [https://service.elsevier.com/app/answers/detail/a\\_id/14883/kw/scimago/supporthub/scopus/](https://service.elsevier.com/app/answers/detail/a_id/14883/kw/scimago/supporthub/scopus/)  
Best Regards, SCImago Team

V **Vani** 5 months ago

good evening, whether this journal is Q4 or Q2 ?

reply



**Melanie Ortiz** 5 months ago

SCImago Team

Dear Vani,  
Thank you for contacting us.  
As said below, we calculate the SJR data for all the publication's types, but the Quartile's data are only calculated for Journals and Book Series.  
Best regards, SCImago Team

F **FEROSKHAN M** 10 months ago

IOP Conference Series: Earth and Environmental Science - Volume 573 is not available in Scopus. But later volumes are available. May I know when will they publish in Scopus?

reply



**Melanie Ortiz** 10 months ago

SCImago Team

Dear Sir/Madam,

thank you very much for your comment, unfortunately we cannot help you with your request. We suggest you contact Scopus support: [https://service.elsevier.com/app/answers/detail/a\\_id/14883/kw/scimago/supporthub/scopus/](https://service.elsevier.com/app/answers/detail/a_id/14883/kw/scimago/supporthub/scopus/)  
Best Regards, SCImago Team

**N** **Natt** 1 year ago

I would like to know the quartile of this journal. Why isn't it showing on the website?

reply



**Melanie Ortiz** 1 year ago

SCImago Team

Dear Natt,  
Thank you for contacting us. We calculate the SJR data for all the publication's types, but the Quartile's data are only calculated for Journals and Book Series.  
Best regards, SCImago Team

**N** **Nurgustaana** 1 year ago

Dear SCImago Team!  
I want to know previous quartiles of journal (for 2018 and 2019 years). I have tried find information about a quartile, but discovered just SJR for 2018. Could you please provide information about it?  
Yours sincerely, Nurgustaana

reply



**Melanie Ortiz** 1 year ago

SCImago Team

Dear Nurgustaana,  
Thank you for contacting us. We calculate the SJR data for all the publication types, but the Quartile data are only calculated for Journal type's publications. Best regards,  
SCImago Team

**M** **Mora** 1 year ago

hello, how to search one of journal who publised by IOP, because when i find it by the title, they are not able in scimagojr but the publisher is available in here, thank you for the respond it means a lot

reply



**Melanie Ortiz** 1 year ago

SCImago Team

Dear Mora,  
  
thank you for contacting us. Could you provide us the Title of the journal? We remember that SCImago Journal & Country Rank shows all the information have been provided by Scopus. If you didn't localize the journal in the search engine, it means that Scopus / Elsevier has not provided us the corresponding data.

Best Regards, SCImago Team

D **Dr. Yousif** 2 years ago

Dear Sir,

I have published a paper in Earth and Environmental Science Journal (only myself, single author) I am trying to withdraw it after 28 days of publishing online, is it possible? Could you please tell me the procedure of withdrawing a paper?

Thank you,

reply



**Melanie Ortiz** 2 years ago

SCImago Team

Dear Yousif,

thank you for contacting us.

We are sorry to tell you that SCImago Journal & Country Rank is not a journal. SJR is a portal with scientometric indicators of journals indexed in Elsevier/Scopus.

Unfortunately, we cannot help you with your request, we suggest you to contact the journal's editorial staff , so they could inform you more deeply. Best Regards, SCImago Team

A **Agustinus Kastanya** 2 years ago

need information about renking of the Journal on Scopus

reply



**Melanie Ortiz** 2 years ago

SCImago Team

Dear Agustinus, thank you very much for your comment, unfortunately we cannot help you with your request. We suggest you to consult the Scopus database directly. Remember that the SJR is a static image of a database (Scopus) which is changing every day. Best regards, SCImago Team

M **Mahipal** 2 years ago

Dear Admin,

How could our journal include in your IOP?

reply



**Melanie Ortiz** 2 years ago

SCImago Team

Dear Mahipal,

thank you for contacting us.

We suggest you to contact the IOP's editorial staff , so they could inform you more deeply.

If you would like to make an application to Scopus, please contact them to help you with this issue here: <https://www.elsevier.com/solutions/scopus/content/content-policy-and-selection>

<http://suggestor.step.scopus.com/suggestTitle/step1.cfm>

Best Regards, SCImago Team

M **Mursalin** 2 years ago

Dear SCImago Team

My name is mursalin from Jambi City, Indonesia. I have published my article titled The Effect of Temperature on MDAG Purification Using Creaming Demulsification Technique at the IOP Conference Series: Earth and Environmental Science, Volume 309, conference 1 and could be accessed at: <https://iopscience.iop.org/article/10.1088/1755-1315/309/1/012068>. But why until now it does not appear into Google Scholar and my account.

Please help me to resolve the issue. Thank you for your kindness. I am waiting for good news from you.

Sincerelly,  
Mursalin

reply



**Melanie Ortiz** 2 years ago

SCImago Team

Dear Mursalin,

Thank you for contacting us. Unfortunately, we can not help you with your request. Maybe other users can help you. Best Regards, SCImago Team

N **Танзиля Созаева** 2 years ago

Доброго времени суток! Интересует информация по квартилю журнала (издания)

reply



**Melanie Ortiz** 2 years ago

SCImago Team

Dear user, thank you very much for your request. You can consult that information in SJR website. Best Regards, SCImago Team

S **syaiful** 3 years ago

I am very interested to send my paper to this conference

best regards

syaiful

reply



**Elena Corera** 3 years ago

SCImago Team

Dear user, in the link below you will find the information corresponding to the author's instructions of this journal. Best regards, SCImago Team  
<https://publishingsupport.iopscience.iop.org/author-guidelines-for-conference-proceedings/>

**Leave a comment**

Name

Email

(will not be published)

 I'm not a robot reCAPTCHA  
Privacy - Terms**Submit**

The users of Scimago Journal & Country Rank have the possibility to dialogue through comments linked to a specific journal. The purpose is to have a forum in which general doubts about the processes of publication in the journal, experiences and other issues derived from the publication of papers are resolved. For topics on particular articles, maintain the dialogue through the usual channels with your editor.

Developed by:



Powered by:



Follow us on @ScimagoJR

Scimago Lab, Copyright 2007-2020. Data Source: Scopus®

**EST MODUS IN REBUS**  
Horatio (Saturni 1, 1, 106)



# Source details

## IOP Conference Series: Earth and Environmental Science

Scopus coverage years: from 2010 to Present

ISSN: 1755-1307 E-ISSN: 1755-1315

- Subject area:
- Environmental Science: General Environmental Science
  - Earth and Planetary Sciences: General Earth and Planetary Sciences
  - Physics and Astronomy: General Physics and Astronomy

Source type: Conference Proceeding

CiteScore 2020 **0.5**

SJR 2020 **0.179**

SNIP 2020 **0.436**

[View all documents >](#)

[Set document alert](#)

[Save to source list](#) [Source Homepage](#)

[CiteScore](#) [CiteScore rank & trend](#) [Scopus content coverage](#)

### i Improved CiteScore methodology

CiteScore 2020 counts the citations received in 2017-2020 to articles, reviews, conference papers, book chapters and data papers published in 2017-2020, and divides this by the number of publications published in 2017-2020. [Learn more >](#)

CiteScore 2020

$$0.5 = \frac{25,463 \text{ Citations } 2017 - 2020}{49,883 \text{ Documents } 2017 - 2020}$$

Calculated on 05 May, 2021

CiteScoreTracker 2021

$$0.5 = \frac{32,770 \text{ Citations to date}}{65,646 \text{ Documents to date}}$$

Last updated on 04 September, 2021 • Updated monthly

### CiteScore rank 2020

Category	Rank	Percentile
Environmental Science		
General Environmental Science	#183/220	17th
Earth and Planetary Sciences		
General Earth and Planetary Sciences	#157/186	15th

[View CiteScore methodology >](#) [CiteScore FAQ >](#) [Add CiteScore to your site](#)



## About Scopus

[What is Scopus](#)  
[Content coverage](#)  
[Scopus blog](#)  
[Scopus API](#)  
[Privacy matters](#)

## Language

[日本語に切り替える](#)  
[切换到简体中文](#)  
[切换到繁體中文](#)  
[Русский язык](#)

## Customer Service

[Help](#)  
[Contact us](#)

**ELSEVIER**

[Terms and conditions](#) ↗ [Privacy policy](#) ↗

Copyright © Elsevier B.V. ↗. All rights reserved. Scopus® is a registered trademark of Elsevier B.V.

We use cookies to help provide and enhance our service and tailor content. By continuing, you agree to the use of cookies.

 RELX

## Table of contents

### Volume 293

2019

[◀ Previous issue](#)   [Next issue ▶](#)

The 2nd International Conference on Natural Resources and Life Sciences (NRLS-2018) 23–24 August 2018, Ibis Styles Hotel, Surabaya, Indonesia

Accepted papers received: 17 May 2019

Published online: 01 July 2019

[Open all abstracts](#)

### Preface

**OPEN ACCESS** 011001

[The 2nd International Conference on Natural Resources and Life Sciences \(NRLS-2018\)](#)

[+ Open abstract](#)   [View article](#)   [PDF](#)

**OPEN ACCESS** 011002

[Conference Photos](#)

[+ Open abstract](#)   [View article](#)   [PDF](#)

**OPEN ACCESS** 011003

[Conference Sponsors](#)

[+ Open abstract](#)   [View article](#)   [PDF](#)

**OPEN ACCESS** 011004

[Scientific & Editorial Boards the 2<sup>nd</sup> NRLS-2018](#)

[+ Open abstract](#)   [View article](#)   [PDF](#)

**OPEN ACCESS** 011005

[Peer review statement](#)

[+ Open abstract](#)   [View article](#)   [PDF](#)

### Papers

**OPEN ACCESS** 012001

[Android based rice pest detection system using learning vector quantization method](#)

A Budiman, P Utomo and S Rahayu

[+ Open abstract](#)   [View article](#)   [PDF](#)

**OPEN ACCESS** 012002

[Community behavior and single-use plastic bottle consumption](#)

A Khoironi, S Anggoro and S Sudarmo

[+ Open abstract](#)   [View article](#)   [PDF](#)

- 
- OPEN ACCESS** 012003  
Genetic variability of soybean (*Glycine max* L. Merrill) genotypes for pod shattering resistance  
A Krisnawati and M M Adie  
[+ Open abstract](#) [View article](#) [PDF](#)
- 
- OPEN ACCESS** 012004  
Review on biogas from palm oil mill effluent (POME): Challenges and opportunities in Indonesia  
A Rajani, Kusnadi, A Santosa, A Saepudin, S Gobikrishnan and D Andriani  
[+ Open abstract](#) [View article](#) [PDF](#)
- 
- OPEN ACCESS** 012005  
Early detection of somaclonal variation in oil palm callus culture through cytological and SDS-PAGE protein analysis  
A Sahara, Reflini, C Utomo and T Liwang  
[+ Open abstract](#) [View article](#) [PDF](#)
- 
- OPEN ACCESS** 012006  
Genetic diversity of Indonesian soybean (*Glycine max* L. Merrill) germplasm based on morphological and microsatellite markers  
A Sulisty, F C Indriani, M J Mejaya, A N Sugiharto and J Agranoff  
[+ Open abstract](#) [View article](#) [PDF](#)
- 
- OPEN ACCESS** 012007  
Fermentation quality of *Pennisetum purpureum* cv. Mott ensiled with *Lactobacillus plantarum* and sugarcane molasses in tropic  
A Wahyudi, L Hendraningsih, Sutawi, R H Setyobudi and M Mel  
[+ Open abstract](#) [View article](#) [PDF](#)
- 
- OPEN ACCESS** 012008  
The evaluation of estimated breeding value and the most probable producing ability for the basis selection of Ettawa crossbred goat (*Capra hircus* sp.) at Malang, East Java, Indonesia  
A Winaya, Suyatno, P Coy and N Fauzi  
[+ Open abstract](#) [View article](#) [PDF](#)
- 
- OPEN ACCESS** 012009  
Menthol from the stem and leaf *in-vitro* *Mentha piperita* Linn.  
B K Wijaya, P H Hardjo and S Emantoko  
[+ Open abstract](#) [View article](#) [PDF](#)
- 
- OPEN ACCESS** 012010  
A review: Biomass-based fuel pellet usage in biomass gasifier-an optimization on non-wood material  
D Andriani, T D Atmaja, M Arifin, A Rajani and Kusnadi  
[+ Open abstract](#) [View article](#) [PDF](#)
- 
- OPEN ACCESS** 012011  
Effectiveness of the use of organic waste as fertilizer and physical scarification of seeds on growth of seeds nila plants (*Indigofera* sp.)  
D Roeswitawati, M Huda, D Indratmi and M Mel  
[+ Open abstract](#) [View article](#) [PDF](#)

- 
- OPEN ACCESS** 012012  
Physiology and genotyping of adaptive and sensitive oil palm progenies under unwatered stress condition  
D Yono, E Purwanti, A Sahara, Y A Nugroho, Z A Tanjung, R Aditama, C Dewi, A E Sihotang, C Utomo and T Liwang  
[+ Open abstract](#) [View article](#) [PDF](#)
- 
- OPEN ACCESS** 012013  
Effect of phosphoric acid pretreatment on characterization of gelatin from broiler chicken (*Gallus gallus domesticus* L.) bones  
D Yuliani, A Maunatin, A Jannah and H H Fauziyyah  
[+ Open abstract](#) [View article](#) [PDF](#)
- 
- OPEN ACCESS** 012014  
Selection for drought-resistant rice (*Oryza sativa* L.) using polyethylene glycol  
E D Purbajanti, F Kusmiyati, E Fuskhah, R Rosyida, P G Adinurani and Z Vincēviča-Gaile  
[+ Open abstract](#) [View article](#) [PDF](#)
- 
- OPEN ACCESS** 012015  
Effectiveness the source of nitrogen from  $\text{NO}_3$  and  $\text{NH}_4$  for *Panicum maximum* Jacq. growth in saline soil  
E D Purbajanti, P G Adinurani, T Turkadze, Z Vincēviča-Gaile and R H Setyobudi  
[+ Open abstract](#) [View article](#) [PDF](#)
- 
- OPEN ACCESS** 012016  
Anticancer potential from ethanol extract of *Zanthoxylum acanthopodium* DC. seed to against MCF-7 cell line  
E V Arsita, D E Saragih and K Aldrin  
[+ Open abstract](#) [View article](#) [PDF](#)
- 
- OPEN ACCESS** 012017  
Long-term Proton Pump Inhibitors induces recurrent Urinary Tract Infections: A case study  
F Cokro and S T Arrang  
[+ Open abstract](#) [View article](#) [PDF](#)
- 
- OPEN ACCESS** 012018  
Molecular docking, drug-likeness, and ADMET study of 1-benzyl-3-benzoylurea and its analogs against VEGFR-2  
F Suhud, D H Tjahjono, T A Yuniarta, G S Putra and J Setiawan  
[+ Open abstract](#) [View article](#) [PDF](#)
- 
- OPEN ACCESS** 012019  
Microcontroller application for pH and temperature control system in liquid sugar liquefaction process made from cassava (*Manihot esculenta* Crantz.)  
H Santosa, Yuliaty, I J Mulyana, D S Sirait and S D Novitasari  
[+ Open abstract](#) [View article](#) [PDF](#)
- 
- OPEN ACCESS** 012020  
The effect of fermentation process on physical and chemical characteristics of pitaya (*Hylocereus polyrhizus* [F.A.C. Weber] Britton & Rose) stem flour  
H Soedjatmiko, R Chrisnasari and P H Hardjo  
[+ Open abstract](#) [View article](#) [PDF](#)