

# Thin Layer Chromatography Fingerprinting and Clustering of *Orthosiphon stamineus* Benth. from Different Origins

Kartini Kartini<sup>1,\*</sup>, Ervina Rustiana Dewi<sup>1</sup>, Fandi Achmad<sup>1</sup>, Nikmatul Ikhrom Eka Jayani<sup>1</sup>, Mochammad Arbi Hadiyat<sup>2</sup>, Christina Avanti<sup>3</sup>

Kartini Kartini<sup>1,\*</sup>, Ervina Rustiana Dewi<sup>1</sup>, Fandi Achmad<sup>1</sup>, Nikmatul Ikhrom Eka Jayani<sup>1</sup>, Mochammad Arbi Hadiyat<sup>2</sup>, Christina Avanti<sup>3</sup>

<sup>1</sup>Department of Pharmaceutical Biology, Faculty of Pharmacy, University of Surabaya, Raya Kalirungkut Road, Surabaya 60293, INDONESIA.

<sup>2</sup>Department of Industrial Engineering, Faculty of Engineering, University of Surabaya, Raya Kalirungkut Road, Surabaya 60293, INDONESIA.

<sup>3</sup>Department of Pharmaceutic, Faculty of Pharmacy, University of Surabaya, Raya Kalirungkut Road, Surabaya 60293, INDONESIA.

## Correspondence

Kartini Kartini

Department of Pharmaceutical Biology, Faculty of Pharmacy, University of Surabaya, Raya Kalirungkut Road, Surabaya 60293, INDONESIA.

E-mail: kartini@staff.ubaya.ac.id

## History

- Submission Date: 14-11-2019;
- Review completed: 29-11-2019;
- Accepted Date: 09-12-2019.

DOI : 10.5530/pj.2020.12.257

## Article Available online

<http://www.phcogj.com/v12/i1>

## Copyright

© 2019 Phcogj.Com. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International license.

## ABSTRACT

**Introduction:** *Orthosiphon stamineus* has been widely used across Asian countries for the treatment of various diseases. The quality of herbal medicine determine its safety as well as efficacy; and geographical origin is important factor contributing on the quality of herb and its products. **Materials and Methods:** Thin Layer Chromatography (TLC) method combined with chemometric, Principal Component Analysis (PCA), has been employed to evaluate the quality of *Orthosiphon stamineus* leaves collected from eleven origins in Indonesia. **Results:** The results showed that mobile phase suitable for *Orthosiphon stamineus* was chloroform, dichloromethane, ethyl acetate (7:4:1). The method used has met the requirements of TLC system stability and precision. TLC-fingerprints analyzed with chemometrics showed an ability to discriminate *Orthosiphon stamineus* from various origins. PCA score plot of the first two principal components (PC) clearly distinguished 3 clusters of samples, whereas the loading plot of the first two PC showed that compounds with the R<sub>f</sub> values of 0.0-0.1, 0.1-0.2, 0.2-0.3, and 0.9-1.0 are the most important compounds for clustering of samples. **Conclusions:** TLC-fingerprint combined with the PCA was able to discriminate among the leaves of *Orthosiphon stamineus* originated from various locations. TLC-fingerprints analyzed with chemometrics can be used as an alternative of marker-oriented method to evaluate the quality of *Orthosiphon stamineus*.

**Key words:** Geographical origin; Herbal medicine; Marker; Principal component analysis; Quality; TLC.

## INTRODUCTION

*Orthosiphon stamineus* (Indonesia: *Kumis kucing*) is a member of the Lamiaceae family. This plant has been empirically used in several Southeast Asian countries, such as Indonesia, Malaysia, Thailand, Vietnam, and Myanmar. *Orthosiphon stamineus* (OS) in Indonesia is traditionally used for diuretic, rheumatic, diabetes, hypertension, tonsillitis, epilepsy, menstrual disorders, gonorrhea, syphilis, kidney stones, nephritis, gout arthritis, and antipyretics.<sup>1</sup> Various studies on OS and its compounds indicated that this plant has antioxidant, cytotoxic, diuretic, nephroprotective, antidiabetic, antihypertensive, anti-inflammatory, antimicrobial, anti-obesity, and hepatoprotective activities, as well as activity on the cardiovascular system.<sup>1-5</sup> Bioactivities of OS has been attributed by its various chemical constituents classified as monoterpenes, diterpenes, triterpenes, essential oil, sterols, saponins, flavonoids, and organic acids.<sup>1,6</sup>

*Orthosiphon stamineus* is an herbaceous plant and it has two types of flower, white and purple colour. This variety may cause the differences in chemical contents as well as biological activities.<sup>7</sup> The purple variety has more bioactive compounds than the white one. However, most scientific investigations have used the white variety.<sup>8</sup> Moreover, solvent, duration, temperature, and method of extraction as well as geographical location may also influence the

chemical compounds of OS.<sup>6,9-13</sup> Geographical origin of plants is important to be recognized since it affects the levels of bioactive compounds, a general criterion in the selection of raw materials. Therefore, a simple and accurate analytical method is needed to identify the geographical origin of OS.

Quality evaluation of OS crude drugs and its extract can be carried out using two approaches, the concentration of marker compound(s) and the profile of fingerprint. A groups of marker, rosmarinic acid (RA), 3'-hydroxy-5,6,7,4'-tetramethoxyflavone (TMF), sinensetin (SIN), and eupatorin (EUP) have been determined using HPLC<sup>11</sup> and HPTLC<sup>13,14</sup> methods to evaluate OS extract obtained from various origins, solvent, and method of extraction. Gas chromatography (GC) and FT-IR have been applied to develop the fingerprint of OS collected from various origins and prepared with different methods of extraction.<sup>7,10-12</sup>

Application of fingerprint reveals several superiorities compared to those of marker compounds. Fingerprint is a distinctive profile or pattern of sample which chemically reflects its composition in which as much information as possible is presented.<sup>15,16</sup> Therefore, almost all of compounds in a plant are considered in the analysis. This allows quality evaluation of a plant to be more objective than using only certain compound(s). TLC-fingerprint has been widely used in herbal medicines assessment. It is simpler, faster, and more cost effective compared to GC,

**Cite this article:** Kartini K, Dewi ER, Achmad F, Jayani NIE, Hadiyat MA, Avanti C. Thin Layer Chromatography Fingerprinting and Clustering of *Orthosiphon stamineus* Benth. from Different Origins. Pharmacogn J. 2020;12(1):1683-91.

HPLC, and FT-IR. TLC is able to analyze many samples in one running time. It is also possible to visually analyze TLC-chromatograms; however, this technique is subjective and not quantitative. Moreover, fingerprint chromatograms are complex multivariate data sets which cause difficulty in evaluation of very similar chromatograms. Thus, chemometrics should be taken into consideration. This approach, although more difficult, is based on objective mathematical methods and treats the chromatogram as a unique signal, without a need to identify and interpret the peaks. Therefore, it provides a good possibility for mining more useful chemical information from original-rich data.<sup>16-18</sup> In this research we developed TLC-fingerprint combined with chemometrics to differentiate OS collected from various origins. Although several works on fingerprinting of OS have been reported before, none of them involved comparison between OS from different origins using TLC method.

## MATERIALS AND METHODS

### Chemicals

The following chemicals were procured from Merck (Darmstadt, Germany): TLC plate pre-coated with silica gel 60 F<sub>254</sub>, n-hexane, diethyl ether, tetrahydrofuran, acetic acid, n-propanol, dichloromethane (DCM), ethyl acetate (EA), dioxane, toluene, chloroform (CHCl<sub>3</sub>), methanol, ethanol, anisaldehyde, and sulfuric acid.

### Plant materials

*Orthosiphon stamineus* were collected from eleven cultivation area in East and Central of Java, Indonesia (Table 1, Figure 1). Plant parts used were 5 leaves from the shoots that have not flowered yet. Leaves were harvested and cleaned with tap water. All samples were authenticated

by Center for Traditional Medicine Information and Development, Faculty of Pharmacy, University of Surabaya, Indonesia (certificate number: 1400/D.T/V/2019). Leaves were then dried and ground into mesh 60.

### Preparation of extracts

One gram of powdered *Orthosiphon stamineus* leaves was extracted with 10 ml of methanol, using Ultrasound-Assisted Extraction (UAE) method for 15 minutes. Extracts were subsequently filtered and kept in a tightly closed bottle.

### TLC-fingerprint analysis


#### TLC condition

A Camag TLC system comprising of Linomat 5 sample applicator, twin-through chamber, and TLC-Visualizer with 12 bit CCD camera and Camag VideoScan 1.02 software serial number 2503D001 (Camag, Muttenz, Switzerland) were used. Chromatography was performed on Merck TLC plates (Art. No.: 1.05554.0001, silica gel 60 F<sub>254</sub> pre-coated, 20 × 20 cm, 175-225 µm layer thickness, aluminium-backed, particle size distribution: 9.5-11.5 µm) with a 100-µl Camag syringe. Samples were spotted under a flow of nitrogen as 8 mm bands, 10 mm from the left edge, 15 mm from the bottom edge and 20 mm of track distance. Development was carried out in a chamber previously equilibrated (for 30 min at room temperature) with mobile phase (see: *selection of mobile phase*) and migration distance was 80 mm. The plates were dried under room temperature and then dipped in anisaldehyde-sulfuric acid reagent. TLC plates were then dried in fume hood and heated for 10 min at 100 °C. TLC plate were subsequently illuminated under short-wave UV (254 nm), long-wave UV (366 nm), and white light by using

**Table 1: Geographical origins of *Orthosiphon stamineus* from Indonesia.**

Code	District (City)	Height (mamsl)*	Latitude, Longitude	Time of collection
A	Lamongan	6	7°11' S; 112°42' E	May 2019
B	Pasuruan	800	7°41' S; 112°39' E	May 2019
C	Malang	445	7°43' S; 112°40' E	May 2019
D	Gresik	3	7°10' S; 112°40' E	May 2019
E	Surabaya	2	7°15' S; 112°42' E	May 2019
F	Sidoarjo	3	7°29' S; 112°43' E	May 2019
G	Jombang	44	7°20' S; 112°45' E	May 2019
H	Mojokerto 1	700	7°40' S; 112°36' E	May 2019
I	Karanganyar	1200	7°42' S; 111°80' E	February 2018
J	Batu	875	7°42' S; 112°42' E	February 2018
K	Mojokerto 2	700	7°40' S; 112°36' E	May 2019

\*Metres above mean sea level



**Figure 1:** *Orthosiphon stamineus* from Lamongan (a), Pasuruan (b), Malang (c), Gresik (d), Surabaya (e), Sidoarjo (f), Jombang (g), Mojokerto 1 (h), Karanganyar (i), Batu (j), and Mojokerto 2 (k).

TLC-Visualizer. Quantitative evaluation of digitized images was carried out with Camag VideoScan 1.02 software.

### Selection of the mobile phase

Ten ml of each solvent i.e. diethyl ether, 2-propanol, ethanol, tetrahydrofuran, acetic acid, dichloromethane, ethyl acetate, dioxane, toluene, and chloroform were put into a twin-through chamber and saturated for 30 minutes. Five  $\mu$ l of OS leaves extract was applied on TLC plate. The plate was then developed with the migration distance of 80 mm. After development, the plates were removed and dried at room temperature. In addition to the single mobile phase, the selection of mobile phase was also carried out using mixture of two or three solvents with various comparisons to obtain a chromatogram with the highest number and the best separations of zones.

### Stability of the chromatograms

The stability of analytes in solution and on the plate was assessed by leaving the extract at room temperature and on the plate for 3 hours. As a standard, the new extract was prepared and applied just before the elution is started. The analytes were considered to be stable in the solution and on the plate before the chromatographic process if the difference of their Rf value is not more than 0.05. Stability of the analytes during chromatography was evaluated by two-dimensional (2D) elution. The analytes were concluded to be stable during the chromatographic process if all zones located on a diagonal connecting the initial position of the application with the intersection of the two mobile phase fronts. Moreover, to investigate the stability of the chromatographic result, the sample was chromatographed according to the method and derivatized with anisaldehyde-sulfuric acid reagent. The chromatogram was evaluated using TLC-visualizer repeatedly after 5, 10, 30, and 60 minutes. The chromatogram was concluded to be stable if there are no significant changes in Rf value.<sup>19</sup>

### Precision of the chromatograms

*Orthosiphon stamineus* leaves were extracted in triplicate. Each extract was chromatographed three times on 3 different plates. Intraday precision was analyzed by calculating the Rf values of several prominent zones then calculating its mean and relative standard deviation. A reasonable acceptance criterion would be that Rf values of the same compounds do not vary more than 0.02 from plate to plate. To determine the interday precision, the extract was applied three times on 3 different plates and on 3 different days. Interday precision was acceptable if Rf values of the same substances vary not more than 0.05 between plates from different days.<sup>19</sup>

### TLC of samples from different origins

OS leaves extracts from 11 different locations were chromatographed on a TLC plate. The TLC plate was then eluted as described before. TLC plate was subsequently illuminated under short-wave UV (254 nm), long-wave UV (366 nm), and white light by using TLC-Visualizer. Videodensitogram of samples were then analyzed using chemometric method.

### Chemometric analysis

Rf value, height, and area of each peak (compound) obtained from videodensitogram was tabulated and then analyzed using Principal Component Analysis (PCA). PCA is an exploratory data analysis. This method is based on the information available in the fingerprints only. PCA reduces the complexity of the multivariate data set by explaining the correlation amongst a large number of variables in terms of a smaller number of underlying factors (principal components or PCs) without losing much information. The projections of the n objects from the original data on PCs are called the scores plots, whereas the contribution of each original variable to the score is presented by its

loading, which detects the variables responsible for the clustering<sup>15</sup>. PCA was carried out using Minitab v.16 (Minitab Inc., USA).


## RESULTS AND DISCUSSION

### Selection of the mobile phase


Methanol extracts of OS leaves collected from 11 origins are presented at Figure 2. All of extracts show dark green colour, except OS extract from Batu district (sample j). This is an initial indicator that sample from Batu may differ from the others. One of these extracts was then chosen for the mobile phase selection. At the first step, 10 single mobile phases were used and chloroform, dichloromethane, as well as ethyl acetate were the best mobile phase. These single mobile phases exhibited the most number and the best separation of zones on the TLC plate. These solvents were then mixed using simplex centroid with axial design.<sup>20</sup> Ten types of combination in different ratio were deduced and CHCl<sub>3</sub>, DCM, EA (4:1:1) showed the optimum solvent system (Figure 3). However, the separation of zones with this solvent ratio was not so good, therefore modified ratios were applied i.e. 5:2:1 and 7:4:1. Finally, CHCl<sub>3</sub>, DCM, EA in the ratio of 7:4:1 was then selected as a solvent system for the development of OS TLC fingerprinting. It exhibited 11 zones after derivatized with anisaldehyde-sulfuric acid reagent under white light.

### Stability test

The stability of the analytes before and during the chromatography process is important to be established due to off-line nature of TLC system. The stability of the analyte before chromatography was determined by developing two extracts prepared at different times. The Rf values of the prominent zone or marker (\*, yellowish-blue) at tracks I and II were compared to those at tracks III and IV to evaluate the stability of the analytes on the plate, while the zones at tracks V and VI were compared to those at tracks III and IV to assess the stability of the analytes in the solution.<sup>19</sup> The results showed that all tracks afforded the same pattern (Figure 4) and Rf value, with the  $\Delta R_f < 0.05$  (Table 2). These indicated that the analytes remained stable for three hours before chromatography began, both on the plates and in the extract solution.




**Figure 2:** *Orthosiphon stamineus* leaves extracts from Lamongan (a), Pasuruan (b), Malang (c), Gresik (d), Surabaya (e), Sidoarjo (f), Jombang (g), Mojokerto 1 (h), Karanganyar (i), Batu (j), and Mojokerto 2 (k).



**Figure 3:** Total zones of OS leaves extract. Mobile phase (MP): CHCl<sub>3</sub>, DCM, EA in different ratios. Detection (D): anisaldehyde-sulfuric acid reagent, white light.





**Figure 4:** Stability of the analytes on plate and in solution. I, II: extracts on the plate for 3 hours; III, IV: extracts on the plate for less than 5 minutes; V, VI: extracts in solution for 3 hours. MP:  $\text{CHCl}_3$ , DCM, EA (7:4:1). D: anisaldehyde-sulfuric acid reagent, UV 366 nm.

**Table 2: Stability of analyte on the plate and in solution.**

Marker (*)	Track I	Track II	Track III	Track IV	Track V	Track VI
Colour	Yellowish-blue	Yellowish-blue	Yellowish-blue	Yellowish-blue	Yellowish-blue	Yellowish-blue
R <sub>f</sub>	0.418	0.394	0.388	0.384	0.391	0.403
$\Delta R_f^1$	0.032	0.008	0	0	0.005	0.017

<sup>1</sup>calculated to the marker compound at tracks III and IV

Two-dimensional (2D) development was carried out to determine the stability of the analytes during chromatography. A stable compounds will have the same R<sub>f</sub> value in the first and second elution.<sup>19</sup> Figure 5 exhibited that the analytes were stable during the chromatographic process indicated by a diagonal line on 2D chromatogram, including marker compound (\*).

Stability of the chromatographic results was determined by observing the color of the zones for 60 minutes. Figure 6 and Table 3 showed that the number and color of the zones were stable for 60 minutes. Differences of R<sub>f</sub> value for the marker zone was not more than 0.05.

### Precision test

Intraday precision test was conducted by chromatography of three OS leaves extracts prepared at the same time on one plate. Chromatography was replicated on 3 plates on the same day. Figure 7 and Table 4 show that the intraday precision of the method is acceptable since the  $\Delta R_f$  values of the marker zone (\*) from plate to plate is 0.02.<sup>19</sup>

The interday precision test was carried out similar to those of intraday precision, however the development of plates I, II, and III was conducted on 3 different days. Figure 8 and Table 5 exhibited that the interday precision of the method is acceptable because the difference of the R<sub>f</sub> value of marker (\*) is 0.02.<sup>19</sup>


### TLC-fingerprints of OS leaves from different origins

TLC profiles of OS leaves from 11 origins before and after derivatization are shown in Figure 9 and 10. Visually, sample 7 (Jombang) and 10 (Batu) showed significant differences compared to the others. Sample 7 has more zones and is more intensive compared to the others, while sample 10 is the opposite. This profile correlates with the colour of extract as presented at Figure 2.


TLC-fingerprint of samples after derivatization and visualized under UV 366 nm was then transferred into videodensitogram (Figure 11) to show their R<sub>f</sub> value, height, and area. Total peak number of each sample varies between 27 and 34 peaks, all show major peak with R<sub>f</sub> value 0.2-0.3. This peak can be further studied and can be considered as marker compound for OS. Moreover, height and peak area of sample 7 is more prominent compared to the others. This can be an initial indicator that the quality of sample 7 is better than the others. To compare the quality of samples more objectively, chemometric analysis was then performed on these TLC-fingerprints.

### Principal component analysis (PCA)

Videodensitogram of OS leaves extracts from 11 different origins were analyzed with chemometric using PCA method. The height and area of each peak detected on the videodensitogram were then tabulated based on the origin of the sample (table is not shown). PCA with full cross validation was applied to the data set of the 11 fingerprints of OS from 11 origins, 3 replications respectively. Analysis was conducted



**Figure 5:** Stability of the analytes during chromatography. MP:  $\text{CHCl}_3$ , DCM, EA (7:4:1). D: anisaldehyde-sulfuric acid reagent, UV 366 nm.



**Figure 6:** Stability of the chromatographic result after 5 (a), 10 (b), 30 (c), and 60 minutes (d). MP:  $\text{CHCl}_3$ , DCM, EA (7:4:1). D: anisaldehyde-sulfuric acid reagent, UV 366 nm.

**Table 3: Stability of the chromatographic result.**





Marker (*)	Track a	Track b	Track c	Track d
Colour	Yellowish-blue	Yellowish-blue	Yellowish-blue	Yellowish-blue
Rf	0.367	0.366	0.376	0.378
$\Delta Rf^1$	0	0.001	0.009	0.011

<sup>1</sup>calculated to the marker compound at track a**Table 4: Intraday precision test.**

Marker (*)	Track								
	I.1	I.2	I.3	II.1	II.2	II.3	III.1	III.2	III.3
Colour	Yellowish-blue			Yellowish-blue			Yellowish-blue		
Rf	0.404	0.388	0.388	0.431	0.409	0.406	0.415	0.402	0.393
Mean of Rf	0.39			0.41			0.40		
$\Delta Rf$				0.02					

**Table 5: Interday precision test.**

Marker (*)	Track								
	I.1	I.2	I.3	II.1	II.2	II.3	III.1	III.2	III.3
Colour	Yellowish-blue			Yellowish-blue			Yellowish-blue		
Rf	0.404	0.388	0.388	0.431	0.409	0.406	0.420	0.377	0.367
Mean Rf	0.39			0.41			0.38		
$\Delta Rf$				0.02					

**Figure 7:** Intraday precision test. 1, 2, 3 are OS leaves extracts prepared separately, while I, II, III show the different plates. MP: CHCl<sub>3</sub>, DCM, EA (7:4:1). D: anisaldehyde-sulfuric acid reagent, UV 366 nm.**Figure 8:** Interday precision test. 1, 2, 3 are OS leaves extracts prepared separately, while I, II, III show the plates developed in 3 different days. MP: CHCl<sub>3</sub>, DCM, EA (7:4:1). D: anisaldehyde-sulfuric acid reagent, UV 366 nm.**Figure 9:** TLC-fingerprint OS leaves from Lamongan (1), Pasuruan (2), Malang (3), Gresik (4), Surabaya (5), Sidoarjo (6), Jombang (7), Mojokerto (8), Karanganyar (9), Batu (10), and Mojokerto (11). MP: CHCl<sub>3</sub>, DCM, EA (7:4:1). D: none, white light (I), UV 254 nm (II), UV 366 nm (III).**Figure 10:** TLC-fingerprint of OS leaves from Lamongan (1), Pasuruan (2), Malang (3), Gresik (4), Surabaya (5), Sidoarjo (6), Jombang (7), Mojokerto (8), Karanganyar (9), Batu (10), and Mojokerto (11). MP: CHCl<sub>3</sub>, DCM, EA (7:4:1). D: anisaldehyde-sulfuric acid reagent, white light (I), UV 366 nm (II).




on the peak height and area of the full fingerprints without any preprocessing. The score plot of the first two PC (Figure 12) clearly distinguished 3 clusters of samples. The first cluster consists of OS from Mojokerto 1, Mojokerto 2, Malang, Jombang, and Sidoarjo. OS from Gresik, Pasuruan, Lamongan, Surabaya, and Karang Anyar gathered as second cluster, whereas third cluster consists of a sample from Batu. Samples in the cluster 1 show that the compounds with Rf value 0.2-0.3 are higher than those in clusters 2 and 3. However, these results indicate that the clustering of OS samples is not directly related to the height of the original location. Other factors such as soil type, rainfall,


lighting, fertilizing, etc. are predicted to be responsible for the samples grouping.<sup>21</sup>

To estimate which compounds are responsible for the grouping, PCA was conducted by projecting each origin variable called as loading plot. The loading plot of the first PC (Figure 13) showed that compounds with the Rf values of 0.0-0.1, 0.1-0.2, 0.2-0.3, and 0.9-1.0 are the most important compounds for clustering of samples.

Previous studies have been carried out to classify OS from different geographical origin. They involved both chromatography (HPLC, GC)



**Figure 11:** Videodensitogram of OS leaves, track 1-11 represent samples from Lamongan, Pasuruan, Malang, Gresik, Surabaya, Sidoarjo, Jombang, Mojokerto, Karanganyar, Batu, and Mojokerto, respectively.



**Figure 12:** PCA score plot of OS from different origins on the first two principal components.




Figure 13: PCA loading plot of OS from different origins on the first two principal components.

as well as spectroscopy (FTIR) method coupled with chemometrics analysis.<sup>10-12</sup> Our finding suggested that TLC fingerprint coupled with chemometrics analysis is a good alternative method to HPLC, GC, and FTIR for quality control of OS as well as the other crude drugs. In addition to its simplicity and low cost, by using TLC we can simultaneously analysis up to 20 samples under identical conditions.

## CONCLUSION

TLC-fingerprint combined with the PCA was able to discriminate among the leaves of *Orthosiphon stamineus* originated from various locations. Compounds with the Rf values 0.0-0.1, 0.1-0.2, 0.2-0.3, and 0.9-1.0 are the most important compounds for the clustering of samples. TLC-fingerprints analyzed with chemometrics is powerful method to evaluate the quality of *Orthosiphon stamineus*.


## ACKNOWLEDGEMENTS

The authors are thankful to Ministry of Research, Technology, and Higher Education of the Republic of Indonesia for financing this research under Fundamental Research Scheme with the contract number: 004/SP2H/LT/MULTI/L7/2019. This work also supported by University of Surabaya under Competitive Research Scheme with the contract number: 017/SP-Lit/LPPM-01/Int/FF/III/2019.

## REFERENCES

- Adnyana IK, Setiawan F, Insanu M. From ethnopharmacology to clinical study of *Orthosiphon stamineus* Benth. studies. 2013;5(3):66-73.
- Akowuah G, Zhari I, Norhayati I, Sadikun A. Radical scavenging activity of methanol leaf extracts of *orthosiphon stamineus*. *Pharmaceutical Biology*. 2005;42(8):629-35.
- Ho CH, Noryati I, Sulaiman SF, Rosma A. *In vitro* antibacterial and antioxidant activities of *Orthosiphon stamineus* Benth. extracts against food-borne bacteria. *Food Chemistry*. 2010;122(4):1168-72.
- Mohamed EAH, Siddiqui MJA, Ang LF, Sadikun A, Chan SH, Tan SC, et al. Potent  $\alpha$ -glucosidase and  $\alpha$ -amylase inhibitory activities of standardized 50% ethanolic extracts and sinensetin from *Orthosiphon stamineus* Benth as anti-diabetic mechanism. *BMC Complementary and Alternative Medicine*. 2012;12(1):176.
- Alshawsh MA, Abdulla MA, Ismail S, Amin ZA, Qader SW, Hadi HA, et al. Free radical scavenging, antimicrobial and immunomodulatory activities of *Orthosiphon stamineus*. *Molecules*. 2012;17(5):5385-95.
- Siddiqui MJA, Ismail Z. Simultaneous analysis of bioactive markers from *Orthosiphon stamineus* Benth leaves extracts by reverse phase high performance liquid chromatography. *Tropical Journal of Pharmaceutical Research*. 2011;10(1):97-103.
- Man S, Kiong LS, Ab'lah NA, Abdullah Z. Differentiation of the white and purple flower forms of *orthosiphon aristatus* (Blume) Miq. BY 1D and 2D correlation IR spectroscopy. *Jurnal Teknologi*. 2015;77(3):81-6.
- Ameer OZ, Salman IM, Asmawi MZ, Ibraheem ZO, Yam MF. *Orthosiphon stamineus*: traditional uses, phytochemistry, pharmacology, and toxicology. *Journal of Medicinal Food*. 2012;15(8):678-90.
- Chew K, Khoo M, Ng S, Thoo Y, Aida WW, Ho C. Effect of ethanol concentration, extraction time and extraction temperature on the recovery of phenolic compounds and antioxidant capacity of *Orthosiphon stamineus* extracts. *International Food Research Journal*. 2011;18(4):1427-35.
- Rafi M, Purwakusumah ED, Ridwan T, Barus B, Sutandi A, Darusman LK. Geographical classification of Java tea (*Orthosiphon stamineus*) from Java Island by FTIR spectroscopy combined with canonical variate analysis. *J Sains Matematika*. 2015;23:25-31.
- Saidan NH, Hamil MSR, Memon AH, Abdelbari MM, Hamdan MR, Mohd KS, et al. Selected metabolites profiling of *Orthosiphon stamineus* Benth leaves extracts combined with chemometrics analysis and correlation with biological activities. *BMC Complementary and Alternative Medicine*. 2015;15(1):350-62.
- Sim C, Ahmad M, Ismail Z, Othman A, Noor N, Zaihidee E. Chemometric classification of herb-*Orthosiphon stamineus* according to its geographical origin using virtual chemical sensor based upon fast GC. *Sensors*. 2003;3(10):458-71.
- Akowuah G, Zhari I, Sadikun A, Norhayati I. HPTLC Densitometric Analysis of *Orthosiphon stamineus*. Leaf Extracts and Inhibitory Effect on Xanthine Oxidase Activity. *Pharmaceutical Biology*. 2006;44(1):65-70.
- Hashim S, Beh H, Hamil M, Ismail Z, Majid A. High-performance thin-layer chromatography method development, validation, and simultaneous quantification of four compounds identified in standardized extracts of *Orthosiphon stamineus*. Gad HA, El-Ahmady SH, Abou-Shoer MI, Al-Azizi MM. Application of chemometrics in authentication of herbal medicines: a review. *Phytochem Anal*. 2013;24(1):1-24.
- Tistaert C, Dejaegher B, Heyden YV. Chromatographic separation techniques and data handling methods for herbal fingerprints: a review. *Analytica Chimica Acta*. 2011;690(2):148-61.
- Komsta Ł. Chemometrics in fingerprinting by means of thin layer chromatography. *Chromatography Research International*. 2011;2012.
- Bansal A, Chhabra V, Rawal RK, Sharma S. Chemometrics: a new scenario in herbal drug standardization. *Journal of Pharmaceutical Analysis*. 2014;4(4):223-33.
- Reich E, Schibli A. High-performance thin-layer chromatography for the analysis of medicinal plants. New York: Thieme; 2007.
- Brereton RG. Chemometrics: data analysis for the laboratory and chemical plant. West Sussex: John Wiley & Sons; 2003.
- Kartini, Azminah. Chromatographic fingerprinting and clustering of *Plantago major* L. from different areas in Indonesia *Asian Journal of Pharmaceutical and Clinical Research*. 2012;5(4):191-5.

## GRAPHICAL ABSTRACT



## ABOUT AUTHORS



**Kartini, Ph.D.** is an Associate Professor in the Department of Pharmaceutical Biology, Faculty of Pharmacy, University of Surabaya, Indonesia. She has completed her Ph.D. in Phytopharmaceutical Sciences from Faculty of Graduate Studies Mahidol University, Thailand. She is currently the Director of Center for Traditional Medicine Information & Development, Faculty of Pharmacy, University of Surabaya. She works on standardization of herbal medicines and its application as wound healing, anticancer, and immunomodulator.



**Ervina Rustiana Dewi** is a pharmacist who graduated from Faculty of Pharmacy, University of Surabaya, Indonesia. Her research focused on pharmacognosy study and standardization of *Orthosiphon stamineus*.






**Fandi Achmad** is a pharmacist who graduated from Faculty of Pharmacy, University of Surabaya, Indonesia. He is interested on research focusing on phytochemistry and quality assurance of *Orthosiphon stamineus*.



**Nikmatul Ikhrom Eka Jayani, M.Pharm.-Clin.** is a lecturer of Pharmaceutical Biology Department, Faculty of Pharmacy, University of Surabaya, Indonesia. She teaches some subjects, including Pharmacognosy and Indonesian Indigenous Herbal Medicine. She focuses her research on standardization of herbal medicine and its activity as antidiabetes as well as its application in community setting.



**M. Arbi Hadiyat, M.Sc.** is a lecturer of Industrial Engineering Department, Faculty of Engineering, University of Surabaya, Indonesia. Some courses have been taught by him, such as Industrial Statistics, Design of Experiment, and Quality Engineering. Currently, he is working on some researches in quality and process design optimization, machine learning in industry, and customer satisfaction mathematical model.



**Christina Avanti, Ph.D.** is an Associate Professor, Vice Rector for Student Affairs and Alumni, Former Dean of Faculty of Pharmacy University of Surabaya, Indonesia. PhD graduated in Pharmacy at the Department of Pharmaceutical Technology and Biopharmacy University of Groningen. She is currently working in several projects developing techniques to improve bioavailability and stability of lipophilic pharmaceuticals and develop modified delivery system of Indonesian medicinal plants.

**Cite this article:** Kartini K, Dewi ER, Achmad F, Jayani NIE, Hadiyat MA, Avanti C. Thin Layer Chromatography Fingerprinting and Clustering of *Orthosiphon stamineus* Benth. from Different Origins. *Pharmacog J.* 2020;12(1):1683-91.

# Pharmacognosy Journal

An Open Access, Peer Reviewed Journal in the field of  
Pharmacognosy

also developed by scimago:



SCIMAGO INSTITUTIONS RANKINGS

SJR

Scimago Journal &amp; Country Rank

Enter Journal Title, ISSN or Publisher Name

[Home](#)[Journal Rankings](#)[Country Rankings](#)[Viz Tools](#)[Help](#)[About Us](#)

Ads by Google

[Stop seeing this ad](#)[Why this ad?](#)

## Pharmacognosy Journal

### COUNTRY

[India](#) Universities and research institutions in India

### SUBJECT AREA AND CATEGORY

[Pharmacology, Toxicology and  
Pharmaceutics](#)  
[Drug Discovery](#)  
[Pharmacology](#)

### PUBLISHER

[EManuscript Technologies](#)

### H-INDEX

**21**

### PUBLICATION TYPE

[Journals](#)

### ISSN

09753575

### COVERAGE

2009-2020

### INFORMATION

[Homepage](#)[How to publish in this journal](#)[editor@phcogj.com](mailto:editor@phcogj.com)

Ads by Google

[Stop seeing this ad](#)[Why this ad?](#)

### SCOPE

Pharmacognosy Journal (Phcog J.) covers different topics in natural product drug discovery, and also publishes manuscripts that describe pharmacognostic investigations, evaluation reports, methods, techniques and applications of all forms of medicinal plant research

[Join the conversation about this journal](#)

### Discovery begins with Scopus

Comprehensive &amp; selected by experts.

Scopus

[Learn More](#)





**Scopus Indexed Journal**

Scopus, Web of Science, esci

Publish Your Paper in International Lowest Pub. Fee, Fast Review & Pub.

[ephjournal.org](http://ephjournal.org)

OPEN

## FIND SIMILAR JOURNALS ?


options :

1	<b>Oriental Pharmacy and Experimental Medicine</b> USA	82% similarity
2	<b>Pharmacognosy Research</b> IND	81% similarity
3	<b>Indian Journal of Natural Products and Resources</b> IND	77% similarity
4	<b>Journal of Herbs, Spices and Medicinal Plants</b> USA	76% similarity
5	<b>Pharmacognosy Magazine</b> IND	76% similarity

## Discovery begins with Scopus

Comprehensive &amp; selected by experts.

Scopus

[Learn M](#)

Pharmacognosy Journal

← Show this widget in your own website

Q3 Drug Discovery

SJR 2020 0.27

powered by scimagojr.com

Just copy the code below and paste within your html code:

`<a href="https://www.scimagojr.com" >`

**SCImago Graphica**

Explore, visually communicate and make sense of data with our new free tool.





# Source details

## Pharmacognosy Journal

Scopus coverage years: from 2009 to Present

Publisher: Pharmacognosy Network Worldwide

ISSN: 0975-3575

Subject area: Pharmacology, Toxicology and Pharmaceutics: Pharmacology

Pharmacology, Toxicology and Pharmaceutics: Drug Discovery

Source type: Journal

CiteScore 2020

1.6



SJR 2020

0.268



SNIP 2020

0.775



View all documents >

Set document alert

Save to source list

CiteScore CiteScore rank & trend Scopus content coverage

### i Improved CiteScore methodology



CiteScore 2020 counts the citations received in 2017-2020 to articles, reviews, conference papers, book chapters and data papers published in 2017-2020, and divides this by the number of publications published in 2017-2020. [Learn more >](#)

CiteScore 2020

$$1.6 = \frac{1,368 \text{ Citations 2017 - 2020}}{852 \text{ Documents 2017 - 2020}}$$

Calculated on 05 May, 2021

CiteScoreTracker 2021

$$1.6 = \frac{1,368 \text{ Citations to date}}{852 \text{ Documents to date}}$$

Last updated on 04 September, 2021 • Updated monthly

## CiteScore rank 2020

Category	Rank	Percentile
Pharmacology, Toxicology and Pharmaceutics	#214/297	28th
Pharmacology		
Pharmacology, Toxicology and Pharmaceutics	#112/145	23rd
Drug Discovery		

[View CiteScore methodology >](#) [CiteScore FAQ >](#) [Add CiteScore to your site](#)

About Scopus

- What is Scopus
- Content coverage
- Scopus blog
- Scopus API
- Privacy matters

Language

- 日本語に切り替える
- 切换到简体中文
- 切换到繁體中文
- Русский язык

Customer Service

- Help
- Contact us

ELSEVIER

[Terms and conditions ↗](#)   [Privacy policy ↗](#)

Copyright © Elsevier B.V. All rights reserved. Scopus® is a registered trademark of Elsevier B.V.  
We use cookies to help provide and enhance our service and tailor content. By continuing, you agree to the use of cookies.





Time to  
read  
less than  
1 minute

## About Journal

New site available from 21 Sep, 2015

Share



Print

Read so  
far

100%

Pharmacognosy Journal (Phcog J.) covers different topics in natural product drug discovery, and also publishes manuscripts that describe pharmacognostic investigations, evaluation reports, methods, techniques and applications of all forms of medicinal plant research

**Distinctions:** The most widely read, cited, and known Pharmacognosy journal and website is well browsed with all the articles published. More than 50,000 readers in nearly every country in the world each month

**ISSN :** 0975-3575 ; Frequency : Rapid at a time publication (6 issues/year)

**Indexed and Abstracted in :** SCOPUS, Scimago Journal Ranking, Chemical Abstracts, Excerpta Medica / EMBASE, Google Scholar, CABI Full Text, Index Copernicus, Ulrich's International Periodical Directory, ProQuest, Journalseek & Genamics, PhcogBase, EBSCOHost, Academic Search Complete, Open J-Gate, SciACCESS.

**Rapid publication:** Average time from submission to first decision is 30 days and from acceptance to In Press online publication is 45 days.

**Open Access Journal:** Pharmacognosy Journal is an open access journal, which allows authors to fund their article to be open access from publication.

# Editorial Board (2020-21)

## Editors & Editorial Board Members (2021)

### **Dr.Djemli Samir**

Department of Biology , Applied Neuroendocrinology Laboratory  
Badji Mokhtar Annaba University  
Algeria

### **Dr. Raghava Naidu, Ph.D**

Department of Human Oncology,  
University of Wisconsin,  
1111, Highland Ave, Madison,  
Wisconsin 53705, USA

### **Dr.Karim Raafat**

Associate Professor of Pharmacognosy and Phytochemistry,  
Pharmaceutical Sciences Department,  
Faculty of Pharmacy,  
Beirut Arab University (BAU),  
Beirut 115020, Lebanon

### **Ourlad Alzeus Tantengco, MD-PhD Molecular Medicine**

College of Medicine, University of the Philippines Manila  
Pedro Gil Street, Ermita, Manila, Philippines, 1000

### **Janib Achmad**

Lecturer of Faculty of Fisheries and Marine Science,  
University of Khairun Ternate  
Kampus 2 Jalan Pertamina, Kelurahan Gambesi,  
Ternate Selatan

### **Muammar Fawwaz, Ph.D**

Department of Pharmaceutical Chemistry  
Faculty of Pharmacy  
Universitas Muslim Indonesia  
Makassar 90231, South Sulawesi, Indonesia

### **Hany Ezzat Khalil**

Associate Professor,  
College of Clinical Pharmacy,  
King Faisal University,  
KSA

### **Emad Yousif**

Department of Chemistry  
College of Science  
Al-Nahrain University  
Baghdad, Iraq

### **Sughosh Upasani**

R.C Patel Institute of pharmacy,  
Shirpur, Dist-Dhule, Maharashtra,  
India.

### **Gurusiddaiah suresh kumar**

Scientist  
Dept of biochemistry  
CSIR-CFTRI  
Mysore, Karnataka, INDIA

**Arjun Patra**

Assistant Professor  
School of Pharmaceutical Sciences  
Guru Ghasidas Central University  
Koni, Bilaspur - 495 009  
Chhattisgarh, India

**Francis O. Atanu, Ph.D**

Department of Biochemistry  
Faculty of Natural Sciences  
Kogi State University  
Anyigba, Nigeria.

**Vijay Kumar Chattu**

Faculty of Medical Sciences  
University of the West Indies  
St. Augustine, Trinidad & Tobago.

**Dr.Kunle Okaiyeto, PhD**

Applied and Environmental Microbiology Research Group (AEMREG)  
Department of Biochemistry and Microbiology  
University of Fort Hare  
Alice campus  
5700, Alice  
South Africa.

**Dr. Srisailam Keshetti, Ph.D**

Principal, University College of Pharmaceutical Sciences, Satavahana University  
Karimnagar 505001  
Telangana  
INDIA

**Dr. Gayathri M Rao**

Associate Professor  
Department of Biochemistry  
Kasturba Medical College, Mangaluru.

**Shuge Tian**

Experimental Teaching Demonstration Center of TCM in Xinjiang Medical University  
Department of traditional medicine, TCM  
Xinjiang Medical University  
Xinjiang CHINA 830054

**Dr. Ramachandra Setty Siddamsetty,**

Professor, Govt College of Pharmacy,  
Mission Road, Bengaluru, INDIA

**Dr. (Mrs.) Sayyada Khatoon**

HOD, Pharmacognosy Division  
CSIR-National Botanical Research Institute,  
Rana Pratap Marg, Post Box 436,  
Lucknow-226001 (U.P.) India

**Dr. A. Sajeli Begum**

Department of Pharmacy  
Birla Institute of Technology & Science  
Hyderabad, India



**Olga Silva**

Department of Pharmacological Sciences,  
Faculdade de Farmácia,  
Universidade de Lisboa, Portugal

**Xinwen Wang**

Department of Clinical Pharmacy  
University of Michigan  
USA

**Roman Lysiuk**


Department of Pharmacognosy and Botany,  
Danylo Halytsky Lviv National Medical University,  
Pekarska,69., Lviv 79010, Ukraine

**Arif Nur Muhammad Ansori**

Universitas Airlangga  
Indonesia

# Pharmacognosy Journal, Vol 12, Issue 1, Jan-Feb 2020

## RECENT ARTICLES



Original Article

## Toxicity Assessment of Vachellia karro (Hayne) Banfi and Galasso Pods using Brine Shrimp Assay

S Maposa,AJ Afolayan,GA Otunola

Pharmacognosy Journal,12(1):1-5

DOI: 10.5530/pj.2020.12.1

Published: Fri, 7-Feb-2020

[Read More](#)

Original Article

## Neuro-protective Effect of Ayurveda Formulation, Saraswatharishtam, on Scopolamine Induced Memory Impairment in Animal Model


Jai Prabhu,K Prabhu,Anathbandhu Chaudhuri,Mudiganti Ram Krishn Rao,VS Kalai Selvi,TK Balaji,Sruthi Dinakar

Pharmacognosy Journal,12(1):6-13

DOI: 10.5530/pj.2020.12.2

Published: Fri, 7-Feb-2020

[Read More](#)



Original Article

## In vitro Cytotoxicity of Hibiscus sabdariffa Linn Extracts on A549 Lung Cancer Cell Line

Qotrunnada Fithrotunnisa,Ade Arsianti,Gerry Kurniawan,Fona Qorina,Nadzila Anindya Tejaputri,Norma Nur Azizah



Pharmacognosy Journal,12(1):14-19

DOI: 10.5530/pj.2020.12.3

Published: Fri, 7-Feb-2020

[Read More](#)

Original Article

## Thin Layer Chromatography Fingerprinting and Clustering of *Orthosiphon stamineus* Benth. from Different Origins

Kartini Kartini, Ervina Rustiana Dewi, Fandi Achmad, Nikmatul Ikhrom Eka Jayani, Mochammad Arbi Hadiyat, Christina Avanti



Pharmacognosy Journal, 12(1):79-87

DOI: 10.5530/pj.2020.12.13

Published: Fri, 7-Feb-2020

[Read More](#)

Original Article

## Phytochemical Composition and Evaluation of Marine Algal *Sargassum polycystum* for Antioxidant Activity and In Vitro Cytotoxicity on Hela Cells

Ade Arsianti, Anton Bahtiar, Vincent Kharisma Wangsaputra, Norma Nur Azizah, Wilzar Fachri, Lince Dameria Nadapdap, Ajeng Megawati Fajrin, Hiroki Tanimoto, Kiyomi Kakiuchi



Pharmacognosy Journal, 12(1):88-94

DOI: 10.5530/pj.2020.12.14

Published: Fri, 7-Feb-2020

[Read More](#)

Research Article

## Phytochemical Analysis and Anti-diabetic Activity of Leaf extract of *Psydrax horizontalis* Schum and Thonn (Rubiaceae)

Onyekere Peculiar Feenna, Odoh Uchenna Estella, Ezugwu Christophe Obodike



Pharmacognosy Journal, 12(1):95-102

DOI: 10.5530/pj.2020.12.15

Published: Fri, 7-Feb-2020

[Read More](#)