

Strategies to implement community training to promote responsible self-medication in Indonesia: a qualitative study of trainers

Adji P. Setiadi^{a,*}, Yosi I. Wibowo^a, Eko Setiawan^a, Ika Mulyono^a, Susilo A. Wardhani^b and Bruce Sunderland^c

^aCentre for Medicines Information and Pharmaceutical Care – Faculty of Pharmacy, Universitas Surabaya, Surabaya 60293, Indonesia;

^bEast Java Provincial Health Office, Ministry of Health Republic of Indonesia, Surabaya 60231, Indonesia; ^cSchool of Pharmacy, Faculty of Health Sciences, Curtin University, Perth, Western Australia 6102, Australia

*Corresponding author: Tel: +62 31 2981170; Email: adj_i_ps@staff.ubaya.ac.id

Received 12 May 2019; revised 29 October 2019; editorial decision 30 October 2019; accepted 30 October 2019

Background: A community-based approach has been identified as one key strategy to improve the health of Indonesians. In 2015, the government initiated the ‘Smart Use of Medications Movement’ (GeMa CerMat) to promote responsible self-medication. This study aims to explore pharmacist/pharmacy staff trainers’ views on strategies to implement GeMa CerMat community training.

Methods: Four focus group discussions were conducted with 38 pharmacist/pharmacy staff trainers in Ngawi, Indonesia and a thematic analysis was used to obtain the key strategies.

Results: The identified key strategies were building community readiness via well-designed training (considering participant characteristics, training methods, and materials and resources); policy, regulation and organizational support; access to training in a cultural context; communication media to promote training and the inclusion of active-learning tools.

Conclusions: These identified multilevel strategies require collaboration among national bodies and the involvement of trusted community members (‘change agents’). A new strategy using ‘active learning tools’ was necessary to build critical thinking and understanding of the use of medications in everyday life. Future research should focus on process, impact and outcome evaluation involving GeMa CerMat training implementation and sustainability in Indonesia.

Keywords: education, Indonesia, pharmacist, self-medication

Introduction

Indonesia is the world’s largest island nation with a population of >200 million.¹ With an increasing ageing population and prevalence of non-communicable diseases, in conjunction with the high prevalence of infectious diseases, health challenges in Indonesia are daunting.² It is a priority of the Indonesian government to improve the quality of life among Indonesians through a ‘Healthy Indonesia’ programme, including promoting a healthy living paradigm, strengthening health services/facilities and implementing *Jaminan Kesehatan Nasional* (JKN), which provides basic healthcare coverage to the entire population.³ Primary care facilities, mainly primary health centres (PHCs, *puskesmas*), are the front-line providers to implement these programmes. However, the wide dispersion of islands making up Indonesia creates problems with supply, mix and distribution of primary health facilities and workforce.^{2,3} Thus community

empowerment could be seen as one key strategy to improve the health of Indonesians.²

In 2015, the Indonesian government initiated a community empowerment programme, the ‘Smart Use of Medications Movement’ (*Gerakan Masyarakat Cerdas Menggunakan Obat* [GeMa CerMat]), which aimed to promote responsible self-medication among Indonesians;⁴ this was followed by an overarching programme, the ‘Healthy Lifestyle Movement’ (*Gerakan Masyarakat Hidup Sehat* [GerMas]) to include self-care.⁵ Self-medication is defined as ‘the selection and use of medicines by individuals to treat self-recognized illnesses or symptoms’.⁶ Due to the overstretched health system in Indonesia,² self-medication plays a pivotal role as the first action taken by Indonesians in managing their healthcare, especially for minor ailments. Minor ailments can be defined as usually self-limiting medical conditions that can be reasonably self-diagnosed and self-managed

© The Author(s) 2020. Published by Oxford University Press on behalf of Royal Society of Tropical Medicine and Hygiene.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact journals.permissions@oup.com

with over-the-counter (OTC) products; this might include headache, coughs, colds and constipation.⁷ Based on a survey in 2014, 61% of the Indonesian population practises self-medication.⁸

While responsible self-medication can move people toward greater independence to treat their minor ailments, potential risks of self-medication have been reported, such as incorrect self-diagnosis, drug interactions, infrequent but severe adverse reactions, incorrect choice of therapy and risk of drug dependence and abuse.^{9,10} Community pharmacists are well established in Indonesia and have traditionally assisted the community with the management of minor ailments and, when appropriate, referred them to other health practitioners; however, many medicines are also available from drugstores, informal outlets and street peddlers.² Hence the GeMa CerMat initiative is expected to improve self-medication literacy, thus empowering Indonesians in the safe use of OTC medicines.

As part of the GeMa CerMat initiative, community training on basic medication information was developed.⁴ Pharmacists, with their educational background in medications and as suppliers of OTC medicines, are appropriate healthcare practitioners for providing community training. In 2016, selected pharmacist-pharmacy staff were invited for a GeMa CerMat trainers' briefing in four cities/districts: Nganjuk, Blitar, Ngawi and Sumenep.¹¹ Little is known, however, regarding pharmacist/pharmacy staff trainers' experiences (barriers/facilitators) while performing the community training in their practice settings. This study aims to explore trainers' views on strategies to implement GeMa CerMat community training to promote responsible self-medication in Indonesia.

Methods

The study was approved by the ethics committee of the Faculty of Medicine, Islamic University of Indonesia (No. 08/Ka.Kom. Et/70/KE/IX/2016).

Setting, design and sample

The study site was Ngawi, a district (*kabupaten*) in the western part of East Java province, Indonesia. Based on statistical data from 2012,¹² Ngawi covers an area of 1296 km² with a population of 911 911 people and is divided into 17 subdistricts (*kecamatan*) consisting of 217 villages (*desa*). Health facilities in Ngawi include 2 hospitals, 24 PHCs and 51 community pharmacies; a PHC is a technical unit of the Region Health Office that provides primary healthcare at a subdistrict level. In 2016, 46 pharmacist/pharmacy staff in Ngawi were invited to attend a GeMa CerMat trainers' briefing. A module was provided to guide trainers to facilitate interactive learning on basic medication information provided on the label/package/leaflet (i.e. drug name, indication, administration, adverse effects and storage).¹¹

This study was to follow up pharmacists/pharmacy staff involved in the prior trainers' briefing in Ngawi to explore their views on strategies to implement community training to promote responsible self-medication in Indonesia. Focus group discussions (FGDs) were chosen since this allowed the trainers, who had similar backgrounds or experiences, to discuss a specific

topic of interest.¹³ Thus FGDs are a good way to provide insights into a range of views/opinions of people involved in the change process, especially where the aim is to improve a targeted service. These were guided by the Consolidated Criteria for Reporting Qualitative Research (COREQ) requirements.¹⁴

Data collection

Invitation letters were sent by the head of the Pharmacy Division of the Ngawi Health Office to a total of 46 pharmacist/pharmacy staff who had participated in the prior GeMa CerMat trainers' briefing to attend a meeting that was scheduled in May 2017. The meeting was conducted in the Ngawi Health Office. The researcher initially introduced the research and asked for the trainers' participation. Those willing to participate were asked to complete a short demographic questionnaire and to provide written informed consent. They were then invited to join one of four FGDs.

Each FGD was facilitated by one of the researchers, who were a master/doctor in pharmacy as well as an academic. All FGDs were conducted in Bahasa Indonesia and were aided by a semi-structured FGD guide. The guide was developed in conjunction with all researchers and was field tested. The guide explored two areas: training experiences (barriers/facilitators) and proposed strategies to implement the community training. Probing questions were asked when necessary to expand on the participants' responses. The FGDs lasted 45–60 min and were audio recorded.

Data analysis

Audio-recorded data from the FGDs were transcribed verbatim by the facilitators. Transcripts were reviewed using inductive thematic analysis to identify salient themes;¹⁵ all transcripts were coded manually by cutting and pasting between documents. The analysis was performed by YW and the emerging themes were validated by discussion with the other research team members (AP, ES) to reach a consensus. The data analysis first involved a process of familiarization; significant statements related to proposed strategies to implement community training from the data were then identified and coded. The codes were categorized into a broader conceptual level (i.e. into themes). The process of theme generation was reviewed and refined by going back and forth between the themes, the codes and the transcripts until the final themes were identified. Finally, examples of responses were selected to illustrate each theme. The final themes and quotes were discussed with the trainers during a follow-up seminar in October 2017 as a means of member-checking, thus ensuring the credibility of the data.¹⁶ Data analysis was conducted in Bahasa Indonesia and the illustrative verbatim quotes and theme labels were translated into English.

Results

Of 46 pharmacists/pharmacy staff who participated in the prior trainers' briefing and were thus invited to participate in this study, a total of 40 consented and attended a meeting held in the Ngawi Health Office in May 2017. Two subsequently withdrew due to emergency work tasks, leaving 38 who were allocated to join four FGDs (their characteristics are summarized in Table 1).

Table 1. Characteristics of the pharmacist/pharmacy staff trainers involved in the FGDs

| Characteristics | FGD 1 (n=10) | FGD 2 (n=9) | FGD 3 (n=10) | FGD 4 (n=9) |
|--|--|---|--|---|
| Age range (years) | 30–48 | 29–39 | 30–51 | 25–41 |
| Gender | 3 males, 7 females | 9 females | 3 males, 7 females | 9 females |
| Position | 2 PHC pharmacists, 3 community pharmacists, 5 PHC pharmacy staff | 2 PHC pharmacists, 2 community pharmacists, 5 PHC pharmacy staff | 2 PHC pharmacists, 8 PHC pharmacy staff | 5 community pharmacists, 4 PHC pharmacy staff |
| Training experience | | | | |
| – Number of training sessions (range per person) | 1–4 | 1–4 | 1–3 | 1–3 |
| – Training participants | | | | |
| Participant characteristics | Seniors (1 time), members of women's organizations (1–3 times), members of religious gathering (1 time), community health representatives/leaders (1 time), general population (1–2 times) | Seniors (1–2 times), members of women's organizations (1–3 times), community health representatives/leaders (1 time), students (1–2 times), general population (1 time) | Seniors (1–2 times), members of women's organizations (1–3 times), community health representatives (1 time), general population (1–2 times) | Community health representatives/leaders (1 time), general population (1–2 times) |
| Number of participants (range) | 30–100 | 15–200 | 15–100 | 15–100 |

Trainers, perceived strategies to implement community training

Pharmacists/pharmacy staff involved in the FGDs had been involved in at least one training session as trainers (range 1–4 sessions per person); while the training sessions were often conducted during community meetings, including meetings of women's organizations, senior gatherings, religious gatherings, educational sessions at school or community health representative/leader meetings (see Table 1). Responses from the FGDs in relation to the trainers' perceived strategies to implement training to promote responsible self-medication in Ngawi identified five main themes: building community readiness via well-designed training; policy, regulation and organizational support; access to training in a cultural context; communications media and active learning tools.

Building community readiness via well-designed training

This study indicated the importance of designing appropriate training to build community readiness to practise responsible self-medication. It was suggested that factors directly contributing to training delivery were participant characteristics, training methods, and materials and training resources.

Participant characteristics

Participant characteristics might be linked with learning capacity and motivation. This should be considered when delivering training to promote responsible self-medication. Some of the trainers believed that it would be best for the training to be

targeted to specific participant groups, thus the training could be designed accordingly. Community health representatives, community leaders, students or mothers could be considered as the initial targets; these groups generally have higher learning capacity and/or influence, thus having the potential to assist pharmacists to implement community training in their local communities ('change agents'). This was illustrated by comments from pharmacist 4.4: *'The first would be community health representatives. Every year, we are given the opportunity to speak at community health representatives meetings – we usually use it to present GeMa CerMat'*; pharmacist 2.9: *'Usually, community leaders would have better capability – even asking extended questions'*; pharmacist 4.3: *'Clearly, I would prioritise mothers; as in general, they were the ones who take the initiative when their children were ill'*; and pharmacist 2.8: *'In my opinion, we need to target high-school students; younger people have higher capability to learn and could assist their family on using medications properly'*.

Training methods and materials

Training methods and materials should be made to fit participant characteristics. Due to the oral traditions among the Ngawi communities, some trainers suggested interactive methods with oral and fewer written tasks for the participants, such as using games, question-and-answer sessions, pictures/videos or simulations using medication packages. This was illustrated by comments from pharmacy staff 3.4: *'Usually the participants are mothers . . . they are reluctant to do writing tasks; they want us to do questions and answers . . . just like discussion'*; and pharmacist

2.9: *'To attract high school students' interests, I am not using slides instead I am adding some videos'.*

Further, a few trainers indicated the importance of developing a range of training guides focusing on specific or local health issues, e.g. anaemia in pregnancy, diarrhoea or the common cold. It was suggested that trainers, to bridge the literacy gap with participants, should use simple language when delivering the material; as pharmacist 1.8 stated: *'In some cases, we need to adjust – using language that could be easily understood, even using local dialects etc.'*

Training resources

Training resources refers to resources needed to perform self-medication training, including trainers, funding and facilities. Some trainers revealed that the current GeMa CerMat community training is not yet a priority activity of the local health office or the Indonesian Pharmacists Association (*Ikatan Apoteker Indonesia [IAI]*), thus permanent funding has not been established. To obtain the facilities required for the training, the training was often conducted along with other established community meetings, such as community health representative meetings, religious gatherings, health education sessions (at school) or neighbourhood meetings, as illustrated by pharmacist 2.8: *'We have routine activities such as providing health education at schools and we usually include GeMa CerMat there'.*

Adequate numbers of trainers should be available to facilitate efficient interactive discussion; as pharmacy staff 3.4 stated: *'Ideally one group is facilitated by one trainer'.* Few trainers also perceived the importance of setting skill qualifications for those who are allowed to be trainers, such as pharmacists or having completed the trainers' briefing, as illustrated by pharmacy staff 4.2: *'I am a diploma graduate, thus my clinical role is limited, sometimes the participants know more than me. It would be better to have a pharmacist'.*

Policy, regulation and organizational supports

Despite the Indonesian government's initiative, GeMa CerMat community training has not been necessarily translated into the local health office and/or IAI policies and priority programmes. This was illustrated by comments from pharmacist 2.8: *'At PHC, 60% are primary activities and 40% are supporting activities. Pharmacy is only included within supporting activities'.*; and pharmacist 4.3: *'Community pharmacists usually involve (to introduce GeMa CerMat) in their local community activities on their own initiatives'.*

Only after the training was viewed as a priority or routine activity within the local health office and/or IAI did some trainers believe that permanent funding would be established for sustained community training. Further, it was suggested that the IAI develop a remuneration system for pharmacists involved as trainers, such as incentives and/or continuing professional development (CPD) credits.

Access to training in a cultural context

Access to training and various cultural influences might affect the implementation of community training. Trainers emphasized the issues of oral language traditions among target communities; this sociocultural context should guide the development of

appropriate training methods and materials (see 'Training methods and material'). In addition, limited access to training—due to isolated and scattered residential areas—was also considered a barrier; as pharmacy staff 1.5 stated: *'I am residing in an isolated area, in the forest, very rural. Gathering 50 people would be very difficult, (as) they have scattered residences, long distances between them, lots of forest'.* Thus it was suggested to involve community health representatives/leaders in local areas to assist in training implementation to reach a broader population.

Communication media

Some trainers indicated the importance of the local health office in collaboration with the IAI to communicate or disseminate the GeMa CerMat programme to increase awareness of the target community; as pharmacist 4.3 stated: *'On a special occasion, such as World Pharmacist Day, the local IAI could promote "GeMa CerMat", such as installing banners at pharmacies, sharing via social media, radio etc.'*

Active learning tools

While participatory and interactive training materials might help participants to learn about medications, some concerns were raised regarding community empowerment beyond the training; as pharmacist 1.10 mentioned: *'Participants need to be subsequently evaluated and continuously encouraged to be careful when using medications'.* This situation might imply the need for the training to provide community members with a range of active learning tools to build their critical thinking on the use of medications in their daily life.

The five key strategies identified from the FGDs of trainers could be interpreted as an in-process evaluation as part of a cyclical process (assessment–implementation–evaluation) that allows for continuous improvement of a health programme as described in the PRECEDE–PROCEED model.¹⁷ Considering the structure of the Indonesian health system,¹⁸ the strategy component of 'policy, regulation and organizational support' could be further explained in terms of a pyramid model to represent levels of policy, regulation and support needed while implementing the community training (Figure 1). At the top would be the national policy/regulation related to GeMa CerMat, which is then required to be translated down to the regional and local health offices, and the IAI, to reach the target communities.

Discussion

Community-based interventions have been identified at the government level as one key strategy to improve the health status of Indonesians. This qualitative study was initiated to follow up pharmacists/pharmacy staff who had previously participated in a trainers' briefing to explore their views on strategies used to perform community training as an intervention to enhance self-medication in their practice settings. The findings identified five key strategies: building community readiness via well-designed training; policy, regulation and organizational support; access to training in a cultural context; communication media and the inclusion of active learning tools to build critical thinking on the use of medications in everyday life (see Figure 1). These strategies




Figure 1. Levels of policy, regulation and organisational support for sustained community training to promote self-medication in Indonesia.

target factors directly related to the training (internal factors) as well as enabling and reinforcing external factors, all of which could produce synergistic effects for a successful implementation.^{17,19}

Building community readiness via a well-designed training model could be seen as a strategy directly related to the training (internal factors). This study indicated that the training should be designed based on group target participant characteristics. Initially the training could target trusted members of the community, such as health representatives/leaders, students or mothers, as they have the potential to assist in the training implementation ('change agents') in the household or community groups. This should be facilitated with the development of appropriate training methods and materials. Although the evidence available in the literature was inconclusive on which methods work best for patient education, audio-visual, written materials (prepared at an appropriate reading level), demonstrations (if appropriate for the situation) and the use of mixed methods were considered to be effective.²⁰

While the current training material developed was at the general information level on self-medication use, this could be modified to focus on specific local health groups, such as mothers, or other target participant needs. In addition to common minor illnesses (such as diarrhoea, cough, cold and joint/muscular pain),^{12,21} the increasing prevalence of diabetes and hypertension in Indonesia also warrants consideration in developing training materials related to a healthy lifestyle. This should provide self-care support to the current priority health programme, GeMa, which was introduced after GeMa CerMat.⁵ In addition to the improved materials, this study implied the need to provide the training participants with active learning tools. The tools should facilitate community members to be actively and critically able

to obtain information from health professionals, for example, 'Asking 5+' when obtaining medications:²²

1. Name and composition—what is the name and composition of this medication?
2. Indication—what is this medication used for?
3. Directions for use—how do I use this medication?
4. Side effects—what are the common side effects of this medication?
5. Storage—how to store this medication?

+ the expiry date.

Past research has suggested a continuous process of discussion, reflection and action, termed 'critical awareness' and 'critical thinking', forming the basis for approaches for learning and social change.²³

Access to training in a cultural context relates to strategies to enable implementation of the training across the district. Considering Indonesia's communal and paternalistic culture as well as geographic proximity between areas,^{24,25} it would be best to integrate the educational programme into local community activities, thus the involvement of trusted members of the community, such as community health representatives/leaders, would be of importance. Studies worldwide have reported that pupils can act as agents for health interventions at both the school and local community levels ('change agents').²⁵⁻²⁷ As part of the community, these change agents would be more accessible and acceptable to the local community.²⁶

Furthermore, this study has suggested strategies related to 'public policy and organizational support' and 'communication media' to reinforce sustained community training (reinforcement factors). Although the Indonesian government initiated GeMa CerMat in 2015,^{4,11} a key challenge lies in translating the

movement into local regulations/policies. Indonesia's decentralized government system provides districts (*kabupaten*) and municipalities (*kota*) with autonomy to manage regional public sectors, including health.²⁸ Hence GeMa CerMat training should become part of the regional health office's responsibility for supporting and funding GeMa CerMat training performed at the local community level. In addition, the IAI (in collaboration with the local health office) would have a pivotal role in arranging for the training as well as establishing trainers' accreditation and remuneration systems.

The importance of communication media to promote community training was raised. Literature has suggested that health programmes may benefit from media campaigns to expose large audiences to focused messages over time;²⁶ however, the decision to use a particular medium should first consider the audiences' socio-economic and cultural backgrounds.^{29,30} Considering the oral traditions of the target community in this study, the health office and IAI should make use of audio-visual media (e.g. television and radio advertisements) and social media for training publicity.

There are some limitations to this study. This study only included views of pharmacist/pharmacy staff trainers, thus it might not include views from other stakeholders, such as community members or policymakers. However, as the main providers in this initiative, the views of pharmacist/pharmacy staff trainers should provide insight regarding the strategies to promote responsible self-medication in Indonesia. As qualitative data are a product of the views, experiences and perceptions of the respondents, bias can occur if respondents are not sharing their true responses.¹³ The FGDs, however, were carefully designed to include participants with similar backgrounds and experiences (pharmacist trainers) and to discuss a common issue (implementation strategy), thus creating a non-intimidating environment where participants should have been able to express their true ideas/opinions. To ensure accuracy, the results of this study were provided to the participating trainers (member checking).¹⁵ In addition, no new themes emerged at the third FGD, suggesting saturation was reached.¹³

Conclusions

This study has proposed a multilevel strategy for sustained community training to promote responsible self-medication within an Indonesian context. There is a need for the training to provide active learning tools to help Indonesians engage in learning activities while obtaining their medications in their daily lives. These findings should be used as a basis to develop a range of interventions to support trainers while implementing community training. Future research should focus on the process, impact and outcome evaluation involving all stakeholders to provide comprehensive and sustainable community training to promote responsible self-medication in Indonesia.

Authors' contributions: AP made contributions to the conception and design of the study. AP, YW, ES and IM facilitated the FGDs and transcribed the data. AP, YW, ES, IM and ASW participated in the data analysis. YW wrote the first draft of the manuscript. BS contributed to the design of the

study and interpretation of data. All authors were involved in the revisions to the manuscript and final approval of the version to be published.

Acknowledgements: We thank all the participants who took part in this study as well as the chief and staff from the East Java Provincial Health Office and Ngawi District Health Office – Ministry of Health Republic of Indonesia and the Centre of Medicines Information and Pharmaceutical Care (CMIPC) – Faculty of Pharmacy Universitas Surabaya for their technical support in making this study possible.

Funding: This work was supported by the Ministry of Research, Technology and Higher Education, Republic of Indonesia (grant no. 27/SP-Lit/LPPM-01/Dikti/FF/V/2017).

Competing interests: None declared.

Ethical approval: The data collection instrument and methodology used in this study were approved by the ethics committee of the Faculty of Medicine, Islamic University of Indonesia (No. 08/Ka.Kom.Et/70/KE/IX/2016).

References

- 1 Statistics Indonesia (Badan Pusat Statistik – BPS), National Population and Family Planning Board (BKKBN), Ministry of Health Republic of Indonesia (Kementerian Kesehatan – Kemenkes RI), ICF International. Indonesia Demographic and Health Survey 2012. Jakarta:Indonesia BPS, BKKBN, Kemenkes RI and ICF International, 2013.
- 2 Ministry of Health Republic of Indonesia (Kementerian Kesehatan Republik Indonesia – Kemenkes RI). General guidance on healthy Indonesia program. Jakarta:Indonesia Ministry of Health, 2016.
- 3 Republic of Indonesia (Pemerintah Republik Indonesia – Pemerintah RI). The road to National Health Insurance (JKN). Jakarta: Pemerintah RI; 2012. <http://www.tnp2k.go.id> [accessed 7 January 2019].
- 4 Ministry of Health Republic of Indonesia (Kementerian Kesehatan Republik Indonesia – Kemenkes RI). SK Menteri Kesehatan RI Nomor HK.02.02/MENKES/427/2015 tentang Gerakan Masyarakat Cerdas Menggunakan Obat (GeMa CerMat). Jakarta:Kemenkes RI, 2015.
- 5 President, Republic of Indonesia (President RI). Instruksi Presiden Republik Indonesia Nomor 1 Tahun 2017: Gerakan Masyarakat Hidup Sehat. Jakarta:President RI, 2017.
- 6 World Health Organization. The role of the pharmacist in self-care and self-medication. Geneva:World Health Organization, 1998.
- 7 Pharmaceutical Society of Australia. Delivering safe and sustainable clinical services. https://www.dhhs.tas.gov.au/data/assets/pdf_file/0003/194934/Pharmaceutical_Society_of_Australia.pdf [accessed 16 September 2019].
- 8 Statistics Indonesia (Badan Pusat Statistik – BPS). Persentase penduduk yang mengobati sendiri selama sebulan terakhir menurut provinsi dan jenis obat yang digunakan, 2000–2014. <https://www.bps.go.id/linkTableDinamis/view/id/926> [accessed 27 December 2018].
- 9 Hughes C, McElroy J, Fleming G. Benefits and risks of self-medication. *Drug Saf.* 2001;24(14):1027–1037.
- 10 Ruiz M. Risks of self-medication practices. *Curr Drug Saf.* 2010;5(4): 315–323.
- 11 Setiadi A, Wibowo Y, Setiawan E, et al. Evaluation of a community-based training to promote responsible self-medication in East Java, Indonesia. *Int J Pharm Pract.* 2019;27(1):63–70.
- 12 Statistics Ngawi (Badan Pusat Statistik – BPS Kabupaten Ngawi). Kabupaten Ngawi dalam angka 2012. Ngawi:BPS Ngawi, 2012.
- 13 Smith F. Research methods in pharmacy practice. London: Pharmaceutical Press, 2002.
- 14 Tong A, Sainsbury P, Craig J. Consolidated criteria for reporting qualitative research (COREQ): a 32-item checklist for interviews and focus groups. *Int J Qual Health Care.* 2007;19(6):349–357.

- 15 Braun V, Clarke V. Using thematic analysis in psychology. *Qual Res Psychol.* 2006;3(2):77–101.
- 16 Shenton A. Strategies for ensuring trustworthiness in qualitative research projects. *Educ Inf.* 2004;22(2):63–75.
- 17 Green L, Kreuter M. *Health promotion planning: an educational and ecological approach.* Mountain View, CA:Mayfield, 2005.
- 18 Ministry of Health Republic of Indonesia (Kementerian Kesehatan Republik Indonesia – Kemenkes RI). *Peraturan Menteri Kesehatan Republik Indonesia Nomor 75 Tahun 2014: Pusat Kesehatan Masyarakat.* Jakarta:Kemenkes RI; p. 2014.
- 19 McLeroy K, Bibeau D, Steckler A, Glanz K. An ecological perspective on health promotion programs. *Health Educ Q.* 1988;15(4):351–377.
- 20 Friedman J, Cosby R, Boyko S, Hatton-Bauer J, Turnbull G. Effective teaching strategies and methods of delivery for patient education: a systematic review and practice guideline recommendations. *J Canc Educ.* 2011;26(1):12–21.
- 21 Statistics Jawa Timur (Badan Pusat Statistik – BPS Provinsi Jawa Timur). *Provinsi Jawa Timur dalam angka 2016.* Jakarta:BPS, 2016.
- 22 Setiadi A, Wibowo Y, Setiawan EI et al. *Modul pengenalan obat swamedikasi edisi 1 (2017), 1st edn.* Surabaya:PIOLK Universitas Surabaya and East Java Provincial Health Office, 2017.
- 23 Shrestha S. A conceptual model for empowerment of the female health volunteers in Nepal. *Educ Health.* 2003;16(3):318–327.
- 24 Moffatt A. *Indonesian cultural profile: An initiative of HACC Multicultural Advisory Service.* West End, QL, Australia: Diversicare; 2012. http://www.diversicare.com.au/wp-content/uploads/2015/10/Profile_Indonesian.pdf.
- 25 Simovska V, Carlson M. Health-promoting changes with children as agents: Findings from a multiple case study research. *Health Educ.* 2012;112(3):292–304.
- 26 Lehman U, Sanders D. *Community health workers: what do we know about them? The state of the evidence on programmes, activities, costs and impact on health outcome of using community health workers.* Geneva:World Health Organization, 2007.
- 27 Clausen L, Schmidt C, Aagaard-Hansen J, Reinbach H, Toft U, Bloch P. Children as visionary change agents in Danish school health promotion. *Health Promo Int.* 2018. <https://doi.org/10.1093/heapro/day049>.
- 28 World Health Organization. *Country health system profile: Indonesia.* Geneva:World Health Organization, 2007.
- 29 Wakefield M, Loken B, Hornik R. Use of mass media campaigns to change health behaviour. *Lancet.* 2010;376(9748):1261–1271.
- 30 O'Mara B. Social media, digital video and health promotion in a culturally and linguistically diverse Australia. *Health Promo Int.* 2013;28(3):466–476.




International Health



Volume 14 Number 4 July 2022 ISSN (ONLINE) 1876-3405

OXFORD
UNIVERSITY PRESS



OXFORD
ACADEMIC

International Health



RSTMH
DEDICATED TO GLOBAL HEALTH SINCE 1907

Editorial Board

Editor-in-Chief:

Professor David Molyneux

Liverpool School of Tropical Medicine, UK

Managing Editor:

Emma Williams

journals@rstmh.org

Royal Society of Tropical Medicine and Hygiene, Northumberland House, 303-306 High Holborn,
WC1V 7JZ London, UK

Associate Editors:

Olatunji Adetokunboh

South Africa

Elena Ambrosino

The Netherlands

Charles Ameh

UK

Frank Baiden

Sierra Leone

Pauline Bakibinga

Kenya

Ross Boyce

USA

Michael Coleman

UK

Wei Du

Australia

Khalifa Elmusharaf

Ireland

Caterina Fanello

UK

Rebecca Grais

France

Michael Head

UK

Derrick Ho

Hong Kong

Leena Inamdar

UK

Zohra Lassi

Australia

David Lawrence

UK and Botswana

Graeme MacLaren

Singapore

Tom Peto

UK and Thailand

Peiyuan Qui

China

Eric Tenkorang

Canada

Editorial Advisors:

Vineta Bhalla

UK

Rishi Caleyachetty

UK

David Durrheim

Australia

Stephanie Meredith

France

Editorial board:

Margaret Baker

Washington, DC, USA

William R Brieger

Baltimore, MD, USA

Paul Courtright

Moshi, Tanzania

Karen Devries

London School of Hygiene and Tropical Medicine, UK

Christopher Duggan

Boston, USA

Christopher Dye

Geneva, Switzerland

Kevin Marsh

Kilifi, Kenya

Melanie Newport

Brighton, UK

Moffat Nyirenda

Edinburgh University, UK

Watcharapong Piyaphanee

Bangkok, Thailand

Eduard Sanders

Kilifi, Kenya

Andrew Tatem

Southampton, UK

Huw Taylor

Brighton, UK

Geordie Woods

West Sussex, United Kingdom

Kaspar Wyss

Basel, Switzerland

Production Editor:

Daniel Moore

Oxford University Press



| Latest | Most Read | Most Cited |
|---|-----------|------------|
| <p data-bbox="220 1486 1357 1562">Trends in the prevalence of cognitive impairment in Chinese older adults: based on the Chinese Longitudinal Healthy Longevity Survey cohorts from 1998 to 2014</p> <hr/> <p data-bbox="220 1625 1256 1701">2030 Countdown to combating malnutrition in Burundi: comparison of proactive approaches for case detection and enrolment into treatment</p> <hr/> <p data-bbox="220 1764 1334 1881">A comparison of individual and ecological indicators of social deprivation and their association with hospital efficiency in the context of infectious diseases in two French general paediatric departments</p> <hr/> | | |

Evaluation of iron intake in preschool children in a setting with high anemia burden

Strategies to implement community training to promote responsible self-medication in Indonesia: a qualitative study of trainers



[About International Health](#)

[Editorial Board](#)

[Policies](#)

[Author Guidelines](#)

[Contact RSTMH](#)

[Purchase](#)

[Recommend to your Library](#)

[Advertising and Corporate Services](#)

[Journals Career Network](#)

[RSTMH Blog](#)

[Facebook](#)

[Twitter](#)

[YouTube](#)

[LinkedIn](#)

International Health

Online ISSN 1876-3405

Print ISSN 1876-3413

Copyright © 2020 Royal Society of Tropical Medicine and Hygiene

[About Us](#)

[Contact Us](#)

[Careers](#)

[Help](#)

[Access & Purchase](#)

[Rights & Permissions](#)

[Open Access](#)

Connect

[Join Our Mailing List](#)

[OUPblog](#)

[Twitter](#)

[Facebook](#)

[YouTube](#)

[Tumblr](#)

Resources

[Authors](#)

[Librarians](#)

[Societies](#)

[Sponsors & Advertisers](#)

[Press & Media](#)

[Agents](#)

Explore

[Shop OUP Academic](#)

[Oxford Dictionaries](#)

[Oxford Index](#)

[Epigeum](#)

[OUP Worldwide](#)

[University of Oxford](#)

Oxford University Press is a department of the University of Oxford. It furthers the University's objective of excellence in research, scholarship, and education by publishing worldwide

OXFORD
UNIVERSITY PRESS

Copyright © 2020 Oxford University Press

[Legal Notice](#)

[Site Map](#)

[Accessibility](#)

[Cookie Policy](#)

[Privacy Policy](#)

[Get Adobe Reader](#)

Browse issues

Year Issue [Browse by volume](#)[Browse supplements](#)

Volume 14, Issue 4

July 2022

[Cover image](#)

EISSN 1876-3405

[< Previous](#)

Volume 14, Issue 4, July 2022

ORIGINAL ARTICLES

Metabolic syndrome and related risk factors among adults in the northern West Bank, a cross-sectional study

[Basma Damiri, Luna Badran, Deya Safadi, Ahmad Sawalha, Younis Yasin ...](#)*International Health*, Volume 14, Issue 4, July 2022, Pages 339–345, <https://doi.org/10.1093/inthealth/ihz093>[Abstract](#) ▼ [View article](#)

Socioeconomic status and education level are associated with dyslipidemia in adults not taking lipid-lowering medication: a population-based study

[Luçandra R Espírito Santo, Thaís O Faria, Carla Silvana O Silva, Lorena A Xavier, Vivianne C Reis ...](#)*International Health*, Volume 14, Issue 4, July 2022, Pages 346–353, <https://doi.org/10.1093/inthealth/ihz089>[Abstract](#) ▼ [View article](#) [Supplementary data](#)

Investigating the prevalence of and factors associated with depressive symptoms among urban and semi-urban school adolescents in Bangladesh: a pilot study

Afifa Anjum, Sahadat Hossain, Tajuddin Sikder, Md Elias Uddin, Dewan Abdur Rahim

International Health, Volume 14, Issue 4, July 2022, Pages 354–362, <https://doi.org/10.1093/inthealth/ihz092>

[Abstract](#) ▼ [View article](#)

Social determinants of violence against women in Panama: results from population-based cross-sectional studies and a femicide registry

Ana Santamaría, Carmen Gerald, Fermina Chamorro, Víctor Herrera, Haydee Flores ...

International Health, Volume 14, Issue 4, July 2022, Pages 363–372, <https://doi.org/10.1093/inthealth/ihz116>

[Abstract](#) ▼ [View article](#) [Supplementary data](#)

Healthcare-seeking behaviour, barriers to care and predictors of symptom improvement among patients with cardiovascular disease in northern Tanzania

Julian T Hertz, Francis M Sakita, Godfrey L Kweka, Zak Loring, Nathan M Thielman ...

International Health, Volume 14, Issue 4, July 2022, Pages 373–380, <https://doi.org/10.1093/inthealth/ihz095>

[Abstract](#) ▼ [View article](#)

Household structure, maternal characteristics and children's stunting in sub-Saharan Africa: evidence from 35 countries

Sanni Yaya, Olanrewaju Oladimeji, Emmanuel Kolawole Odusina, Ghose Bishwajit

International Health, Volume 14, Issue 4, July 2022, Pages 381–389, <https://doi.org/10.1093/inthealth/ihz105>

[Abstract](#) ▼ [View article](#)

Prevalence and determinants of poor glycaemic control: a cross-sectional study among Moroccan type 2 diabetes patients

Ahmed Chetoui, Kamal Kaoutar, Soufiane Elmoussaoui, Kaltoum Boutahar, Abdesslam El Kardoudi ...

International Health, Volume 14, Issue 4, July 2022, Pages 390–397, <https://doi.org/10.1093/inthealth/ihz107>

[Abstract](#) ▼ [View article](#)

Strategies to implement community training to promote responsible self-medication in Indonesia: a qualitative study of trainers

Adji P Setiadi, Yosi I Wibowo, Eko Setiawan, Ika Mulyono, Susilo A Wardhani ...

International Health, Volume 14, Issue 4, July 2022, Pages 398–404, <https://doi.org/10.1093/inthealth/ihz115>

[Abstract](#) ▼ [View article](#)

A comparison of individual and ecological indicators of social deprivation and their association with hospital efficiency in the context of infectious diseases in two French general paediatric departments

Aphaia Roussel, Albert Faye, Alain Lefevre-Utile, Loic De Pontual, Karine Chevreul ...

International Health, Volume 14, Issue 4, July 2022, Pages 405–412, <https://doi.org/10.1093/inthealth/ihz102>

[Abstract](#) ▼ [View article](#)

2030 Countdown to combating malnutrition in Burundi: comparison of proactive approaches for case detection and enrolment into treatment

Emmanuel Nene Odjidja, Cathryn Christensen, Ghislaine Gatasi, Sonia Hakizimana, Honorine Murorunkwere ...

International Health, Volume 14, Issue 4, July 2022, Pages 413–420, <https://doi.org/10.1093/inthealth/ihz119>

[Abstract](#) ▼ [View article](#)

Inter-relationships among key reproductive health indicators in sub-Saharan Africa focusing on the central role of maternal literacy

Mulu Abraha Woldegiorgis, Denny Meyer, Janet E Hiller, Wubegzier Mekonnen, Jahar Bhowmik

International Health, Volume 14, Issue 4, July 2022, Pages 421–433, <https://doi.org/10.1093/inthealth/ihz117>

[Abstract](#) ▼ [View article](#)

A qualitative narrative review of protocols for women's health on short-term medical missions in Latin America and the Caribbean

Christopher Dainton, Charlene H Chu

International Health, Volume 14, Issue 4, July 2022, Pages 434–441, <https://doi.org/10.1093/inthealth/ihz109>

[Abstract](#) ▼ [View article](#)

Trends in type 1 diabetes diagnosis in Ghana

[Osei Sarfo-Kantanka](#), [Michael Asamoah-Boaheng](#), [Joshua Arthur](#), [Martin Agyei](#), [Nana Ama Barnes](#) ...

International Health, Volume 14, Issue 4, July 2022, Pages 442–446, <https://doi.org/10.1093/inthealth/ihz096>

[Abstract](#) ▼ [View article](#)

SHORT COMMUNICATIONS

Delivery of home-based post-partum contraception in rural Guatemalan women: feasibility, recruitment and retention in a cluster-randomized trial

[Margo S Harrison](#), [Saskia Bunge-Montes](#), [Claudia Rivera](#), [Andrea Jimenez-Zambrano](#), [Gretchen Heinrichs](#) ...

International Health, Volume 14, Issue 4, July 2022, Pages 447–449, <https://doi.org/10.1093/inthealth/ihz098>

[Abstract](#) ▼ [View article](#)

Evaluation of iron intake in preschool children in a setting with high anemia burden

[Maria G Dunn](#), [Ryan M Close](#), [Steven G McKee](#), [Ramona Cordero](#), [Ingrid F Japa](#) ...

International Health, Volume 14, Issue 4, July 2022, Pages 450–451, <https://doi.org/10.1093/inthealth/ihz106>

[Abstract](#) ▼ [View article](#)



[Front Matter](#)



[Table of Contents](#)



[Editorial Board](#)

[All issues](#)



International Health

Country [United Kingdom](#) - SIR Ranking of United Kingdom

24

H Index

Subject Area and Category

[Medicine](#)

[Medicine \(miscellaneous\)](#)

[Public Health, Environmental and Occupational Health](#)

[Social Sciences](#)

[Health \(social science\)](#)

Publisher

[Oxford University Press](#)

Publication type

[Journals](#)

ISSN

18763413


Coverage

2009-ongoing



[Join the conversation about this journal](#)


Quartiles



SJR




Citations per document





Total Cites

Self-Cites






External Cites per Doc Cites per Doc +




% International Collaboration +

Citable documents Non-citable documents +



Cited documents Uncited documents +



International Health

Q2 Health (social science) best quartile

SJR 2018 0.71

powered by scimagojr.com

← Show this widget in your own website

Just copy the code below and paste within your html code:

```
<a href="https://www.scimag
```

Leave a comment

Name

Email

(will not be published)

I'm not a robot reCAPTCHA
Privacy - Terms

Submit

The users of Scimago Journal & Country Rank have the possibility to dialogue through comments linked to a specific journal. The purpose is to have a forum in which general doubts about the processes of publication in the journal, experiences and other issues derived from the publication of papers are resolved. For topics on particular articles, maintain the dialogue through the usual channels with your editor.

Developed by:



Powered by:



Follow us on @ScimagoJR

Scimago Lab, Copyright 2007-2019. Data Source: Scopus®





Source details

International Health

Scopus coverage years: from 2009 to Present

Publisher: Oxford University Press

ISSN: 1876-3413 E-ISSN: 1876-3405

Subject area: Social Sciences: Health (social science) Medicine: Public Health, Environmental and Occupational Health

Source type: Journal

[View all documents >](#)

[Set document alert](#)

[Save to source list](#) [Source Homepage](#)

[CiteScore](#) [CiteScore rank & trend](#) [Scopus content coverage](#)

CiteScore 2021 ▼

$$3.9 = \frac{1,337 \text{ Citations 2018 - 2021}}{339 \text{ Documents 2018 - 2021}}$$

Calculated on 05 May, 2022

CiteScoreTracker 2022 ⓘ

$$3.3 = \frac{1,010 \text{ Citations to date}}{305 \text{ Documents to date}}$$

Last updated on 06 June, 2022 • Updated monthly

CiteScore rank 2021 ⓘ

| Category | Rank | Percentile |
|--|----------|------------|
| Social Sciences | | |
| Health (social science) | #72/323 | 77th |
| Medicine | | |
| Public Health, Environmental and Occupational Health | #174/562 | 69th |

About Scopus

[What is Scopus](#)

[Content coverage](#)

[Scopus blog](#)

[Scopus API](#)

[Privacy matters](#)

Language

[日本語版を表示する](#)

[查看简体中文版本](#)

[查看繁體中文版本](#)

[Просмотр версии на русском языке](#)

Customer Service

[Help](#)

[Tutorials](#)

[Contact us](#)

ELSEVIER

[Terms and conditions ↗](#) [Privacy policy ↗](#)

Copyright © [Elsevier B.V ↗](#). All rights reserved. Scopus® is a registered trademark of Elsevier B.V.

We use cookies to help provide and enhance our service and tailor content. By continuing, you agree to the [use of cookies ↗](#).

