


<https://scialert.net/archivedetails.php?issn=1682-3974&issueno=114>

Inbox (1,761) - poppy_hardjo@... x Asian Journal of Plant Sciences - x Asian Network for Scientific Info: x +

Not secure | scialert.net/archivedetails.php?issn=1682-3974&issueno=114

Asian Journal of Plant Sciences

Publisher: Asian Network for Scientific Information



Asian Journal of Plant Sciences is a high quality scientific journal publishes the original research in all areas of plant science and botany. Scope of the journal includes: Plant biotechnology, plant cell and tissue culture, plant gene transfer, development, growth regulation, molecular cell biology and genetics, signal transduction, photosynthesis, pathogen resistance, nutrition, water relations and gas exchange, symbiosis, stress physiology, population genetics, ecology and molecular systematic.

Asian Journal of Plant Sciences now accepting new submissions. Submit your best paper via [online submission system](#).

eISSN: 1812-5697
pISSN: 1682-3974
Editor-in-Chief: **Rajarshi Kumar Gaur**

Volume 18, Number 2, 2019

Effect of Nitrogen and Phosphorus on the Growth and Redox Homeostasis of Salt-stressed Mustard Plants
Naveed Gulzar, Mohammad Yaseen Mir and Saima Hamid
Asian Journal of Plant Sciences Volume 18, Number 2, 52-59, 2019
[\[Abstract\]](#) [\[Fulltext PDF\]](#) [\[Fulltext HTML\]](#) [\[XML: Abstract + References\]](#) [\[References\]](#)

Improvement of Blast Resistance and Yield Productivity in Rice (*Oryza sativa* L.) Using Tissue Culture and Combining Ability
Mohammed I. Al-Daej, Adel A. Rezk and Mohamed M. El-Malky
Asian Journal of Plant Sciences Volume 18, Number 2, 60-74, 2019
[\[Abstract\]](#) [\[Fulltext PDF\]](#) [\[Fulltext HTML\]](#) [\[XML: Abstract + References\]](#) [\[References\]](#)

Impact of Habitat Heterogeneity on Growth Dynamics and Physiological Responses of *Dipterygium glaucum*
Neelam Sherwani and Sardar Abulfazal Farooq
Asian Journal of Plant Sciences Volume 18, Number 2, 75-84, 2019
[\[Abstract\]](#) [\[Fulltext PDF\]](#) [\[Fulltext HTML\]](#) [\[XML: Abstract + References\]](#) [\[References\]](#)


Shoots Culture of *Gynura procumbens* (Lour.) Merr. in Balloon-Type Bubble-bioreactor Influenced by Sucrose Concentration and Inoculum Density
Ira Nailas Saadah, Afinda Novi Kristanti, Popy Hartatie Hardjo and Yosephine Sri Wulan Manuhara
Asian Journal of Plant Sciences Volume 18, Number 2, 85-90, 2019
[\[Abstract\]](#) [\[Fulltext PDF\]](#) [\[Fulltext HTML\]](#) [\[XML: Abstract + References\]](#) [\[References\]](#)

Induction of Banana Autotetraploids "Klutuk Sukun" and their Reproductive Function for Producing Triploid Hybrids
Yuyu Suryasari Poerba, Diyah Martanti, Tri Handayani and Witjaksono
Asian Journal of Plant Sciences Volume 18, Number 2, 91-100, 2019
[\[Abstract\]](#) [\[Fulltext PDF\]](#) [\[Fulltext HTML\]](#) [\[XML: Abstract + References\]](#) [\[References\]](#)

Navigation


- Online First
- Current Issue
- Previous Issues
- Editorial Board
- Submit a Manuscript
- Guide to Authors
- Article Processing Charges
- Subscribe to E-alerts

Asian Journal of Plant Sciences
Agronomy and Crop Science
Q4
best quartile
SJR 2018
0.18
powered by scimagojr.com



Indexed In

- AGRIIS
- ASCI-Database
- Asian Digital Library
- Cambridge Scientific Abstract
- Chemical Abstract Services
- ISI Web of Knowledge
- SCIMAGO
- Thomson ISI



Asian Journal of Plant Sciences

Publisher: Asian Network for Scientific Information



Asian Journal of Plant Sciences is a high quality scientific journal publishes the original research in all areas of plant science and botany. Scope of the journal includes: Plant biotechnology, plant cell and tissue culture, plant gene transfer, development, growth regulation, molecular cell biology and genetics, signal transduction, photosynthesis, pathogen resistance, nutrition, water relations and gas exchange, symbiosis, stress physiology, population genetics, ecology and molecular systematic.

Asian Journal of Plant Sciences now accepting new submissions. Submit your best paper via [online submission system](#).

eISSN: 1812-5697
pISSN: 1682-3974

Editor-in-Chief: [Rajarshi Kumar Gaur](#)

Editor-in-Chief



Rajarshi Kumar Gaur
Mody Institute of Technology and Science, India

REGIONAL EDITORS



Guno Sindhu Chakraborty
Parul Institute of Pharmacy & Research,
India

ASSOCIATE EDITORS



Hala Mohamed Safwat El-Bassiouny
National Research Center, Egypt



Mohammad Moneruzzaman Khandaker
Universiti Sultan Zaimal Abidin, Malaysia



Mirza Hasanuzzaman
Sher-e-Bangla Agricultural University,
Bangladesh



Mohd Rafiq Wani
Abdul Azad Memorial Degree
College, India



Mitra Noori
Arak University, Iran



Mitalkumar Jayantilal Kaneria
Saurashtra University, India



Hongxin Cao
Institute of Agricultural Resources and
Environment, China



Shah Fahad
Huazhong University of Science and
Technology, China




Vinay Kumar
Modern College, India

Navigation

- Online First
- Current Issue
- Previous Issues
- Editorial Board
- Submit a Manuscript
- Guide to Authors
- Article Processing Charges
- Subscribe to E-alerts

Asian Journal of Plant Sciences



Indexed In

- AGRIS
- ASCI Database
- Asian Digital Library
- Cambridge Scientific Abstract
- Chemical Abstract Services
- ISI Web of Knowledge
- SCIMAGO
- Thomson ISI



James Nte Nwite
Ebonyi State University, Abakaliki,
Nigeria



V. Vadivel
V.O. Chidambaram College, India



Osama H. Sayed
Jazan University, Saudi Arabia



Ekta Menghani
JECRC University, India



Mohd Sayeed Akhtar
Gandhi Fauz-E-Aam College,
Shahjahanpur, India



khalid khalid
National Research Centre, Egypt

TECHNICAL EDITORS



Natchimuthu Karmegam
Government Arts College, India



Irina Neta Gostin
Alexandru Ioan Cuza University of Iasi,
Romania



Kamuran Aktas
Celal Bayar University, Turkey



Ravindran Chandran
Agricultural College and Research
Institute, Kilikulam, India



Md. Shafiqul Islam
Bangladesh Agricultural University,
Bangladesh



Madhu Kamle
Central Institute for Subtropical
Horticulture, Lucknow, India



G.K. Ramegowda
University of Horticultural Sciences, India



Atul Bhargava
Amity University, India



Qurban Ali Panhwar
Universiti Putra Malaysia, Malaysia



Tanweer Azam
Haramiabo Agricultural College, Eritrea



Devendra Kumar Payasi
Jawaharlal Nehru Agricultural University,
India



Amal Mohamed Eliwa Abdel-Hamid
Ain Shams University, Egypt



Sarika Gupta
Banasthali Vidyapeeth, India



Ilknur Ak
Canakkale Onsekiz Mart University,
Turkey



Zehra Ekin
Van Yuzuncu Yil University, Turkey



Bhupinder Dhir
Indira Gandhi National Open University,
Delhi, India



Md. Amir Hossain
Bangladesh Agricultural University,
Bangladesh



Ugur Cakicioglu
Munzur University, Turkey



Mandaliya Viralkumar Bhikkhalal
Gujarat National Law University, India



Keshav Kant Sahu
Pt. Ravishankar Shukla University, India



Md. Rostom Ali
Bangladesh Agricultural University,
Bangladesh



Hameda El Sayed Ahmed El Sayed
Umm Al-Qura University, Saudi Arabia



**Magda Magdy Mohamed Fahmy
Abd El-Salam**
Alexandria University, Egypt



Engin Eroglu
Duzce University, Turkey



Ebru Yuce Babacan
Munzur University, Turkey



Demir Kok
Namik Kemal University, Turkey



Rakesh Kumar
Vjay Singh Pathik Govt. (PG) College,
India



**Riad Sedki Riad Ibrahiem El-
Mohamedy**
National Research Center, Egypt



Udhav Narba Bhale
Arts, Science and Commerce College
Naldurg, India



Ahmad AlSayed Mohamad Omar
University of Florida, USA



Burhan Arslan
Namik Kemal University, Turkey



Canan Ozdemir
Celal Bayar University, Turkey



Thomas Edward Smith
Ave Maria University, USA



**Amany Mohamed Abd El-Fatah
Haroon**
National Institute Oceanography and
Fisheries, Cairo, Egypt



Turan Karadeniz
Karadeniz Technical University, Turkey



**Saad Farouk Mohamed Hussien
Gadalla**
Mansoura University, Egypt



Ram Cahndra Suthar
Pramukh Swami Science & H.D.Patel
Arts College, India



Kemal Yildiz
Celal Bayar University, Turkey



Rafiq Ahmad Lone
University of Kashmir, Srinagar Jammu
and Kashmir



Kiomars Sefidi
University of Mohaghegh Ardabili, Iran



Kala R.G.
Rubber Research Institute of India, India



Chandrasekaran Rajasekaran
VIT University, India



**Preeya Puangsomlee
Wangsomuk**
University of East Anglia, Norwich, UK



Mohammad Israil Ansari
University of Lucknow, India



Hayet Lahmar Edziri
High institute of Biotechnology,Tunesia



**Dina Zein El Abdin Abdel Kader
Saleh**
Taibah University, Saudi Arabia



Nurdan Tuna Gunes
Ankara University, Turkey



L.P. Yadava
Central Institute for Subtropical
Horticulture, India



Mahmoud Hussien Abou Delf
National Research Centre, Egypt

[Home](#) · [Journals](#) · [For Authors](#) · [For Subscribers](#) · [ASCI](#)

© Science Alert. All Rights Reserved

Search SciAlert website











Asian Journal of Plant Sciences

Country	Pakistan - SJR Ranking of Pakistan
Subject Area and Category	Agricultural and Biological Sciences Agronomy and Crop Science Plant Science
Publisher	Asian Network for Scientific Information
Publication type	Journals
ISSN	16823974, 18125697
Coverage	2006-ongoing
Scope	Information not localized
	Homepage
	How to publish in this journal
	Contact
	Join the conversation about this journal

24

H Index





Asian Journal of Plant Sciences

Q4 Agronomy and Crop Science
best quartile

SJR 2018
0.18

powered by scimagojr.com

← Show this widget in your own website

Just copy the code below and paste within your html code:

```
<a href="https://www.scimagojr.com" style="border: 1px solid #ccc; padding: 2px 5px; display: inline-block;">https://www.scimagojr.com
```



Asian Journal of Plant Sciences

ISSN 1682-3974

science
alert

ANSI*net*
an open access publisher
<http://ansinet.com>



Research Article

Shoots Culture of *Gynura procumbens* (Lour.) Merr. in Balloon-Type Bubble-bioreactor Influenced by Sucrose Concentration and Inoculums Density

¹Ira Nailas Saadah, ²Alfinda Novi Kristanti, ³Popy Hartatie Hardjo and ¹Yosephine Sri Wulan Manuhara

¹Department of Biology, Faculty of Science and Technology, Airlangga University, Surabaya, Indonesia

²Department of Chemistry, Faculty of Science and Technology, Airlangga University, Surabaya, Indonesia

³Faculty of Biotechnology, Surabaya University, Surabaya, Indonesia

Abstract

Background and objectives: *Gynura procumbens* has been widely used as a traditional medicine in Indonesia. Micropropagation of this plant was important to fulfill the need of pharmaceutical industry. The aims of this study were to increasing the number of *G. procumbens* shoots in balloon-type bubble bioreactor influenced by sucrose concentration and inoculums density. **Materials and Methods:** Nodal segments of *G. procumbens* were used as explants which were grown in balloon type bubble bioreactor in MS liquid medium supplemented with 2 mg L⁻¹ IAA and 4 mg L⁻¹ BAP. Shoot culture were conducted under treatment of various concentration of sucrose (10, 30 and 50 g L⁻¹) and various inoculums density (5, 10 and 15 explants). **Results:** The highest multiplication of shoots and growth index were obtained at 50 g L⁻¹ sucrose treatment and 5 explants as initial inoculums; there are 9.36 shoots/explants and 13.8 shoots/explants, respectively. **Conclusion:** The study found that *G. procumbens* shoots cultured in balloon-type bubble bioreactors increased shoots multiplication significantly.

Key words: Balloon-type, bubble bioreactor, *Gynura procumbens*, inoculums density, sucrose, shoot culture

Citation: Ira Nailas Saadah, Alfinda Novi Kristanti, Popy Hartatie Hardjo and Yosephine Sri Wulan Manuhara, 2019. Shoots culture of *Gynura procumbens* (Lour.) Merr. in balloon-type bubble-bioreactor influenced by sucrose concentration and inoculums density. Asian J. Plant Sci., 18: 85-90.

Corresponding Author: Yosephine Sri Wulan Manuhara, Department of Biology, Faculty of Science and Technology, Airlangga University, Surabaya, Indonesia Tel: 62817484872

Copyright: © 2019 Ira Nailas Saadah *et al.* This is an open access article distributed under the terms of the creative commons attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

Competing Interest: The authors have declared that no competing interest exists.

Data Availability: All relevant data are within the paper and its supporting information files.

INTRODUCTION

Gynura procumbens is a plant that has been traditionally used as an herbal medicine and scientifically proven its extract benefits to treat various health problems such as antimicrobial¹⁻³, antioxidants⁴⁻⁷ and anticancer⁸⁻¹⁰. Micropropagation of this plant was important to fulfill the need of pharmaceutical industry. Tissue culture is the best solution for growing plants in large quantities and faster than conventional cultivation. In recent years, diverse bioreactor design has been used for cultivation of differentiated plant organ cultures^{11,12}. One of the bioreactor that was used to growing cell and organ culture is balloon-type bubble bioreactor. The application of balloon-type bubble bioreactors has been widely developed for various plant cell and organ culture such as *Morinda citrifolia* (L.) cell culture¹³, *Eurycoma longifolia* adventitious root culture¹⁴, *Panax ginseng* CA Meyer hairy root culture¹⁰, *Cyclopia genistoides* (L.) Vent. shoot culture¹⁵, *Hypericum perforatum* shoot culture¹⁶, *Aloe barbadensis*¹⁷, *Dendrobium candidum* Wall ex Lindl. protocorm culture¹⁸, *G. procumbens* adventitious roots culture¹⁹.

Sucrose is a source of energy to build cell organelles that make up plant organs affecting the growth of plant organ cultures and the production of secondary metabolite. Increased levels of sucrose up to 6% could increase biomass of *Talinum paniculatum* hairy root culture²⁰, 5% sucrose also increased biomass of *G. procumbens* adventitious roots that were grown in liquid culture with agitation²¹ and in a temporary immersion bioreactor²².

The density of explants in bioreactor affected the growth and accumulation of secondary metabolite that was showed in production of ginsenosides from adventitious root suspension cultures of ginseng²³, adventitious roots culture of *Eleutherococcus koreanum*²⁴ and microtuber production of potato in modified nutrient spray bioreactor system²⁵. During the process of culture, explants produced CO₂, which was the result of catabolism process. Increasing the amount of explants will multiply the CO₂ content in the medium, so it affects the growth of explants in the bioreactor²⁴.

Shoot culture of *G. procumbens* has been done in solid medium and liquid culture in temporary immersion system; the highest growth of shoots was achieved in MS (Murashige and Skoog) solid medium supplemented with kinetin and combination of IAA (indol acetic acid) and BA (benzyl adenine) but in temporary immersion system, number of shoots lower than in solid medium²⁶. This study aimed to increase the growth of *G. procumbens* shoots in balloon-type bubble bioreactor (BTBB) influenced by sucrose concentration and inoculums density.

MATERIALS AND METHODS

Gynura procumbens (Lours.) Merr. was obtained from the Botanical Garden Purwodadi, Pasuruan, East Java, Indonesia. This research was conduct at January, 2017 until September, 2018. Stem nodes of *G. procumbens* were used as an explant. Stem nodes sterilization were performed using Clorox 1% (Bayclin, Johnson, Indonesia) for 10 min and rinsed with sterile distilled water three times. The stem nodes were grown in MS²⁷ supplemented with 30 g L⁻¹ sucrose, 7 g L⁻¹ agar and combination of 2 mg L⁻¹ IAA and 4 mg L⁻¹ BAP. The medium used has been adjusted to pH 5.8 and autoclaved for 15 min (1 atm and 121°C). Cultures were maintained at room temperature (25±3°C) with fluorescent lighting (General electric cool white fluorescent tubes) 1900 lux. On the 3rd day, explants was transferred into MS liquid medium using the same combination of plant growth regulator as solid medium and cultured in a shaker incubator (110 rpm) for 3 days. Explants were ready for use to culture in balloon-type bubble bioreactors.

Shoots culture in bioreactor: Seven shoots as an initial inoculum were grown on 1 L bioreactor using 400 mL of MS liquid medium supplemented with 2 mg L⁻¹ IAA and 4 mg L⁻¹ BAP with each sucrose concentrations of 10, 30 and 50 g L⁻¹. The inoculums density treatment were done using 5, 10 and 15 shoots explants with 50 g L⁻¹ sucrose in each bioreactor. The bioreactor was equipped with an explants retaining wire (net) to reduce hyperhydricity that can interfere explants growth by causing abnormal explants growth²⁸. Aeration rate was adjusted to 0.2 vvm. Cultures were maintained by fluorescent light (General electric cool white fluorescent tubes) continuously at 1900 lux and incubation temperature was 25±3°C for 4 weeks.

Data were recorded after 4 week of culture period including total fresh weight, total dry weight, number of shoots per explants, number of leaves per explants and growth index. Dry weight was obtained from the oven-dried shoots at 60°C for 48 h. The pH value and the medium conductivity were measured every week. The growth index of *G. procumbens* shoots was calculated using the following equation²⁹:

$$\text{Growth index} = \frac{\text{Final dry weight of planlet} - \text{dry weight of initial inoculum}}{\text{Dry weight of initial inoculum}}$$

Statistical analysis: Biomass of shoots (fresh weight and dry weight), number of shoots, number of leaves and length of shoots data were analyzed using statistical software program (SPSS 19). Each mean value represented the replicate of two

determinations were analyzed using one-way ANOVA test ($p < 0.05$). To determine the significant difference between treatments, Duncan's test was done.

RESULTS

Effect of sucrose concentration on shoots multiplication:

The highest multiplication of *G. procumbens* shoots in balloon type bubble-bioreactor was obtained in medium supplemented with sucrose 50 g L^{-1} ; these results were powered by data of number of shoots/explants, number of leaves, total fresh weight, dry weight and growth index (Table 1). The higher of sucrose concentration in medium, more the number of shoots per explants formed. The highest length of shoots were achieved by treatment of 10 g L^{-1} sucrose followed by sucrose treatment of 50 and 30 g L^{-1} , respectively. The length of shoot was almost the same in the treatment of 10 and 50 g L^{-1} sucrose. Morphology of *G. procumbens* shoots in various concentration of sucrose was showed at Fig. 1a-c. Although the number of shoots and number of leaves in the treatment of 10 and 30 g L^{-1} sucrose were not significantly different, the length of shoots produced was significantly different.

Effect of initial inoculums density on shoots multiplication:

The initial inoculums density affected the growth of *G. procumbens* shoots in balloon-type bubble bioreactors (Fig. 1d-e). The fastest shoot multiplication was obtained in the treatment of inoculums density 5 explants, that produced 13.8 shoots/explants and number of leaves

272.2 but length of shoots were not significantly different with the other treatments. Fresh weight and dry weight increased along with the increasing inoculums density planted. The highest growth index was achieved at the treatment of inoculums density 5 explants but the highest length of shoots was obtained in inoculums density 15 explants, although based on statistical analysis was not significantly different (Table 2).

Physical condition of medium: The physical condition of medium during shoots culture in balloon type bubble bioreactors was showed by pH values and electrical conductivity. The pH of the medium had a tendency to decrease during 4th week culture period (Fig. 2). In the treatment of various sucrose concentrations, the decrease of pH at the end of culture was not significantly different, that was 4.6-4.8 but in the treatment of 5 explants as inoculums density, there was a significant decrease of pH at the end of culture, which was 4. The decrease in pH was most prevalent in the inoculums density of 15 explants. The available nutrition in the medium can be utilized for plant growth. This was supported by the pH and electrical conductivity of the medium that decreased gradually from week to week. In the treatment of 5 explants, the electrical conductivity of medium increased from week to week although it is not significant (Fig. 3).

The electrical conductivity of the medium decreased until the end of the 4th week of culture, except in the initial inoculums density of 5 explants. In the treatment of 5 explants, the medium conductivity showed an increase but not significant (Fig. 3).




Fig. 1(a-f): Shoot culture of *G. procumbens* in BTBB, (a) Sucrose treatment 10 g L^{-1} , (b) 30 g L^{-1} , (c) 50 g L^{-1} and inoculums density (d) 5 explants, (e) 10 explants and (f) 15 explants

Bars: 1.4 cm

Table 1: Effect of sucrose concentrations on multiplication of *G. procumbens* shoots culture in balloon type bubble-bioreactor

Sucrose concentration (g L ⁻¹)	No. of shoot/explants	No. of leaves	Length of shoots (cm)	Total FW (g)	Total DW (g)	Growth index
10	4.00±1.57 ^b	151.21±49.71 ^{ab}	4.34±1.10 ^a	16.85	0.46	0.93
30	5.64±3.27 ^b	118.36±68.32 ^{ab}	3.15±1.16 ^b	22.94	1.22	4.10
50	9.36±5.050 ^a	195.50±71.5 ^a	4.03±0.95 ^{ab}	26.95	2.10	7.80

Mean values within a column followed by the same letters are not significantly different at p = 0.05 according to Duncan's Multiple Range test

Table 2: Effect of inoculums density on multiplication of *G. procumbens* shoots cultured in balloon type bubble-bioreactor

Inoculums density (explants)	No. of shoot/explants	No. of leaves	Length of shoots (cm)	Total FW (g)	Total DW (g)	Growth index
5	13.80±2.39 ^a	272.20±109.94 ^a	3.94±0.40 ^a	17.79	0.94	12.80
10	7.60±3.27 ^b	102.80±24.68 ^c	3.89±0.49 ^a	27.39	2.23	6.60
15	6.67±3.85 ^b	178.33±73.17 ^b	4.94±1.23 ^a	49.09	3.13	5.60

Mean values within a column followed by the same letters are not significantly different at p = 0.05 according to Duncan's Multiple Range test




Fig. 2: The pH of medium during *G. procumbens* shoot culture period in balloon-type bubble-bioreactor

S10: 10 g L⁻¹ sucrose, S30: 30 g L⁻¹ sucrose, S50: 50 g L⁻¹ sucrose, T05: 5 explants, T10: 10 explants, T15: 15 explants




Fig. 3: Electrical conductivity medium during *G. procumbens* shoot culture period in balloon type bubble-bioreactors

S10: 10 g L⁻¹ sucrose, S30: 30 g L⁻¹ sucrose, S50: 50 g L⁻¹ sucrose, T05: 5 explants, T10: 10 explants, T15: 15 explants

DISCUSSION

The higher sucrose concentration supplemented in the culture medium, the higher carbon source was available in the medium, so it increased the multiplication of shoots. Sucrose will be hydrolyzed into free glucose and fructose by invertase enzyme before being used³⁰. Some of the sucrose in the tissue culture medium is hydrolyzed into free fructose and glucose during the process of sterilization³¹. Hydrolyzed

sucrose is easier used by the plant for the process of cell metabolism and cell organelles builders. Sucrose also plays an important role in the formation of secondary metabolite compounds used as medicinal ingredients³². In this study higher sucrose concentration in medium could enhance shoot multiplication. Optimum results were also obtained at higher sucrose addition in *A. barbadensis* shoot culture¹⁷, *Talinum paniculatum* adventitious root²⁰ and *Dendrobium candidum* protocorm¹⁸.

The number of shoots increased along with the concentration level of sucrose. That was also reported in *A. barbadensis* shoot culture¹⁷. It indicated that the availability of high sucrose in the medium is utilized well for the growth process. Sucrose also affected the growth index of *G. procumbens* shoots in this study. Sucrose not only acts as a carbon source for build plant organs but also has a function as osmotic control of plant tissues³⁰. The difference in concentration gradient between plant cells will affect the polarity of plant cells that will affect the morphogenesis process and the growth index of the plant. However, the concentration gradient difference is not the only factor affecting the polarity process that affects plant morphogenesis. There are other factors that affect the process of plant morphogenesis, such as genetic factors of plants³².

The number of shoots obtained was getting lower along with the higher inoculums density planted. The highest number of leaves were obtained at the treatment of initial inoculum of 5 explants. The more number of initial inoculums, the tighter competition to get nutrition and space to grow. The tight competition affected the growth of plants. It was also reported on the production of potato microtubers²⁵. In this study the best shoot height was obtained at the density treatment of 15 explants. This phenomenon proved that more explants planted, the plants will compete to get the light so that trigger the plant to grow higher towards the light to support the process of photosynthesis. But, the

treatment of 5 explants and supplementation of sucrose 50 g L⁻¹ was the best treatment to obtain the high multiplication of shoots (13.8 shoots/explants and 9.38 shoots/explants). This result showed that *G. procumbens* shoots cultured in balloon-type bubble-bioreactor increased multiplication of shoots better than in solid medium or in temporary immersion system. In solid medium, maximum number of shoots was 7 shoots/explants, whereas in temporary immersion system was 8 shoots/explant²⁶.

The total fresh weight and total dry weight increased along with the high inoculums density but the growth index was decreased. Cui *et al.*¹⁸ reported that fresh weight and dry weight were higher along with the higher inoculums density grown in bioreactors. Lee *et al.*²⁴ also reported that fresh weight and dry weight obtained were higher and the growth index obtained was lower along with the high inoculums density on the product ion of *Eleutherococcus koreanum* Nakai root in bioreactors. In this study, the best dry weight was obtained in the initial inoculums treatment of 15 explants. This proved that the plants formed the cell mass well, so the percentage of moisture content in the cell was the lowest compared to other treatments.

CONCLUSION

From the results obtained, it can be inferred that supplementation of 50 g L⁻¹ sucrose produce the highest number of shoots and growth index of *G. procumbens* shoot culture, whereas the highest number of shoot and growth index was obtained at inoculums density 5 explants.

SIGNIFICANCE STATEMENT

This study discovered the shoot multiplication of *G. procumbens* in balloon-type bubble bioreactor can increased significantly, so that this method can be beneficial to produce biomass in large scale. This study will help the researchers to uncover the critical areas of shoot multiplication in bioreactor that many researchers were not able to explore. Thus a new method to speed up micropropagation of this plant maybe arrived at to support the need of pharmaceutical industry.

ACKNOWLEDGMENT

This study was supported by Mandatory Research Grant from Airlangga University, Surabaya, Indonesia with grant No. 1922/UN3.1.8/LT/2018.

REFERENCES

1. Jarikasem, S., S. Charuwichitratana, S. Siritantikorn, W. Chantratita, M. Iskander, A.W. Frahm and W. Jiratchariyakul, 2013. Antiherpetic effects of *Gynura procumbens*. Evidence-Based Complement. Altern. Med., Vol. 2013. 10.1155/2013/394865.
2. Vejanan, V., J. Latip, L.P. Chin, N. Embi and H.M. Sidek, 2012. *In vitro* and *in vivo* anti-plasmodial activities of *Gynura procumbens*. Sains Malaysiana, 41: 1535-1542.
3. Zheng, G.D., L.Q. Shuai, D.M. Li and Y.T. Zhu, 2014. Extraction and antibacterial effects of *Gynura procumbens* leaves. Food Sci. Technol., 39: 218-221.
4. Afandi, A., A. Sadikun and S. Ismail, 2014. Antioxidant properties of *Gynura procumbens* extracts and their inhibitory effects on two major human recombinant cytochrome P450s using a high throughput luminescence assay. Asian J. Pharm. Clin. Res., 7: 36-41.
5. Kaewseejan, N. and S. Siriamornpun, 2015. Bioactive components and properties of ethanolic extract and its fractions from *Gynura procumbens* leaves. Ind. Crops Prod., 74: 271-278.
6. Krishnan, V., S. Ahmad and M. Mahmood, 2015. Antioxidant potential in different parts and callus of *Gynura procumbens* and different parts of *Gynura bicolor*. BioMed Res. Int., Vol. 2015. 10.1155/2015/147909.
7. Rosidah, M. Yam, A. Sadikun and M. Asmawi, 2008. Antioxidant potential of *Gynura procumbens*. Pharm. Biol., 46: 616-625.
8. Meiyanto, E. and R.I. Jenie, 2007. Co-chemotherapy of sambung nyawa (*Gynura procumbens* (Lour.) Merr.) leaves ethanolic extract and doxorubicin on breast cancer cell. Indonesian J. Pharm., 18: 81-87.
9. Shwter, A.N., N.A. Abdullah, M.A. Alshawsh, A. Alsalahi and M. Hajrezaei *et al.*, 2014. Chemoprevention of colonic aberrant crypt foci by *Gynura procumbens* in rats. J. Ethnopharmacol., 151: 1194-1201.
10. Wang, J., S. Man, W. Gao, L. Zhang and L. Huang, 2013. Cluster analysis of ginseng tissue cultures, dynamic change of growth, total saponins, specific oxygen uptake rate in bioreactor and immuno-regulative effect of ginseng adventitious root. Ind. Crops Prod., 41: 57-63.
11. Shukla, A.A. and U. Gottschalk, 2013. Single-use disposable technologies for biopharmaceutical manufacturing. Trends Biotechnol., 31: 147-154.
12. Trejo-Tapia, G., A. Jimenez-Aparicio, L. Villarreal and M. Rodriguez-Monroy, 2001. Broth rheology and morphological analysis of *Solanum chrysostrichum* cultivated in a stirred tank. Biotechnol. Lett., 23: 1943-1946.

13. Jang, Y.S., M.A. Baque, M.H.K. Shiragi, S.H. Moh, E.J. Lee and K.Y. Paek, 2013. CO₂-enriched microenvironment induces biosynthesis of anthraquinones, phenolics and flavonoids in bioreactor cell suspension cultures of *Morinda citrifolia* (L.): The role of antioxidants and enzymes. Aust. J. Crop Sci., 7: 1606-1616.
14. Lulu, T., S.Y. Park, R. Ibrahim and K.Y. Paek, 2015. Production of biomass and bioactive compounds from adventitious roots by optimization of culturing conditions of *Eurycoma longifolia* in balloon-type bubble bioreactor system. J. Biosci. Bioeng., 119: 712-717.
15. Kokotkiewicz, A., A. Bucinski and M. Luczkiewicz, 2015. Xanthone, benzophenone and bioflavonoid accumulation in *Cyclopia genistoides* (L.) Vent. (honeybush) shoot cultures grown on membrane rafts and in a temporary immersion system. Plant Cell Tissue Org. Cult., 120: 373-378.
16. Kwiecien, I., A. Szydłowska, B. Kawka, L. Beerhues and H. Ekiert, 2015. Accumulation of biologically active phenolic acids in agitated shoot cultures of three *Hypericum perforatum* cultivars: 'Elixir', 'Helos' and 'Topas'. Plant Cell Tissue Org. Cult., 123: 273-281.
17. Mariateresa, C., C.S.C. Maria and C. Giuseppe, 2014. Influence of ozone treatments on *in vitro* propagation of *Aloe barbadensis* in continuous immersion bioreactor. Ind. Crops Prod., 55: 194-201.
18. Cui, H.Y., H.N. Murthy, S.H. Moh, Y. Cui, E.J. Lee and K.Y. Paek, 2014. Protocorm culture of *Dendrobium candidum* in balloon type bubble bioreactors. Biochem. Eng. J., 88: 26-29.
19. Manuhara, Y.S.W., D.Y. Kusuma, R.L.K. Sari and A.N. Kristanti, 2017. Biomass production of *Gynura procumbens* adventitious roots in different type of liquid culture. Biosaintifika: J. Biol. Biol. Edu., 9: 523-529.
20. Manuhara, Y.S.W., A.N. Kristanti and E.S.W. Utami, 2015. Optimization of culture conditions of *Talinum paniculatum* Gaertn. Adventitious roots in balloon type bubble bioreactor using aeration rate and initial inoculum density. Asian J. Biol. Sci., 8: 83-92.
21. Noviyanti, R., R.L.K. Sari, A.N. Kristanti, A. Yachya and Y.S.W. Manuhara, 2017. Biomass and flavonoid production of *Gynura procumbens* adventitious roots induced by sucrose, phenylalanine and tyrosine. Biosci. Res., 14: 934-941.
22. Kusuma, D.Y., A.N. Kristanti and Y.S.W. Manuhara, 2017. Effect of sucrose and immersion frequency on production of adventitious roots and secondary metabolites of *Gynura procumbens* (Lour.) merr in temporary immersion bioreactors. Asian J. Plant Sci., 16: 24-36.
23. Jeong, C.S., H.N. Murthy, E.J. Hahn, H.L. Lee and K.Y. Paek, 2009. Inoculum size and auxin concentration influence the growth of adventitious roots and accumulation of ginsenosides in suspension cultures of ginseng (*Panax ginseng* C.A. Meyer). Acta Physiol. Plant., 31: 219-222.
24. Lee, E.J., S.H. Moh and K.Y. Paek, 2011. Influence of inoculum density and aeration volume on biomass and bioactive compound production in bulb-type bubble bioreactor cultures of *Eleutherococcus koreanum* Nakai. Bioresour. Technol., 102: 7165-7170.
25. Rahman, M.Z., S.M.S. Islam, A.N. Chowdhury and S. Subramaniam, 2015. Efficient microtuber production of potato in modified nutrient spray bioreactor system. Sci. Hortic., 192: 369-374.
26. Pramita, A.D., A.N. Kristanti, E.S.W. Utami and Y.S.W. Manuhara, 2018. Production of biomass and flavonoid of *Gynura procumbens* (Lour.) Merr shoots culture in temporary immersion system. J. Genet. Eng. Biotechnol., 16: 639-643.
27. Murashige, T. and F. Skoog, 1962. A revised medium for rapid growth and bio assays with tobacco tissue cultures. Physiol. Planta., 15: 473-497.
28. Debnath, S.C., 2008. Developing a scale-up system for the *in vitro* multiplication of thidiazuron-induced strawberry shoots using a bioreactor. Can. J. Plant Sci., 88: 737-746.
29. Baque, M.A., M.H.K. Shiragi, S.H. Moh, E.J. Lee and K.Y. Paek, 2013. Production of biomass and bioactive compounds by adventitious root suspension cultures of *Morinda citrifolia* (L.) in a liquid-phase airlift balloon-type bioreactor. *In vitro* Cell. Dev. Biol.-Plant, 49: 737-749.
30. Salisbury, B.F. and C.W. Ross, 1992. Plant Physiology. 4th Edn., Wadsworth Publishing Company, Belmont, CA., USA., ISBN-13: 9780534151621, Pages: 682.
31. Ball, E., 1953. Hydrolysis of sucrose by autoclaving media, a neglected aspect in the technique of culture of plant tissues. Bull. Torrey Bot. Club, 80: 409-411.
32. Medvedev, S.S., 2012. Mechanisms and physiological role of polarity in plants. Russian J. Plant Physiol., 59: 502-514.