

PAPER • OPEN ACCESS

Selection of potential Indonesian plant species for antioxidant

To cite this article: K Kartini *et al* 2020 *IOP Conf. Ser.: Earth Environ. Sci.* **457** 012040

View the [article online](#) for updates and enhancements.

A promotional banner for the PRIME 2020 meeting. The background features a blue and white globe. On the left, there are three circular logos: the top one is 'ECS' in a white circle, the middle one is 'The Electrochemical Society' with a stylized 'ECS' logo, and the bottom one is 'THE KOREAN ELECTROCHEMICAL SOCIETY'. The main text in the center reads 'Joint International Meeting PRIME 2020 October 4-9, 2020' in white and blue. Below this, a blue bar contains the text 'Attendees register at NO COST!' in white. On the right side, there is a large white logo for 'PRIME' with a stylized 'P' and 'M' shape above it, followed by 'PACIFIC RIM MEETING ON ELECTROCHEMICAL AND SOLID STATE SCIENCE' and '2020' in white. At the bottom right, a blue bar contains the text 'REGISTER NOW' in white with a white arrow pointing right.

ECS

Joint International Meeting

PRIME 2020

October 4-9, 2020

Attendees register at NO COST!

PRIME
PACIFIC RIM MEETING
ON ELECTROCHEMICAL
AND SOLID STATE SCIENCE

2020

REGISTER NOW ▶

Selection of potential Indonesian plant species for antioxidant

K Kartini^{1*}, F Setiawan¹, J Sukweenadhi², O Yunita¹ and C Avanti³

¹Department of Pharmaceutical Biology, Faculty of Pharmacy, University of Surabaya, Surabaya, Indonesia

²Faculty of Technobiology, University of Surabaya, Surabaya, Indonesia

³Department of Pharmaceutic, Faculty of Pharmacy, University of Surabaya, Surabaya, Indonesia

*Corresponding author: kartini@staff.ubaya.ac.id

Abstract. Diabetic foot ulcer (DFU) is commonly complication of diabetes mellitus. Antimicrobial and antioxidant herbs can be used as complementary therapy in DFU and these properties will be strengthened by silver nanoparticles (AgNPs). This study was performed for screening of Indonesian herbs promising as antioxidant which can be subsequently involved in the green synthesis of AgNPs. Antioxidant of eight herbs extracts were evaluated using 1,1-diphenyl-2-picrylhydrazyl method. Their total group of compounds were quantified as well. *Phyllanthus niruri* and *Orthosiphon stamineus* exhibited the strongest antioxidant power with IC₅₀ of 102 and 133 µg/ml, followed by *Curcuma domestica*, *Stelechocarpus burahol*, and *Curcuma xanthorrhiza* with the IC₅₀ of 363, 481, and 540 µg/ml, respectively. *Sonchus arvensis*, *Apium graveolens*, *Centella asiatica* did not have antioxidants activity. Total phenolics from the highest is: *O. stamineus* > *C. domestica* > *C. xanthorrhiza* > *P. niruri* > *S. burahol* > *S. arvensis* > *A. graveolens* > *C. asiatica*, whereas total flavonoids order is *O. stamineus* > *S. burahol* > *P. niruri* > *A. graveolens* > *C. asiatica* > *S. arvensis* > *C. xanthorrhiza* > *C. domestica*. We concluded that *O. stamineus* was the most suitable herb to be used as a bioreductor in the green synthesis of AgNPs.

Keywords: Antioxidant, silver nanoparticles, diabetic wound healing, medicinal plants

1. Introduction

A high prevalence of diabetes causes the emergence of comorbidities. The critical effects of hyperglycemia include micro-vascular and macro-vascular complications [1]. Diabetes is the cause of non-traumatic lower limb amputation, which is preceded by wounds that do not heal [2]. The risk of wound in diabetics is 15-20% [3]. About 15-27% of diabetic patients experience amputation due to infection [4]. 45-60% of diabetic foot ulcers (DFU) are estimated due to neuropathy [5]. Emerging studies show high infections of *Pseudomonas* [6], *E. coli* [7], and *S. aureus* [8] in DFU. Diabetic foot ulcers treatment requires handling of wounds as well as giving appropriate antibiotics [1]. Silver/sulphadiazine dressing can be considered in DFU therapy. Silver and polyherbal dressing shows promising results in wound healing [9].

Various plant species have been used empirically in Indonesia for wound healing [10]. Some potential species for DFU therapy are *Curcuma xanthorrhiza* (*temulawak*), *Curcuma domestica* (*kunyit*), *Centella asiatica* (*pegagan*), *Stelechocarpus burahol* (*kepel*), *Orthosiphon stamineus* (*kumis kucing*), *Phyllanthus niruri* (*meniran*), *Sonchus arvensis* (*tempuyung*), and *Apium graveolens* (*seledri*)



[11]. These plants contain antioxidant and antimicrobial compounds, as well as being able to reduce metals [12-14].

Nanoparticles are the current technology and have been used in many fields, one of which is metal nanoparticles [2]. It has been widely used in biomedical and physicochemical fields [2]. Silver nanoparticles are also widely used for antimicrobial, anticancer, anti-inflammatory, and wound treatment [15, 16]. Green synthesis of nanoparticles is carried out using microorganisms, plants, yeast, or viruses. Nanoparticle synthesis using plants, phytonanotechnology, has a lot of advantages such as biocompatibility, fast, stable, environmentally friendly, affordable, and easy to scale-up [2]. Various chemical compounds including primary metabolites such as protein and polysaccharides as well as secondary metabolites i.e. terpenoids, phenolics, flavonoids, and alkaloids can be applied in the green synthesis of metal nanoparticles including silver nanoparticles (AgNPs) [17]. They work as reducing and capping agents [18]. The green synthesis of AgNPs involving plant extracts can be an option to increase the effectiveness of topical preparations for DFU treatment [19]. As an initial step, this research was conducted to select native Indonesian plant species having potential as antioxidants. Selected plants will be further developed for the green synthesis of silver nanoparticles as DFU treatment.

2. Material and methods

2.1. Plant species

This research used eight plant species as described at table 1. All crude drugs were obtained from and authenticated by Center for Research and Development of Medicinal Plants and Traditional Medicines (B2P2TOOT) located at Tawangmangu, Indonesia.

Table 1. Selected Indonesian plant species for antioxidant.

Plant species	Part used
<i>Orthosiphon stamineus</i>	Leaves
<i>Curcuma domestica</i>	Rhizome
<i>Curcuma xanthorrhiza</i>	Rhizome
<i>Phyllanthus niruri</i>	Herbs
<i>Sonchus arvensis</i>	Leaves
<i>Apium graveolens</i>	Herbs
<i>Centella asiatica</i>	Herbs
<i>Stelechocarpus burahol</i>	Leaves

2.2. Extraction

Dried plants were milled into 2 mm before used. Each powdered plant was then refluxed with aqueous ethanol for 3 x 2.5 h. The ethanol were then removed from the extracts by using rotary evaporator at the temperature of 40°C, continued by evaporation on waterbath to yield a thick extract. The thick extracts were subsequently stored in a desiccator.

2.3. Quantification of total phenolics

Concentration of the total phenolics in the crude extracts was determined by using Folin-Ciocalteu (FC) reagent according to the previous studies [20, 21]. Analysis was carried out on 96-well microplates by using gallic acid as a standard. It was prepared in various concentrations started from 2 to 10 µg/ml. Reaction was initiated by adding FC reagent (0.1 mol/l, 150 µl), followed by 30 µl of each ethanol extract or standard solution. After 10 min, solution of Na₂CO₃ (7.5%, 120 µl) was added and incubated for two hours. The absorbance was subsequently read at the wavelength of 760 nm. Total phenolics concentration of the extracts was calculated by interpolation of the absorbance to the calibration curve of standard. The total phenolics concentration was presented as gram of gallic acid equivalents (GAE) per 100 g of thick extract.

2.4. Quantification of total flavonoids

Evaluation of total flavonoids concentration was conducted by established colorimetric methods as detailed in previous study [22]. Thick extract was dissolved and diluted in 96% ethanol. One ml of extract was transferred into a volumetric flask, supplemented with 4 ml of demineralized water and solution of NaNO_2 (5%, 0.3 ml), followed by incubation at room temperature for 5 minutes. Solution of AlCl_3 (10%, 0.3 ml) was then added at the 6th minute. Sodium hydroxide (1 M, 2 ml) and demineralized water were finally added up to 10.0 ml. The absorbance was subsequently read at the wavelength of 400.2 nm. Total flavonoids concentration of the extracts was calculated by interpolation of the absorbance to the calibration curve of standard (quercetin, 5 to 25 $\mu\text{g/ml}$). The total flavonoids concentration was expressed as gram of quercetin equivalents (QE) per 100 g of thick extract.

2.5. Antioxidant activity

Antioxidant activity test referred to the previous work with slight modification [23]. It was based on the capability of the extract to scavenge free radical of DPPH. In summary, the method is as follows: solution of DPPH in methanol (0.026%, 50 μl) was pipetted to 100 μl of serial dilution of extracts in methanol (total volume 150 μl). It was left in darkness for 15 min and the absorbance was then measured at the wavelength of 517 nm using a microplate reader. Free radical scavenging activity was calculated using the following formula:

$$\% \text{ Inhibition} = \frac{A_B - A_A}{A_B} \times 100$$

where, A_B = absorption of DPPH solution ($t = 0$ min); A_A = absorption of tested samples ($t = 15$ min). IC_{50} value, the concentration of extract, which exhibited 50% radical scavenging activity, was deduced from the linear regression of concentration versus percentage of inhibition [24].

2.6. Statistical analysis

Both total phenolics/flavonoids and IC_{50} values were presented as average \pm SD, $n = 3$ or 4. Data were statistically analysed using ANOVA followed with post hoc Tukey test. All analysis was conducted with the statistical software, GraphPad Prism (GraphPad Software Inc. Windows Version 5.01). Differences were concluded to be significant at P value of less than 0.05.

3. Results and discussion

3.1. Total phenolics and flavonoid concentration

Group of compounds, phenolics and flavonoids, have been determined in this research to evaluate the quality of each herb material. Total phenolics was analysed with the Folin-Ciocalteu method, whereas total flavonoids was evaluated with the aluminum chloride (AlCl_3) colorimetric method using spectrophotometer. A series concentration of gallic acid (figure 1) as well as quercetin (figure 2) were prepared to set up the standard calibration curves, which were then applied to quantify the total phenolics and flavonoid contents in each crude extract. Table 2 presents total phenolics content as well as total flavonoid compounds calculated for each plant extract. It could be observed that phenolics and flavonoid contents of *Orthosiphon stamineus* extract were the highest.

Table 2 demonstrate that total phenolics content from the highest is as follow: *O. stamineus* > *C. domestica* > *C. xanthorrhiza* > *P. niruri* > *S. burahol* > *S. arvensis* > *A. graveolens* > *C. asiatica*, whereas the total flavonoid content order is *O. stamineus* > *S. burahol* > *P. niruri* > *A. graveolens* > *C. asiatica* > *S. arvensis* > *C. xanthorrhiza* > *C. domestica*. *Orthosiphon stamineus* contains the highest phenolics and flavonoids.




Figure 1. Calibration curve of gallic acid




Figure 2. Calibration curve of quercetin

Table 2. Total phenolics and flavonoids concentration of selected Indonesian plant species extracts.

Plant species	Total phenolics concentration (g of GAE/100 g of extract)*	Total flavonoids concentration (g of QE/100 g of extract)**
<i>Orthosiphon stamineus</i>	44.91 ± 0.24 ^a	20.03 ± 0.68 ^a
<i>Curcuma domestica</i>	13.10 ± 0.67 ^b	1.24 ± 0.48 ^b
<i>Curcuma xanthorrhiza</i>	7.38 ± 0.40 ^c	2.77 ± 1.58 ^{b,c}
<i>Phyllanthus niruri</i>	7.22 ± 0.06 ^c	8.34 ± 0.74 ^d
<i>Sonchus arvensis</i>	2.62 ± 0.15 ^d	3.78 ± 0.09 ^c
<i>Apium graveolens</i>	2.16 ± 0.05 ^d	6.67 ± 0.31 ^d
<i>Centella asiatica</i>	0.67 ± 0.03 ^e	4.06 ± 0.27 ^c
<i>Stelechocarpus burahol</i>	5.85 ± 0.18 ^f	9.74 ± 0.27 ^e

*,**Values in the column represent average ± SD of 3 determinations. Averages followed by different notation within same column show significant differences ($P < 0.05$).

Flavonoids such as 3'-hydroxy-5,6,7,4'-tetramethoxyflavone, sinensetin, as well as eupatorin have been identified from *O. stamineus*, in addition to rosmarinic acid, a caffeic acid derivative. Moreover, total phenolics and flavonoids of this crude drug varies according to the geographical origin [25]. Phenolics content of *O. stamineus* correlates with the solvent, duration, and temperature of extraction. The optimum conditions for phenolics extraction from *O. stamineus* were 40% alcohol for 120 min at 65°C [26].

3.2. Antioxidant activity

Antioxidant activity of eight Indonesian plant species has been determined with free radical scavenging method against stable DPPH (1,1-diphenyl-2-picrylhydrazyl). DPPH is a commonly used free radical in the antioxidant screening assay. It has an unpaired electron from nitrogen atom. When a

compound or material with capability to donate hydrogen is reacted with the DPPH, it will reduce DPPH become DPPH-H (1,1-diphenyl-2-picrylhydrazyn) (figure 3) [27].




Table 3. IC₅₀ value of DPPH scavenging activity by selected extracts of Indonesian plant species.

Extract of plant species	IC ₅₀ (µg/ml)
<i>Orthosiphon stamineus</i>	133 ± 6.6 ^a
<i>Curcuma domestica</i>	363 ± 9.7 ^b
<i>Curcuma xanthorrhiza</i>	540 ± 12.8 ^c
<i>Phyllanthus niruri</i>	102 ± 1.7 ^a
<i>Sonchus arvensis</i>	1135 ± 7.1 ^d
<i>Apium graveolens</i>	2263 ± 61.8 ^e
<i>Centella asiatica</i>	ND
<i>Stelechocarpus burahol</i>	481 ± 2.7 ^f
Ascorbic acid*	31 ± 1.6
Trolox*	76 ± 1.1

*standard compound; ND: not detected; Values in the column represent average ± SD of IC₅₀ (inhibitory concentration 50, $n = 4$). Averages followed by different notation within same column show significant differences ($P < 0.05$).

Flavonoids and phenolics are ubiquitous compounds which can be found in many species of plants. They are main source of natural antioxidants in plants, fruits, and vegetables. In addition to flavonoids especially lipophilic flavones as well as caffeic acid derivatives, *O. stamineus* contains terpenoids and essential oil. Lipophilic flavones and caffeic acid ester of 3-(3,4-dihydroxyphenyl)lactic acid were main phenolics compounds found in the crude extract of *O. stamineus* and their antioxidant activities have been well determined [29, 30]. However, in our research, there was no linear correlation between the antioxidant capacity and the total phenolics concentration ($R^2 = 0.238$) as well as between antioxidant power and the total flavonoids concentration ($R^2 = 0.079$). The same pattern on the correlation between antioxidant capacity of plant extracts and the phenolics concentration have been previously investigated ($R_{xy} = 0.048$) [25, 31]. Phenolics is not the only phytochemical with antioxidant properties. The existence of various other anti-oxidant compounds in the plant extracts, which are able to donate hydrogen may erroneously contribute to the content of the total phenols determined with Folin–Ciocalteu reagent. Therefore, it is no simple correlation between the level of total phenolics and the antioxidant activity when compare among herb materials [25].

4. Conclusion

Phyllanthus niruri and *Orthosiphon stamineus* showed the most potent antioxidant with an IC₅₀ value of 102 and 133 µg/ml, followed by *C. domestica*, *S. burahol*, and *C. xanthorrhiza* with the IC₅₀ value of 363, 481, and µg/ml ppm, respectively. *Sonchus arvensis*, *A. graveolens*, and *C. asiatica* exhibited poor antioxidant capacity with the IC₅₀ value > 1000 ppm. Total phenolics content from the highest is as follow: *O. stamineus* > *C. domestica* > *C. xanthorrhiza* > *P. niruri* > *S. burahol* > *S. arvensis* > *A. graveolens* > *C. asiatica*, whereas the total flavonoids content order is *O. stamineus* > *S. burahol* > *P. niruri* > *A. graveolens* > *C. asiatica* > *S. arvensis* > *C. xanthorrhiza* > *C. domestica*. *Orthosiphon stamineus* contains the highest concentration phenolics and flavonoids. *Orthosiphon stamineus* was the most suitable herbs to be used as a source of bioreductors in the green synthesis of silver nanoparticles for foot ulcer.

Acknowledgements

We thank to the Ministry of Research, Technology and Higher Education of the Republic of Indonesia who has supported this work under Fundamental Research Scheme with the contract number: 004/SP2H/LT/MULTI/L7/2019.

References

- [1] Clayton Jr W and Elasy T A 2009 *Clin. Diabetes* **27** 52
- [2] Kavitha K V, Tiwari S, Purandare V B, Khedkar S, Bhosale S S and Unnikrishnan A G 2014 *World J. Diabetes* **5** 546
- [3] Singh P, Kim Y-J, Zhang D and Yang D-C 2016 *Trends Biotechnol.* **34** 588
- [4] Reiber G E 2002 *The Diabetic Foot* (Netherland: Springer Netherlands)
- [5] Richard J-L, Sotto A and Lavigne J-P 2011 *World J. Diabetes* **2** 24
- [6] Shanmugam P and Jeya M 2013 *J. Clin. Diagn. Res.* **7** 441
- [7] Tiwari S, Pratyush D D, Dwivedi A, Gupta S K, Rai M and Singh S K 2012 *J. Infect. Dev. Ctries.* **6** 329
- [8] Abdulrazak A, Bitar Z I, Al-Shamali A A and Mobasher L A 2005 *J. Diabetes Complications* **19** 138
- [9] Viswanathan V, Kesavan R, Kavitha K and Kumpatla S 2011 *Indian J. Med. Res.* **134** 168
- [10] Elfahmi, Woerdenbag H J and Kayser O 2014 *J. Herb. Med.* **4** 51
- [11] Wientarsih I, Winarsih W and Sutardi L N 2013 *Jurnal Veteriner* **13** 251 (In Indonesian Language)
- [12] Shameli K, Ahmad M B, Zamanian A, Sangpour P, Shabanzadeh P, Abdollahi Y, et al 2012 *Int. J. Nanomedicine* **7** 5603
- [13] Logeswari P, Silambarasan S and Abraham J 2015 *J. Saudi Chem. Soc.* **19** 311
- [14] Singh K, Panghal M, Kadyan S and Yadav J P 2014 *JNMNT* **5** 1
- [15] Singh N, Armstrong D G and Lipsky B A 2005 *Jama* **293** 217
- [16] Keat C L, Aziz A, Eid A M and Elmarzugi N A 2015 *Bioresour. Bioprocess.* **2** 47
- [17] Ahmed S, Ahmad M, Swami B L and Ikram S 2016 *J. Adv. Res.* **7** 17
- [18] Duan H, Wang D and Li Y 2015 *Chem. Soc. Rev.* **44** 5778
- [19] Mishra M, Kumar H and Tripathi K 2008 *Dig. J. Nanomater. Bios.* **3** 49
- [20] Beara I N, Lesjak M M, Jovin E Đ, Balog K J, Anačkov G T, Orčić D Z et al 2009 *J. Agric. Food Chem.* **57** 9268
- [21] Singleton V L, Orthofer R and Lamuela-Raventos R M 1999 *Methods Enzymol.* **299C** 152
- [22] Chang C-C, Yang M H, Wen H M and Chern J C 2002 *J. Food Drug Anal.* **10** 178
- [23] Galvez M, Martin-Cordero C, Houghton P J and Ayuso M J 2005 *J. Agric. Food Chem.* **53** 1927
- [24] Chen Z, Bertin R and Frolidi G 2013 *Food Chem.* **138** 414
- [25] Akowuah G, Zhari I, Norhayati I, Sadikun A and Khamsah S 2004 *Food Chem.* **87** 559
- [26] Chew K, Khoo M, Ng S, Thoo Y, Aida WW and Ho C 2011 *Int. Food Res. J.* **18** 1427
- [27] Liang N and Kitts D 2014 *Molecules* **19** 19180
- [28] Jun M, Fu H Y, Hong J, Wan X, Yang C and Ho CT 2003 *J. Food Sci.* **68** 2117
- [29] Brand-Williams W, Cuvelier M E and Berset C 1995 *LWT - Food Sci. Technol.* **28** 25
- [30] Cuvelier M-E, Richard H and Berset C 1992 *Biosci. Biotechnol. Biochem.* **56** 324
- [31] Maillard M-N and Berset C 1995 *J. Agric. Food Chem.* **43** 1789



PAPER • OPEN ACCESS

LIST OF COMMITTEE ICOBIO 2019

To cite this article: 2020 *IOP Conf. Ser.: Earth Environ. Sci.* **457** 011002

View the [article online](#) for updates and enhancements.

You may also like

- [Design and Optimization of Static Characteristics for a Steering System in an ATV](#)

L. Rajeshkumar, V. Bhuvanewari, B. Pradeepraj et al.

- [Quantum steering: a review with focus on semidefinite programming](#)

D Cavalcanti and P Skrzypczyk

- [Evaluation of Acceptability of Steering Support Method Using Impedance Control with Multiple HMs](#)

Takuya Hirata, Soichiro Hayakawa, Shigeyoshi Tsutsumi et al.



ECS Membership = Connection

ECS membership connects you to the electrochemical community:

- Facilitate your research and discovery through ECS meetings which convene scientists from around the world;
- Access professional support through your lifetime career;
- Open up mentorship opportunities across the stages of your career;
- Build relationships that nurture partnership, teamwork—and success!

Join ECS!

Visit electrochem.org/join



LIST OF COMMITTEE ICOBIO 2019

Steering Committee

Dr. Miftahudin (IPB University, Indonesia)
Dr. Sri Nurdiati (IPB University, Indonesia)
Prof. Dr. Aris Tri Wahyudi (IPB University, Indonesia)
Prof. Dr. Anja Meryandini (IPB University, Indonesia)
Dr. Chatchawan Jantasuriyarat (Kasetsart University, Thailand)
Prof. Dr. Ahmad Ismail, FASc (Universiti Putra Malaysia, Malaysia)
Dr. Martyna M. Kotowska (Georg-August-Universität Göttingen, German)
Dr. Jun Kohyama (Keio University, Japan)

Organizing Committee (IPB University)

Dr. Ence D.J .Supena (Chairman)
Dr. Berry Juliandi (Vice-Chairman)
Dr. Rika Indri Astuti (Secretary)
Dr. Puji Rianti (Secretary)
Dr. Nina Ratna Djuita (Secretary)
Dr. Sulistijorini (Treasurer)
Prof. Dr. YM Diah Ratnadewi
Dr. Triadiati
Dr. Sri Budiarti
Dr. Nampiah Sukarno
Dr. Ibnul Qayim
Mafrikhul Muttaqin, MSi
Ivan Permana Putra, MSi
Windra Priawandiputra, PhD
Dr. Rika Rafiuddin
Dra. Hilda Akmal, MSi
Hirmas Fuady Putra, MSi

Scientific Editors

Dr. Miftahudin
Dr. Berry Juliandi
Windra Priawandiputra, PhD
Mafrikhul Muttaqin, MSi



PAPER • OPEN ACCESS

Editorial Board Members

To cite this article: 2016 *IOP Conf. Ser.: Mater. Sci. Eng.* **120** 011002

View the [article online](#) for updates and enhancements.

Related content

- [Editorial board](#)

- [Editorial Board](#)

- [Editorial board](#)



The banner features a background image of Earth from space. On the left, there are three circular logos: the top one is 'ECS' in a circle, the middle one is 'The Electrochemical Society' with a stylized 'ECS' logo, and the bottom one is 'THE KOREAN ELECTROCHEMICAL SOCIETY'. The main text in the center reads 'Joint International Meeting PRIME 2020 October 4-9, 2020' in white and blue. Below this, a blue bar contains the text 'Attendees register at NO COST!' in white. On the right side, there is a large 'PRIME' logo with 'PACIFIC RIM MEETING ON ELECTROCHEMICAL AND SOLID STATE SCIENCE' underneath, and '2020' in large white numbers. At the bottom right, a blue bar contains the text 'REGISTER NOW' with a white arrow pointing right.

Editorial Board Members

Subramaniam Ananthakrishnan (Mumbai, India)

Pavel Belov (Russia)

Charles Cavalcante Casimiro (Brazil)

Sergio Colafrancesco (Italy)

Mérouane Debbah (France)

Lars Jacob Foged (Denmark)

Debatosh Guha (India)

Sébastien Lalléchère (France)

Jean-Daniel Lan Sun Luk (France)

Dominique Lesselier (France)

André de Lustrac (France)

James McLean (USA)

Eric Mokole (USA)

Vikass Monebhurrin (France)

Shailendra Oree (Mauritius)

Lionel Pichon (France)

Blaise Ravelo (China)

Tapan Sarkar (India)

Russell Taylor (USA)

Bernard Veyret (France)



NOTICE: There are currently some performance issues with IOPscience, which may also cause error messages to appear. Apologies for the inconvenience..

Table of contents

Volume 457

2020

◀ Previous issue Next issue ▶

The 3rd International Conference on Biosciences 8 August 2019, IPB International Convention Centre, Bogor, Indonesia

Accepted papers received: 10 February 2020

Published online: 26 March 2020

[Open all abstracts](#)

Preface

OPEN ACCESS 011001

The 3rd International Conference on Biosciences (The 3rd ICoBio)

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS 011002

LIST OF COMMITTEE ICOBIO 2019

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS 011003

Logo

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS 011004

Peer review statement

[+ Open abstract](#) [View article](#) [PDF](#)

Biodiversity, Conservation Ecology, Climate Change

OPEN ACCESS 012001

Synurbic avian species in Greater Jakarta Area, Indonesia

A. Mardiasuti, Y. A. Mulyani, D. Rinaldi, W. Rumbat, L. K. Dewi, A. Kaban and H. Sastranegara

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more,

[see our Privacy and Cookies policy](#) [PDF](#)



[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012038

Selection and characterization yeast potential from pineapple for bioethanol production using some sugar sources

N Chasanah, A Meryandini and T C Sunarti

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012039

The toxicity of the extract of tobacco leaf *Nicotiana tabacum* L, marigold leaf *Tithonia diversifolia* (HAMSLEY) and citrus japonsche citroen peel *Citrus limonia* against citrus psyllid (*Diaphorina citri* Kuwayama), the vector of citrus HLB disease

S Wuryantini, Harwanto and R A Yudistira

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012040

Selection of potential Indonesian plant species for antioxidant

K Kartini, F Setiawan, J Sukweenadhi, O Yunita and C Avanti

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012041

Antimicrobial activity of biosynthesized silver nanoparticles using *Diospyros celebica* Bakh. (Eboni) extra

W Handayani, H Zukhrufa, Yasman and C Imawan

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012042

Biological synthesis and characterization of silver nanoparticles synthesized from *Pometia pinnata* and *Diospyros discolor* Fruits

W Handayani, I Nolia, R M Sundari and C Imawan

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012043

Isolation and characterization endophytic bacteria as biological control of fungus *Colletotrichum* sp. on onion plants (*Allium cepa* L.)

A Murtado, N R Mubarik and A Tjahjoleksono

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012044

Ethnopharmacology and chemical screening of piper from ketambe research station, Gunung Leuser National Park, Southeast Aceh

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our [Privacy and Cookies policy](#).





Test Potensi Akademik Online

Tingkatkan kemungkinan anda diterima bekerja dgn melampirkan hasil Test Potensi Akademik

tokopedia.com

BUKA

IOP Conference Series: Earth and Environmental Science

Country [United Kingdom](#) - [SIR Ranking of United Kingdom](#)

Subject Area and Category [Earth and Planetary Sciences](#)
[Earth and Planetary Sciences \(miscellaneous\)](#)

[Environmental Science](#)
[Environmental Science \(miscellaneous\)](#)

Publisher [IOP Publishing Ltd.](#)

Publication type Conferences and Proceedings

ISSN 17551315, 17551307

Coverage 2010-2020

Scope The open access IOP Conference Series: Earth and Environmental Science (EES) provides a fast, versatile and cost-effective proceedings publication service.

[Homepage](#)

[How to publish in this journal](#)

[Contact](#)

[Join the conversation about this journal](#)

18








H Index

Test Potensi Akademik Online

Tingkatkan kemungkinan anda diterima bekerja dgn melampirkan hasil Test Potensi Akademik

tokopedia.com

BUKA



IOP Conference Series: Earth and Environmental...

← Show this widget in your own website

Not yet assigned quartile

Just copy the code below and paste within your html code:

SJR 2019
0.18

powered by scimagojr.com

```
<a href="https://www.scimagojr.com/journalsearch.php?q=19900195068&tip=sid&clean=0" style="border: 1px solid #ccc; padding: 2px 5px; display: inline-block;"></pre>

```



Source details

IOP Conference Series: Earth and Environmental Science

Scopus coverage years: from 2010 to Present

ISSN: 1755-1307 E-ISSN: 1755-1315

Subject area: Environmental Science: General Environmental Science

Earth and Planetary Sciences: General Earth and Planetary Sciences

CiteScore 2019

0.4



SJR 2019

0.175



SNIP 2019

0.514



[View all documents >](#)

[Set document alert](#)

[Save to source list](#) [Journal Homepage](#)

[CiteScore](#) [CiteScore rank & trend](#) [Scopus content coverage](#)

Improved CiteScore methodology

CiteScore 2019 counts the citations received in 2016-2019 to articles, reviews, conference papers, book chapters and data papers published in 2016-2019, and divides this by the number of publications published in 2016-2019. [Learn more >](#)

CiteScore 2019

$$0.4 = \frac{11,544 \text{ Citations 2016 - 2019}}{32,872 \text{ Documents 2016 - 2019}}$$

Calculated on 06 May, 2020

CiteScoreTracker 2020

$$0.4 = \frac{17,363 \text{ Citations to date}}{42,393 \text{ Documents to date}}$$

Last updated on 07 September, 2020 • Updated monthly

CiteScore rank 2019

Category	Rank	Percentile
Environmental Science	#176/210	16th
General Environmental Science		
Earth and Planetary Sciences	#164/187	12th
General Earth and Planetary Sciences		

[View CiteScore methodology >](#) [CiteScore FAQ >](#) [Add CiteScore to your site](#)

About Scopus

- [What is Scopus](#)
- [Content coverage](#)
- [Scopus blog](#)
- [Scopus API](#)
- [Privacy matters](#)

Language

- [日本語に切り替える](#)
- [切换到简体中文](#)
- [切换到繁體中文](#)
- [Русский язык](#)

Customer Service

- [Help](#)
- [Contact us](#)



The 3rd International Conference on
Biosciences (IcoBio) 2019
IPBICC, Indonesia, 8 August 2019

CERTIFICATE

This is to certify that

Kartini, S.Si., M.Si., Apt., Ph.D.

has attended as


Oral Presenter

in the 3rd International Conference on Biosciences (ICoBio) 2019
organized by the Department of Biology, IPB University on 8 August 2019
in IPB International Convention Center (IPBICC), Bogor, Indonesia

Bogor, 8 August 2019



Dr. Ir. Sri Nurdianti, MSc
Dean of Faculty of Mathematics
and Natural Sciences, IPB University



Dr. Ence Darmo Jaya Supena
Organizing Committee Chairman
The 3rd IcoBio 2019

The Committee of The 3rd International Conference on Biosciences (ICoBio) 2019
Department of Biology, Faculty of Mathematics and Natural Sciences, IPB University
Gedung Biologi, Jl. Agatis-Kampus IPB Dramaga Bogor 16680 Indonesia
<https://biologi.ipb.ac.id/web/en/page/icobio> | social media: bioipb

BIOLOGI.
IPB.AC.ID