

A PROSPECTIVE MODIFICATION STRUCTURE: THE EFFECT OF LIPOPHILIC AND ELECTRONIC PROPERTIES OF *N*-(PHENYLCARBAMOTHYOIL)BENZAMIDE DERIVATIVES ON CYTOTOXIC ACTIVITY BY IN SILICO AND IN VITRO ASSAY WITH T47D CELLS

D. Kesuma¹, A.L. Nasyanka^{2,*}, M. Rudyanto³, Siswandono³, B.T. Purwanto³ and I.G.A. Sumartha¹

¹Department of Pharmaceutical Chemistry, University of Surabaya, Surabaya-60293, (East Java) Indonesia

²Departement of Pharmacy, University of Muhammadiyah Gresik, Gresik-61111, (East Java) Indonesia

³Department of Medicinal Chemistry, University of Airlangga, Surabaya-60286, (East Java) Indonesia

*E-mail: anindilupita@umg.ac.id

ABSTRACT

Topliss aromatic modification structure considers two important properties, lipophilic and electronic. This study is to prove lipophilic and electronic properties effect of an anticancer lead compound *N*-(phenylcarbamothyoil) benzamide using in silico and in vitro method. We used in silico method that docked two derivatives (4-nitro and 4-methyl moieties) in the sirtuin-1 receptor (PDB ID: 4i5i) using autodock tools. They were screened for cytotoxic activity on T47D cells with MTT Method. The result showed that there were different activities in docking prediction and cytotoxic activity on T47D. 4-Nitro PCTB had more increased activity than 4-methyl. It was implied that electronic activity is more affecting cytotoxic activity than lipophilic properties.

Keywords: Lipophilic Effect, Electronic Effect, *N*-(phenylcarbamothyoil)Benzamide, T47D Cell, *In-silico*

© RASĀYAN. All rights reserved

INTRODUCTION

Since a long ago, modification of drug structure is one of the promising methods to find new active drug compounds. A lead compound must be developed first with this method¹. Lead compound structure modification varies, but changing its physical and chemical properties by adding substituent makes it more active as a drug². Every substituent has its effect on pharmacokinetics and pharmacodynamics³.

Topliss proposed guidelines for structural modification of non-mathematical, non-statistical and non-computerized parent compounds using the basic principles of the structural relationship approach and the activities of the Hansch model⁴. The structural modification of Topliss method includes groups that have certain lipophilic, electronic, and steric properties at certain positions in the lead compound structure, which are predicted to produce compounds with higher, equal, or lower activity than the lead compound, and lastly to find the most profitable synthesis pathway⁵.

In this study, we had structural modification of two *N*-(phenylcarbamothyoil)benzamide (PCTB) derivatives based on their lipophilic and electronic effects. *N*-(phenylcarbamothyoil)benzamide is known as cytotoxic thiourea derivatives compound that has better activity than hydroxyurea⁶. Tenovin 1 is a name given to thiourea derivatives 1-(4 acetamidophenyl)3-4-tertbutylbenzoyl) thiourea. It has been recognized to possess anticancer activity through synthesis and evaluation as suggested by another research. Also, it

owns a reputation to elevate the levels of p53 protein in vitro (which inhibits SIRT1)⁷. Moreover, N-(allylcarbamothioyl) benzamide is also evident to have activity as breast anticancer (T47D cells)⁸.

Aromatic ring modification in Topliss method only considers two properties, lipophilic and electronic⁹. Lipophilic properties affect drug penetration through the cell membrane and increase the amount of drug bound into a receptor, which increases its activity. While electronic properties have a role in drug solubility at the distribution and interaction of drug-receptor¹⁰.

To prove lipophilic and electronic effects of N-(phenylcarbamothioyl)benzamide on cytotoxic activity, we used 4-Nitro and 4-Methyl to represent each effect, respectively. Nitro has strong electronic effect ($\sigma = 0.78$) and methyl has strong lipophilic effect ($\pi = 0.56$)¹¹. Later on, we used two different cytotoxic activity methods on T47D, in silico and in vitro, to imply the effect on N-(phenylcarbamothioyl) benzamide.

EXPERIMENTAL

Molecular Modeling

The chemical structures were sketched using Marvin sketch software (Version 19.17.0) and converted to 3D Form for docking studies using Avogadro (Version 1.2.0). Merck Molecular Force Field (MMFF94), which optimizes the geometry's structures, was also utilized with Avogadro principles saved in .mol2.

In-silico Cytotoxic Activity Prediction

Molecular modeling of N-(phenylcarbamothioyl) benzamide was conducted to predict the activity. A molecular docking method with Histone deacetylase SIRT1 (ID= 4i5i) was performed by utilizing Autodock software. Retrieved from PDB bank server was the Sirtuin-1 (SIRT1) enzyme. As to provide protein molecule ligand, AutoDock Tool (version 1.5.6) was applied. Based on experimental conducts, grid box values were acquired to assign the perfect grid parameters in the ligand.

Synthesis

Synthesis had been done by reacting 0.03 mol N-phenylthiourea, 0.025 mol triethylamine, and 0.025 mol 4-Nitro/4-Methylbenzoyl chloride solution in tetrahydrofuran, followed by *refluxing* and stirring the mixture placed in a water bath and TLC, once per hour. Just as any spot appeared, the reaction was halted. Afterward, THF evaporation on the rotary was continued with recrystallization. The compound structure identification was conducted based on infrared spectroscopy, ¹H-NMR, ¹³C-NMR, and HRMS result.¹²⁻¹⁴

Cytotoxic Activity on T47D Cell

By employing T47D, the breast cancer cells, anticancer activity is conducted in vitro. To produce 50.000 µg/mL concentration, the test compound (5 mg) was fused in DMSO (100µl). To derive a group of standard working solution, the mother liquor was steadily diluted both by adding RPMI culture medium for T47D cells and M199 for Vero cells¹⁵. In a 96-wells microplate, a culture of T47D cancer cell was set in the contour of cell suspension. It was employed with 10,000 cells / pitting in terms of density for this study. For media control, the plate was added with empty wells followed by CO₂ 5% incubator, left for 24 hours. After the incubation period was over, the plate was taken away. With 180 degrees of reversion, the media was removed. Every single well was rinsed using 100µl PBS. Afterward, the plate was rotated 180 degrees and PBS was discarded. Set in microplate wells were the concentration of a standard working solution (100µL), positive control and solvent control. Each of the concentrations was then duplicated three times. As for media control, it was derived from the sums containing no T47D cancer cells, incubated 24 hours with 5% CO₂ incubator at 37°. Both the plate and the incubator were then discarded after that 24 hours. Added to every well was 0.5 mg/ml of MTT up to 100µL/well, incubated for three hours¹⁶. After the incubation and the microplate were removed, terminating the reaction was proceeded by putting 100 µL SDS 10% in 0.1 N HCl inside the hole. Covered with aluminum paper, the microplate was nurtured with 5% CO₂ in an incubator for 24 hours and inserted in Elisa Reader at $\lambda = 595$ nm. IC₅₀ values were obtained using probit analysis out of the test compound.¹⁷




RESULTS AND DISCUSSION

In-silico Cytotoxic Activity Prediction

In silico study, the result showed that both 4-NO₂-PCTB and 4-CH₃-PCTB has *Binding Score* (BS) lower than hydroxyurea (HU) as reference. Hence, it was predicted that they have better activity in sirtuin-1

receptor. According to Table-1, 4-NO₂-PCTB (-9.69 kcal/mol) has BS lower than other derivatives, 4-CH₃-PCTB (-8.34 kcal/mol). It inferred that 4-NO₂ substituent donates a strong and stable ligand-protein bond than 4-CH₃. Nitro substituent in para position has a stronger electronic effect compared to its lipophilic effect ($\sigma = 0.78$), while methyl in para position had stronger lipophilic effect than its electronic effect in QSAR parameter ($\pi = 0.56$)¹⁸. This was following previous research which states the electronic properties play a role in the stability of interactions between drugs and receptors.¹⁹

Table-1: Binding Score and Interaction Binding of 4-CH₃-PCTB, 4-NO₂-PCTB, and Hydroxyurea

| Substituent | Binding Score (Kcal/mol) | Interaction Binding |
|-------------------------|--------------------------|---|
| 4-CH ₃ -PCTB | -8.34 |  <p>Interactions</p> <ul style="list-style-type: none"> Conventional Hydrogen Bond Pi-Donor Hydrogen Bond Pi-Sigma Pi-Sulfur Pi-Pi T-shaped Pi-Alkyl |
| 4-NO ₂ -PCTB | -9.69 |  <p>Interactions</p> <ul style="list-style-type: none"> Conventional Hydrogen Bond Pi-Donor Hydrogen Bond Pi-Sigma Pi-Sulfur Pi-Pi T-shaped Pi-Alkyl |
| HU | -2.26 |  <p>Interactions</p> <ul style="list-style-type: none"> van der Waals Water Hydrogen Bond Conventional Hydrogen Bond Unfavorable Donor-Donor |

Another important result from Table-1 was the interaction of 4-NO₂-PCTB and 4-CH₃-PCTB with sirtuin-1. It showed that lower BS is affected by the quantity and the quality of occurring bond²⁰. In 4-NO₂-PCTB strengthened with 13 bonds, there are 6 hydrogen conventional bonds (GLNA345 (two bond in receptor), LYSA44, SERA422, HISA363, ASNA346), 3 *pi*-alkyl bonds (ILEA411, ALAA262, VALA445), 1 *pi*-*pi* T-shaped bond (PHE A297), 1 *pi*-sulfur bond (PHEA273), 1 *pi*-sigma bond (ILEA347), and 1 *pi* donor hydrogen bond (GLNA345). However, 4-CH₃-PCTB has higher BS because it was strengthened with 8 bonds; They are 2 hydrogen conventional bonds (GLNA345, HIS A363), 1 *pi* donor hydrogen bond (GLN A345), 1 *pi* sigma bond(ILE A347), 1 *pi* sulfur bond(PHEA273), 1 *pi*-*pi* T shaped bond (PHEA297), and 2 *pi*-alkyl bonds (ILEA411 and ALA A262). Therefore, in silico study result predicted that more bonds in ligand-receptor interaction create a stronger bond with lower binding score²¹.

Synthesis

The synthesis began with R-benzoyl Chloride (R=4-CH₃ and 4-NO₂) combined with N-phenylthiourea – a yellow light crystals luster and insoluble in water - in one step, from which the 4-CH₃-PCTB and 4-NO₂-PCTB compounds were derived. The structure identification of the synthesized compounds was conducted by IR, H NMR, C NMR and HRMS spectroscopy as mentioned in the following details:

N-(Phenylcarbamothioyl)-4-methyl benzamide

White crystal, yield 77%, m.p. 115–116 °C. ¹H NMR (DMSO-*d*₆, 500 MHz). δ 2.45 (s, 3H,CH₃); δ 7.29 (t, *J*=7.8 Hz, 1H,Ar-H); δ 7.33 (d, *J*=8.1 Hz, 2H, Ar-H); δ 7.42 (t, *J*=7.8 Hz, 2H, Ar-H); δ 7.71 (d, *J*=7.8 Hz, 2H, Ar-H); δ 7.79 (d, *J*=8.1 Hz, 2H, Ar-H); δ 9.10 (s, 1H,O=C-NH-C=S); δ 12.64 (s, 1H,S=C-NH-Ar). ¹³C NMR (DMSO-*d*₆, 125 MHz). δ 21.8 (1C, CH₃); δ 124.3 (2C, Ar); δ 127.0 (2C, Ar); δ 127.7 (1C, Ar); 128.8 (2C, Ar); δ 129.0 (2C, Ar); 130.0 (1C, Ar); δ 137.8 (1C, Ar); δ 145.0 (1C, Ar); δ 167.0 (1C, C=O); δ 178.6 (1C, C=S). IR (KBr), ν maks(cm⁻¹): 1672 (C=O amide); 1603 dan 1496 (C=C Aromatic); 3313 and 1603 (NH stretch sec. amides); 1075 dan 807 (C=S). HRMS (*m/z*) C₁₅H₁₃N₂OS: (M-H)⁻ = 269.0757 and Calc. Mass = 269.0749.

N-(Phenylcarbamothioyl)-4-nitro benzamide

Yellow crystal, yield 65 %, m.p. 129–130 °C. ¹H NMR (DMSO-*d*₆, 500 MHz) δ 6.94 (t, *J*= 8.0, 1H, Ar-H); δ 7.12 (t, *J*=8.0 Hz, 2H, Ar-H); δ 7.37 (d, *J*=8.0 Hz, 2H, Ar-H); δ 7.76 (d, *J*=8.0 Hz, 2H, Ar-H); δ 8.35 (d, *J*=8.0 Hz, 2H, Ar-H); δ 9.56 (s, 1H, O=C-NH-C=S); δ 10.54 (s, 1H, S=C-NH-Ar). ¹³C NMR (DMSO-*d*₆, 125 MHz) δ 118.7 (2C, Ar); δ 121.1 (2C, Ar); δ 122.4 (1C, Ar); δ 122.5 (2C, Ar); δ 124.1 (2C, Ar); δ 124.9 (1C, Ar); δ 129.6 (1C, Ar); δ 140.3 (1C, Ar); δ 160.3 (1C, C=O); δ 171.2 (1C, C=S). IR (KBr), ν maks(cm⁻¹): 1651 (C=O amide); 1635 and 1497 (C=C aromatic); 3322 and 1635 (NH stretch sec amides); 1104 and 830 (C=S). HRMS (*m/z*) C₁₄H₁₂N₃O₃S: (M+H)⁺ = 302.0536 and Calc. Mass = 302.0534.

Cytotoxic Activity

In vitro MTT Assay in T47D result showed the same activity to in silico study. IC₅₀ 4-NO₂-PCTB (0.12 mM) was lower than 4-CH₃-PCTB (1.08 mM). In other words, 4-NO₂-PCTB is more potent than 4-CH₃-PCTB as an anticancer agent in T47D. The effect of the electron withdrawal group is stronger than the electron contributor group.

Both of 4-NO₂-PCTB and 4-CH₃-PCTB had better activity than hydroxyurea as reference. They also had cancer cell-targeted mechanisms from the cytotoxic assay in Vero cell as a normal cell. The result was both of them toxic in T47D cell but not in vero cell. Thus, they are worthy to develop as an anticancer agent in T47D cells with mechanism prediction of sirtuin-1 receptor inhibition.

Table-2: RS, IC₅₀ T47D, and Vero Cells Values of Test Compounds and Reference Compounds

| Compounds | RS (Kcal/mol) | IC ₅₀ T47D Cells(mM) | IC ₅₀ Vero Cells(mM) |
|-------------------------|---------------|---------------------------------|---------------------------------|
| 4-CH ₃ -PCTB | -8.34 | 1.08 ± 0.013 | 53.58±0.004 |
| 4-NO ₂ -PCTB | -9.69 | 0.12± 0.014 | 65.63±0.003 |
| HU | -2.26 | 4.58± 0.019 | 369.88±0.015 |

To develop another derivative of *N*-(phenylcarbamothioyl)benzamide, structural modification in the aromatic compound must be considered in the selection of moiety with another electron withdrawal group with a higher electronic parameter σ (+) but slightly higher lipophilic parameter (π). It should still use the Lipinski rule because the activity is affected by the physicochemical characteristic of the moiety. Furthermore, in silico study in both derivatives supports invitro cytotoxic activity assay. Hence, in silico study can be used as activity prediction before synthesis and in vitro study.

CONCLUSION

In *N*-(phenylcarbamothioyl)benzamide derivatives compounds, nitro substituent has a stronger cytotoxic effect than methyl. It can be concluded that the *N*-(phenylcarbamothioyl) benzamide derivatives compound has a stronger electronic effect than the lipophilic effect. This can be used to develop more anticancer agent compounds from this lead compound.

ACKNOWLEDGEMENT

Funding support for this study was from Indonesia's Doctorate General of Resources for Science, Technology and Higher Education under the Ministry of Research, Technology and Higher Education (KEMENRISTEK DIKTI) following the scholarship funding scheme for PhD program at University of Airlangga, Surabaya Indonesia.

REFERENCES

1. J.M. Beale and J.H. Block, *Wilson and Gisvolds Textbook of Organic Medicinal and Pharmaceutical Chemistry 12th ed*, Lippincott Williams & Wilkins, Philadelphia, (2011)
2. S. Siswandono, *Pengembangan Obat Baru*, Airlangga University Press, Surabaya, (2014)
3. G. Thomas, *Medicinal Chemistry 2nd ed*, John Wiley & Sons, New York, (2007)
4. J.G. Topliss, *Journal of Medicinal Chemistry*, **15**(10), 1006(1972), DOI:10.1021/jm00280a002
5. G.L. Patrick, *An Introduction to Medicinal Chemistry 5thed*, Oxford University Press, New York, (2013)
6. D. Kesuma, Siswandono, B.T. Purwanto and M. Rudyanto, *Journal Chinese Pharmaceutical Science*, **29**(2), 123(2020), DOI:10.5246/jcps.2020.02.010
7. A.R. McCarthy, L. Pirrie, J.J. Hollick, S. Ronseaux, J.Campbell, M. Higgins, O.D.Staples, F. Tran, A.M.Z. Slawin, S.Lain and N.J.Westwodd, *Bioorganic & Medicinal Chemistry*, **20**,1779(2012), DOI:10.1016/j.bmc.2012.01.001
8. T. Widiandani, S. Siswandono, E. Meiyanto, M. Ika, BT. Purwanto and S. Hardjono, *Tropical Journal of Pharmaceutical Research*, **17**(8), 1607(2018), DOI:10.4314/tjpr.v17i8.20
9. D. Abraham, *Burger's Medicinal Chemistry and Drug Discovery 6thed Volume 1: Drug Discovery*, John Wiley and Sons, New York, (2003)
10. A.L. Nasyanka, S. Siswandono and S. Hardjono, *Thai Journal Pharmaceutical Sciences*, **41**(3), 99(2017)
11. S. Siswandono, *Kimia Medisinal I*, Airlangga University Press, Surabaya, (2016)
12. J. Clayden, N. Greeves, S. Warren and P. Wothers, *Organic Chemistry 2nded*, Oxford University Press, New York, (2012)
13. J.M. McMurry, *Fundamental of Organic Chemistry 7thed*, Brooks/Cole, Belmont, (2011)
14. D.L. Pavia, G.M. Lampman, G.S. Kriz, R. James and J.R. Vyvyan, *Spectroscopy 4thed*, Brooks/Cole, Belmont, (2009)
15. D. Satria, J. Silalahi, G. Haro, S. Ilyas and P.A.Z. Hasibuan, *Rasayan Journal Chemistry*, **12**(2), 803(2019), DOI:10.31788/RJC.2019.1225180
16. V. Venkataramireddy, M. Shankaraiah, A. Tejeswara Rao, C. Kalyani, M. L.Narasu, R.Varala and A. Jayashree, *Rasayan Journal Chemistry*, **9**(1), 31(2016)
17. <http://ccrc.farmasi.ugm.ac.id/wp-content/uploads/03.010.02-uji-sitotoksik-MTT.pdf>
18. C. Avendano and J.C. Menendes, *Medicinal Chemistry of Anticancer Drugs 2nd ed.*, Elsevier, Amsterdam, (2015)
19. A. Bendjeddou, T. Abbaz, S. Maache, R. Rehamnia, A. K. Gouasmia, and D. Villemin, *Rasayan Journal Chemistry*, **9**(1), 18(2016)
20. X. Du, Y. Li, Y. L. Xia, S. M. Ai, J. Liang, P. Sang, and S. Q. Liu, *International Journal of Molecular Sciences*, **17**(2), 144(2016), DOI: 10.3390/ijms17020144
21. P. L. Kastritisand A. M. J. J.Bonvin, *Journal of Royal Society Interface*, **10**(20120835), 1(2012), DOI: 10.1098/rsif.2012.0835

[RJC-5694/2020]

RASAYAN

JOURNAL OF CHEMISTRY

An International Quarterly Research Journal of Chemical Sciences

Rasayan J. Chem. ISSN: 0974-1496(Print) ISSN:
0976-0083(Online) CODEN: RJCABP Frequency: March,
June, September & December

SCOPUS
Indexed Since, 2008
&
UGC
Approved

[HOME](#) | [ABOUT US](#) | [EDITORIAL BOARD](#) | [SUBSCRIPTION](#) | [ABSTRACTED](#) | [CONTACT](#) | [PUBLICATION PROCESS](#) | [MANUSCRIPT SUBMISSION](#)

Welcome to RJC

RASAYAN means Chemical, which is an incredibly reactive and self explanatory term in itself. **RASAYAN** is an open access as well as print journal and widely covers all branches of Chemistry including : Organic, Inorganic, Physical, Analytical, Biological, Pharmaceutical, Industrial, Environmental, Agricultural & Soil, Petroleum, Polymers, Nanotechnology, Green Chemistry, Forensic, Phytochemistry, Synthetic Drugs, Computational, as well as Chemical Physics and Chemical Engineering. In online version of **RASAYAN** all figures and photographs are colored with printable clarity.

RJC is Abstracted / Indexed in



ELSEVIER
SCOPUS



A division of the American Chemical Society



CAB
INTERNATIONAL




ILLUMINA

Deep Indexing added to selected databases



Google
scholar



[Current Issue](#) →

[Archive Issues](#) →

SCIRUS
www.scirus.com

ULRICHSWEB™
GLOBAL SERIALS DIRECTORY

SJR SCImago
Journal & Country
Rank

EBSCO
INFORMATION SERVICES

ROAD DIRECTORY
OF OPEN ACCESS
SCIENTIFIC
RESOURCES

crossref

Plagiarism Check Partner



1.8 CiteScore²⁰¹⁹

56th percentile

Powered by Scopus

[Downloads](#) →

[RJC Conferences](#) →

SJR SCImago
Journal & Country
Rank

Rasayan Journal of
Chemistry

Q2

Pharmacology,
Toxicology and
Pharmaceutics...
best quartile

SJR 2019
0.27

powered by scimagojr.com

Editorial Board



Editor-in-Chief:

Prof. (Dr.) Sanjay K. Sharma

Managing Editor:

Dr. Pratima Sharma

Editorial Office:

23 'Anukampa', Janakpuri,
Opp. Heerapura Power Stn.,
Ajmer Road, Jaipur-302024 (India)

Mobile; 09414202678, 9001699997

E-mail: rasayanjournal@gmail.com

Nnabuk Okon EDDY

Professor, Department of Chemistry,
Ahmadu Bello University, Zaria, Kaduna
State, Nigeria.

Goutam BRAHMACHARI

Professor, Chemistry Department, Visva-
Bharati University, Santiniketan-731235,
India.

Man SINGH

Professor and Dean, school of Chemical
sciences, Gujrat central University,
Gandhinagar, Gujrat, India

Susheel MITTAL

Senior Professor, School of Chemistry
& Biochemistry, Thapar Institute of
Engineering & Technology (Deemed
to be University), Bhadson Road,
Patiala-147004, India

V.K. GARG

Professor and Dean Centre for
Environmental Science and Technology
School of Environment and Earth
Sciences Central University of Punjab,
Bathinda- 151001, India

S. P.MOULIK

Emeritus Professor & Hony Scientist (INSA),
Centre for Surface Science, Department of
Chemistry, Jadavpur university, Kolkata –
700 032, India

Fernando de Carvalho DA SILVA

Professor Adjunto III, UNIVERSIDADE
FEDERAL FLUMINENSE Instituto de
Química, Departamento de Química
Orgânica, Sala 509, Centro, Niterói-RJ,
Brazil.

Abdulahakim A. AHMED Department of
Chemistry, Faculty of Science, University of
Benghazi- Libya, P.O.Box 1308

Roberto BALLINI

School of Science and Technology,
University of Camerino – Italy

Marei Mailoud EL-AJAILY

University of Benghazi, Faculty of Science ,
Department of Chemistry, Benghazi, Libya

Soro YAYA

Laboratoire des Procédés Industriels
de Synthèse, de l'Environnement et des
Energies Nouvelles (LAPISEN), Institut
National Polytechnique (INP-
HB), Yamoussoukro, BP 991
Yamoussoukro(Côte d'Ivoire)

William E. ACREE

Department of Chemistry, 1155 Union
Circle Drive #305070, University of
North Texas, Denton, TX 76203-5017
(USA)

Junfang ZHAO

Division of Pathology and Laboratory
Medicine, Cincinnati Children's
Hospital Medical Center, 3333 Burnet
Ave. Cincinnati, OH, USA 45208

Catalina PISOSCHI

Department of Biochemistry, University
of Medicine and Pharmacy, Craiova,
Romania

Hakan ARSLAN

Department of Chemistry, Faculty of Arts and Science, Mersin University, Mersin, TR-33343, Turkey

R.V. SINGH

Ex Professor, Department of Chemistry, University of Rajasthan, Jaipur, India

Dong CHEN

Department of Engineering, Indiana University - Purdue University Fort Wayne, 2101 E Coliseum Blvd Fort Wayne, IN 46805, USA

Giusy LOFRANO

Department of Environment, University of Salerno, Salerno, Italy

Essam Khamis Ibrahim ALL-HANASH

City of Scientific Research and Technological Applications (SRTA-City), Universities and Research Center District, New Borg El- Arab, Egypt. P.O. Box: 21934 ALEX

Willian Aperador CHAPARRO

School of Engineering, Universidad Militar Nueva Granada, Bogotá-111121, Colombia

Ishmael MASESANE

Professor, Department of Chemistry, University of Botswana, Botswana

Eno E. EBENSO

Professor, North-West University Gauteng, South Africa

Pankaj KUMAR

Professor and Head, Department of Chemistry, University of Energy and Petroleum studies, Dehradun, India

Virendra GOMASE

Department of Bioinformatics, Padmashree Dr. D.Y. Patil University, Navi Mumbai, 400614, India

Pavel MOKREJS

Department of Polymeric Engineering, Tomas Bata University, 762 72, Zlin, The Czech Republic

Nicola D'ANATONA

National Research Council of Italy, Institute of Biomolecular Chemistry, Italy

Sadanandan E. VELU

Department of Chemistry, The University of Alabama at Birmingham, 901, 14th Street South, Birmingham, AL - 35294 - 1240

Y. Frank CHENG

Corrosion and Pipeline Engineering Laboratory, Dept. of Mechanical and Manufacturing Engineering, Schulich School of Engineering, University of Calgary, Calgary, Alberta, Canada T2N 1N4

Tingyue GU

Dept. of Chemical and Biomolecular Engineering Ohio University Athens, Ohio 45701, USA

Surendra PRASAD

School of Biological and Chemical Sciences, The University of the South Pacific, Suva, FIJI

Mika SILLANPÄÄ

Department of Chemical Engineering, School of Mining, Metallurgy and Chemical Engineering, University of Johannesburg, Doornfontein 2028, South Africa

Current Issue



Volume 13, Number 3, July - September (2020)

REMOVAL OF ARSENIC FROM WATER VIA SOLAR ENERGY HARVESTING TECHNIQUE WITH INTEGRATED SYSTEM OF ACTIVATED ALUMINA ABSORBENT BED AND DOUBLE SLOPE SOLAR STILL

Suresh Kumar Patel, Pradeep Kumar, G. Srinivasan and Dhananjay Singh

Rasayan J. Chem, 13 (3), 1293 - 1307 (2020)

Keywords: Arsenic, Adsorption, Activated Alumina, Solar Distillation, Solar Energy Harvesting.

DOI: <http://dx.doi.org/10.31788/RJC.2020.1335725>

HUMATES FROM COAL MINING WASTE: SYNTHESIS, STUDY OF COMPOSITION AND RADIOACTIVITY

M. Kambatyrov, U. Nazarbek, P. Abdurazova, S. Nazarbekova and Y. Raiymbekov

Rasayan J. Chem, 13 (3), 1308 - 1312 (2020)

Keywords: Coal Mining Waste, Humate, Radioactivity, Feed Additives, Calcium Humate.

DOI: <http://dx.doi.org/10.31788/RJC.2020.1335729>

ATENOLOL IMPRINTED POLYMER WITH BUTANOL AS A POROGENIC SOLVENT: USING TWO POLYMERIZATION METHODS

A. N.Hasanah, J. Tristi and D. Rahayu

Rasayan J. Chem, 13 (3), 1313 - 1320 (2020)

Keywords: Atenolol, Molecular Imprinting Polymer, Solid-phase Extraction, Methyl Methacrylate, Butanol

DOI: <http://dx.doi.org/10.31788/RJC.2020.1335765>

MOLECULAR INTERACTION OF THE DOWNSTREAM EXECUTIONER CYSTEINE ASPARTYL PROTEASES (CASPASE-3 AND CASPASE-7) WITH CORILAGIN, QUERCETIN, RUTIN, KAEMPFEROL, GALLIC ACID, AND GERANIIN OF *Acalypha wilkesiana* Müll.Arg.

S. Megantara, M. Mutakin, E. Halimah, E. Febrina and J. Levita

Rasayan J. Chem, 13 (3), 1321 - 1329 (2020)

Keywords: Apoptosis, Aspartic Acid, Flavonoids, Caspase, Kaempferol, Quercetin, Rutin

DOI: <http://dx.doi.org/10.31788/RJC.2020.1335766>

ISOLATION AND CHARACTERISATION OF FORCED DEGRADATION PRODUCTS OF A CALCIUM CHANNEL BLOCKER BY LC/MS TECHNIQUE

L. Sawaikar and P. Kapupara

Rasayan J. Chem, 13 (3), 1330 - 1334 (2020)

Keywords: Cilnidipine, LC/MS, Forced Degradation

DOI: <http://dx.doi.org/10.31788/RJC.2020.1335855>

SYNTHESIS AND CHARACTERIZATION OF CERAMIC MEMBRANE FROM FLY ASH AND CLAY PREPARED BY SINTERING METHOD AT LOW TEMPERATURE

Selvie Diana, Reza Fauzan, Nasrul Arahman, Fachrul Razi and Muhammad Roil Bilad

Rasayan J. Chem, 13 (3), 1335 - 1341 (2020)

Keywords: Clay, Fly Ash, Ceramic Membrane, Low Temperature, Sintering.

DOI: <http://dx.doi.org/10.31788/RJC.2020.1335707>

COMPOSITE GALVANIC COATINGS BASED ON ZINC

N. A. Vysotskaya, B. N. Kabyzbekova, K.A. Bekzhigitova, L.D. Aikozova, G.M.Adyrbekova, L.A. Zhurhabayeva and E. N. Abdulova

Rasayan J. Chem, 13 (3), 1342 - 1346 (2020)

Keywords: Composite Zinc Coatings, Thiourea, Gelatin, BGU-12, Energy Dispersion Analysis

DOI: <http://dx.doi.org/10.31788/RJC.2020.1335841>

FORMULATION AND CHARACTERIZATION OF QUERCITRIN NANOEMULSION ISOLATED FROM *Dendrophoe falcata* AND ITS ANTIOXIDANT ACTIVITY TEST

S.Atun, R. Arianingrum, L. Cahyaningsih, F. A. Pratiwi, R. Kusumaningrum, and M. Khairuddean

Rasayan J. Chem, 13 (3), 1347 - 1356 (2020)

Keywords: Antioxidant, *Dendrophoe falcata*, Nanoemulsion, Quercitrin

DOI: <http://dx.doi.org/10.31788/RJC.2020.1335868>

A PROSPECTIVE MODIFICATION STRUCTURE: THE EFFECT OF LIPOPHILIC AND ELECTRONIC PROPERTIES OF N-(PHENYLCARBAMOTHYOIL)BENZAMIDE DERIVATIVES ON CYTOTOXIC ACTIVITY BY IN SILICO AND IN VITRO ASSAY WITH T47D CELLS

D. Kesuma, A.L. Nasyanka, M. Rudyanto, Siswandono, B.T. Purwanto and I.G.A. Sumartha

Rasayan J. Chem, 13 (3), 1914 - 1918 (2020)

Keywords: Lipophilic Effect, Electronic Effect, N-(phenylcarbamothyoil)Benzamide, T47D Cell, In-silico

DOI: <http://dx.doi.org/10.31788/RJC.2020.1335694>

THE OPERATIONAL PARAMETERS EFFECT ON PHOTOCATALYTIC DEGRADATION OF DIAZINON USING CARBON AND NITROGEN MODIFIED TiO₂

K. Khoiriah, S. Safni, S. Syukri, and J. Gunlazuardi

Rasayan J. Chem, 13 (3), 1919 - 1925 (2020)

Keywords: C,N-codoped TiO₂, Diazinon, Photocatalysis, Mineralization

DOI: <http://dx.doi.org/10.31788/RJC.2020.1335743>

ANTIBACTERIAL STUDIES OF NEWLY SYNTHESIZED 3d MACROCYCLES

S. Sharma, R. Mehani ne'e Chopra and V. Chugh

Rasayan J. Chem, 13 (3), 1926 - 1932 (2020)

Keywords: Hexane-1, 6-diamine, Metal Halide, 2, 2-Dihydroxyindane-1, 3-dione, Physico-chemical Techniques, Antibacterial Screening.

DOI: <http://dx.doi.org/10.31788/RJC.2020.1335629>
