

## EFFECT OF RED MULBERRY JUICE (*MORUS RUBRA*) ON THE INFLAMMATORY RESPONSE IN MALE WISTAR RATS DUE TO EXPOSURE TO CIGARETTE SMOKE

Rivan Virlando Suryadinata<sup>1\*</sup>, Merryana Adriani<sup>2</sup>, Stefani Cornelia Sardjono<sup>3</sup>

<sup>1</sup>Faculty of Medicine, University of Surabaya (UBAYA), Surabaya

<sup>2</sup>Department of Nutrition, Faculty of Public Health, Universitas Airlangga, Surabaya

<sup>3</sup>Students of the Medical Study Program, University of Surabaya (UBAYA), Surabaya

\*E-mail: rivan.virlando.suryadinata@gmail.com

### ABSTRACT

The increasing number of tobacco users will directly impact to an increase in the number of passive smokers. Various harmful substances that enter the respiratory tract can cause health problems. The airway will recognize these substances as pathogen and activate immune response accordingly. Inflammatory response becomes one of the effects of body compensation due to an increased pathogen. This effect is indicated by the elevation of C-Reactive Protein as one systemic inflammatory response parameters in the body. This study aimed to determine the efficacy of red mulberry juice to reduce levels of C-Reactive Protein in Wistar rats because exposure to cigarette smoke. This is an experimental study with a post-test control group design. Animals were divided into negative control group, positive control group, and 3 treatment groups with different red mulberry doses. The positive control group showed the highest increase in C-Reactive Protein levels compared with the red mulberry intake group ( $p < 0,001$ ). In conclusion, the administration of red mulberry juice can reduce inflammatory response in rats because of exposure to cigarette smoke.

**Keywords:** c-reactive protein, cigarette, inflammation, red mulberry

### INTRODUCTION

Tobacco cigarettes are the most common type of cigarette used worldwide (Gowing et al., 2015). Various studies have shown the dangers of using tobacco cigarettes both as active smokers and passive smokers (Suryadinata et al., 2017). Smoking has caused 6 million people dies each year and it is estimated will reach 8 million in 2030. Globally, the number of tobacco smokers has reached 1.1 billion people and almost 70% are residents of developing countries (Nasser et al, 2018). The prevalence of active smokers in developed countries has shown a significant decrease. Estimated in 2015, number of smokers in developed countries like America and Europe was around 15% of total population (West, 2017). In developing countries, the number of smokers increases every year, especially in the productive age (Hossain et al., 2017).

Smoke contains more than 4000 types various ingredients, including nicotine, ammonia, phenols, polycyclic aromatics, carbon monoxide, nitrogen oxides, and various metals (Kamceva et al., 2016). Two main phases that can be identified in cigarette smoke are tar phase and gas phase. Both phases

contain high free radicals such as superoxide ( $O_2^{\bullet-}$ ) and hydroxyl ( $^{\bullet}OH$ ) (Suryadinata, 2018). Increased free radicals in body due to exposure to smoke cigarettes entering airways will cause oxidative stress and trigger lipid peroxidation. This will potentially cause cell damage and necrosis (Suryadinata et al., 2019). Normally, free radicals are mitochondrial metabolism and can be neutralized by enzymatic antioxidants in body (Suryadinata et al, 2019). In pathological conditions, excessive free radicals will directly stimulate inflammatory response through phagocytic process of macrophages (Arulselvan et al., 2016).

Inflammatory process is a form of body defence mechanism against pathogenic reactions from outside. Various pro-inflammatory cytokines are secreted into body microenvironment such as interleukin-1, interleukin-6, interleukin-8, and tumor necrosis factor-alpha ( $TNF-\alpha$ ). Inflammatory mediator will induce liver to release C-Reactive Protein (CRP) into the blood (Phaniendra et al., 2015). Prevention of inflammatory process due to free radical increase in the body can be done with administration of antioxidants from the outside.

Various types of non-enzymatic antioxidants can be obtained from food especially flavonoids (Pratiwi et al., 2018).

Mulberry fruit is one type of fruit that high in antioxidants. Mulberries originated from genus *Morus* of Moraceae family, to date there are 24 species and 100 varieties (Zhang et al., 2018). Various studies have shown the provision of mulberry fruit has potential impact as antioxidant (Zhang and Shi, 2010). This study used red mulberry juice (*Morus rubra*) juice to reduce body inflammatory response due to exposure to tobacco smoke. Besides, red mulberries are also widely planted and consumed by Indonesian people so that at the same time can preserve local food. This study purpose is to analyze the efficacy of red mulberry juice to reduce levels of C-Reactive Protein in Wistar rats exposed to cigarette smoke.

## METHODS

This research was an experimental study with post-test control group design performed in 25 male Wistar rats (*Rattus norvegicus*) which were divided into 5 groups for 30 days. Inclusion criteria in this study were experimental animals aged 3-4 months with weight around 200 g. While the exclusion criteria are animals get sick or die during study. Maintenance and care of the animals during study are based on 3R principles (Replacement, Reduction, and Refinement). This research has been obtained Ethical Clearance Certificate from Institutional Ethical Committee, University of Surabaya (No:135/KE/VI/2020).

Red mulberries (*Morus rubra*) that have been obtained, sorted and cleaned in order to get fruit in good condition. Red mulberry juice that has been obtained then performed flavonoid levels measurement ( $\pm 38$  mg/g). Results of flavonoids levels that have been obtained then converted from daily needs of adults ( $\pm 190$  mg/kg) into daily dose intake of Wistar rats with 200g body weight ( $\pm 240$  mg/rats) using *Laurence and Bacharach* table. The amount of red mulberry juice given orally to the treatment group was 3.25 g, 6.5 g, and 13 g.

The results of CRP level measurement are based on agglutination process in blood through intracardiac puncture. Dilution process show the results of C-Reactive Protein measurement i.e., >

6 mg/L, 12 mg/L, 24 mg/L, 48 mg/L and 96 mg/L. Measurement results were classified as positive if C-Reactive protein levels obtained more than 6 mg/L, and negative if C-Reactive Protein level obtained were less than 6 mg/L.

This research was conducted for 30 days by dividing rats into 5 groups. The first was the negative control group, where animals did not received any treatment. In positive control group, animals received daily intake and exposure to 2 tobacco cigarettes smoke per day in smoking chamber. While last three groups were exposed to 2 tobacco cigarettes smoke per day, daily intake, and 3.25 g, 6.5 g, or 13 g red mulberry juice.

Results of study were ordinal data in the form of CRP levels in each group in mg/L units. CRP measurement method uses dilution to get a multiple of the result. CRP levels of >6 mg/L, 12 mg/L, 24 mg/L and 48 mg/L will be carried out for coding. Data analysis was performed using Kruskal Wallis analysis with SPSS version 22 to observe differences between groups.

## RESULTS AND DISCUSSIONS

Results of study were carried out by comparing C-Reactive Protein levels between groups. Table 1. shows results of C-Reactive Protein level measurement in blood between group. Group I, IV, and V had the same levels of C-Reactive Protein. The highest level of C-Reactive Protein obtained in Group II which is the Positive Control Group.

Kruskal Wallis test in Table 1 shows p-value of C-Reactive Protein levels between group was <0,001 (*p-value* <0.05), so it can be concluded that there are significant differences of C-Reactive Protein levels between groups.

Exposure to free radicals from tobacco smoke that enters respiratory tract causes body develop defence mechanisms to prevent the impact of cell damage due to oxidative stress (Indraswari et al., 2018). Antioxidant produced in body are enzymatic antioxidants (superoxide dismutase [SOD], glutathione peroxidase [GSH-Px], and catalase), while antioxidant from outside are non-enzymatic antioxidant derived from food intake such as flavonoids (Suryadinata et al., 2016). The balance between number of free radicals and

**Table 1.** Levels of C-Reactive Protein in research

Group		C-Reactive Protein levels (mg / L) in Wistar Rats					Kruskal Wallis Test
		1 <sup>st</sup>	2 <sup>nd</sup>	3 <sup>rd</sup>	4 <sup>th</sup>	5 <sup>th</sup>	
I	The negative control group without exposure to cigarette smoke and intake of red mulberry juice	<6	<6	<6	<6	<6	
II	The positive control group received exposure to smoke cigarettes and no intake of red mulberry juice	48	48	48	24	24	
III	Treatment group I was exposed to cigarette smoke and red mulberry juice intake of 3.25 g	24	12	12	<6	<6	<0.001
IV	Treatment group II received smoke exposure cigarettes and red mulberry juice intake of 6.5 g	<6	<6	<6	<6	<6	
V	Treatment group III received smoke exposure cigarettes and red mulberry juice intake of 13 g	<6	<6	<6	<6	<6	

antioxidants is needed for the stability of body normal metabolic process (Biswas et al., 2017).

Reactive Oxygen Species (ROS) are the most reactive and radical types that directly respond to inflammation in body. These radicals can be neutralized with superoxide dismutase (SOD) into hydrogen peroxidase (H2O2) then will be neutralized with glutathione peroxidase (GSH-Px) and catalase into water (H2O) and oxygen (O2) (Nimse and Pal, 2015).

However, an increase in excessive free radicals can result in cell damage or necrosis. This process will cause cells to become lysis and is called debris cells (Vénéreau et al., 2015). This will stimulate the movement of macrophages as one mechanism of body defence. Phagocytic process carried by macrophages will trigger secretion of pro-inflammatory cytokines such as interleukin-1, interleukin-6, and tumor necrosis factor-alpha (TNF-α) (Thomsen et al., 2016). Excessive release of proinflammatory cytokines will induce liver to secrete C-Reactive Protein (Sproston and Ashworth, 2018).

Inflammatory processes in the body have positive and negative effects. In acute and low condition, Inflammatory processes will function as a defence and healing process for cell tissue, while if the condition is chronic, then it will have impact on cell damage. Exposure to cigarette smoke that given regularly and continuously will create chronic inflammatory conditions that have a negative impact on health (Srivastava and Kumar, 2015). Therefore, the administration of antioxidants through food intake is expected to

be able to prevent the inflammatory process by decreasing free radicals.

Administration of red mulberry juice (*Morus rubra*) which rich in flavonoids shows its efficacy on decreasing levels of C-Reactive Protein in blood due to exposure to tobacco smoke. Therefore, the use of red mulberries as an additional intake of antioxidants from outside body can be considered as one way to prevent cell damage due to inflammatory process.

CONCLUSION

Exposure to tobacco smoke can increase body inflammatory response through increased levels of C-Reactive Protein in the blood. Giving red mulberry juice can reduce the response of inflammation caused by exposure to tobacco smoke in rats.

ACKNOWLEDGEMENT

The researcher is grateful to all laboratory staff who have assisted the research process.

REFERENCES

Arulselvan, P., Fard, M. T., Tan, W. S., Gothai, S., Fakurazi, S., Norhaizan, M. E., ... Kumar, S.S. (2016). Role of antioxidants and natural products in inflammation. *Oxidative Medicine and Cellular Longevity*, 2016, 1–15. doi: 10.1155/2016/5276130

Biswas, S., Das, R., & Ray Banerjee, E. (2017). Role of free radicals in human inflammatory

- diseases. *AIMS Biophysics*, 4(4), 596–614. doi: 10.3934/biophy.2017.4.596
- Gowing, L. R., Ali, R. L., Allsop, S., Marsden, J., Turf, E. E., West, R., & Witton, J. (2015). Global statistics on addictive behaviours: 2014 status report. *Addiction*, 110(6), 904–919. doi: 10.1111/add.12899.
- Hossain, S., Hossain, S., Ahmed, F., Islam, R., Sikder, T., & Rahman, A. (2017). Prevalence tobacco smoking and factors associated with the initiation of smoking among university students in Dhaka, Bangladesh. *Central Asian Journal of Global Health*, 6(1). doi: 10.5195/cajgh.2017.244
- Indraswari, P. I. I., Lorensia, A., and Suryadinata, R. V. (2018). Analysis effect of nutrition intake on lung function of active smoker and non smoker. *Jurnal Kesehatan Masyarakat*, 14(2), 247–253. Retrieved from <https://journal.unnes.ac.id/nju/index.php/kemas/article/view/14947>
- Kamceva, G., Arsova-Sarafinovska, Z., Ruskovska, T., Zdravkovska, M., Kamceva-Panova, L., & Stikova, E. (2016). Cigarette smoking and oxidative stress in patients with coronary artery disease. *Open Access Macedonian Journal of Medical Sciences*, 4(4), 636. doi: 10.3889/oamjms.2016.117
- Nasser, A., Salah, B., Regassa, L., Alhakimy, A., & Zhang, X. (2018). Smoking prevalence, attitudes and associated factors among students in health-related Departments of Community College in rural Yemen. *Tobacco Induced Diseases*, 16(July). doi: 10.18332/tid/92547
- Nimse, S. B. & Pal, D. (2015). Free radicals, natural antioxidants, and their reaction mechanisms. *RSC Advances*, 5(35), 27986–28006. doi: 10.1039/C4RA13315C
- Phaniendra, A., Jestadi, D. B., & Periyasamy, L. (2015). Free radicals: properties, sources, targets, and their implication in various diseases. *Indian Journal of Clinical Biochemistry*, 30(1), 11–26. doi: 10.1007/s12291-014-0446-0
- Pratiwi, S. R., Lorensia, A., & Suryadinata, R. V. (2018). Vitamin C and E Intake with SQ-FFQ towards Smokers' and Non-Smokers' Lung Function. *Media Kesehatan Masyarakat Indonesia*, 14(2), 101. Retrieved from <http://journal.unhas.ac.id/index.php/mkmi/issue/view/559>
- Sproston, N. R., & Ashworth, J. J. (2018). Role of c-reactive protein at sites of inflammation and infection. *Frontiers in Immunology*, 9. doi: 10.3389/fimmu.2018.00754
- Srivastava, K. K., & Kumar, R. (2015). Stress, oxidative injury and disease. *Indian Journal of Clinical Biochemistry*, 30(1), 3–10. doi: 10.1007/s12291-014-0441-5
- Suryadinata, Rivan Virlando, Wirjatmadi, B. (2019). The role of selenium micronutrients as antioxidants in exposure to e-cigarette smoke. *Asian Journal of Pharmaceutical and Clinical Research*, 12(8), 265–268. Retrieved from <https://innovareacademics.in/journals/index.php/ajpcr/issue/view/466>
- Suryadinata, R. V., Wirjatmadi, B., & Adriani, M. (2016). Pengaruh perubahan hiperplasia sel goblet selama 28 hari paparan asap rokok dengan pemberian antioksidan superoxide dismutase. *The Indonesian Journal of Public Health*, 11(1), 60. Retrieved from 10.20473/ijph.v11i1.2016.60-68
- Suryadinata, R. V., Wirjatmadi, B., & Adriani, M. (2017). Efektivitas penurunan malondialdehyde dengan kombinasi suplemen antioksidan superoxide dismutase melon dan gliadin akibat paparan rokok. *Global Medical And Helath Communication*, 5(2), 79–83. Retrieved from
- Suryadinata, R. V., Wirjatmadi, B., Adriani, M., and Sumarmi, S. (2019). The effects of exposure duration to electronic cigarette smoke on differences in superoxide dismutase and malondialdehyde in blood of wistar rats. *International Journal of Current Pharmaceutical Research*, 11(3), 13–16. doi: 10.22159/ijcpr.2019v11i3.34084
- Thomsen, M., Kersten, C., Sorbye, H., Skovlund, E., Glimelius, B., Pfeiffer, P., et al. (2016). Interleukin-6 and C-reactive protein as prognostic biomarkers in metastatic colorectal cancer. *Oncotarget*, 7(46), 75013–75022. doi: 10.18632/oncotarget.12601.
- Vénéreau, E., Ceriotti, C., and Bianchi, M. E. (2015). DAMPs from cell death to new life. *Frontiers in Immunology*, 6.
- Virlando Suryadinata, R. (2018). Effect of free radicals on inflammatory process in chronic obstructive pulmonary disease (COPD). *Amerta Nutrition*, 2(4), 317–324. doi: 10.20473/amnt.v2i4.2018.317-324
- West, R. (2017). Tobacco smoking: Health impact, prevalence, correlates and interventions. *Psychology & Health*, 32(8), 1018–1036. doi: 10.1080/08870446.2017.1325890

Zhang, H., Ma, Z., Luo, X., & Li, X. (2018). Effects of mulberry fruit (*morus alba* l.) consumption on health outcomes: a mini-review. *Antioxidants*, 7(5), 69. doi: 10.3390/antiox7050069

Zhang, Z. & Shi, L. (2010). Anti-inflammatory and analgesic properties of cis-mulberroside A from *Ramulus mori*. *Fitoterapia*, 81(3), 214–218. doi: 10.1016/j.fitote.2009.09.005






# Jurnal Ilmiah **MEDIA GIZI** *Indonesia*



Accredited by SINTA Indonesia (SINTA 2)  
SK DITJEN RISSANG KEMENRISTEKDIKTI RI NO 3/E/KTP/2019  
(National Nutrition Journal)





Home > Vol 16, No 3 (2021)

## MEDIA GIZI INDONESIA

*Media Gizi Indonesia* (MGI) / National Nutrition Journal (p-ISSN 1693-7228 ; e-ISSN 2540-8410) is a scientific journal on nutrition. It is periodically published every 3 months, in January, May, and September. MGI also collaborates with *Perhimpunan Pakar Gizi dan Pangan Indonesia* (PERGIZI PANGAN Indonesia). MGI serves as communication media for disseminating information on research results and literature reviews which focus on community nutrition, clinical nutrition, institutional nutrition, food service management, food technology, and current issues on food and nutrition.



## ANNOUNCEMENTS

### ARTICLE PROCESSING CHARGE (APC)

Dear respected authors and readers,

We are announcing that starting March 29, 2021, Article Processing Charge (APC) payment via Bank BNI No.Virtual Account 9883030300000331. Name: Pembayaran Article Publications Charge (APC) Jurnal FKM Unair.

Thank you

Posted: 2021-04-05

### CALL FOR PAPER

National Nutrition Journal, Faculty of Public Health Universitas Airlangga announced a \*Call for Paper for International author\*

The article must be an \*original research or literature review\* that fits to our \*Focus and Scope\* in nutrition fields.

Submit your articles by \*July, 15 2020\* through <https://e-journal.unair.ac.id/MGI>

\*Selected articles will be published on Vol. 15 Issue 3, by September 2020 after peer reviewed.

Please contact us for more information:

+6282230069797

[jmggi@fkm.unair.ac.id](mailto:jmggi@fkm.unair.ac.id)

### Accreditation & Certificate



### Download Certificate

### Instruction for Author

Guide for authors

Online Submission

Document Template

### Journal Policy

Focus and Scope

Publication Ethics

Article Processing Charge

Peer Reviewers

Peer Reviewers Process

Editorial Team

Open Access Statement

Plagiarism

Copyright

Contact

Old Website

### USER

Username

Password

☐ Remember me

Login

### NOTIFICATIONS

[View](#)  
[Subscribe](#)

### JOURNAL CONTENT

Search

Search Scope

All

Search



Posted: 2020-06-15

## ENGLISH PROOFREAD CERTIFICATE

Dear

Authors of Media Gizi Indonesia

We would like to announce that starting from March 2020, for those who would like to submit manuscript in english must attach the english proofread to our editor.

Manuscript submission without english proofread certificate will not be accepted.

Thank you for your cooperation

Posted: 2020-03-20

[More Announcements...](#)

## VOL 16, NO 3 (2021): JURNAL MEDIA GIZI INDONESIA (NATIONAL NUTRITION JOURNAL)

### TABLE OF CONTENTS

#### ARTICLES

##### THE ACCEPTANCE OF BAKED GETUK (CASSAVA CAKE) WITH RICE BRAN AND RED BEANS SUBSTITUTION AS HIGH FIBER SNACK FOR ELDERLY

PDF  
200-  
206

10.20473/mgi.v16i3.200-206

*Ali Iqbal Tawakal, Annis Catur Adi, Dominikus Raditya Atmaka*

##### CORRELATION OF SNACKING FREQUENCY, HEMOGLOBIN LEVELS, PHYSICAL ACTIVITY AND POCKET MONEY WITH NUTRITIONAL STATUS IN FEMALE ADOLESCENTS

PDF  
207-  
214

10.20473/mgi.v16i3.207-214

*Dinar Putri Rahmawati, Dono Indarto, Diffah Hanim*

##### THE ASSOCIATION OF FAMILY SUPPORT IN FULFILLING HEALTHY NUTRITIOUS FOODS, PROTEIN AND MICRONUTRIENT INTAKE WITH HEMOGLOBIN LEVELS AMONG ELDERLY

PDF  
215-  
223

10.20473/mgi.v16i3.215-223

*Luberta Ebta Wiyani, Diffah Hanim, Dono Indarto*

##### FACTORS RELATED TO BODY IMAGE AND IT CORRELATION WITH NUTRITIONAL STATUS AMONG FEMALE ADOLESCENTS: A LITERATURE REVIEW

PDF  
224-  
232

10.20473/mgi.v16i3.224-232

*Lusiana Pradana Hariyanti, Nila Reswari Haryana*

##### EFFECT OF RED MULBERRY JUICE (MORUS RUBRA) ON THE INFLAMMATORY RESPONSE IN MALE WISTAR RATS DUE TO EXPOSURE TO CIGARETTE SMOKE

PDF  
233-  
237

10.20473/mgi.v16i3.233-237

*Rivan Virlando Suryadinata, Merryana Adriani, Stefani Cornelia Sardjono*

##### DIETARY INTAKE AND SUN EXPOSURE RELATED TO VITAMIN D CONCENTRATION IN THALASSEMIA PATIENTS: A LITERATURE REVIEW

PDF  
238-  
247

10.20473/mgi.v16i3.238-247

Browse

[By Issue](#)

[By Author](#)

[By Title](#)

[Other Journals](#)

### INFORMATION

[For Readers](#)

[For Authors](#)

[For Librarians](#)



**RELATIONSHIP OF FORMER HISTORY OF DIARRHEA AND SANITATION WITH THE PREVALENCE OF STUNTING AMONG CHILDREN AGED 1-5 YEARS IN SIDOTOPO WETAN, SURABAYA**

PDF  
248-  
255

 10.20473/mgi.v16i3.248-255

Naisya Azalia Samsuddin, Emyr Reisha Isaura, Sri Sumarmi

**ASAM KLOROGENAT PADA KOPI DAN OBESITAS: A SYSTEMATIC REVIEW Chlorogenic Acid in Coffee and Obesity: A Systematic Review**

PDF  
256-  
266

 10.20473/mgi.v16i3.256-266

Kamila Dwi Febrianti, Stefania Widya Setyaningtyas

**PENGARUH DOSIS DAN LAMA PEMBERIAN EKSTRAK DAUN ASAM JAWA (TAMARINDUS INDICA LINN) TERHADAP HOMA-B PADA TIKUS MODEL DIABETES MELLITUS TIPE 2 The effect of Tamarind Leaf (tamarindus indica linn) Extract on HOMA-β in Rats with Type 2 Diabetes Mellitus Model**

PDF  
267-  
272

 10.20473/mgi.v16i3.267-272

Devi Novia, Sugiarto Sugiarto, Yulia Lanti Dewi

**PENINGKATAN PENGETAHUAN DAN SELF-EFFICACY UPAYA PENCEGAHAN COVID-19 MELALUI EDUKASI GIZI KONVENSIONAL Increased Knowledge and Self-Efficacy of COVID-19 Prevention through Conventional Nutrition Education**

PDF  
273-  
279

 10.20473/mgi.v16i3.273-279

Qonita Rachmah, Triska Susila Nindya, Arif Sabta Aji, Sitti Pattimah, Nabilla Rachmah, Nabiil Ikbaar Maulana, Asri Meidyah Agustin, Junaida Astina

**KORELASI TINGKAT ASUPAN LEMAK DAN AKTIVITAS FISIK DENGAN KUALITAS HIDUP LANJUT USIA Correlation between Fat Intake and Physical Activity with Quality of Life in Elderly**

PDF  
280-  
286

 10.20473/mgi.v16i3.280-286

Roni Nurdianto, Diffah Hanim, Eti Poncorini Pamungkasari

**PENGARUH ASUPAN SAYUR KANGKUNG (IPOMEA REPTANS POIR.) DAN OLAHRAGA RENANG TERHADAP INTENSITAS KECEMASAN PADA REMAJA LAKI-LAKI The Effect of Kangkong Vegetables (Ipomea reptans Poir.) and Swimming on Anxiety Intensity in Adolescent Male**

PDF  
287-  
295

 10.20473/mgi.v16i3.287-295

Sigit Yudhistira, Adi Magna Patriadi Nuhriawangsa, Mohammad Fanani

**UJI EFEKTIFITAS SINBIOTIK KEFIR TEPUNG PISANG BATU TERHADAP KADAR GLUKOSA DARAH DAN KADAR PROFIL LIPID TIKUS MODEL SINDROM METABOLIK Effectiveness Test of banana (Musa balbisiana) Flour Kefir Synbiotic on Blood Glucose Level and Lipid Profile Level in Metabolic Syndrome Rats Model**

PDF  
296-  
305

 10.20473/mgi.v16i3.296-305

Taufik Maryusman, Christine Diane Dien, Santi Herlina Mail

**HUBUNGAN ANTARA STATUS GIZI TB/U DENGAN KEMAMPUAN KOGNITIF, FISIK, DAN SOSIO-EMOSIONAL ANAK USIA PRA-SEKOLAH Association of Height-For-Age Nutritional Status with Cognitive, Physical, and Socio-Emotional Ability of Pre-School Children**

PDF  
306-  
312

 10.20473/mgi.v16i3.306-312

Vidya Anggarini Rahmasari, Lailatul Muniroh





**In Collaboration**



**00413635**

[View MGI Stats](#)



Pusat Pengembangan Jurnal dan Publikasi Ilmiah  
**Universitas Airlangga**  
copyright©2017 Template PPJPI



[Home](#) > [About the Journal](#) > [Editorial Team](#)

## EDITORIAL TEAM

### CHIEF EDITOR

[Dr. Annis Catur Adi, Ir., M.Si.](#), (SCOPUS ID: 57200582013 ; H-Index : 3), Department of Nutrition, Faculty of Public Health, Universitas Airlangga, Indonesia

### EDITORIAL BOARD

[Assoc. Prof. Wantanee Kriengsinyos, Ph.D.](#), (SCOPUS ID: 6506938692 ; H-Index : 10), Human Nutrition Division, Institute of Nutrition, Mahidol University, Thailand

[Assoc. Prof. Hazreen Bin Abdul Majid, Bsc. Dietetic., M.Nut.Dietetic., Ph.D.](#), (SCOPUS ID: 54893067500 ; H-Index : 15), Department Social and Preventive Medicine, Faculty of Medicine, University of Malaya, Malaysia

[Prof. Dr. Ir. Dodik Briawan, MCN.](#), (SCOPUS ID: 56157628600 ; H-Index : 5), Department of Community Nutrition, Human Ecology Faculty, Bogor Agricultural University, Indonesia

[Prof. R. Bambang W, dr, MS, MCN, Ph.D, SpGK.](#), (SCOPUS ID: 56403330300 ; H-Index : 4), Department of Nutrition, Faculty of Public Health, Universitas Airlangga, Indonesia

[Prof. Dr. Luthfiyah Nurlaela, M.Pd.](#), (SCOPUS ID: 57193699945 ; H-Index : 1), State University of Surabaya, Indonesia

[Dr. Beben Benyamin, PhD.](#), (SCOPUS ID: 16314927500 ; H-Index : 27), School of Health Sciences, University of South Australia, Australia

[Loh Su Peng, PhD.](#), (SCOPUS ID: 23972729900 ; H-Index : 15), Department of Nutrition & Dietetics, Faculty of Medicine & Health Sciences, Universiti Putra Malaysia

[Sueppong Gowachirapant, Ph.D.](#), (SCOPUS ID: 12764961000), Institute of Nutrition, Mahidol University, Thailand

[Dr. Ir. Tri Dewanti Widyaningsih, M.Kes.](#), (SCOPUS ID: 55900709900 ; H-Index : 5), Department of Food Science and Technology, Faculty of Agriculture Technology, University of Brawijaya, Indonesia, Indonesia

[Dr. Toto Sudargo, M.Kes.](#), (SCOPUS ID: 56158010900 ; H-Index : 3), Undergraduate School of Nutrition and Health, Faculty of Medicine, Gadjah Mada University, Indonesia

[Prof. Dian Handayani, SKM., MKes., Ph.D.](#), (SCOPUS ID: 56035896000 ; H-Index : 3), Nutrition Department, Faculty of Medicine, University of Brawijaya, Indonesia

### MANAGING EDITORS

[Qonita Rachmah, S.Gz, M.Sc \(Nutr & Diet\).](#), (SCOPUS ID: 60069383 ; H-Index : 3), Department of Nutrition, Faculty of Public Health, Universitas Airlangga, Indonesia

[Stefania Widya Setyaningtyas, S.Gz, MPH.](#), (SCOPUS ID: 57208284644 ; H-Index : 1) Department of Nutrition, Faculty of Public Health, Universitas Airlangga, Indonesia

[Dominikus Raditya Atmaka, S.Gz., MPH.](#), (SCOPUS ID: 57222577092), Department of Nutrition, Faculty of Public Health, Universitas Airlangga, Indonesia

### ADMINISTRATIVE ASSISTANT

[Hadiar Huriyah Rahma, S.KM.](#), Department of Nutrition, Faculty of Public Health, Universitas Airlangga, Indonesia

#### Accreditation & Certificate



#### Download Certificate

#### Instruction for Author

[Guide for authors](#)

[Online Submission](#)

[Document Template](#)

#### Journal Policy

[Focus and Scope](#)

[Publication Ethics](#)

[Article Processing Charge](#)

[Peer Reviewers](#)

[Peer Reviewers Process](#)

[Editorial Team](#)

[Open Access Statement](#)

[Plagiarism](#)

[Copyright](#)

[Contact](#)

[Old Website](#)

#### USER

Username

Password

☐ Remember me

#### NOTIFICATIONS

[View](#)  
[Subscribe](#)

#### JOURNAL CONTENT

Search

Search Scope

All



In Collaboration

# MGI

*by* Rivan Suryadinata

---

**Submission date:** 30-Sep-2021 07:35PM (UTC+0700)

**Submission ID:** 1661510835

**File name:** Media\_Gizi\_Indonesia\_September\_2021.pdf (189.66K)

**Word count:** 2676

**Character count:** 14807

## EFFECT OF RED MULBERRY JUICE (*MORUS RUBRA*) ON THE INFLAMMATORY RESPONSE IN MALE WISTAR RATS DUE TO EXPOSURE TO CIGARETTE SMOKE

Rivan Virlando Suryadinata<sup>1\*</sup>, Merryana Adriani<sup>2</sup>, Stefani Cornelia Sardjono<sup>3</sup>

<sup>1</sup>Faculty of Medicine, University of Surabaya (UBAYA), Surabaya

<sup>2</sup>Department of Nutrition, Faculty of Public Health, Universitas Airlangga, Surabaya

<sup>3</sup>Students of the Medical Study Program, University of Surabaya (UBAYA), Surabaya

\*E-mail: rivan.virlando.suryadinata@gmail.com

### ABSTRACT

The increasing number of tobacco users will directly impact to an increase in the number of passive smokers. Various harmful substances that enter the respiratory tract can cause health problems. The airway will recognize these substances as pathogen and activate immune response accordingly. Inflammatory response becomes one of the effects of body compensation due to an increased pathogen. This effect is indicated by the elevation of C-Reactive Protein as one systemic inflammatory response parameters in the body. This study aimed to determine the efficacy of red mulberry juice to reduce levels of C-Reactive Protein in Wistar rats because exposure to cigarette smoke. This is an experimental study with a post-test control group design. Animals were divided into negative control group, positive control group, and 3 treatment groups with different red mulberry doses. The positive control group showed the highest increase in C-Reactive Protein levels compared with the red mulberry intake group ( $p < 0.001$ ). In conclusion, the administration of red mulberry juice can reduce inflammatory response in rats because of exposure to cigarette smoke.

**Keywords:** c-reactive protein, cigarette, inflammation, red mulberry

### INTRODUCTION

Tobacco cigarette are the most common type of cigarette used worldwide (Gowing et al., 2015). Various studies have shown the dangers of using tobacco cigarettes both as active smokers and passive smokers (Suryadinata et al., 2017). Smoking has caused 6 million people dies each year and it is estimated will reach 8 million in 2030. Globally, the number of tobacco smokers has reached 1.1 billion people and almost 70% are residents of developing countries (Nasser et al, 2018). The prevalence of active smokers in developed countries has shown a significant decrease. Estimated in 2015, number of smokers in developed countries like America and Europe was around 15% of total population (West, 2017). In developing countries, the number of smokers increases every year, especially in the productive age (Issain et al., 2017).

Smoke contains more than 4000 types various ingredients, including nicotine, ammonia, phenols, polycyclic aromatics, carbon monoxide, nitrogen oxides, and various metals (Kamceva et al., 2016). Two main phases that can be identified in cigarette smoke are tar phase and gas phase. Both phases

contain high free radicals such as superoxide ( $O_2^{\bullet-}$ ) and hydroxyl ( $\bullet OH$ ) (Suryadinata, 2018). Increased free radicals in body due to exposure to smoke cigarettes entering airways will cause oxidative stress and trigger lipid peroxidation. This will potentially cause cell damage and necrosis (Suryadinata et al., 2019). Normally, free radicals are mitochondrial metabolism and can be neutralized by enzymatic antioxidants in body (Suryadinata et al, 2019). In pathological conditions, excessive free radicals will directly stimulate inflammatory response through phagocytic process of macrophages (Arulselvan et al., 2016).

Inflammatory process is a form of body defence mechanism against pathogenic reactions from outside. Various pro-inflammatory cytokines are secreted into body microenvironment such as interleukin-1, interleukin-6, interleukin-8, and tumor necrosis factor-alpha (TNF- $\alpha$ ). Inflammatory mediator will induce liver to release C-Reactive Protein (CRP) into the blood (Phaniendra et al., 2015). Prevention of inflammatory process due to free radical increase in the body can be done with administration of antioxidants from the outside.



Various types of non-enzymatic antioxidants can be obtained from food especially flavonoids (Pratiwi et al., 2018).

Mulberry fruit is one type of fruit that high in antioxidants. Mulberries originated from genus *Morus* of Moraceae family, to date there are 24 species and 100 varieties (Zhang et al., 2018). Various studies have shown the provision of mulberry fruit has potential impact as antioxidant (Zhang and Shi, 2010). This study used red mulberry juice (*Morus rubra*) juice to reduce body inflammatory response due to exposure to tobacco smoke. Besides, red mulberries are also widely planted and consumed by Indonesian people so that at the same time can preserve local food. This study purpose is to analyze the efficacy of red mulberry juice to reduce levels of C-Reactive Protein in Wistar rats exposed to cigarette smoke.

## METHODS

This research was an experimental study with post-test control group design performed in 25 male Wistar rats (*Rattus norvegicus*) which were divided into 5 groups for 30 days. Inclusion criteria in this study were experimental animals aged 3-4 months with weight around 200 g. While the exclusion criteria are animals get sick or die during study. Maintenance and care of the animals during study are based on 3R principles (Replacement, Reduction, and Refinement). This research has been obtained Ethical Clearance Certificate from Institutional Ethical Committee, University of Surabaya (No:135/KE/VI/2020).

Red mulberries (*Morus rubra*) that have been obtained, sorted and cleaned in order to get fruit in good condition. Red mulberry juice that has been obtained then performed flavonoid levels measurement ( $\pm 38$  mg/g). Results of flavonoids levels that have been obtained then converted from daily needs of adults ( $\pm 190$  mg/kg) into daily dose intake of Wistar rats with 200g body weight ( $\pm 240$  mg/rats) using Laurence and Bacharach table. The amount of red mulberry juice given orally to the treatment group was 3.25 g, 6.5 g, and 13 g.

The results of CRP level measurement are based on agglutination process in blood through intracardiac puncture. Dilution process show the results of C-Reactive Protein measurement i.e., >

8  
6 mg/L, 12 mg/L, 24 mg/L, 48 mg/L and 96 mg/L. Measurement results were classified as positive if C-Reactive protein levels obtained more than 6 mg/L, and negative if C-Reactive Protein level obtained were less than 6 mg/L.

This research was conducted for 30 days by dividing rats into 5 groups. The first was the negative control group, where animals did not received any treatment. In positive control group, animals received daily intake and exposure to 2 tobacco cigarettes smoke per day in smoking chamber. While last three groups were exposed to 2 tobacco cigarettes smoke per day, daily intake, and 3.25 g, 6.5 g, or 13 g red mulberry juice.

Results of study were ordinal data in the form of CRP levels in each group in mg/L units. CRP measurement method uses dilution to get a multiple of the result. CRP levels of >6 mg/L, 12 mg/L, 24 mg/L and 48 mg/L will be carried out for coding. Data analysis was performed using Kruskal Wallis analysis with SPSS version 22 to observe differences between groups.

## RESULTS AND DISCUSSIONS

Results of study were carried out by comparing C-Reactive Protein levels between groups. Table 1. shows results of C-Reactive Protein level measurement in blood between group. Group I, IV, and V had the same levels of C-Reactive Protein. The highest level of C-Reactive Protein obtained in Group II which is the Positive Control Group.

Kruskal Wallis test in Table 1 shows p-value of C-Reactive Protein levels between group was <0,001 ( $p\text{-value} < 0.05$ ), so it can be concluded that there are significant differences of C-Reactive Protein levels between groups.

Exposure to free radicals from tobacco smoke that enters respiratory tract causes body develop defence mechanisms to prevent the impact of cell damage due to oxidative stress (Indraswari et al., 2018). Antioxidant produced in body are enzymatic antioxidants (superoxide dismutase [SOD], glutathione peroxidase [GSH-Px], and catalase), while antioxidant from outside are non-enzymatic antioxidant derived from food intake such as flavonoids (Suryadinata et al., 2016). The balance between number of free radicals and

**Table 1.** Levels of C-Reactive Protein in research

	Group	C-Reactive Protein levels (mg / L) in Wistar Rats					Kruskal Wallis Test
		1 <sup>st</sup>	2 <sup>nd</sup>	3 <sup>rd</sup>	4 <sup>th</sup>	5 <sup>th</sup>	
I	The negative control group without exposure to cigarette smoke and intake of red mulberry juice	<6	<6	<6	<6	<6	
II	The positive control group received exposure to smoke cigarettes and no intake of red mulberry juice	48	48	48	24	24	
III	Treatment group I was exposed to cigarette smoke and red mulberry juice intake of 3.25 g	24	12	12	<6	<6	<0.001
IV	Treatment group II received smoke exposure cigarettes and red mulberry juice intake of 6.5 g	<6	<6	<6	<6	<6	
V	Treatment group III received smoke exposure cigarettes and red mulberry juice intake of 13 g	<6	<6	<6	<6	<6	

antioxidants is needed for the stability of body normal metabolic process (Biswas et al., 2017).

Reactive Oxygen Species (ROS) are the most reactive and radical types that directly respond to inflammation in body. These radicals can be neutralized with superoxide dismutase (SOD) into hydrogen peroxidase (H<sub>2</sub>O<sub>2</sub>) then will be neutralized with glutathione peroxidase (GSH-Px) and catalase into water (H<sub>2</sub>O) and oxygen (O<sub>2</sub>) (Nimse and Pal, 2015).

However, an increase in excessive free radicals can result in cell damage or necrosis. This process will cause cells to become lysis and is called debris cells (Vénéreau et al., 2015). This will stimulate the movement of macrophages as one mechanism of body defence. Phagocytic process carried by macrophages will trigger secretion of pro-inflammatory cytokines such as interleukin-1, interleukin-6, and tumor necrosis factor-alpha (TNF- $\alpha$ ) (Thomsen et al., 2016). Excessive release of proinflammatory cytokines will induce liver to secrete C-Reactive Protein (Sproston and Ashworth, 2018).

Inflammatory processes in the body have positive and negative effects. In acute and low condition, Inflammatory processes will function as a defence and healing process for cell tissue, while if the condition is chronic, then it will have impact on cell damage. Exposure to cigarette smoke that given regularly and continuously will create chronic inflammatory conditions that have a negative impact on health (Srivastava and Kumar, 2015). Therefore, administration of antioxidants through food intake is expected to

be able to prevent the inflammatory process by decreasing free radicals.

Administration of red mulberry juice (*Morus rubra*) which rich in flavonoids shows its efficacy on decreasing levels of C-Reactive Protein in blood due to exposure to tobacco smoke. Therefore, the use of red mulberries as an additional intake of antioxidants from outside body can be considered as one way to prevent cell damage due to inflammatory process.

## CONCLUSION

Exposure to tobacco smoke can increase body inflammatory response through increased levels of C-Reactive Protein in the blood. Giving red mulberry juice can reduce the response of inflammation caused by exposure to tobacco smoke in rats.

## ACKNOWLEDGEMENT

The researcher is grateful to all laboratory staff who have assisted the research process.

## REFERENCES

- Arulselvan, P., Fard, M. T., Tan, W. S., Gothai, S., Fakurazi, S., Norhaizan, M. E., ... Kumar, S.S. (2016). Role of antioxidants and natural products in inflammation. *Oxidative Medicine and Cellular Longevity*, 2016, 1–15. doi: 10.1155/2016/5276130
- Biswas, S., Das, R., & Ray Banerjee, E. (2017). Role of free radicals in human inflammatory

- diseases. *AIMS Biophysics*, 4(4), 596–614. doi: 10.3934/biophy.2017.4.596
- Gowing, L. R., Ali, R. L., Allsop, S., Marsden, J., Turf, E. E., West, R., & Witton, J. (2015). Global statistics on addictive behaviours: 2014 status report. *Addiction*, 110(6), 904–919. doi: 10.1111/add.12899.
- Hossain, S., Hossain, S., Ahmed, F., Islam, R., Sikder, T., & Rahman, A. (2017). Prevalence tobacco smoking and factors associated with the initiation of smoking among university students in Dhaka, Bangladesh. *Central Asian Journal of Global Health*, 6(1). doi: 10.5195/cajgh.2017.244
- Indraswari, P. I. I., Lorensia, A., and Suryadinata, R. V. (2018). Analysis effect of nutrition intake on lung function of active smoker and non smoker. *Jurnal Kesehatan Masyarakat*, 14(2), 247–253. Retrieved from <https://journal.unnes.ac.id/nju/index.php/kemas/article/view/14947>
- Kamceva, G., Arsova-Sarafinovska, Z., Ruskovska, T., Zdravkovska, M., Kamceva-Panova, L., & Stikova, E. (2016). Cigarette smoking and oxidative stress in patients with coronary artery disease. *Open Access Macedonian Journal of Medical Sciences*, 4(4), 636. doi: 10.3889/oamjms.2016.117
- Nasser, A., Salah, B., Regassa, L., Alhakimy, A., & Zhang, X. (2018). Smoking prevalence, attitudes and associated factors among students in health-related Departments of Community College in rural Yemen. *Tobacco Induced Diseases*, 16(July). doi: 10.18332/tid/92547
- Nimse, S. B. & Pal, D. (2015). Free radicals, natural antioxidants, and their reaction mechanisms. *RSC Advances*, 5(35), 27986–28006. doi: 10.1039/C4RA13315C
- Phaniendra, A., Jestadi, D. B., & Periyasamy, L. (2015). Free radicals: properties, sources, targets, and their implication in various diseases. *Indian Journal of Clinical Biochemistry*, 30(1), 11–26. doi: 10.1007/s12291-014-0446-0
- Pratiwi, S. R., Lorensia, A., & Suryadinata, R. V. (2018). Vitamin C and E Intake with SQ-FFQ towards Smokers' and Non-Smokers' Lung Function. *Media Kesehatan Masyarakat Indonesia*, 14(2), 101. Retrieved from <http://journal.unhas.ac.id/index.php/mkmi/issue/view/559>
- Sproston, N. R., & Ashworth, J. J. (2018). Role of c-reactive protein at sites of inflammation and infection. *Frontiers in Immunology*, 9. doi: 10.3389/fimmu.2018.00754
- Srivastava, K. K., & Kumar, R. (2015). Stress, oxidative injury and disease. *Indian Journal of Clinical Biochemistry*, 30(1), 3–10. doi: 10.1007/s12291-014-0441-5
- Suryadinata, Rivan Virlando, Wirjatmadi, B. (2019). The role of selenium micronutrients as antioxidants in exposure to e-cigarette smoke. *Asian Journal of Pharmaceutical and Clinical Research*, 12(8), 265–268. Retrieved from <https://innovareacademics.in/journals/index.php/ajpcr/issue/view/466>
- Suryadinata, R. V., Wirjatmadi, B., & Adriani, M. (2016). Pengaruh perubahan hiperplasia sel goblet selama 28 hari paparan asap rokok dengan pemberian antioksidan superoxide dismutase. *The Indonesian Journal of Public Health*, 11(1), 60. Retrieved from 10.20473/ijph.v11i1.2016.60-68
- Suryadinata, R. V., Wirjatmadi, B., & Adriani, M. (2017). Efektivitas penurunan malondialdehyde dengan kombinasi suplemen antioksidan superoxide dismutase melon dan gliadin akibat paparan rokok. *Global Medical And Helath Communication*, 5(2), 79–83. Retrieved from
- Suryadinata, R. V., Wirjatmadi, B., Adriani, M., and Sumarmi, S. (2019). The effects of exposure duration to electronic cigarette smoke on differences in superoxide dismutase and malondialdehyde in blood of wistar rats. *International Journal of Current Pharmaceutical Research*, 11(3), 13–16. doi: 10.22159/ijcpr.2019v11i3.34084
- Thomsen, M., Kersten, C., Sorbye, H., Skovlund, E., Glimelius, B., Pfeiffer, P., et al. (2016). Interleukin-6 and C-reactive protein as prognostic biomarkers in metastatic colorectal cancer. *Oncotarget*, 7(46), 75013–75022. doi: 10.18632/oncotarget.12601.
- Vénéreau, E., Ceriotti, C., and Bianchi, M. E. (2015). DAMPs from cell death to new life. *Frontiers in Immunology*, 6.
- Virlando Suryadinata, R. (2018). Effect of free radicals on inflammatory process in chronic obstructive pulmonary disease (COPD). *Amerta Nutrition*, 2(4), 317–324. doi: 10.20473/amnt.v2i4.2018.317-324
- West, R. (2017). Tobacco smoking: Health impact, prevalence, correlates and interventions. *Psychology & Health*, 32(8), 1018–1036. doi: 10.1080/08870446.2017.1325890

Zhang, H., Ma, Z., Luo, X., & Li, X. (2018). Effects of mulberry fruit (*morus alba* l.) consumption on health outcomes: a mini-review. *Antioxidants*, 7(5), 69. doi: 10.3390/antiox7050069

Zhang, Z. & Shi, L. (2010). Anti-inflammatory and analgesic properties of cis-mulberroside A from *Ramulus mori*. *Fitoterapia*, 81(3), 214–218. doi: 10.1016/j.fitote.2009.09.005

## ORIGINALITY REPORT

13%

SIMILARITY INDEX

11%

INTERNET SOURCES

7%

PUBLICATIONS

7%

STUDENT PAPERS

## PRIMARY SOURCES

1	e-journal.unair.ac.id Internet Source	3%
2	id.123dok.com Internet Source	2%
3	innovareacademics.in Internet Source	2%
4	Submitted to British College of Osteopathic Medicine Student Paper	1%
5	link.springer.com Internet Source	1%
6	Repositori.Usu.Ac.Id Internet Source	1%
7	www.ncbi.nlm.nih.gov Internet Source	1%
8	Submitted to University of Hull Student Paper	1%
9	www.jove.com Internet Source	1%



10

[www.mjms.mk](http://www.mjms.mk)

Internet Source

1 %

11

[worldwidescience.org](http://worldwidescience.org)

Internet Source

&lt;1 %

12

Lun-Hsien Chang, Jue-Sheng Ong, Jiyuan An, Karin J.H. Verweij et al. "Investigating the genetic and causal relationship between initiation or use of alcohol, caffeine, cannabis and nicotine", Drug and Alcohol Dependence, 2020

Publication

&lt;1 %

13

[cbt2olympicad6.unimus.ac.id](http://cbt2olympicad6.unimus.ac.id)

Internet Source

&lt;1 %

14

Jing Qu, Yue Cheng, Wenchao Wu, Lixing Yuan, Xiaojing Liu. "Glycocalyx Impairment in Vascular Disease: Focus on Inflammation", Frontiers in Cell and Developmental Biology, 2021

Publication

&lt;1 %

Exclude quotes On

Exclude matches &lt; 5 words

Exclude bibliography On

FINAL GRADE

/10

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5