



Relationship of Knowledge on Respiratory Disorders with Lung Function in Masons in East Surabaya

Amelia Lorensia^{1,*}, Rivan Virlando Suryadinata², Wilma Adib Gardiawan³

¹Department of Clinical-Community Pharmacy, Faculty of Pharmacy, Universitas Surabaya (UBAYA)

²Department of Public Health, Faculty of medicine, Universitas Surabaya (UBAYA)

³Bachelor Student of Faculty of Pharmacy, Faculty of Pharmacy, Universitas Surabaya (UBAYA)

*Corresponding author: amelia.lorensia@staff.ubaya.ac.id

Abstract

Masons are regularly exposed to air pollution in the workplace. The pollution exposure received by them is a risk factor for respiratory problems. Most of the masons had insufficient knowledge that is very important for the management of a person's illness. The purpose of this study was to determine relationship of knowledge on respiratory disorders with lung function in masons in east surabaya. This research was an observational study with a cross-sectional design with a purposive method. The research was conducted from April to July 2019 in the East Surabaya area. The variables that will be observed in this study are the level of knowledge of risk factors, symptoms, and treatment and therapy regarding respiratory disorders. In this study, there were 158 respondents consisting of 79 groups with impaired lung function and 79 groups without lung function disorders. Most of the level of knowledge of both groups was good in risk factors and treatment of respiratory disease, but instead on the symptoms regarding respiratory disease. The research showed that there was a relationship between the level of knowledge about risk factors for respiratory disease ($p(0.223) < 0.05$) and lung function, but there was no difference in knowledge about symptoms ($p(0.745) < 0.05$) and treatment ($p(0.741) < 0.05$) respiratory disease with pulmonary function. Therefore, it was necessary to plan further educational programs in increasing masons' knowledge of respiratory symptoms so that they could catch respiratory diseases earlier.

Keywords: knowledge, lung function, masons, respiratory disorders

Submitted: 03 May 2021

Accepted: 25 December 2021

DOI: <https://doi.org/10.25026/jtpc.v5i4.342>

1 Introduction

Surabaya is the second metropolitan city in Indonesia with a total area of 326.36 km². In 2017 the city of Surabaya has a population of 2.87 million and is predicted to increase every year [1]. The efforts of the Surabaya City government in realizing development planning require labor, one of which is a builder. A builder is a person who works by relying on his physical strength and has skills in handwork (digger, bricklayer, blacksmith, carpenter, painter) in making buildings in general (such as buildings, houses, shops) [2]. Masons have a very high risk of being exposed to air pollution [3,4]. According to research data in 2010, more than 50% of masons are regularly exposed to air pollution such as steam, gas, dust, or smoke in the workplace [5].

The pollution exposure received by masons is a risk factor for respiratory problems [6]. A work environment full of air pollution will have a negative impact on lung health [7]. Air pollution can have a direct or indirect effect on environmental quality degradation, namely air pollution [4,8]. The impact of air pollutants on the respiratory system has been consistently reported in recent years. Adverse effects include decreased lung function, increased infection, increased respiratory symptoms, acute exacerbations, onset of asthma, and increased respiratory mortality [6].

Decreased lung function in the form of respiratory disorders that occur in the respiratory tract and lungs that affect the human respiratory system [10]. Respiratory disorders can also be seen from spirometry measurements by looking at the FEV1/FVC value, where if the FEV1/FVC (Forced Expiratory Volume 1/ Forced Vital Capacity) value is <0.7 then it can be stated that you have respiratory problems [11]. In 2017, respiratory system diseases were also included in the top 10 most diseases in Surabaya and even ranked first [12]. Most of the masons had insufficient knowledge. Based on the results of the study, where most of the mason respondents were at the elementary education level and dropped out of school [13]. Lack of knowledge can affect the prevention of respiratory disorders. According to research conducted by Bhandari et al., it

shows that 88.0% of respondents have a below-average level of knowledge to prevent respiratory problems. Likewise, 41.0% of respondents had an average level of knowledge regarding the prevention of respiratory problems [14]. Knowledge of respiratory health shows that 65% of people have insufficient knowledge, 35% of people have moderate knowledge and no one has sufficient knowledge [15].

Knowledge is very important for the management of a person's illness, self-management in order to avoid a disease [16,17]. According to research, if a person has more knowledge of the disease, it is more likely that someone will obey the treatment [18]. Therefore, pharmacists are required to provide good health services to patients and pharmacists must have knowledge, skills and behavior that are always improved [19]. The purpose of this study was to determine differences in knowledge regarding (1) risk factors for respiratory disorders; (2) respiratory symptoms; and (3) treatment and therapy of respiratory disorders in masons with impaired lung function and without pulmonary function disorders.

2 Materials and Methods

2.1 Research design

The research was conducted from April to July 2019 in the East Surabaya area. This study was an observational study using a cross sectional study design to determine the profile of knowledge about respiratory disorders in builders based on lung function. Retrieval of research data by means of direct question and answer (interview) between the interviewer and the respondent.

2.2 Research variable

This study has two variables, namely the independent variable and the dependent variable. The independent variable of this study was impaired lung function. While the dependent variable in this study was the level of knowledge related to respiratory disorders.

The builder referred to in this research was a person who works by relying on his physical strength and has skill in a handwork (digger, bricklayer, bricklayer, carpenter, painter) in making buildings (such as buildings, houses, shops). Lung function disorders were diseases and disorders that occur in the respiratory tract and lungs that affect human respiration [10]. A person's lung function can be known through spirometric measurements, if the measurement using spirometry shows the results of the FEV1 value <70%, it can be said that you have lung function disorders [20]. The knowledge referred to in this research was knowledge about risk factors, symptoms, and treatment and therapy that the respondent has regarding respiratory disorders.

2.3 Population and Research Sample

The population in this study were masons who were working on a project or were active in the East Surabaya area. The sample in this study were masons who were in the East Surabaya area who were included in the inclusion and exclusion criteria. Respondents who are willing to participate in this research and have filled in the informed consent, are male, b. Age 17 - 65 years, active smokers, minimum length of work for 5 years, and no mental or physical disorders. The sampling technique used was random sampling (nonprobability sampling) with purposive sampling and consecutive sampling methods. The number of samples taken in this study using the Lemeshow formula with an unknown or infinite population. The following is the formula used: $n = Z^2 [P(1-P)] / d^2$ (n = number of samples; Z =standard normal value (if $\alpha = 0.05$ then $Z = 1.960$); $P(1-P)$ =estimated population proportion (if $P=0.1$ then $P(1-P)=0.09$); d^2 : tolerant deviation (10%). So the minimum number of samples needed in this study is 62 respondents for each group.

2.4 Method of collecting data

Respondents will be given a questionnaire which was one of the instruments, important in research, the questionnaire given was about knowledge related to respiratory disorders. The questionnaire in this study consisted of 25 questions, of which 11 questions about knowledge of risk factors for respiratory disorders, 6 questions about knowledge of

respiratory symptoms, and 8 questions about knowledge about treatment and therapy.

Pulmonary function measurements were carried out using a Handled Spirometer where BMI data in the form of body weight and height as well as respondent data such as gender, age, smokers or non-smokers were also entered into the spirometer. The results of the spirometer measurement are used to determine respondents who have respiratory problems or do not experience respiratory problems based on FEV1, if FEV1 <70% indicates respiratory problems [20]. So if FEV1 <70% it can be categorized in the group that has impaired lung function and if FEV1 >70% can be categorized in the group that does not experience lung function disorders.

2.5 Data analysis method

Data processing was carried out in stages, starting with the data collected in the field until the data was ready for analysis. The data that has been collected in the field was checked, if there was a lack of data, it will be supplemented with repeated interviews with research respondents. The answers from the respondents will be coded to make the data input process easier. Then the data were analyzed descriptively (minimum, maximum, average and standard deviation values) and statistically using SPSS version 24 software.

The expected value in the chi-square should not be less than 5 (maximum 20% of expected frequencies <5). If the expected value was not met (20% expected frequencies <5), then the chi-square must be replaced with an alternative test and in determining the P value, if the P value was greater than 0.05 then there was no significant difference, but if the P value was more smaller than 0.05, there was a significant difference.

The level of knowledge of risk factors for respiratory disorders was measured using a questionnaire about respiratory disorders in which there were 11 questions. Every question that was answered correctly will be given a score of 1, while questions that are answered incorrectly or don't know will be given a score of 0. Then the level of knowledge will be classified as good and bad. It was said to be good if it gets a score of ≥ 8 and it is said to be bad if it gets a score of <8. Then the chi-square test was

carried out and the P_{value} was obtained of 0.223, which means that there was no significant difference between the group with pulmonary function disorders and the group without lung function disorders.

The level of knowledge of respiratory symptoms was measured using a questionnaire about respiratory disorders in which there are 6 questions. Every question that was answered correctly will be given a score of 1, while questions that were answered incorrectly or don't know will be given a score of 0. Then the level of knowledge will be classified as good and bad. It was said to be good if it gets a score of ≥ 4 and it was said to be bad if it gets a score of <4 . Then performed the chi-square test and obtained a P_{value} of 0.745 which means none a significant difference between the group with impaired lung function and the group without pulmonary function impairment.

The level of knowledge of respiratory symptoms was measured using a questionnaire about respiratory disorders in which there were 8 questions. Every question that was answered correctly will be given a score of 1, while questions that were answered incorrectly or don't know will be given a score of 0. Then the level of knowledge will be classified as good and bad. It was said good if it gets a score of ≥ 6 and it was said to be bad if it gets a score <6 . Then the chi-square test is carried out and the P_{value} was obtained of 0.741 which means no a significant difference between the group with impaired lung function and the group without pulmonary function impairment.

3 Results and Discussion

This research was conducted in April-July 2019 by conducting direct interviews with respondents using a questionnaire about respiratory disorders to determine the knowledge profile of risk factors, symptoms, and treatment and therapy for respiratory disorders based on lung function in masons in East Surabaya. Respondent data obtained in this study were 158 masons consisting of 79 people with pulmonary function disorders and 79 people without pulmonary function disorders. There were 20 respondents who refused to fill out the questionnaire, where 16 people did not want their rest time to be disturbed and 4 people did not meet the criteria.

3.1 Validity and Reliability of the Questionnaire

The results of the validity test of the risk factor knowledge questionnaire were declared valid. The questionnaire can be said to be valid if $r_{\text{count}} > r_{\text{table}}$ product moment, where the value of r_{table} for 30 respondents was 0.361 and all question items from the questionnaire have a value greater than the value of r_{table} product moment. The results of the reliability test of the risk factor knowledge questionnaire were declared reliable because they had a Cronbach's Alpha value equal to or greater than 0.61 (Cronbach's Alpha ≥ 0.61). The results of the reliability test of the risk factor knowledge questionnaire that have been obtained are declared reliable. The results of the validity of the questionnaire for knowledge of respiratory symptoms were declared valid because the value of $r_{\text{count}} > r_{\text{table}}$. Then then the questionnaire will be tested for reliability. The results of the questionnaire reliability test were declared reliable because Cronbach's Alpha value was ≥ 0.61 (Table 1).

3.2 Characteristics of Research Respondents

Respondents in this study were grouped into 2 groups, namely the group without lung function disorders and the group with lung function disorders. Respondents used in both groups were 158 respondents with 79 respondents in each group. Characteristics of respondents based on age consisted of late adolescents (17-25 years), early adults (26-35 years), late adults (36-45 years), early seniors (46-55 years), and late elderly (56-65 years old). Respondents without pulmonary function disorders in late adolescence were 20 respondents, early adults 33 respondents, late adults 13 respondents, early elderly 10 respondents, and late elderly by 3 respondents. Meanwhile, respondents with lung function disorders consisted of 15 respondents as late as adolescence, 45 respondents for early adulthood, 15 respondents for late adults, 3 respondents for early elderly, and 1 respondent for late elderly. Based on the age characteristics after testing the differences in the two groups showed that there was no significant difference between the two groups (Table 2).

Table 1. Validity and Reliability Test

Domain	Validity Test			Reliability Test		
	No.	Question	Corrected Item-Total Correlation	Conclusion	Cronbach's Alpha	Conclusion
Knowledge of Risk Factors for Respiratory Disorders	1	Environments with high pollution levels have greater risk of respiratory distress	0.457	Valid	0.614	Valid
	2	The smoke from the combustion/ factory /mine is a pollution which can cause respiratory distress	0.417	Valid		
	3	Smoking is not a cause of respiratory distress	0.419	Valid		
	4	The more the number of cigarettes smoked then further increase the risk factors for the disorder respiration	0.551	Valid		
	5	People working in dusty environments not experiencing respiratory distress	0.509	Valid		
	6	People who work in industrial areas have poor lung condition due to pollution in the area industry	0.411	Valid		
	7	People living in cities are more at risk experiencing respiratory distress than staying in the village	0.503	Valid		
	8	People living in industrial areas are more risk of experiencing respiratory distress	0.411	Valid		
	9	Increasing age does not increase the risk respiratory disorders	0.539	Valid		
	10	Respiratory disorders cannot be inherited or derived from the biological family	0.400	Valid		
	11	History of respiratory disorders in infants and children can present with disabilities throughout their lifetime adult	0.374	Valid		
Knowledge of Respiratory Disorders Symptoms	1	Shortness of breath is not a symptom of the disorder inhalation	0.481	Valid	0.612	Valid
	2	Shortness of breath in respiratory disorders can getting worse when doing	0.518	Valid		
	3	Cough that is continuous/ not persistent healing is a symptom of the disorder respiration	0.564	Valid		
	4	Increased respiratory rate is common in patients with respiratory disorders	0.707	Valid		
	5	Wheezing (such as a whistling sound or a 'ngic' sound during breathing) is a symptom of respiratory distress	0.520	Valid		
	6	The respiratory rate is increased in respiratory distress not a sign of worsening condition	0.704	Valid		
Knowledge of Medicine and Respiratory Disorders Therapy	1	Cough in respiratory distress may reduce with treatment (medicine) given	0.502	Valid	0.631	Valid
	2	Treatment (medication) of respiratory disorders is given only for symptom relief	0.476	Valid		
	3	Treatment (drugs) given can cure respiratory disorders	0.722	Valid		
	4	Quitting smoking has no effect reduction of respiratory distress symptoms	0.565	Valid		
	5	Breathing exercises aim to improve effort to breathe	0.434	Valid		
	6	Quitting smoking can make you breathe for the better	0.528	Valid		
	7	Breathing muscle training is not a viable exercise reduce shortness of breath in sufferers of the disorder respiration	0.567	Valid		
	8	Body activity program (sports) for increase tolerance of physical activity	0.435	Valid		

Table 2. Respondent Characteristics Frequency Distribution

Characteristics of Respondents	Total Number of Respondents (n: 158)				Different Test	
	Group Without Lung Function Disorders (n: 79)		Group With Lung Function Disorders (n: 79)		Pvalue	Conclusion
	Frequency	Percentage (%)	Frequency	Percentage (%)		
Age (year)	Late adolescents (17-25 years)	20	25.32	15	18.99	0.113 no difference
	Early adults (26-35 years)	33	41.77	45	56.96	
	Late adults (36-45 years)	13	16.46	15	18.99	
	Early seniors (46-55 years)	10	12.66	3	3.79	
	Late elderly (56-65 years old)	3	3.79	1	1.26	
BMI (Kg/m²)	Thin (<18.5)	7	8.86	6	7.59	0.485 no difference
	Normal (18.5-25.0)	59	74.68	66	83.54	
	Overweight (25.0-27.0)	8	10.13	5	6.33	
	Obesity	5	6.33	2	2.53	
Spirometry value [11]	<i>Mild</i> (FEV1 >80% predicted)	0	0	66	83.54	-
	<i>Worsening</i> (50%<FEV1<80% predicted)	0	0	13	16.46	-

Characteristics of respondents based on BMI were classified into 4, namely thin (<18.5), normal (18.5-25.0), overweight (25.0-27.0), and obesity (≥ 27.0). Respondents without pulmonary function disorders with a thin BMI were 7 respondents, normal was 59 respondents, overweight was 8 respondents, and obesity was 5 respondents. Meanwhile, 6 respondents with lung function disorders with a thin BMI, 66 normal respondents, 5 respondents with overweight, and 2 respondents with obesity. After testing the differences in the two groups, it was found that there was no significant difference in the two groups. Characteristics of respondents based on spirometric values to see lung function in groups with respiratory disorders will be classified based on GOLD [11] into 2, namely mild and worsening (**Table 2**).

Chronic lung disease was a major public health problem in the world. Although the chronic lung disease database on these outcome parameters is inadequate and often less reliable than mortality data, limited data suggest that chronic lung disease morbidity increases with age. Although the development of chronic lung disease comorbidities may occur at a younger age. Chronic lung disease was also more prevalent at age >40 years than <40 years, and was more common in men than women. Much of the increase in COPD mortality was due to the progression of the epidemic from smoking, the decrease in mortality from other common causes of death such as ischemic heart disease, infectious diseases [11]. We can broadly divide

the structural changes of aging lungs into three categories:

1. Changes in the structure of the lungs. Changes in the structure of the lungs are mainly associated with an increase in the size of the alveolar spaces in the absence of inflammation or damage to the alveolar walls, which is called "senile emphysema". This microscopic emphysema increased linearly with increasing age in non-smokers, but in smokers a more progressive increase in alveolar space size can be observed in certain (susceptible) individuals only. Senile emphysema can be a consequence of the loss of the supporting structures of the lung parenchyma. In addition, it has been observed that the elasticity of the lungs decreases with age. It has been postulated that this phenomenon is due more to the reduced surface tension force of the alveoli due to an increase in the size of the individual diameter compared to changes in elastin and collagen in the pulmonary parenchyma [21].
2. Changes in the chest wall. With increasing age, chest wall adherence progressively decreases. First, the shape of the thorax may change with age due to reduced thickness of the intervertebral discs, leading to reduced intrathoracic volume. Second, age-related osteoporosis can cause vertebral fractures, resulting in a change in the shape of the chest. Third, rib stiffness increases with age, thereby increasing the force required for chest movement [21].
3. Changes in the respiratory muscles. In general, muscle strength decreases with age. This loss of muscle strength is also reflected in

the diaphragm, the most important respiratory muscle, and thus affects the breathing pattern [21].

Malnutrition can be categorized by calculating the body mass index (BMI). BMI was often defined as body weight (kilograms) divided by height (meters) squared (kg/m^2). BMI is related to body fat which can be used to categorize obesity and malnutrition in both adults and children. BMI is also used to assess nutritional deficiencies, where a BMI $<18.5 \text{ kg}/\text{m}^2$ indicates nutritional deficiency [22]. Malnutrition was a common and serious problem in chronic lung disease patients, as COPD (chronic obstructive pulmonary disease) especially those with emphysema. COPD patients with emphysema have a lower body mass index (BMI) than COPD patients with chronic bronchitis [23,24,25]. Malnutrition in COPD was associated with complications and increased mortality. COPD patients with low body weight have a lower diffusion capacity and exercise than normal weight COPD patients. Reduced body cell mass was associated with reduced diaphragm mass and respiratory muscle mass. Malnutrition is also associated with decreased immune status, resulting in unwanted complications, such as nosocomial lung infections and hypercapnic lung failure [26].

The spirometric value that has been obtained from each respondent will be calculated the predictive value by comparing the spirometric value of each respondent with the normal FEV1 value, which was 3.2 liters. Then get the predicted value of each

respondent, then classified. In this characteristic, the chi-square test cannot be carried out because data was not obtained in the group without lung function disorders, so that it cannot be compared between the group without pulmonary function disorders and the group with lung function disorders. Average value data in the group without lung function disorders was 77.51, the highest value was 117, the lowest value was 70, the standard deviation was 7.19, while the average value in the group with pulmonary function disorders of 63.42, the highest value of 69, the lowest value of 37, the standard deviation was 6.32.

3.3 Research Results Knowledge

At the level of knowledge of risk factors for respiratory disorders, the two groups are mostly at a good level, and there are differences in the level of knowledge of risk factors for lung function disorders among respondents with and without pulmonary function disorders. At the level of knowledge of respiratory disorders symptoms, the two groups are mostly at a bad level, and there is no difference in the level of knowledge of risk factors for lung function disorders among respondents with and without pulmonary function disorders. Meanwhile, in the knowledge of medicine and respiratory disorders therapy, the two groups were mostly at a good level, and there was no difference in the level of knowledge of risk factors for lung function disorders among respondents with and without pulmonary function disorders (**Table 3**).

Table 3. Results of Knowledge of Risk Factors for Respiratory Disorders

Domain	Knowledge Level	Total Number of Respondents (n: 158)			Different Test	
		Group Without Lung Function Disorders (n: 79)	Group With Lung Function Disorders (n: 79)	Total	Pvalue	Conclusion
Knowledge of Risk Factors for Respiratory Disorders	Good	52	59	111	0.223	There was a difference in the level of knowledge of risk factors for lung function disorders among respondents with and without pulmonary function disorders
	Poor	27	20	47		
	Total	79	79	158		
Knowledge of Respiratory Disorders Symptoms	Good	32	30	62	0.745	there was no difference in the level of knowledge among respondents with and without pulmonary function disorders
	Poor	47	49	96		
	Total	79	79	158		
Knowledge of Medicine and Respiratory Disorders Therapy	Good	51	49	62	0.741	there was no difference in the level of knowledge among respondents with and without pulmonary function disorders
	Poor	28	30	96		
	Total	79	79	158		

The knowledge profile about the risk factors for respiratory disorders shows that there is no difference in the level of knowledge of the respondents with or without respiratory disorders. The knowledge profile about respiratory symptoms shows that there is no difference in the level of knowledge of respondents with or without respiratory disorders. And the knowledge profile about the treatment and therapy of respiratory disorders shows that there is no difference in the level of knowledge of respondents with or without respiratory disorders. Table 3 showed that there was a relationship between the level of knowledge about risk factors for respiratory disease ($p(0.223) < 0.05$) and lung function, but there was no difference in knowledge about symptoms ($p(0.745) < 0.05$) and treatment ($p(0.741) < 0.05$) respiratory disease with pulmonary function (**Table 3**).

The lung function test was a combination of studies carried out in clinical practice to determine lung capacity and possible decrease in mechanical function of the lungs, respiratory muscles, and chest wall. In clinical guidelines, pulmonary function interpretation has historically been based on the most important anthropometric factors, including weight, height, sex, and age [27]. In Lorensia et al. [28], in 148 respondents pedicab drivers as high risk chronic pulmonary disease (135 with respiratory disorder and 13 without respiratory disorder). The results of this study found the most common risk factor was air pollution exposure. In 145 respondents (97.97%), the average lung function was 52.56%. The most frequent early symptom was dyspnea with percentage of 7.43% and average lung function value of 26.20%. Air pollution was the most common risk factor and dyspnea was the most common early symptom in pedicab drivers with respiratory disorders in Surabaya.

4 Conclusions

The knowledge profile about the risk factors for respiratory disorders shows that there is no difference in the level of knowledge of the respondents with or without respiratory disorders. The knowledge profile about respiratory symptoms shows that there is no

difference in the level of knowledge of respondents with or without respiratory disorders. And the knowledge profile about the treatment and therapy of respiratory disorders shows that there is no difference in the level of knowledge of respondents with or without respiratory disorders.

5 Acknowledgments

Funding sources: this research was funded by the Institute of Research and Community Service of the Universitas Surabaya.

6 Author Contribution

All authors contributed starting from data collection, data analysis, and writing papers.

7 Ethical Clearance

No. 002-OL/KE/IV/2019 from Health Research Ethics Committee in Universitas Surabaya.

8 Conflicts of Interest

The authors declare no conflict of interest.

9 References

- [1] Badan Pusat Statistik, 2018, Statistik Daerah Kota Surabaya. Surabaya: Badan Pusat Statistik Kota Surabaya.
- [2] Moynihan, M.C., Allwood, J.M., 2014. Utilization of structural steel in buildings. *Proc Math Phys Eng Sci*, 470(2168), pp.20140170. doi:10.1098/rspa.2014.0170
- [3] Bauleo, L., Bucci, S., Antonucci, C., et al., 2019. Long-term exposure to air pollutants from multiple sources and mortality in an industrial area: a cohort study. *Occup Environ Med*, 76(1), pp.48-57. doi:10.1136/oemed-2018-105059
- [4] Manisalidis, I., Stavropoulou, E., Stavropoulos, A., Bezirtzoglou, E., 2020. Environmental and Health Impacts of Air Pollution: A Review. *Front Public Health*, 8, pp.14. doi:10.3389/fpubh.2020.00014
- [5] Wiston, M., 2017. Status of Air Pollution in Botswana and Significance to Air Quality and Human Health. *J Health Pollut*, 7(15), pp.8-17. doi:10.5696/2156-9614-7.15.8
- [6] Jiang, X.Q., Mei, X.D., Feng, D., 2016. Air pollution and chronic airway diseases: what should

- people know and do?. *J Thorac Dis*, 8(1), pp.E31-E40. doi:10.3978/j.issn.2072-1439.2015.11.50
- [7] Kurt, O.K., Zhang, J., Pinkerton, K.E., 2016. Pulmonary health effects of air pollution. *Curr Opin Pulm Med*, 22(2), pp.138-143. doi:10.1097/MCP.0000000000000248
- [8] Ghorani-Azam, A., Riahi-Zanjani, B., Balali-Mood, M., 2016. Effects of air pollution on human health and practical measures for prevention in Iran. *J Res Med Sci*, 21, pp.65. doi:10.4103/1735-1995.189646
- [9] Kim, D., Chen, Z., Zhou, L.F., Huang, S.X., 2018. Air pollutants and early origins of respiratory diseases. *Chronic Dis Transl Med*, 4(2), pp.75-94. doi:10.1016/j.cdtm.2018.03.003
- [10] Alvarado, A., Arce, I., 2016. Metabolic Functions of the Lung, Disorders and Associated Pathologies. *J Clin Med Res*, 8(10), pp.689-700. doi:10.14740/jocmr2668w
- [11] GOLD (Global Initiative For Chronic Lung Disease), 2018. Global Strategy for the Diagnosis, Management, and Prevention of Chronic Obstructive Pulmonary Disease (2018 Report) Global Initiative for Chronic Obstructive Disease, Inc, USA.
- [12] Dinas Kesehatan Surabaya. 2016. Profil Kesehatan Kota Surabaya. Surabaya: Dinas Kesehatan Surabaya.
- [13] Supriadi, L.S.R., Sui-Pheng, L., 2017. The Indonesian Construction Industry. *Business Continuity Management in Construction*, pp.117-154. doi:10.1007/978-981-10-5487-7_6
- [14] Aryal-Bhandari, A., Gautam, R., Bhandari, S., 2015. Knowledge and Practice on Prevention of Respiratory Health Problems among Traffic Police in Kathmandu, Nepal. *Int Sch Res Notices*, 2015, pp.716257. doi:10.1155/2015/716257
- [15] Missriya, S., Subashini, S., 2017. Assess The Prevalence of Respiratory Problem and Knowledge on Health Effects of People Residing Near Quarry. *Int J Pharm Pharm Sci*, 9(2), pp.250-253.
- [16] Grady, P.A., Gough, L.L., 2014. Self-management: a comprehensive approach to management of chronic conditions. *Am J Public Health*, 104(8), pp.e25-e31. doi:10.2105/AJPH.2014.302041
- [17] Ebrahimi-Belil, F., Alhani, F., Ebadi, A., Kazemnejad, A., 2018. Self-Efficacy of People with Chronic Conditions: A Qualitative Directed Content Analysis. *J Clin Med*, 7(11), pp.411. doi:10.3390/jcm7110411
- [18] Settineri, S., Frisone, F., Merlo, E.M., Geraci, D., Martino, G., 2019. Compliance, adherence, concordance, empowerment, and self-management: five words to manifest a relational maladjustment in diabetes. *J Multidiscip Healthc*, 12, pp.299-314. doi:10.2147/JMDH.S193752
- [19] Khan, F.U., Waqas, N., Ihsan, A.U., Khongorzul, P., Wazir, J., Gang, W., Mengqi, Y., Xiaoqian, L., Han, L., Zhou-Xiaohui, Z., 2019. Analysis of the Qualities Matching New Classification of Clinical Pharmacist. *Indian J Pharm Sci*, 81(1), pp.2-10.
- [20] Ranu, H., Wilde, M., Madden, B., 2011. Pulmonary function tests. *Ulster Med J*, 80(2), pp.84-90.
- [21] Brandsma, C.A., Vries, M.D., Costa, R., Woldhuis, R.R., Konigshoff, M., Timens, W., 2017. Lung ageing and COPD: is there a role for ageing in abnormal tissue repair?. *European Respiratory Review*, 26, pp.170073.
- [22] Nuttall, F.Q., 2015. Body Mass Index: Obesity, BMI, and Health: A Critical Review. *Nutr Today*, 50(3), pp.117-128. doi:10.1097/NT.0000000000000092
- [23] Rawal, G., Yadav, S., 2015. Nutrition in chronic obstructive pulmonary disease: A review. *J Transl Int Med*, 3(4), pp.151-154. doi:10.1515/jtim-2015-0021
- [24] Collins, P.F., Yang, I.A., Chang, Y.C., Vaughan, A., 2019. Nutritional support in chronic obstructive pulmonary disease (COPD): an evidence update. *J Thorac Dis*, 11(Suppl 17), pp.S2230-S2237. doi:10.21037/jtd.2019.10.41
- [25] Lorensia, A., Suryadinata, R.V., Sidabutar., 2021. Effect Analysis of Protein Intake of Pedicab Driver in Surabaya. *Journal of Tropical Pharmacy and Chemistry*, 5(3), pp.188-193.
- [26] Walson, J.L., Berkley, J.A., 2018. The impact of malnutrition on childhood infections. *Curr Opin Infect Dis*, 31(3), pp.231-236. doi:10.1097/QCO.0000000000000448
- [27] Barroso, A.T., Martin, E.M., Romero, L.M.R., Ruiz, F.O., 2018. Factors Affecting Lung Function: A Review of the Literature. *Arch Bronconeumol*, 54(6), pp.327-332.
- [28] Lorensia, A., Suryadinata, R.V., Diputra. I.N.Y., 2019. Risk Factors and Early Symptoms Related to Respiratory Disease in Pedicab Drivers in Surabaya. *Jurnal Kesehatan Masyarakat*, 15(2), pp.224-235.



[Current](#) [Archives](#) [Focus and Scope](#) [Article processing charge](#) [Author Guidelines](#)
[Announcements](#) [Scopus Citation Analysis](#) [About ▾](#)

[Search](#)



Journal of Tropical Pharmacy and Chemistry (J. Trop. Pharm. Chem)

[Journal of Tropical Pharmacy and Chemistry](#) (p-ISSN: 2087-7099, e-ISSN: [2407-6090](#)) is a Biannual, international, open access (Indexed in [SINTA 4](#) since 2016, and start from Volume 4 No. 4 2018 indexed in [SINTA 3](#) (SK No. 10/E/KPT/2019) by [Ministry of Research Technology and Higher Education, Indonesia](#), [GARUDA](#), [DOAJ](#), [Index Copernicus](#), [Crossref](#), [CiteFactor](#), [PKP Index](#), [BASE](#), [Moraref](#), [Google Scholar](#)), journal dedicated to various disciplines of pharmaceutical and allied sciences. The journal had been established in 2010, and online publication was

begun in 2014. Since 2018, the journal has been published in English by [Faculty of Pharmacy, Universitas Mulawarman](#) (UNMUL), Samarinda, Indonesia.

[Journal of Tropical Pharmacy and Chemistry](#) only receives manuscripts in English.

[Journal of Tropical Pharmacy and Chemistry](#) is a Six monthly, international, open access, journal dedicated to various disciplines of pharmaceutical and allied sciences. Journal of Tropical Pharmacy and Chemistry publishes manuscripts (Original research Article, review articles, Mini-reviews, and Short communication) on original work, either experimental or theoretical in the following areas: Pharmaceutics & Biopharmaceutics, Novel & Targeted Drug Delivery, Nanotechnology & Nanomedicine, Pharmaceutical Chemistry, Pharmacognosy & Ethanobotany, Phytochemistry, Pharmacology & Toxicology, Pharmaceutical Biotechnology & Microbiology, Pharmacy practice & Hospital Pharmacy, Pharmacogenomics, Pharmacovigilance, Natural Product Research, Drug Regulatory Affairs, Case Study & Full clinical trials, Biomaterials & Bioactive polymers, Analytical Chemistry, Organic Chemistry, Physical Pharmacy, Clinical Pharmacy.

Submission and publication in [Journal of Tropical Pharmacy and Chemistry](#) is **free of charge**.

[Journal of Tropical Pharmacy and Chemistry](#) Indexed by:



Journal Template



Hal yang harus diperhatikan oleh calon penulis

📅 2021-12-17

perhatian untuk calon penulis

[Read More](#) ➔

Author Contribution

 2020-07-16

Current Issue

Vol 5 No 4 (2021): Journal of Tropical Pharmacy and Chemistry

Published: 2021-12-31

Articles

Test of the Aphrodisiac Effect of Ethanol Extract of Roots, Stems and Leaves of Packaged Plants (*Smilax rotundifolia*) on Male White Rats (*Sprague Dawley*)

Kiki Rizki Handayani, Ayu Wulandari

280-285

 PDF

Proximate Composition, In vitro Antioxidant and Anti-inflammatory Properties of *Adansonia digitata* and *Belanites aegyptiaca* Seeds

Mercy Badu, Mary-Magdalene Pedavoah, Nathaniel O Boadi, Irene Y Dzaye

286-298

 PDF

Isolation and Evaluation of Antibacterial Potential Test of Plant *Carthamus oxycantha*

Amir Hassan, Ibad Ullah, Waqas Ahmad

299-308

 PDF

Quality Control and Anti-Inflammatory Activity of the Stem Bark of *Chlorophora regia* A. Chev. (*Moraceae*)

Quality Control of Herbal Products

Isaiah Amoo, James Oppong-Kyekyeku, Eric Boakye-Gyasi, Samuel Asare-Nkansah, Joseph Kwasi Adu

309-323

 PDF

Quality Test of Extemporaneously Prepared Tramadol and Paracetamol Capsules Combination Derived From a Private Hospital in Semarang

Paulus Unggul Wikan Prabandono, Michael Raharja Gani, Sri Hartati Yuliani

324-329



PDF

Evaluation of Antibacterial and Antifungal Activity of Herbs Used in Treatment of Diabetes, Malaria and Pneumonia in Kisii and Nyamira Counties Region, Kenya

Moses A. Guto Maobe, Leonard Gitu, Erastus Gatebe

330-337



PDF

Spectrophotometric Methods for the Determination of Caffeine in Beverages Use Solvent Extraction Techniques and Adsorption of Activated Carbon

Edy Agustian Yazid, Abdul Wafi, Agustin Eka Wulandari

338-346



PDF

Analysis of SARS-CoV-2 Spike Protein as The Key Target in the Development of Antiviral Candidates for COVID-19 through Computational Study

Taufik Muhammad Fakhri, Mentari Luthfika Dewi

347-352



PDF

Analysis of Ethyl p-Methoxycinnamate from Kaempferia galanga L. Extract by High Performance Liquid Chromatography

Wiwin Winingsih, Sri Gustini Husein, Rozalia Putri Neno Ramdhani

353-358



PDF

Analysis of Flavonoids and Terpenoids in Ethanol Extract of Colocasia esculenta L. (Schoot) Stalk and Leaves

Erni Rustiani, Aulia Fitriani, Sri Wardatun

359-364



PDF

Antioxidant Activity Test of Effervescent Granules Morinda citrifolia L Leaf Extract with DPPH Free Radical Absorption Method

Muthmainah Tuldjanah, Niluh Puspita Dewi, Dian Rahmawati
365-368

[PDF](#)

Physical Evaluation of Transfersome that Contains Pandan Leaves Extract (*Pandanus amaryllifolius* R.)

Rini Ambarwati, Yulianita Yulianita
369-374

[PDF](#)

Antibacterial Activity of Salam (*Syzygium polyanthum*) Leaves 70% Ethanolic Extract on *Staphylococcus aureus* and *Staphylococcus epidermidis*

Ayu Nala El Muna Haerussana, Wulan Putri Dwiastuti, Cindi Arwan Sukowati
375-380

[PDF](#)

Identification and Determination of Berberine from *Arcangelisia Flava*, East Borneo

Riski Sulistiarini
381-386

[PDF](#)

Relationship of Knowledge on Respiratory Disorders with Lung Function in Masons in East Surabaya

Amelia Lorensia, Rivan Virlando Suryadinata, Wilma Adib Gardiawan
387-395

[PDF](#)

[View All Issues >](#)

[Make a Submission](#)

Information

[For Readers](#)

[For Authors](#)

[For Librarians](#)

JOURNAL TEMPLATE

1. [RTF Version](#)
2. [Doc Version](#)
3. [Docx Version](#)

Reference Style

[1] Author, Year. Title. Journal. Volume. (Issue). Pages. (reference style for journal)

[2] Author, Year. Title. Publisher. Place (reference style for book)

Accepted Articles

[Article in Press](#)

PEOPLE

[Contact](#)

[Editorial Boards](#)

[Reviewer](#)

EDITORIAL INFORMATION

[Author Fees](#)

[Author Guidelines](#)

[Focus and Scope](#)

[Publication Ethics](#)

[About J.Trop.Pharm.Chem.](#)

Journal of Tropical Pharmacy and Chemistry Indexed by:

1. [DOAJ](#)
2. [Microsoft Academic](#)
3. [Index Copernicus](#)

4. [Sinta](#)
5. [Crossref](#)
6. [Google Scholar](#)
7. [PKP Index](#)
8. [Citefactor](#)
9. [Base](#)
10. [Keio](#)
11. [Worldcat](#)
12. [Dimensions](#)
13. [Scilit](#)
14. [Hinari](#)
15. [Road](#)
16. [Moraref](#)
17. [Garuda](#)

Accreditation

[SK Akreditasi](#)

[Accreditation Certificate](#)

TOOLS

[Grammarly](#)

[Endnote](#)

[Mendeley](#)

[Zotero](#)

[Turnitin](#)

STATISTIC

 [Web Analytics Made Easy - StatCounter View My Stats](#)

 [Flag Counter](#)

ADDRESS

Gedung Administrasi Fakultas Farmasi Jl. Penajam, Kampus Unmul Gunung Kelua, Kota Samarinda,
Kalimantan Timur, Indonesia, 75119

p-ISSN: 2087-7099; *e*-ISSN: [2407-6090](https://doi.org/10.24070/2407-6090)

Email: admin@farmasi.unmul.ac.id

Address: Gedung Administrasi Fakultas Farmasi Universitas Mulawarman, Jalan Penajam, Kampus UNMUL Gn. Kelua, Samarinda, 75119. Indonesia.



This work is licensed under a [Creative Commons Attribution-NonCommercial 4.0 International License](https://creativecommons.org/licenses/by-nc/4.0/).

Platform &
workflow by
OJS / PKP

Journal Of Tropical Pharmacy And Chemistry

eISSN : 24076090 | pISSN : 20877099

[Health](#) [Science](#) [Agriculture](#) [Engineering](#)

[Universitas Mulawarman](#)



S3

Sinta Score



Indexed by GARUDA

14

H-Index

13

H5-Index

668

Citations

629

5 Year Citations



Penerbit:

Fakultas Farmasi Universitas Mulawarman

[Website](#) | [Editor URL](#)

Address:

Gedung Administrasi Fakultas Farmasi Jl. Penajam Kampus UNMUL Gn. Kelua
Samarinda

Email:

jtpc_unmul@farmasi.unmul.ac.id

Phone:

0541-739491

Last Updated :

2021-12-04

2017

2018

2019



Search..



Page 1 of 23 | Total Records : 224

Publications

Citation

<u>Formulasi dan optimasi basis gel HPMC (hidroxy propyl methyl cellulose) dengan berbagai variasi konsentrasi</u> M Ardana, V Aeyni, A Ibrahim Journal of Tropical Pharmacy and Chemistry 3 (2), 101-108	53
<u>Hubungan antara tempat perindukan nyamuk Aedes aegypti dengan kasus demam berdarah dengue di Kelurahan Penajam Kecamatan Penajam Kabupaten Penajam Paser Utara</u> S Badrah, N Hidayah Journal of Tropical Pharmacy and Chemistry 1 (2), 150-157	32
<u>Aktivitas ekstrak daun salam (eugenia polyantha) sebagai antiinflamasi pada tikus putih (rattus norvegicus)</u> R Agustina, DT Indrawati, MA Masruhim Journal Of Tropical Pharmacy And Chemistry 3 (2), 120-123	31
<u>Optimalisasi ekstraksi dan uji metabolit sekunder tumbuhan liho (Ficus variegata Blume)</u> L Febrina, R Rusli, F Mufliah Journal of Tropical Pharmacy and Chemistry 3 (2), 74-81	29
<u>Potensi Tumbuhan Liho (Ficus variegata Blume) sebagai Sumber Bahan Farmasi Potensial</u>	

Publications

Journal of Tropical Pharmacy and Chemistry 2 (3), 166-179

Profil kromatografi senyawa aktif antioksidan dan antibakteri fraksi n-heksana daun Libo (Ficus variegata Blume)

R Rusli, MP Hardina, F Muflifah, A Rahmadani

Journal of Tropical Pharmacy and Chemistry 3 (2), 124-130

22

Jamur Endofit, Biodiversitas, Potensi Dan Prospek Penggunaannya Sebagai Sumber Bahan Obat Baru

H Kuncoro, NE Sugijanto

Journal of Tropical Pharmacy and Chemistry 1 (3), 247-262

19

Identifikasi metabolit sekunder dan aktivitas antibakteri ekstrak daun sungkai (Peronema canescens Jack.) terhadap beberapa bakteri patogen

A Ibrahim, H Kuncoro

Journal of Tropical Pharmacy and Chemistry 2 (1), 8-18

16

Pengaruh Ekstrak Etanol Daun Jambu Air (Syzygium aqueum (Burm.f.) Alston) terhadap Glukosa Darah, Ureum dan Kreatinin Tikus Putih (Rattus norvegicus)

J Tandi

Journal of Tropical Pharmacy and Chemistry 4 (2), 43-51

15

Pengembangan sistem nanostructured lipid carriers (NLC) meloxicam dengan lipid monostearin dan miglyol 808 menggunakan metode emulsifikasi

R Annisa, E Hendradi, D Melani

Journal of Tropical Pharmacy and Chemistry 3 (3), 156-169


15

Page 1 of 23 | Total Records : 224



Citation Statistics





Copyright © 2017

Kementerian Pendidikan, Kebudayaan, Riset, dan Teknologi

(The Ministry of Education, Culture, Research, and Technology)

All Rights Reserved.



[Current](#) [Archives](#) [Focus and Scope](#) [Article processing charge](#) [Author Guidelines](#)
[Announcements](#) [Scopus Citation Analysis](#) [About ▾](#)

[Search](#)

[Home](#) / [Editorial Team](#)

Editorial In Chief:

Dr. [Hadi Kuncoro](#), M.Farm., Apt. Faculty of Pharmacy, Universitas Mulawarman; [google scholar profile](#), [researchgate profile](#), & [orcid ID](#)

Co Editor:

1. Prof. Dr. [Laode Rijai](#), M.Si., Drs. Faculty of Pharmacy, Universitas Mulawarman; [google scholar profile](#)
2. Dr. [Rolan Rusli](#), M.Si. Faculty of Pharmacy, Universitas Mulawarman; [google scholar profile](#), [researchgate profile](#), [orcid ID](#), & [Publons](#)

Advisory Editor:

1. Prof. Dr. [Deddi Prima Putra](#), Apt. (Universitas Andalas)
2. [Pinus Jumaryatno](#), M.Phill., Ph.D., Apt. (Universitas Islam Indonesia)
3. [Nasrul Wathoni](#), Ph. D., Apt. (Universitas Padjadjaran)

4. Prof. Dr. [Yoshihito Shiono](#) (Yamagata University, Japan)
5. Prof. [Muchtari M](#), Ph. D., M.Si., Apt. (Universitas Padjadjaran)
6. Prof. Dr. [Unang Supratman](#) (Universitas Padjadjaran)
7. Dr. [Aty Widawaruyanti](#), MS., Apt. (Universitas Airlangga)
8. Prof. Dr. [Taslim Ersam](#), MS. (ITS)

[Make a Submission](#)

Information

[For Readers](#)

[For Authors](#)

[For Librarians](#)

JOURNAL TEMPLATE

1. [RTF Version](#)
2. [Doc Version](#)
3. [Docx Version](#)

Reference Style

[1] Author, Year. Title. Journal. Volume. (Issue). Pages. (reference style for journal)

[2] Author, Year. Title. Publisher. Place (reference style for book)

Accepted Articles

[Article in Press](#)

PEOPLE

[Contact](#)

[Editorial Boards](#)

[Reviewer](#)

EDITORIAL INFORMATION

[Author Fees](#)

[Author Guidelines](#)

[Focus and Scope](#)

[Publication Ethics](#)

[About J.Trop.Pharm.Chem.](#)

Journal of Tropical Pharmacy and Chemistry Indexed by:

1. [DOAJ](#)
2. [Microsoft Academic](#)
3. [Index Copernicus](#)
4. [Sinta](#)
5. [Crossref](#)
6. [Google Scholar](#)
7. [PKP Index](#)
8. [Citefactor](#)
9. [Base](#)
10. [Keio](#)
11. [Worldcat](#)
12. [Dimensions](#)
13. [Scilit](#)
14. [Hinari](#)
15. [Road](#)
16. [Moraref](#)
17. [Garuda](#)

Accreditation

[SK Akreditasi](#)

[Accreditation Certificate](#)

TOOLS

[Grammarly](#)

[Endnote](#)

[Mendeley](#)

[Zotero](#)

[Turnitin](#)

STATISTIC

 [Web Analytics Made Easy - StatCounter View My Stats](#)

 [Flag Counter](#)

ADDRESS

Gedung Administrasi Fakultas Farmasi Jl. Penajam, Kampus Unmul Gunung Kelua, Kota Samarinda, Kalimantan Timur, Indonesia, 75119

p-ISSN: 2087-7099; *e*-ISSN: [2407-6090](#)

Email: admin@farmasi.unmul.ac.id

Address: Gedung Administrasi Fakultas Farmasi Universitas Mulawarman, Jalan Penajam, Kampus UNMUL Gn. Kelua, Samarinda, 75119. Indonesia.



This work is licensed under a [Creative Commons Attribution-NonCommercial 4.0 International License](#).

Platform &
workflow by
OJS / PKP

[Current](#)[Archives](#)[Focus and Scope](#)[Article processing charge](#)[Author Guidelines](#)[Announcements](#)[Scopus Citation Analysis](#)[About ▾](#)[Search](#)

[Home](#) / [Archives](#) / Vol 5 No 4 (2021): Journal of Tropical Pharmacy and Chemistry

DOI: <https://doi.org/10.25026/jtpc.v5i4>

Published: 2021-12-31

Articles

Test of the Aphrodisiac Effect of Ethanol Extract of Roots, Stems and Leaves of Packaged Plants (*Smilax rotundifolia*) on Male White Rats (*Sprague Dawley*)

Kiki Rizki Handayani, Ayu Wulandari

280-285

[PDF](#)

Proximate Composition, In vitro Antioxidant and Anti-inflammatory Properties of *Adansonia digitata* and *Belanites aegyptiaca* Seeds

Mercy Badu, Mary-Magdalene Pedavoah, Nathaniel O Boadi, Irene Y Dzaye

286-298

[PDF](#)

Isolation and Evaluation of Antibacterial Potential Test of Plant *Carthamus oxycantha*

Amir Hassan, Ibad Ullah, Waqas Ahmad

299-308



PDF

Quality Control and Anti-Inflammatory Activity of the Stem Bark of *Chlorophora regia* A. Chev. (Moraceae)

Quality Control of Herbal Products

Isaiah Amoo, James Oppong-Kyekyeku, Eric Boakye-Gyasi, Samuel Asare-Nkansah, Joseph Kwasi Adu

309-323



PDF

Quality Test of Extemporaneously Prepared Tramadol and Paracetamol Capsules Combination Derived From a Private Hospital in Semarang

Paulus Unggul Wikan Prabandono, Michael Raharja Gani, Sri Hartati Yuliani

324-329



PDF

Evaluation of Antibacterial and Antifungal Activity of Herbs Used in Treatment of Diabetes, Malaria and Pneumonia in Kisii and Nyamira Counties Region, Kenya

Moses A. Guto Maobe, Leonard Gitu, Erastus Gatebe

330-337



PDF

Spectrophotometric Methods for the Determination of Caffeine in Beverages Use Solvent Extraction Techniques and Adsorption of Activated Carbon

Edy Agustian Yazid, Abdul Wafi, Agustin Eka Wulandari

338-346



PDF

Analysis of SARS-CoV-2 Spike Protein as The Key Target in the Development of Antiviral Candidates for COVID-19 through Computational Study

Taufik Muhammad Fakhri, Mentari Luthfika Dewi

347-352



PDF

Analysis of Ethyl p-Methoxycinnamate from *Kaempferia galanga* L. Extract by High Performance Liquid Chromatography

Wiwin Winingsih, Sri Gustini Husein, Rozalia Putri Neno Ramdhani

353-358



Analysis of Flavonids and Terpenoids in Ethanol Extract of *Colocasia esculenta* L. (Schoot) Stalk and Leaves

Erni Rustiani, Aulia Fitriani, Sri Wardatun

359-364



Antioxidant Activity Test of Effervescent Granules *Morinda citrifolia* L Leaf Extract with DPPH Free Radical Absorption Method

Muthmainah Tuldjanah, Niluh Puspita Dewi, Dian Rahmawati

365-368



Physical Evaluation of Transfersome that Contains Pandan Leaves Extract (*Pandanus amaryllifolius* R.)

Rini Ambarwati, Yulianita Yulianita

369-374



Antibacterial Activity of Salam (*Syzygium polyanthum*) Leaves 70% Ethanolic Extract on *Staphylococcus aureus* and *Staphylococcus epidermidis*

Ayu Nala El Muna Haerussana, Wulan Putri Dwiastuti, Cindi Arwan Sukowati

375-380



Identification and Determination of Berberine from *Arcangelisia Flava*, East Borneo

Riski Sulistiarini

381-386



Relationship of Knowledge on Respiratory Disorders with Lung Function in Masons in East Surabaya

Amelia Lorensia, Rivan Virlando Suryadinata, Wilma Adib Gardiawan

387-395



[Make a Submission](#)

Information

[For Readers](#)

[For Authors](#)

[For Librarians](#)

JOURNAL TEMPLATE

1. [RTF Version](#)
2. [Doc Version](#)
3. [Docx Version](#)

Reference Style

[1] Author, Year. Title. Journal. Volume. (Issue). Pages. (reference style for journal)

[2] Author, Year. Title. Publisher. Place (reference style for book)

Accepted Articles

[Article in Press](#)

PEOPLE

[Contact](#)

[Editorial Boards](#)

[Reviewer](#)

EDITORIAL INFORMATION

[Author Fees](#)

[Author Guidelines](#)

[Focus and Scope](#)

[Publication Ethics](#)

[About J.Trop.Pharm.Chem.](#)

Journal of Tropical Pharmacy and Chemistry Indexed by:

1. [DOAJ](#)
2. [Microsoft Academic](#)
3. [Index Copernicus](#)
4. [Sinta](#)
5. [Crossref](#)
6. [Google Scholar](#)
7. [PKP Index](#)
8. [Citefactor](#)
9. [Base](#)
10. [Keio](#)
11. [Worldcat](#)
12. [Dimensions](#)
13. [Scilit](#)
14. [Hinari](#)
15. [Road](#)
16. [Moraref](#)
17. [Garuda](#)

Accreditation

[SK Akreditasi](#)

[Accreditation Certificate](#)

TOOLS

[Grammarly](#)

[Endnote](#)

[Mendeley](#)

[Zotero](#)

[Turnitin](#)

STATISTIC

 [Web Analytics Made Easy - StatCounter View My Stats](#)

 [Flag Counter](#)

ADDRESS

Gedung Administrasi Fakultas Farmasi Jl. Penajam, Kampus Unmul Gunung Kelua, Kota Samarinda, Kalimantan Timur, Indonesia, 75119

p-ISSN: 2087-7099; *e*-ISSN: [2407-6090](#)

Email: admin@farmasi.unmul.ac.id

Address: Gedung Administrasi Fakultas Farmasi Universitas Mulawarman, Jalan Penajam, Kampus UNMUL Gn. Kelua, Samarinda, 75119. Indonesia.



This work is licensed under a [Creative Commons Attribution-NonCommercial 4.0 International License](#).

Platform &
workflow by
OJS / PKP

Relationship of Knowledge on Respiratory Disorders with Lung Function in Masons in East Surabaya

by Amelia Lorensia

Submission date: 25-Dec-2021 10:02PM (UTC+0700)

Submission ID: 1735616496

File name: 342-Article_Text-992-1-10-20211225_1.pdf (292.49K)

Word count: 5615

Character count: 29826



Relationship of Knowledge on Respiratory Disorders with Lung Function in Masons in East Surabaya

Amelia Lorensia^{1,*}, Rivan Virlando Suryadinata², Wilma Adib Gardiawan³

¹Department of Clinical-Community Pharmacy, Faculty of Pharmacy, Universitas Surabaya (UBAYA)

²Department of Public Health, Faculty of medicine, Universitas Surabaya (UBAYA)

³Bachelor Student of Faculty of Pharmacy, Faculty of Pharmacy, Universitas Surabaya (UBAYA)

*Corresponding author: amelia.lorensia@staff.ubaya.ac.id

Abstract

Masons are regularly exposed to air pollution in the workplace. The pollution exposure received by them is a risk factor for respiratory problems. Most of the masons had insufficient knowledge that is very important for the management of a person's illness. The purpose of this study was to determine relationship of knowledge on respiratory disorders with lung function in masons in east surabaya. This research was an observational study with a cross-sectional design with a purposive method. The research was conducted from April to July 2019 in the East Surabaya area. The variables that will be observed in this study are the level of knowledge of risk factors, symptoms, and treatment and therapy regarding respiratory disorders. In this study, there were 158 respondents consisting of 79 groups with impaired lung function and 79 groups without lung function disorders. Most of the level of knowledge of both groups was good in risk factors and treatment of respiratory disease, but instead on the symptoms regarding respiratory disease. The research showed that there was a relationship between the level of knowledge about risk factors for respiratory disease ($p(0.223) < 0.05$) and lung function, but there was no difference in knowledge about symptoms ($p(0.745) < 0.05$) and treatment ($p(0.741) < 0.05$) respiratory disease with pulmonary function. Therefore, it was necessary to plan further educational programs in increasing masons' knowledge of respiratory symptoms so that they could catch respiratory diseases earlier.

Keywords: knowledge, lung function, masons, respiratory disorders

Submitted: 03 May 2021

Accepted: 25 December 2021

DOI: <https://doi.org/10.25026/jtpc.v5i4.342>

1 Introduction

Surabaya is the second metropolitan city in Indonesia with a total area of 326.36 km². In 2017 the city of Surabaya has a population of 2.87 million and is predicted to increase every year [1]. The efforts of the Surabaya City government in realizing development planning require labor, one of which is a builder. A builder is a person who works by relying on his physical strength and has skills in handwork (digger, bricklayer, blacksmith, carpenter, painter) in making buildings in general (such as buildings, houses, shops) [2]. Masons have a very high risk of being exposed to air pollution [3,4]. According to research data in 2010, more than 50% of masons are regularly exposed to air pollution such as steam, gas, dust, or smoke in the workplace [5].

The pollution exposure received by masons is a risk factor for respiratory problems [6]. A work environment full of air pollution will have a negative impact on lung health [7]. Air pollution can have a direct or indirect effect on environmental quality degradation, namely air pollution [4,8]. The impact of air pollutants on the respiratory system has been consistently reported in recent years. Adverse effects include decreased lung function, increased infection, increased respiratory symptoms, acute exacerbations, onset of asthma, and increased respiratory mortality [6].

Decreased lung function in the form of respiratory disorders that occur in the respiratory tract and lungs that affect the human respiratory system [10]. Respiratory disorders can also be seen from spirometry measurements by looking at the FEV1/FVC value, where if the FEV1/FVC (Forced Expiratory Volume 1/ Forced Vital Capacity) value is <0.7 then it can be stated that you have respiratory problems [11]. In 2017, respiratory system diseases were also included in the top 10 most diseases in Surabaya and even ranked first [12]. Most of the masons had insufficient knowledge. Based on the results of the study, where most of the mason respondents were at the elementary education level and dropped out of school [13]. Lack of knowledge can affect the prevention of respiratory disorders. According to research conducted by Bhandari et al., it

shows that 88.0% of respondents have a below-average level of knowledge to prevent respiratory problems. Likewise, 41.0% of respondents had an average level of knowledge regarding the prevention of respiratory problems [14]. Knowledge of respiratory health shows that 65% of people have insufficient knowledge, 35% of people have moderate knowledge and no one has sufficient knowledge [15].

Knowledge is very important for the management of a person's illness, self-management in order to avoid a disease [16,17]. According to research, if a person has more knowledge of the disease, it is more likely that someone will obey the treatment [18]. Therefore, pharmacists are required to provide good health services to patients and pharmacists must have knowledge, skills and behavior that are always improved [19]. The purpose of this study was to determine differences in knowledge regarding (1) risk factors for respiratory disorders; (2) respiratory symptoms; and (3) treatment and therapy of respiratory disorders in masons with impaired lung function and without pulmonary function disorders.

2 Materials and Methods

2.1 Research design

The research was conducted from April to July 2019 in the East Surabaya area. This study was an observational study using a cross sectional study design to determine the profile of knowledge about respiratory disorders in builders based on lung function. Retrieval of research data by means of direct question and answer (interview) between the interviewer and the respondent.

2.2 Research variable

This study has two variables, namely the independent variable and the dependent variable. The independent variable of this study was impaired lung function. While the dependent variable in this study was the level of knowledge related to respiratory disorders.

The builder referred to in this research was a person who works by relying on his physical strength and has skill in a handwork (digger, bricklayer, bricklayer, carpenter, painter) in making buildings (such as buildings, houses, shops). Lung function disorders were diseases and disorders that occur in the respiratory tract and lungs that affect human respiration [10]. A person's lung function can be known through spirometric measurements, if the measurement using spirometry shows the results of the FEV1 value <70%, it can be said that you have lung function disorders [20]. The knowledge referred to in this research was knowledge about risk factors, symptoms, and treatment and therapy that the respondent has regarding respiratory disorders.

2.3 Population and Research Sample

The population in this study were masons who were working on a project or were active in the East Surabaya area. The sample in this study were masons who were in the East Surabaya area who were included in the inclusion and exclusion criteria. Respondents who are willing to participate in this research and have filled in the informed consent, are male, b. Age 17 - 65 years, active smokers, minimum length of work for 5 years, and no mental or physical disorders. The sampling technique used was random sampling (nonprobability sampling) with purposive sampling and consecutive sampling methods. The number of samples taken in this study using the Lemeshow formula with an unknown or infinite population. The following is the formula used: $n = Z^2 [P(1-P)] / d^2$ (n= number of samples; Z=standard normal value (if $\alpha = 0.05$ then $Z = 1.960$); $P(1-P)$ =estimated population proportion (if $P=0.1$ then $P(1-P)=0.09$); d^2 : tolerant deviation (10%)). So the minimum number of samples needed in this study is 62 respondents for each group.

2.4 Method of collecting data

Respondents will be given a questionnaire which was one of the instruments, important in research, the questionnaire given was about knowledge related to respiratory disorders. The questionnaire in this study consisted of 25 questions, of which 11 questions about knowledge of risk factors for respiratory disorders, 6 questions about knowledge of

respiratory symptoms, and 8 questions about knowledge about treatment and therapy.

Pulmonary function measurements were carried out using a Handled Spirometer where BMI data in the form of body weight and height as well as respondent data such as gender, age, smokers or non-smokers were also entered into the spirometer. The results of the spirometer measurement are used to determine respondents who have respiratory problems or do not experience respiratory problems based on FEV1, if FEV1 <70% indicates respiratory problems [20]. So if FEV1 <70% it can be categorized in the group that has impaired lung function and if FEV1 >70% can be categorized in the group that does not experience lung function disorders.

2.5 Data analysis method

Data processing was carried out in stages, starting with the data collected in the field until the data was ready for analysis. The data that has been collected in the field was checked, if there was a lack of data, it will be supplemented with repeated interviews with research respondents. The answers from the respondents will be coded to make the data input process easier. Then the data were analyzed descriptively (minimum, maximum, average and standard deviation values) and statistically using SPSS version 24 software.

The expected value in the chi-square should not be less than 5 (maximum 20% of expected frequencies <5). If the expected value was not met (20% expected frequencies <5), then the chi-square must be replaced with an alternative test and in determining the P value, if the P value was greater than 0.05 then there was no significant difference, but if the P value was more smaller than 0.05, there was a significant difference.

The level of knowledge of risk factors for respiratory disorders was measured using a questionnaire about respiratory disorders in which there were 11 questions. Every question that was answered correctly will be given a score of 1, while questions that are answered incorrectly or don't know will be given a score of 0. Then the level of knowledge will be classified as good and bad. It was said to be good if it gets a score of ≥ 8 and it is said to be bad if it gets a score of <8. Then the chi-square test was

carried out and the P_{value} was obtained of 0.223, which means that there was no significant difference between the group with pulmonary function disorders and the group without lung function disorders.

The level of knowledge of respiratory symptoms was measured using a questionnaire about respiratory disorders in which there are 6 questions. Every question that was answered correctly will be given a score of 1, while questions that were answered incorrectly or don't know will be given a score of 0. Then the level of knowledge will be classified as good and bad. It was said to be good if it gets a score of ≥ 4 and it was said to be bad if it gets a score of <4 . Then performed the chi-square test and obtained a P_{value} of 0.745 which means none a significant difference between the group with impaired lung function and the group without pulmonary function impairment.

The level of knowledge of respiratory symptoms was measured using a questionnaire about respiratory disorders in which there were 8 questions. Every question that was answered correctly will be given a score of 1, while questions that were answered incorrectly or don't know will be given a score of 0. Then the level of knowledge will be classified as good and bad. It was said good if it gets a score of ≥ 6 and it was said to be bad if it gets a score <6 . Then the chi-square test is carried out and the P_{value} was obtained of 0.741 which means no a significant difference between the group with impaired lung function and the group without pulmonary function impairment.

3 Results and Discussion

This research was conducted in April-July 2019 by conducting direct interviews with respondents using a questionnaire about respiratory disorders to determine the knowledge profile of risk factors, symptoms, and treatment and therapy for respiratory disorders based on lung function in masons in East Surabaya. Respondent data obtained in this study were 158 masons consisting of 79 people with pulmonary function disorders and 79 people without pulmonary function disorders. There were 20 respondents who refused to fill out the questionnaire, where 16 people did not want their rest time to be disturbed and 4 people did not meet the criteria.

3.1 Validity and Reliability of the Questionnaire

The results of the validity test of the risk factor knowledge questionnaire were declared valid. The questionnaire can be said to be valid if $r_{\text{count}} > r_{\text{table}}$ product moment, where the value of r_{table} for 30 respondents was 0.361 and all question items from the questionnaire have a value greater than the value of r_{table} product moment. The results of the reliability test of the risk factor knowledge questionnaire were declared reliable because they had a Cronbach's Alpha value equal to or greater than 0.61 (Cronbach's Alpha ≥ 0.61). The results of the reliability test of the risk factor knowledge questionnaire that have been obtained are declared reliable. The results of the validity of the questionnaire for knowledge of respiratory symptoms were declared valid because the value of $r_{\text{count}} > r_{\text{table}}$. Then then the questionnaire will be tested for reliability. The results of the questionnaire reliability test were declared reliable because Cronbach's Alpha value was ≥ 0.61 (Table 1).

3.2 Characteristics of Research Respondents

Respondents in this study were grouped into 2 groups, namely the group without lung function disorders and the group with lung function disorders. Respondents used in both groups were 158 respondents with 79 respondents in each group. Characteristics of respondents based on age consisted of late adolescents (17-25 years), early adults (26-35 years), late adults (36-45 years), early seniors (46-55 years), and late elderly (56-65 years old). Respondents without pulmonary function disorders in late adolescence were 20 respondents, early adults 33 respondents, late adults 13 respondents, early elderly 10 respondents, and late elderly by 3 respondents. Meanwhile, respondents with lung function disorders consisted of 15 respondents as late as adolescence, 45 respondents for early adulthood, 15 respondents for late adults, 3 respondents for early elderly, and 1 respondent for late elderly. Based on the age characteristics after testing the differences in the two groups showed that there was no significant difference between the two groups (Table 2).

Table 1. Validity and Reliability Test

Domain	Validity Test		Corrected Item-Total Correlation	Conclusion	Reliability Test	
	No.	Question			Cronbach's Alpha	Conclusion
Knowledge of Risk Factors for Respiratory Disorders	1	Environments with high pollution levels have greater risk of respiratory distress	0.457	Valid	0.614	Valid
	2	The smoke from the combustion/ factory /mine is a pollution which can cause respiratory distress	0.417	Valid		
	3	Smoking is not a cause of respiratory distress	0.419	Valid		
	4	The more the number of cigarettes smoked then further increase the risk factors for the disorder respiration	0.551	Valid		
	5	People working in dusty environments not experiencing respiratory distress	0.509	Valid		
	6	People who work in industrial areas have poor lung condition due to pollution in the area industry	0.411	Valid		
	7	People living in cities are more at risk experiencing respiratory distress than staying in the village	0.503	Valid		
	8	People living in industrial areas are more risk of experiencing respiratory distress	0.411	Valid		
	9	Increasing age does not increase the risk respiratory disorders	0.539	Valid		
	10	Respiratory disorders cannot be inherited or derived from the biological family	0.400	Valid		
	11	History of respiratory disorders in infants and children can present with disabilities throughout their lifetime adult	0.374	Valid		
Knowledge Respiratory Disorders Symptoms	1	Shortness of breath is not a symptom of the disorder inhalation	0.481	Valid	0.612	Valid
	2	Shortness of breath in respiratory disorders can getting worse when doing	0.518	Valid		
	3	Cough that is continuous/ not persistent healing is a symptom of the disorder respiration	0.564	Valid		
	4	Increased respiratory rate is common in patients with respiratory disorders	0.707	Valid		
	5	Wheezing (such as a whistling sound or a 'ngic' sound during breathing) is a symptom of respiratory distress	0.520	Valid		
	6	The respiratory rate is increased in respiratory distress not a sign of worsening condition	0.704	Valid		
Knowledge Medicine and Respiratory Disorders Therapy	1	Cough in respiratory distress may reduce with treatment (medicine) given	0.502	Valid	0.631	Valid
	2	Treatment (medication) of respiratory disorders is given only for symptom relief	0.476	Valid		
	3	Treatment (drugs) given can cure respiratory disorders	0.722	Valid		
	4	Quitting smoking has no effect reduction of respiratory distress symptoms	0.565	Valid		
	5	Breathing exercises aim to improve effort to breathe	0.434	Valid		
	6	Quitting smoking can make you breathe for the better	0.528	Valid		
	7	Breathing muscle training is not a viable exercise reduce shortness of breath in sufferers of the disorder respiration	0.567	Valid		
	8	Body activity program (sports) for increase tolerance of physical activity	0.435	Valid		

Table 2. Respondent Characteristics Frequency Distribution

Characteristics of Respondents	Total Number of Respondents (n: 158)				Different Test		
	Group Without Lung Function Disorders (n: 79)		Group With Lung Function Disorders (n: 79)		Pvalue	Conclusion	
	Frequency	Percentage (%)	Frequency	Percentage (%)			
Age (year)	Late adolescents (17-25 years)	20	25.32	15	18.99	0.113	no difference
	Early adults (26-35 years)	33	41.77	45	56.96		
	Late adults (36-45 years)	13	16.46	15	18.99		
	Early seniors (46-55 years)	10	12.66	3	3.79		
	Late elderly (56-65 years old)	3	3.79	1	1.26		
BMI (Kg/m²)	Thin (<18.5)	7	8.86	6	7.59	0.485	no difference
	Normal (18.5-25.0)	59	74.68	66	83.54		
	Overweight (25.0-27.0)	8	10.13	5	6.33		
	Obesity	5	6.33	2	2.53		
Spirometry value [11]	Mild (FEV1 >80% predicted)	0	0	66	83.54	-	-
	Worsening (50% < FEV1 < 80% predicted)	0	0	13	16.46		

Characteristics of respondents based on BMI were classified into 4, namely thin (<18.5), normal (18.5-25.0), overweight (25.0-27.0), and obesity (≥ 27.0). Respondents without pulmonary function disorders with a thin BMI were 7 respondents, normal was 59 respondents, overweight was 8 respondents, and obesity was 5 respondents. Meanwhile, 6 respondents with lung function disorders with a thin BMI, 66 normal respondents, 5 respondents with overweight, and 2 respondents with obesity. After testing the differences in the two groups, it was found that there was no significant difference in the two groups. Characteristics of respondents based on spirometric values to see lung function in groups with respiratory disorders will be classified based on GOLD [11] into 2, namely mild and worsening (Table 2).

Chronic lung disease was a major public health problem in the world. Although the chronic lung disease database on these outcome parameters is inadequate and often less reliable than mortality data, limited data suggest that chronic lung disease morbidity increases with age. Although the development of chronic lung disease comorbidities may occur at a younger age. Chronic lung disease was also more prevalent at age >40 years than <40 years, and was more common in men than women. Much of the increase in COPD mortality was due to the progression of the epidemic from smoking, the decrease in mortality from other common causes of death such as ischemic heart disease, infectious diseases [11]. We can broadly divide

the structural changes of aging lungs into three categories:

1. Changes in the structure of the lungs. Changes in the structure of the lungs are mainly associated with an increase in the size of the alveolar spaces in the absence of inflammation or damage to the alveolar walls, which is called "senile emphysema". This microscopic emphysema increased linearly with increasing age in non-smokers, but in smokers a more progressive increase in alveolar space size can be observed in certain (susceptible) individuals only. Senile emphysema can be a consequence of the loss of the supporting structures of the lung parenchyma. In addition, it has been observed that the elasticity of the lungs decreases with age. It has been postulated that this phenomenon is due more to the reduced surface tension force of the alveoli due to an increase in the size of the individual diameter compared to changes in elastin and collagen in the pulmonary parenchyma [21].
2. Changes in the chest wall. With increasing age, chest wall adherence progressively decreases. First, the shape of the thorax may change with age due to reduced thickness of the intervertebral discs, leading to reduced intrathoracic volume. Second, age-related osteoporosis can cause vertebral fractures, resulting in a change in the shape of the chest. Third, rib stiffness increases with age, thereby increasing the force required for chest movement [21].
3. Changes in the respiratory muscles. In general, muscle strength decreases with age. This loss of muscle strength is also reflected in

the diaphragm, the most important respiratory muscle, and thus affects the breathing pattern [21].

Malnutrition can be categorized by calculating the body mass index (BMI). BMI was often defined as body weight (kilograms) divided by height (meters) squared (kg/m^2). BMI is related to body fat which can be used to categorize obesity and malnutrition in both adults and children. BMI is also used to assess nutritional deficiencies, where a BMI $<18.5 \text{ kg}/\text{m}^2$ indicates nutritional deficiency [22]. Malnutrition was a common and serious problem in chronic lung disease patients, as COPD (chronic obstructive pulmonary disease) especially those with emphysema. COPD patients with emphysema have a lower body mass index (BMI) than COPD patients with chronic bronchitis [23,24,25]. Malnutrition in COPD was associated with complications and increased mortality. COPD patients with low body weight have a lower diffusion capacity and exercise than normal weight COPD patients. Reduced body cell mass was associated with reduced diaphragm mass and respiratory muscle mass. Malnutrition is also associated with decreased immune status, resulting in unwanted complications, such as nosocomial lung infections and hypercapnic lung failure [26].

The spirometric value that has been obtained from each respondent will be calculated the predictive value by comparing the spirometric value of each respondent with the normal FEV1 value, which was 3.2 liters. Then get the predicted value of each

respondent, then classified. In this characteristic, the chi-square test cannot be carried out because data was not obtained in the group without lung function disorders, so that it cannot be compared between the group without pulmonary function disorders and the group with lung function disorders. Average value data in the group without lung function disorders was 77.51, the highest value was 117, the lowest value was 70, the standard deviation was 7.19, while the average value in the group with pulmonary function disorders of 63.42, the highest value of 69, the lowest value of 37, the standard deviation was 6.32.

3.3 Research Results Knowledge

At the level of knowledge of risk factors for respiratory disorders, the two groups are mostly at a good level, and there are differences in the level of knowledge of risk factors for lung function disorders among respondents with and without pulmonary function disorders. At the level of knowledge of respiratory disorders symptoms, the two groups are mostly at a bad level, and there is no difference in the level of knowledge of risk factors for lung function disorders among respondents with and without pulmonary function disorders. Meanwhile, in the knowledge of medicine and respiratory disorders therapy, the two groups were mostly at a good level, and there was no difference in the level of knowledge of risk factors for lung function disorders among respondents with and without pulmonary function disorders (Table 3).

Table 3. Results of Knowledge of Risk Factors for Respiratory Disorders

Domain	Knowledge Level	Total Number of Respondents (n: 158)		Total	Different Test	
		Group Without Lung Function Disorders (n: 79)	Group With Lung Function Disorders (n: 79)		Pvalue	Conclusion
Knowledge of Risk Factors for Respiratory Disorders	Good	52	59	111	0.223	There was a difference in the level of knowledge of risk factors for lung function disorders among respondents with and without pulmonary function disorders
	Poor	27	20	47		
	Total	79	79	158		
Knowledge of Respiratory Disorders Symptoms	Good	32	30	62	0.745	there was no difference in the level of knowledge among respondents with and without pulmonary function disorders
	Poor	47	49	96		
	Total	79	79	158		
Knowledge of Medicine and Respiratory Disorders Therapy	Good	51	49	62	0.741	there was no difference in the level of knowledge among respondents with and without pulmonary function disorders
	Poor	28	30	96		
	Total	79	79	158		

The knowledge profile about the risk factors for respiratory disorders shows that there is no difference in the level of knowledge of the respondents with or without respiratory disorders. The knowledge profile about respiratory symptoms shows that there is no difference in the level of knowledge of respondents with or without respiratory disorders. And the knowledge profile about the treatment and therapy of respiratory disorders shows that there is no difference in the level of knowledge of respondents with or without respiratory disorders. Table 3 showed that there was a relationship between the level of knowledge about risk factors for respiratory disease ($p(0.223)<0.05$) and lung function, but there was no difference in knowledge about symptoms ($p(0.745)<0.05$) and treatment ($p(0.741)<0.05$) respiratory disease with pulmonary function (Table 3).

The lung function test was a combination of studies carried out in clinical practice to determine lung capacity and possible decrease in mechanical function of the lungs, respiratory muscles, and chest wall. In clinical guidelines, pulmonary function interpretation has historically been based on the most important anthropometric factors, including weight, height, sex, and age [27]. In Lorensia et al. [28], in 148 respondents pedicab drivers as high risk chronic pulmonary disease (135 with respiratory disorder and 13 without respiratory disorder). The results of this study found the most common risk factor was air pollution exposure. In 145 respondents (97.97%), the average lung function was 52.56%. The most frequent early symptom was dyspnea with percentage of 7.43% and average lung function value of 26.20%. Air pollution was the most common risk factor and dyspnea was the most common early symptom in pedicab drivers with respiratory disorders in Surabaya.

4 Conclusions

The knowledge profile about the risk factors for respiratory disorders shows that there is no difference in the level of knowledge of the respondents with or without respiratory disorders. The knowledge profile about respiratory symptoms shows that there is no

difference in the level of knowledge of respondents with or without respiratory disorders. And the knowledge profile about the treatment and therapy of respiratory disorders shows that there is no difference in the level of knowledge of respondents with or without respiratory disorders.

5 Acknowledgments

Funding sources: this research was funded by the Institute of Research and Community Service of the Universitas Surabaya.

6 Author Contribution

All authors contributed starting from data collection, data analysis, and writing papers.

7 Ethical Clearance

No. 002-OL/KE/IV/2019 from Health Research Ethics Committee in Universitas Surabaya.

8 Conflicts of Interest

The authors declare no conflict of interest.

9 References

- [1] Badan Pusat Statistik, 2018, Statistik Daerah Kota Surabaya. Surabaya: Badan Pusat Statistik Kota Surabaya.
- [2] Moynihan, M.C., Allwood, J.M., 2014. Utilization of structural steel in buildings. *Proc Math Phys Eng Sci*, 470(2168), pp.20140170. doi:10.1098/rspa.2014.0170
- [3] Bauleo, L., Bucci, S., Antonucci, C., et al., 2019. Long-term exposure to air pollutants from multiple sources and mortality in an industrial area: a cohort study. *Occup Environ Med*, 76(1), pp.48-57. doi:10.1136/oemed-2018-105059
- [4] Manisalidis, I., Stavropoulou, E., Stavropoulos, A., Bezirtzoglou, E., 2020. Environmental and Health Impacts of Air Pollution: A Review. *Front Public Health*, 8, pp.14. doi:10.3389/fpubh.2020.00014
- [5] Wiston, M., 2017. Status of Air Pollution in Botswana and Significance to Air Quality and Human Health. *J Health Pollut*, 7(15), pp.8-17. doi:10.5696/2156-9614-7.15.8
- [6] Jiang, X.Q., Mei, X.D., Feng, D., 2016. Air pollution and chronic airway diseases: what should

- people know and do?. *J Thorac Dis*, 8(1), pp.E31-E40. doi:10.3978/j.issn.2072-1439.2015.11.50
- [7] Kurt, O.K., Zhang, J., Pinkerton, K.E., 2016. Pulmonary health effects of air pollution. *Curr Opin Pulm Med*, 22(2), pp.138-143. doi:10.1097/MCP.0000000000000248
- [8] Ghorani-Azam, A., Riahi-Zanjani, B., Balali-Mood, M., 2016. Effects of air pollution on human health and practical measures for prevention in Iran. *J Res Med Sci*, 21, pp.65. doi:10.4103/1735-1995.189646
- [9] Kim, D., Chen, Z., Zhou, L.F., Huang, S.X., 2018. Air pollutants and early origins of respiratory diseases. *Chronic Dis Transl Med*, 4(2), pp.75-94. doi:10.1016/j.cdtm.2018.03.003
- [10] Alvarado, A., Arce, I., 2016. Metabolic Functions of the Lung, Disorders and Associated Pathologies. *J Clin Med Res*, 8(10), pp.689-700. doi:10.14740/jocmr2668w
- [11] GOLD (Global Initiative For Chronic Lung Disease), 2018. Global Strategy for the Diagnosis, Management, and Prevention of Chronic Obstructive Pulmonary Disease (2018 Report) Global Initiative for Chronic Obstructive Disease, Inc, USA.
- [12] Dinas Kesehatan Surabaya. 2016. Profil Kesehatan Kota Surabaya. Surabaya: Dinas Kesehatan Surabaya.
- [13] Supriadi, L.S.R., Sui-Pheng, L., 2017. The Indonesian Construction Industry. *Business Continuity Management in Construction*, pp.117-154. doi:10.1007/978-981-10-5487-7_6
- [14] Aryal-Bhandari, A., Gautam, R., Bhandari, S., 2015. Knowledge and Practice on Prevention of Respiratory Health Problems among Traffic Police in Kathmandu, Nepal. *Int Sch Res Notices*, 2015, pp.716257. doi:10.1155/2015/716257
- [15] Missriya, S., Subashini, S., 2017. Assess The Prevalence of Respiratory Problem and Knowledge on Health Effects of People Residing Near Quarry. *Int J Pharm Pharm Sci*, 9(2), pp.250-253.
- [16] Grady, P.A., Gough, L.L., 2014. Self-management: a comprehensive approach to management of chronic conditions. *Am J Public Health*, 104(8), pp.e25-e31. doi:10.2105/AJPH.2014.302041
- [17] Ebrahimi-Belil, F., Alhani, F., Ebadi, A., Kazemnejad, A., 2018. Self-Efficacy of People with Chronic Conditions: A Qualitative Directed Content Analysis. *J Clin Med*, 7(11), pp.411. doi:10.3390/jcm7110411
- [18] Settineri, S., Frisone, F., Merlo, E.M., Geraci, D., Martino, G., 2019. Compliance, adherence, concordance, empowerment, and self-management: five words to manifest a relational maladjustment in diabetes. *J Multidiscip Healthc*, 12, pp.299-314. doi:10.2147/JMDH.S193752
- [19] Khan, F.U., Waqas, N., Ihsan, A.U., Khongorzul, P., Wazir, J., Gang, W., Mengqi, Y., Xiaoqian, L., Han, L., Zhou-Xiaohui, Z., 2019. Analysis of the Qualities Matching New Classification of Clinical Pharmacist. *Indian J Pharm Sci*, 81(1), pp.2-10.
- [20] Ranu, H., Wilde, M., Madden, B., 2011. Pulmonary function tests. *Ulster Med J*, 80(2), pp.84-90.
- [21] Brandsma, C.A., Vries, M.D., Costa, R., Woldhuis, R.R., Konigshoff, M., Timens, W., 2017. Lung ageing and COPD: is there a role for ageing in abnormal tissue repair?. *European Respiratory Review*, 26, pp.170073.
- [22] Nuttall, F.Q., 2015. Body Mass Index: Obesity, BMI, and Health: A Critical Review. *Nutr Today*, 50(3), pp.117-128. doi:10.1097/NT.0000000000000092
- [23] Rawal, G., Yadav, S., 2015. Nutrition in chronic obstructive pulmonary disease: A review. *J Transl Int Med*, 3(4), pp.151-154. doi:10.1515/jtim-2015-0021
- [24] Collins, P.F., Yang, I.A., Chang, Y.C., Vaughan, A., 2019. Nutritional support in chronic obstructive pulmonary disease (COPD): an evidence update. *J Thorac Dis*, 11(Suppl 17), pp.S2230-S2237. doi:10.21037/jtd.2019.10.41
- [25] Lorensia, A., Suryadinata, R.V., Sidabutar., 2021. Effect Analysis of Protein Intake of Pedicab Driver in Surabaya. *Journal of Tropical Pharmacy and Chemistry*, 5(3), pp.188-193.
- [26] Walson, J.L., Berkley, J.A., 2018. The impact of malnutrition on childhood infections. *Curr Opin Infect Dis*, 31(3), pp.231-236. doi:10.1097/QCO.0000000000000448
- [27] Barroso, A.T., Martin, E.M., Romero, L.M.R., Ruiz, F.O., 2018. Factors Affecting Lung Function: A Review of the Literature. *Arch Bronconeumol*, 54(6), pp.327-332.
- [28] Lorensia, A., Suryadinata, R.V., Diputra. I.N.Y., 2019. Risk Factors and Early Symptoms Related to Respiratory Disease in Pedicab Drivers in Surabaya. *Jurnal Kesehatan Masyarakat*, 15(2), pp.224-235.

Relationship of Knowledge on Respiratory Disorders with Lung Function in Masons in East Surabaya

ORIGINALITY REPORT

12%

SIMILARITY INDEX

14%

INTERNET SOURCES

7%

PUBLICATIONS

2%

STUDENT PAPERS

PRIMARY SOURCES

1	err.ersjournals.com Internet Source	3%
2	docobook.com Internet Source	2%
3	www.scribd.com Internet Source	2%
4	garuda.kemdikbud.go.id Internet Source	2%
5	www.jpmmh.org Internet Source	2%
6	Submitted to Asian Institute of Technology Student Paper	2%

Exclude quotes On

Exclude matches < 51 words

Exclude bibliography On

Relationship of Knowledge on Respiratory Disorders with Lung Function in Masons in East Surabaya

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8

PAGE 9
