

Study of Jamu as Indonesian Herbal Medicine for Covid-19 Treatment

Marisca Evalina Gondokesumo^{1*}, Krisyanti Budipramana¹, Silmi Qurrotu Aini²

¹ Faculty of Pharmacy, University of Surabaya, East Java

² Biology Programs, Education University of Indonesia

*Corresponding author. Email: marisca@staff.ubaya.ac.id

ABSTRACT

The spread of the Severe Acute Respiratory Syndrome (SARS-CoV-2) virus has caused the COVID-19 pandemic for more than a year. Daily cases in Indonesia keep increasing following the high mortality rate, entitling it as the epicentre country of Asia. The government advises patients with mild symptoms to take self-isolation at home. Due to its rich cultures and beliefs in Indonesia, patients on independent isolation tend to take treatment on their own accord, such as using traditional herbal medicine renowned as jamu. This article is aimed to review the use of various types of concoctions in jamu taken as the prevention and additional therapy for COVID-19 patients. The review may advantage for COVID-19 treatment discovery through indigenous compound and medical care in societies. In writing this article, the author collected information from related articles in online sources such as Google Scholar, Pubmed, Science Direct, as well as circulars from the Indonesian Ministry of Health and treatment guidelines from the Food and Drug Supervisory Agency. Polyherbal formulations that have been widely recommended for COVID-19 treatment are 9 jamu formulas within *Zingiber officinale* for the highest herbal compound used and followed by other 21 herbal plants such as *Curcuma zanthorrhiza*, *Cinnomomun verum*, *Citrus aurantifolia* fruit, *Cymbapogon sitratus*, *Curcuma longa*, *Alpinea galangal*, etc. Five of all founded jamu identified for antiviral agent as well as mostly applied in immune booster traditionally. Moreover, all of jamu formulas has antioxidant activities and antiinflammatory effect that would be promising for COVID-19 treatment, especially for Jamu Pahitan and Wedang Secang could be the candidates for COVID-19 therapy through increasing immunity and reducing risk factors for each symptom. No antagonistic interactions were found in the herbal concoctions, while several synergistic interactions that increase the pharmacological activity. This study would contribute a remarkable literacy since the use of jamu in preventing and treating COVID-19 as a new disease requires careful consideration due to the limited scientific data and clinical evidence to support its use.

Keywords: COVID-19, Herbal, Polyherbal, Active Compound Interaction, Pharmacological Activity

1. INTRODUCTION

Coronavirus Disease (COVID-19) was first discovered as a virus causing pneumonia in Wuhan, China. As a disease occurred from the new variant of the coronaviridae virus family, it is known to have gen's sequence similarities with Severe Acute Respiratory Syndrome (SARS) and Middle East Respiratory Syndrome (MERS), 88% and 50% respectively [1]. Since the infection phase may occur without symptoms, it is hard to detect the virus particle; hence COVID-19 has a faster and broader range of infection and dispersion [2]. On January 30, WHO determined COVID-19 as a Public Health Emergency of International Concern (PHEIC). Afterward, on February 11, 2020, the International Committee on Taxonomy of Viruses officially named this new coronavirus as Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) [3].

In Indonesia, the COVID-19 case was first found on March 2, 2020, in Depok, West Java. Cases of positive infection keep on increasing over time until the virus spread across all provinces in Indonesia. Further, there was a second wave of infection when the Delta variant or B1617.2 as the variant of concern entered Indonesia on May 2021. On August 4, 2021, it was confirmed that the positive cases had reached 3,532,567 people with a daily discovery of 35,867 people and an additional daily death of 1,747 cases [4]. The daily addition of cases and daily death rate are quite high, leading to Indonesia's entitlement as the epicentre of COVID-19 in Asia.

COVID-19 treatment therapy is still under continuous research to find the most effective cure and vaccine. This is a necessary effort in mitigating and controlling the COVID-19. The research usually uses evidence-based approach and multifactorial possibilities, namely through the clinical severity study, transmission and infection rate, as well as the efficacy of medication used in the therapy [5].

So far, some general antiviral medications such as remdesivir, favipiravir, and lopinavir-ritonavir, as well as antimalaria medication such as chloroquine and hydroxychloroquine have been used [6]. However, hydroxychloroquine is the first medication that has evidently been considered to have no clinical benefits in a prospective manner, since it does not significantly reduce the possibility of virus incubation period and death rate [7]. The latest report related to the effectiveness of remdesivir showed that it had reduced the duration of recovery from an average of 15 days to 10 days among the hospitalized in-patient adults.

Besides, remdesivir reduces almost half of the mortality rate among the cured patients. [8] Unfortunately, the supply of this therapeutically medication in the various region keeps on decreasing and experience scarcity.

Other alternatives in curing the COVID-19 patients may be taken through convalescent plasma (CP) therapy. However, this type of medication may only be more effective for COVID-19 patients in an initial phase, where the virus has not yet cause severe damages to the lungs [9]. Various vaccines have been discovered through many types of virus sources, yet the continuing development of the variant of virus has incited debate regarding the efficacy and effectiveness of the respective vaccines, particularly in stopping the transmission or mitigating the virus infection.

In facing the COVID-19, China is the pioneer, where the case was first discovered, and thus, it becomes the first epicentre in the world. It has been simultaneously using medical guidelines merged from traditional Chinese medicine (TCM) and modern medication. The Chinese government stated that 91.50% of the total case occurred, or as much as 74,187 cases of COVID-19, was cured with the mentioned combination, and it produced promising results in all phases of infections. The result obtained refers to the management of the significant symptoms, namely in reducing symptoms, drop in body temperature, average in-patient duration in hospitals, an increase of recovery rate, a decrease of mortality rate in moderate symptoms, and may be used to mitigate the disease. Furthermore, when combined with modern biomedical, herbal cures may help relieve hypoxaemia and chronic obstructive pulmonary disease (COPD) [10].

Coronavirus is known to cause variety of symptoms in each patient with different severity and mortality rate according to the strength of their antibodies and comorbidities. The infection character of this virus makes it difficult for virus and infection control. Medication and control management of COVID-19 need to focus on early diagnosis, isolation, supportive treatment, as well as infection mitigation and control [11]. Hence, aside from medication effort, mitigation and countermeasures of several mild and moderate symptoms through the increase of immune system are also necessary to conduct. WHO has been encouraging innovation worldwide to use traditional medications as the new therapy development to find potential in curing and mitigating of COVID-19. Several compounds discovered as potentially holding up the coronavirus and boosting the immune system

are quercetin, myricetin, psoralidin, caffeic acid, tryptanthrin, lycorine scutellarein, silvestrol, saikosaponin, isobavachalone, and griffithsin [12]. Herbs are contributing in decreasing morbidity and mortality through their immunomodulatory function, it also suppresses the inflammation process; thus, it is capable of preventing worsening symptoms, increasing immunity, and protect body organs. [13]

Indonesia has been using herbal medicines in society for generations. The Herbs and Jamu Research (Ristoja) in 2017 alone has found 6,193 types of jamu formulas for 74 complaints or diseases from 505 out of 1,128 ethnics, in 11 provinces in Indonesia, outside of Java, Sumatera, and Bali. [14]. During this COVID-19 pandemic, the Ministry of Health also encourages people to use herbal medication, mainly through increasing immunity for mitigation purposes. The Foods and Drugs Monitoring Agency has also issued a Herbal Medication Pocket Book During COVID-19 as the guidelines to use herbs, specifically jamu. There are three types of home remedies classified based on the standardization of the materials and empirical tests in Indonesia, namely jamu, Standardized Herbal Medication (SHM), and Phytopharmaceuticals. Home remedies may also be found in the form of concoctions, either in powder or liquid. [15]

As Indonesian herbs, there has been no research concerning the material standardization or clinical tests made against jamu as empirical verification. Jamu has been used much by Indonesian society as cultural heritage due to its efficacy and utilization for generations. [16]. Jamu formula often consist of medicinal plants processed in traditional way or consisted of several kinds of herbs or poly-herbs simultaneously processed to produce several types of active compounds. It is necessary to pay attention to the poly-herbs management to still achieve the therapy efficacy without any interaction of active compounds that may lead to fatal side-effects. [17]

Therefore, this article review is aimed to collect researches related to Indonesian jamu formula used as therapeutic medication and mitigation against the COVID-19. The collection of information regarding jamu formula is also based on the publication of jamu's use during the COVID-19 pandemic among the society. The data obtained shall be used to find each active compounds from either mono-herbs or poly-herbs content within the concoctions, to then compare the effectivity between the two. Besides, the interaction among the active compounds in the poly-herbs shall be delivered as a comprehensive literacy in

the use of jamu for the COVID-19 treatment and the interest of further study.

2. METHODS

This review was performed using the search terms "jamu for COVID-19" that found 312 articles in Google Scholar, 3 articles in Science Direct, and manually searched the references of select articles for additional relevant articles such as phytopharmaceuticals of herbs and antiviral prediction for SARS-CoV-2. Herbs component of jamu founded as COVID-19 treatment in society have compared to FORTI (Formularium of Traditional Indonesian Medicines) from the Ministry of Health as a reference for herbal medicines that are safe for consumption as well as an electronic book of Herbal Use Guidelines issued by BPOM. Moreover, the particular herbal component was checked for herbal-herbal interactions through <https://www.drugs.com/condition/herbal-supplementation.html> page and related articles. Additional articles are obtained by tracking citations from selected publications or directly accessing the journal's website. The literature search included all available and published reports for the period 2011 to 2021. Relevant articles were collected and reviewed.

3. RESULT

Indonesian jamu content of polyherbal that are widely used by society to treat and prevent COVID-19 are presented in Table 1. We found 9 jamu formulas with various herbal components. It is known that there are 22 types of herbal plants used and presented in Diagram 1 with 5 main bioactivities, such as immunomodulatory, antioxidants, anti-inflammatory, antiviral, and antibacterial.

The mostly used plant was *Zingiber officiale* (18%), followed by *Curcuma zanthorrhiza* (9%) and *Cinnomun verum*, *Citrus aurantifolia* fruit, *Cymbapogon sitratus*, *Curcuma longa*, and *Alpinia alanga* that occupied at 6%. In traditional applied 9 kinds of jamu are used for increasing the immune system generally. Other benefits are known to be very diverse, such as relieving rheumatic pain, improving digestion and treating diarrhoea, improving blood circulation, increasing appetite, treating fever and colds, treating sore throat, asthma, or lung disease, and treating malaria.




Diagram 1. Herbal Plant in Jamu Utilizing Founded

4. DISCUSSION

4.1. Pharmacological Potential of Jamu for COVID-19

The thousands of Indonesian ethnic that separated throughout all region has different traditions and

indigenous beliefs, including in the medical tradition. Jamu is a traditional Indonesian medicine that proved an effectivity from generation to generation. As the going spread of COVID-19 today, the Indonesian government has provided to encourage and direct public societies for traditional Indonesian medicine usage through the policy of the Ministry of Health and the Food and Drug Administration in handling COVID-19. However, the clinical research publications of herbs have not been completely found in evaluating an effective treatment of COVID-19 patients, but rather the function of maintaining and enhancing the body's immune system. This activity of the immune system is related with stimulator agents contained in herbs as a stimulator or modulator of various aspects of immunity in the body's system, both adaptive and innate [16]. Therefore, bioactive and pharmacological studies on herbs that are widely used and recommended during this pandemic need to be reviewed to determine the potential for the body's immune system. Furthermore, the active substances that have antiviral activity can be potential candidates for the treatment of COVID-19 involved the in silico analysis to predict objectively.

The results of the literature review found 9 herbal formulas that can be used and are often recommended for the treatment and prevention of COVID-19 in Indonesia, consisting of 22 different types of herbal plants. This result is very small when compared to the result of RISTOJA 2017 from the provinces of Kalimantan, Sulawesi, Nusa Tenggara, and Papua which found as many as 15,671 herbs for health and 1,183 healers / traditional remedies of them. From these ingredients, 11,218 information on types of medicinal plants were obtained; 9,516 were identified, consisting of 1,144 species and 187 genera, while 1,285 plants could not be identified. From all the medicinal plants obtained, various types of ingredients were found consisting of 2-5 types of constituent herbal ingredients [14]. Nevertheless, the composition of the herbs found was by the electronic book Guidelines for the Use of Herbs and Health Supplements in Facing COVID-19 in Indonesia issued by the Food and Drug Supervisory Agency and FORTI (Formulation of Traditional Indonesian Medicines) published in the Decree of the Minister of Health number (HK .01.07/MENKES/187/2017). Both publications are the basis for using herbal medicinal ingredients that have been recognized for their safety and health efficacy by the Indonesian government [16], [43], [44].

Table 1. Herbal formulas that are used for COVID-19 therapy and body immune booster

Herbs Formulation	Bioactive Compound	Pharmacologies Activities	Traditional Used
<p>Wedang Jahe: <i>Cinnamomum verum</i>, <i>Citrus aurantifolia</i> (Christm.) fruit Swingle, and <i>Zingiber officinale</i> var. <i>Rubrum</i></p>	<p>Trans-cinnamaldehyde as immunomodulatory from <i>C. verum</i> 6-gingerol, 6-shogaol, and 8-shogaol as antioxidant from <i>Z. officinale</i></p>	<p><i>C. verum</i>: strong antioxidant (DPPH 71.12 ± 0.77%) [18]. <i>C. aurantifolia</i> fruit inhibit the expression of leukocyte CD18/11a which causes cell adhesion defects and increased leukocyte phagocytosis [19]. <i>Z. officinale</i>, DPPH (IC50 0.64 g/ml), [6]-gingerol (50mg/kg) reduced pain relief in experimental rats by 50% as an effective analgesic and anti-inflammatory effect [20].</p>	<p><i>Wedang Jahe</i> has used traditionally to keep body warm, increase appetite, improve digestion system, and help relieve rheumatic pain [21].</p>
<p>Jamu Temulawak-Meniran: <i>Curcuma xanthorrhiza</i>, <i>Phyllanthus niruri</i>, and <i>Zingiber officinale</i> var. <i>Rubrum</i></p>	<p>Xanthorrhizol as antiinflammatory and antioxidant <i>C. xanthorrhiza</i> <i>P. niruri</i> as immunomodulatory from combination of 12 compounds tannin and flavonoid <i>Z. officinale</i>, 6-gingerol, 6-shogaol, and 8-shogaol as antioxidant</p>	<p><i>C. xanthorrhiza</i> in vitro decreased COX-2, iNOS, TNF-α and IL-6 levels; in vivo neutralized the effect of TPA-induced ODC, COX-2 and iNOS activation in mouse skin, and prevented iKβa degradation; blocked neurogenic and inflammatory pain responses in a formalin-induced pain test in mice [20]; <i>P. niruri</i> as immunomodulatory agent inhibit cellular and humoral immune responses in Balb/C mice; inhibits the release of ROS and NO, inhibits fMLP-induced neutrophil migration; inhibits the release of pro-inflammatory cytokines; and inhibits MyD88-dependent signaling pathways (NF-κB and MAPK)[22]</p>	<p>Jamu Temulawak-Meniran is often used traditionally for digestive disorders, jaundice, vaginal discharge, increasing immune and health [23].</p>
<p>Jamu Beras Kencur: <i>Kaempferia galanga</i> (L.), Saripati <i>Oryza sativa</i> (L.), and <i>Zingiber officinale</i> var. <i>Rubrum</i></p>	<p><i>K. galanga</i> contain ethyl-p-methoxycinnamate and curcuminoids as antiinflammatory and antioxidant Polysaccharides from <i>Oryza sativa</i> as immunomodulatory, γ-oryzanol as antiinflammatory and antioxidant <i>Z. officinale</i>, 6-gingerol, 6-shogaol, and 8-shogaol as antioxidant</p>	<p>Ethyl-p-methoxycinnamate has protective effect since savage ROS and inhibits IL-1, and TNF-alpha[24] -oryzanol as anti-inflammatory prevent IL-6, TNF-α, and NF-κB [23].</p>	<p>Jamu Beras Kencur traditionally applied as a general tonic, relieves fatigue, raises blood circulation, and appetite[21]</p>
<p>Wedang Kunyit: <i>Curcuma longa</i>, <i>Alpinia galanga</i>, <i>Citrus aurantifolia</i></p>	<p>Curcumin as antioxidant, antiviral, and antiinflammatory on <i>C. longa</i> Polysaccharide from <i>A. galanga</i> has activity in immunomodulatory effect [25], phydroxycinnamaldehyde as antiinflammatory.</p>	<p><i>C. longa</i> extract showed an increasing macrophage activity, reducing TNF-, IL-1β, IL-6, and MCP-1 production in microglial cells, and inhibiting the inflammatory response [26]. Curcumin has antiviral activity against different viruses such as influenza virus, respiratory syncytial virus (RSV), Herpes simplex 1 (HSV-1) papillomavirus (HPV), and Human norovirus (HuNoV)[27]. The effect of phydroxycinnamaldehyde has been shown to decrease chondrocyte degeneration in osteoarthritis patients [28].</p>	<p>Wedang Kunyit in traditionally medicine practice has been used for the treatment of strep throat, gout, rheumatism, hepatoprotective, and boosting the</p>

<p>Teh Pegagan: <i>Centella asiatica</i>, <i>Zingiber officinale</i>, <i>Curcuma xanthorrhiza</i></p>	<p>Asiaticoside as immunomodulator in <i>C. asiatica</i> and several polyphenolic compounds with anti-inflammatory and antioxidant activity 6-gingerol, 6-shogaol, and 8-shogaol as antioxidant from <i>Z. officinale</i> Xanthorizol as anti-inflammatory and antibacterial</p>	<p>Leaf extract of <i>C. asiatica</i> increased activity of lymphocyte proliferation in mice on the hepatitis-induced Balb/c pathway [29].</p>	<p>immune system [23]. <i>C. asiatica</i> in Teh Pegagan is traditionally used for the treatment fever, and dementia [29]</p>
<p>Jamu Empon-Empon: <i>Zingiber officinale</i> var <i>Rubrum</i>, <i>Curcuma longa</i>, <i>Curcuma xanthorrhiza</i>, <i>Cymbopogon citratus</i></p>	<p>6-gingerol, 6-shogaol, and 8-shogaol as antioxidant from <i>Z. officinale</i> Curcumin as antioxidant, antiviral, and antiinflammatory pada <i>C. longa</i> <i>Xanthorizol</i> as antiinflammatory and antibacterial Minyak atsiri and polifenol <i>C. citratus</i> memiliki aktivitas antioxidant</p>	<p>Adding the boiled water from <i>C. citratus</i> fruit can significantly increase the TAS value (total antioxidant status) in both serum and liver samples of mice induced by diabetes [30].</p>	<p>Jamu Empon-empun has widely used to treat inflammation, bacterial infections, and enhance the body's immune system [31].</p>
<p>Jamu Kuyit Asam: <i>Tamarindus indica</i>, <i>Curcuma longa</i></p>	<p><i>T. indica</i> possess phenol, glycoside, mallic acid, tartaric acid that haave an antioxidant and antiinflammatory activity Curcumin as antioxidant, antiviral, and antiinflammatory in <i>C. longa</i></p>	<p>The flesh of <i>T. indica</i> fruit has found in reducing NO levels and inhibiting pancreatic damage in streptozotocin-induced rats; the results of oral tests on mice also showed analgesic activity and increased recovery in mice induced [32].</p>	<p>Jamu Kuyit Asam is traditional treatment for getting rid of fever, curing constipation, asthma, diabetes, reducing nausea in pregnancy, slimming, and treatment of lung disease [33].</p>
<p>Jamu Pahitan: Daun <i>Andrographis paniculata</i>, <i>Tinospora cordifolia</i>, <i>Phyllanthus urinaria</i>, <i>Cymbopogon citratus</i>, Batang <i>Alistonia scholaris</i>, <i>Curcuma xanthorrhiza</i>, <i>Zingiber zerumbet</i>, <i>Alpinia galanga</i>, <i>Curcuma aeruginosa</i></p>	<p><i>A. paniculata</i> leaves possessed many bioactive andrographolide as immunostimulator and anti-infektion; neoandrographolide proved anti-inflammatory activity; and 14-deoxyandrographolide as immunomodulator; <i>P. urinaria</i> leaves have 9 kind of flavonoids and 17 kind of tannin that affected in widely bioactive such as antioxidant, anti-inflamasi, and antivirus [34]; Alkaloids and triterpen from <i>A. scholaris</i> leaves as immunomodulatory, antiinflammatory, and antioxidant [35]; <i>Z. zerumbet</i> as antioxidant and anti-inflammatory from zerumbone and α-humulene [36]</p>	<p>The methanolic extract of <i>A. paniculata</i> inhibited 65% NO production by macrophages and significantly inhibited the formation of leg edema in induced rats for continually five days applied (50 mg/day) [26]. Andrographolide inhibits the production of oxygen radicals in neutrophils, inhibits macrophage migration, NF-B activity and TNF- and IL-12 production via the protein kinase C pathway, extracellular signaling-regulated kinase1/2 (ERK1/2) or the PI3K/Akt signaling pathway [37]. <i>P. urinaria</i> leaves have been shown to inhibit dengue virus, enteroviruses, hepatitis B virus (HCV) and human immunodeficiency virus (HIV); Antioxidant testing on <i>A. scholaris</i> showed the results of DPPH (63%), FRAP (74.88%), H₂O₂ (72.28%) and ROS (67.66%)[38]; Zingiber zerumbet extract can inhibit the action of cyclooxygenase,</p>	<p>Jamu Pahitan is traditional herbal medicine widely used for beauty treatment, malaria medicine, fever medicine, epilepsy, and asthma[38]</p>

<p>Wedang Secang: <i>Caesalpinia sappan</i>, <i>Syzygium aromaticum</i>, <i>Cinnamomum verum</i>, <i>Amomum compactum</i>, <i>Zingiber officinale</i> Rosc., <i>Cymbopogon nardus</i> (L.) Rend!</p>	<p><i>C. sappan</i> has brazilin as a strong antioxidant, protosappanin, and chalcone as an anti-inflammatory and immunomodulator. <i>S. aromaticum</i> depends a lot of polyphenols, one of which is eugenol that Has antioxidant, anti-inflammatory, immunomodulatory and antiviral activity. Cineole as antibacterial in <i>A. compactum</i> [39].</p>	<p>lipoxigenase, myeloperoxidase and nitric oxide synthase enzymes that play a role in the inflammatory process [37]. The aqueous extract of <i>C. sappan</i> can extend the allograft survival time in murine skin induced by immune tolerance [40]. <i>S. aromaticum</i> exerts an anti-inflammatory and immunomodulatory activity from eugenol by suppressing the action of lipopolysaccharide (LPS) and the nuclear factor-κB (NF-κB) pathway. High antioxidant activity has been tested by various in vitro methods: DPPH, ROS, FRAP, ABTS, and ORAC. Clove extract has an important role in treating memory deficit due to oxidative stress effect which was demonstrated in scopolamine-induced rats. Eugenol which isolated from extracts of <i>S. aromaticum</i> has been known to be antiviral for strains of herpes virus and Hepatitis-C virus by inhibiting viral DNA polymerase enzymes, as well as herpes simplex virus type 1 (HSV-1) and influenza-A virus when combined with acyclovir. [41].</p>	<p>Wedang Secang traditionally used to warm the body, treat colds, treat flatulence and diarrhea, treat sore throat, treat rheumatism and increase stamina[42].</p>
--	---	---	---

Five of the 9 ingredients obtained, namely Wedang Kunyit, Empon-Empon, Kunyit Asam, Jamu Pahitan, and Wedang Secang have antiviral activity that has been clinically proven against enteroviruses, herpes, HIV, Influenza-A, respiratory syncytial virus (RSV), and Human norovirus (HuNoV) so that it can be a potential treatment agent for COVID-19 [13], [21], [45]. Meanwhile, herbs with immunomodulatory, antioxidant, and anti-inflammatory activities have the potential to prevent COVID-19 and reduce clinical symptoms of COVID-19. Anti-inflammatory agent is an important mediator in the treatment of COVID-19 due to its application in one of the dangerous threats from SARS-CoV-2 infection is hemophagocytic syndrome (HPS) also called hemophagocytic lymphohistiocytosis [6] and also, macrophage activation syndrome (MAS), a subtype of HPS in which it develops like an autoimmune disease.

This syndrome shows hyperinflammatory symptoms that have been found in the lungs based on histopathological findings. This syndrome occurs due to an attack carried out by the body's immunity resulting in an extreme increase in pro-inflammatory cytokines. This syndrome can be observed through IL-1 β , IL-2, IL-6, IL-7, G-CSF1, IP-10, MCP-1 α , MIP-1 α , TNF-, and sIL2R. The binding of SARS-CoV-19 to the Toll-Like Receptor (TLR) will be followed by the release of pro-IL-1b which is cleaved by caspase-1 which will induce the inflammation and IL-1b mediator activation of pneumonia. Clinical observations showed that the health of some patients deteriorated 6 - 8 days after infection. According to the disease tolerance theory, the most severe cases of COVID-19 could be due in part to an uncontrolled immune response to SARS-CoV-2 infection. This reduction in the inflammatory immune response could be considered a potential therapeutic target against severe COVID-19. Consequently, the suggestion of treat severe COVID-19 with immunomodulatory therapies such as those carried out in HPS is likely to be beneficial for overcoming hyperinflammation and ultimately ameliorating severe clinical syndromes [9]. The anti-inflammatory prevent IL-1b and IL-6 as modulating pro-inflammatory cytokines where there is a storm of pro-inflammatory cytokines in the alveoli.

In RISTOJA's discovery, there were 391 herbal formulas for immunomodulatory that were dominated by turmeric (*Curcuma longa*), betel leaves (*Piper betle* L.), ginger (*Zingiber officinale*). Accordance on our founded of the herbals compound consist by the composition of *Zingiber officinale* as well as the most widely used plant species in the treatment and prevention of COVID-19, followed by *Curcuma zanthorrhiza* and *Curcuma longa*. The types of herbs found showed compatibility with herbal plants recommended by BPOM for the treatment of COVID-19, such as ginger, curcuma, turmeric, meniran, and bitter. There are other medicinal plants recommended by BPOM but not found in this herbal

review, namely the fruit and leaves of guava (*Psidium guava* L.). As for the exploration of Indonesian herbal medicines through RISTOJA, the most common ingredients found were herbs for pre/post-partum care which had 321 types of ingredients, followed by open wound herbs (315), malaria ingredients (288), diarrhoea ingredients (255) and hot/hot ingredients. fever (251), ingredients for cancer/tumour (223), tuberculosis (106), and HIV/AIDS (22) [14]. Unfortunately, there is no complete report regarding the composition of the ingredients used. Some herbs in certain ethnic groups must be used with certain rituals or prayers.

Another study conducted by the Ministry of Health to determine the preference for herbal medicine use among Indonesian people in 2010 and 2013 showed that there were 59.12% of the Indonesian population who had consumed herbal medicine (age > 15 years), both men and women, in rural and urban areas with 95.60% of users feeling the benefits. The herbal dosage form that is most preferred by the public is the liquid form, followed by steeping/powder, decoction/chopped, and capsule/pill/tablet form. Basic Health Research Data (Riskesmas) 2013 shows that households that use traditional health services are 30.40%, of which 77.80% choose skills without tools and 49.00% potions [43].

4.2. Herbs – Herbs Interaction in Jamu

Jamu as herbal medicine consists of several medicinal plants within complex bioactive content. The multi-herbs combination would enhance the effectiveness of treatment, consisting of the main component as the primary element in the treatment goal, supporting compounds that help strengthen the effect, then complement or balance. However, not all combinations of medicinal plants can achieve treatment effectiveness some can interact with each other differently, including; (1) a positive synergistic effect indicates that the combined compound has a beneficial interaction, (2) a negative effect indicates that the combined compound eliminate the benefits, and (3) does not synergize indicating that the combined compound does not affect the results of the compound benefit [46].

In addition, certainly pharmacological actions of the active herbal constituents are also needed due to significantly increase as well as be able on more strengthened by other plants instead of its single used [47]. This is possible since the molecules in one herb are not necessarily compounds that are naturally contained but in the form of host-specific metabolites or molecular complexes formed after other herbs co-administration [10].

Table 2. The Interaction into Herbs of Jamu

No	Jamu	Compound Combination	Interaction	Affected Pharmacology
1	Jamu Wedang Jahe	<i>Z. officinale</i> – <i>C. verum</i>	Synergistic	+ decrease blood glucose
2	Jamu Temulawak-meniran	<i>C. zanthorrhiza</i> – <i>P. niruri</i>	Synergistic	+ antioxidant , + antivirus
3	Jamu Beras Kencur	<i>Z. officinale</i> – <i>K. galanga</i>	Synergistic	+ antibacterial
4	Weandg Kunyit	No reported yet		
5	Teh Pegagan	No reported yet		
6	Jamu Empon-empon	<i>Z. officinale</i> – <i>K. galanga</i>	Synergistic	+ decrease blood glucose and colessterol
7		<i>C. longa</i> – <i>Z. officinale</i>	Synergistic	+ antibacterial
8	Jamu Kunyit Asam	<i>T. indica</i> – <i>C longa</i>	Synergistic	+ antioxidant
9	Jamu Pahitan	<i>T. cordifolia</i> - <i>Z. officinale</i>	Synergistic	+ antiinflamantory
10	Wedang Secang	<i>C. sappan</i> – <i>Z. zerumbet</i>	Synergistic	+ cytotoxic
11		<i>Z. officinale</i> – <i>S. aromaticum</i>	Synergistic	+ antivirus
12		<i>S. aromaticum</i> – <i>C. verum</i>	Synergistic	+ antioxidant

Notes: (+) show the increasing of pharmacology activities

Jamu Temulawak-meniran consists of *C. zanthorrhiza* or curcuma, *Phyllanthus niruri* or meniran, and ginger. It was found that there was a synergistic reaction between curcuma and meniran on antioxidant activity. Based on the antioxidant activity of all formulas shown in [48], the highest antioxidant activity (lowest IC50 value) was found in a mixed formula of 50% curcuma extract and 50% meniran extract. The activity of the mixed formula (IC50 9.47) was higher than that of curcuma extract alone (IC50 151.6) and meniran extract alone (IC50 101.91), but not as good as the activity of catechins as a positive control. In addition, the antiviral activity of curcuma and meniran also increased with a concentration of each component of 500 ppm which was indicated by the PCR results on viral cDNA decreasing in Simian Retrovirus Serotype-2 (SRV-2) virus [49].

The components of the jamu *Beras Kencur* include *Kaempferia galanga* (L.), *Oryza sativa* (L.) or rice, and *Zingiber officinale* var. Rubrum or red ginger. The combination of red ginger with *K. galanga* shows a synergistic effect that can increase antibacterial activity against *Staphylococcus aureus* bacteria. The results [49] showed that a single extract of *Z. officinale* had a minimum inhibitory concentration (MIC) of 62.5 g/ml and a minimum bactericidal concentration (MBC) of 125 g/ml, a single extract of kencur had a MIC of 500 g/ml and an MBC of 1000 g. /ml. The results of the combination of *Z. officinale* extract and aromatic ginger extract which increased antibacterial activity, all concentrations of the combined extract showed no growth of *Streptococcus pyogenes* at the

lowest concentration of 62.5 g/ml. These results indicated that the combination of *Z. officinale* and *K. galanga* extracts were more effective than single extracts. However, no reports have been found for a combined effect with *Oryza sativa*.

The following herbal formula is a mixture of *Curcuma longa* or turmeric, *Alpinia galanga* or galangal, and *Citrus aurantifolia* or lime. There were no reports related to their interaction. Using turmeric reported could not be allowed in conjunction with other herbal supplements that caused blood clotting, such as capsicum, cloves, garlic, ginger, ginkgo, or red clover. Moreover, combining with lower blood sugar causes, such as alpha-lipoic acid, chromium, or garlic does not permit. In the Wedang turmeric formula, there are no ingredients that will induce an antagonist reaction as well as it is safe to consume. While had piperine from pepper contain bioavailability of curcumin would increase [50].

Centella tea with a mixture of *Centella asiatica*, *Zingiber officinale* or ginger, and *Curcuma zanthorrhiza* or curcuma has not report synergies or antagonist's interaction publication yet. In order to do so, avoid taking *C. asiatica* with other herbal caused drowsiness, such as 5-HTP (5-hydroxytryptophan), jeringa, or valerian. *C. asiatica* has been reported to interfere with blood glucose levels when given concomitantly with hypoglycemic therapy. In addition, the saponoside fraction of plant extracts containing brahmic acid and the derivatives were found to affect infertility in human and rat sperm [51].

There are no compounds found in Centella tea ingredients that cause these negative interactions so they can be safely consumed.

Jamu empon-empon consist of *Zingiber officinale* var Rubrum or red ginger, *Curcuma longa* or turmeric, *Curcuma zanthorrhiza* or curcuma, *Cymbopogon citratus* or lemongrass. There are studies in ginger consumption published that cannot be combined with turmeric due to significantly lower blood sugar levels agent ($P < 0.05$) and control lipid profiles in diabetic-dyslipidemic rats. However, the other report showed the synergistic therapy of both extracts was most significant ($P < 0.05$) controlling all parameters of diabetic dyslipidemia (78.00 ± 1.06 mg/dL FPG, 62.00 ± 0.58 mg/dL TG, 66.50 ± 0.76 mg/dL cholesterol, HDL 32.00 ± 0.36 mg/dL, LDL 22.43 ± 0.64 mg/dL, and VLDL 12.40 ± 0.12 mg/dL). Additionally, reports of relief from liver cirrhosis after carbon tetrachloride-induced poisoning in rats, as ginger alone or either in combination with curcumin at 100 mg/kg ameliorated liver injury in animals. In addition, another report at 200,400 mg/kg strengthened the activity of antioxidant enzymes (superoxide dismutase, catalase, glutathione peroxidase) while decreasing the activity of liver function enzymes (alanine transaminase, aspartate aminotransferase) in acetaminophen-induced liver [52]. Synergistic activity was also found in galangal and lemongrass through their increasing antibacterial activity compared to a single application. The profile of the combination of galangal oil and lemongrass oil (volume ratio 3:7, 1:1, and 7:3) was tested against four pathogenic microorganisms (*Staphylococcus aureus* 0.5% and 4%, v/v, against *Pseudomonas aeruginosa* 40% and > 40% v/v, against *Streptococcus bovis* 0.25% and 0.5%, v/v, and against *Candida albicans* 0.25% and 0.5%, v/v). Synergistic activity was best recorded for only one ratio (volume ratio 3:7). This investigation provides evidence of their interaction as an herbal combination within the minimum effective dose [53].

Jamu tamarind-turmeric is reported to have a very strong antioxidant synergism greater than the synthetic antioxidant BHT which resembles vitamin E which is widely used by the food industry as a preservative [54]. Jamu Pahitan that have the most herbal medicinal compositions are *Andrographis paniculata* or sambiloto leaves, *Tinospora cordifolia* or bratawali leaves, *Phyllanthus urinaria* or meniran leaves, *Cymbopogon citratus* or lemongrass, *Alstonia scholaris* or pulai stems, *Curcuma xanthorrhiza*, *Zingiber zerumbet* or lempuyang, *Alpinia aeruginosa* or temu hitam. Single or synergistic formulations of *Tinospora cordifolia* with *Zingiber officinale* have been traditionally used in rheumatoid arthritis the

treatment. *Tinospora cordifolia* has been reported to influence the proliferation, differentiation, and mineralization of bone-like as the matrix by in vitro osteoblast model system that is referred a potential anti-osteoporosis agent. The alcoholic extract of *Tinospora cordifolia* has been shown to stimulate osteoblast growth, promote cell differentiation into osteoblastic lineages and also increase mineralization of bone-like matrix [55]. The combination between *Tinospora cordifolia* and *Andrographis paniculata* considered increasing its hepatoprotective activity. It was observed that AHPL/AYTAB/0613 significantly reduced levels of serum glutamic-oxaloacetic transaminase, serum glutamic pyruvic transaminase, alkaline phosphatase, total bilirubin, and also significantly increase the total protein level which has been tested in paracetamol-induced rats [50].

Wedang Secang consists of *Caesalpinia sappan* or sappan, *Syzygium aromaticum* or clove, *Cinnamomum verum*, *Amomum compactum* or cardamom, *Zingiber officinale*, *Cymbopogon nardus* (L.) Rendl. Secang extract and lempuyang rhizome showed a cytotoxic effect on MCF-7 cells with IC₅₀ values of 30 and 155 g/mL, respectively. 0.57-0.85. The combination of 15 g/mL cup with 8 and 24 g/mL lempuyang showed cell cycle inhibition in the G₂/M phase. The combination of the two also enhanced the occurrence of apoptosis induction compared to the control and single treatment. The combination of ethanolic extract of secang wood and lempuyang rhizome produces a synergistic cytotoxic effect. Combined synergism occurs through inhibition of the cell cycle in the G₂/M phase and induction of apoptosis [56]. Ginger and clove extracts have been tested for synergistic interactions as antivirals. Indeed, nevermore do detect an antiviral effect when host cells pretreated with the extracts. Clove and ginger extracts significantly decreased feline calicivirus (FCV) infectivity in virus pretreatment and in coinfection and post-infection treatment but not in cell pretreatment. Eugenol may not be the only antiviral compound in clove extract and extracts in combination with ginger may induce viral inactivation than eugenol alone [56]. The combination of clove and cinnamon extracts also showed a synergistic effect in increasing the cytotoxicity of Hep G2 cancer cells and increasing DPPH radical scavenging [56]. All the herbal-herbal interactions found has summarized in Table 2.

4.3. *Jamu* Application for COVID-19 Treatment

Considering aspects in the using herbal medicine for the treatment of COVID-19 published in [21]. First, the product must be sure of its safety. Although for a long time does jamu used no clinical evidence, especially for COVID-19 treatment. The use of jamu for therapeutic purposes in COVID-19 is focused on symptom reduction as they generally show the best benefit in ongoing chronic post-infection symptoms rather than the acute stage [57]. Second, the evidence of the herbal safety and efficacy should obtain based on clinical trials. In traditional medicine in Indonesia, herbal medicines that have been clinically tested are called Fitofarmaka. The herbs are herbs that have not been standardized or clinically tested. Indonesia through the Food and Drug Supervisory Agency has issued a guidebook for the Original Indonesian Modern Medicine Informatorium (OMAI) during the COVID-19 Pandemic, in which there are 15 Leveraged Herbal Medicines for 5 groups to relieve symptoms of COVID-19 and increase immunity, as well as 10 Phytopharmaceutical to maintain immunity [15]. Jamu as a traditional medicine used for generation to generation can be analysed for its feasibility through efficacy data obtained from in-vitro and in-vivo studies as a rationale for use in the prevention and reduction of COVID-19 symptoms.

Third, the molecular mechanisms that underlie the pharmacological effects of herbal medicine in the treatment of COVID-19 remain unclear. So far, the available data were obtained from in-silico analysis. In addition, hesperidin, a flavonoid found in *Citrus* sp. The fruit peel, which is popularly championed as natural prevention of COVID-19 in Indonesia, is reported to have strong binding to human angiotensin-converting enzyme-2 (hACE-2) and RNA-dependent RNA polymerase (RdRp) as potential targets for SARS-CoV-2 infection [58]. The potential danger of delay in treatment due to consumption of herbal medicine must be known because herbal medicines are sold as over-the-counter products and can be accessed without a doctor's prescription, their use as COVID-19 self-medication by patients who experience symptoms is very possible [59]. Jamu and other herbal medicines generally should not be used in emergency conditions, including in the acute phase of viral infection. Delays in getting proper treatment for COVID-19 cases lengthen the recovery process. In fact, in some cases, this delay would fatal if the virus continues to replicate rapidly in the patient's body.

There has been no publication of detailed reports regarding the clinical use of jamu for COVID-19

treatment patients in Indonesia yet, either alone or in combination with modern medicine. In contrast to treatment in China through the guidelines, Traditional Chinese Medicine (TCM) has been used in the treatment of COVID-19 patients. There are 23 provinces in China that have adopted guidelines for the use of TCM and issued programs on COVID-19 prevention, while 26 provinces have officially established integrative TCM that combines TCM with clinical treatment for COVID-19 patients [21]. More than 85% of the total confirmed COVID-19 patients are reported to have been treated with TCM. Treatment with integrative medicine was given to all COVID-19 cases in Shanghai, while the first patient who recovered in Beijing was also treated with this combination drug [15]. In the Chinese Guidelines, TCM is used for the treatment of mild, moderate, severe, and convalescent stages of COVID-19. As of January 2020, the total confirmed cases treated by TCM have reached 60,107. In 102 cases of mild symptoms that were treated with TCM, the time of onset of clinical symptoms was 2 days faster, body temperature recovery time was 1.7 days shorter, the average length of hospitalization was 2.2 days faster, and the rate of CT images increased by 22%. In addition, in the treatment of acute patients, the mean length of hospitalization and time to PCR to negative has been 2 days shorter [60].

The research results and predictions of the interaction of active compounds from herbal plants in the composition of herbal ingredients with the COVID-19 virus from the types of herbs found from information on herbal formulas for COVID-19 are presented in Table 3. Based on the in silico test, all herbs contain herbal plants that have the potential to be a candidate for COVID-19 drugs because they can bind to the target of drug action on SAR-CoV-2 (Mpro, Spike Protein, Spike glycoprotein, RdRp, and Nonstructural Polyprotein) or the target site of viral infection to humans (ACE2). The herbal medicine with the highest number of herbal plants found to be able to bind to the target of the SAR-CoV-2 drug is Jamu Pahitan, which has 6 types of herbal plants that have a high affinity for the target. Then, herbal medicine wedang secang has 3 types of plants that are capable of affinity. Antiviral activity in Jamu Wedang Secang is likely to work more effectively it is due to the synergistic effect between ginger and clove which can increase antiviral activity. However, there is no clinical evidence regarding the use of herbal medicine for COVID-19 patients.

Table 3. Prediction of Phytochemical Activity of Herbal Plants from Jamu

Herbal Plan	Bioactive Compound	Prediction of Bioactive – SAR-CoV-2	Ref
<i>Cinnamomum verum</i>	Bergamol	Docking was carried out on protein targets including the major protease SARS-CoV-2 (SARS-CoV-2 M ^{pro}), SARS-CoV-2 endoribonuclease (SARS-CoV-2 Nsp15/NendoU), SARS-CoV-2 ADP-ribose-1" phosphatase (SARS-CoV-2 ADRP), SARS-CoV-2 RNA-dependent RNA polymerase (SARS-CoV-2 RdRp), SARS-CoV-2 spike protein binding domain (SARS-CoV-2 rS), and human angiotensin converting enzyme (hACE ₂) to bergamol essential oil. The docking results showed the affinity of bergamol to each target protein in sequence: 98.0kJ/mol, 91.2kJ/mol, 105.9kJ/mol, 71.9kJ/mol, -63.6kJ/mol, and 83.6kJ/mol respectively. However, this docking score is considered too high or too low affinity when compared to the comparison molecule so it is not feasible. The presence of binding to the target protein indicates that cinnamon can still prevent COVID-19, not as a candidate that can be used as a drug.	[61]
<i>Citrus aurantifolia</i>	Naringin	Naringin pretreatment significantly inhibited the expression of HMGB1 protein and LPS-induced proinflammatory cytokines (COX-2, iNOS, IL-1 β and IL-6) in in vitro expression assays. HMGB1 is a ubiquitous DNA-binding core protein and can be actively released by immune cells, such as macrophages and monocytes, following inflammatory stimuli. HMGB1 also acts as a pro-inflammatory cytokine and regulates cytokine storm, increasing cytokines such as TNF-, IL-6, IL-1 β , and IL-8.	[62]
<i>Zingiber officinale</i>	6 – Gingerol	In the molecular docking test between gingerol and COVID-19 virus RNA binding protein (6W4B), N-Terminal RNA Binding Protein (6VSB), spike glycoprotein (6M3M) showed binding affinities of -11.4082 KJ/mol, -12.9523 KJ/mol and -12.8835 KJ/mol. Molecular interactions between the spike glycoprotein COVID-19 virus and Gingerol make the hydrogen bond interactions with Val42, Pro58, Ser60, Thr68 change form an unbond interaction with Arg40, Phe41, Val42, Phe57, Pro58, Lys59, Ser60, Ile66, Thr68, Ile92 residues from spike COVID-19 glycol protein.	[63]
<i>Curcuma xanthorrhiza</i>	Xanthorrhizol	Xanthorrhizol has anti-inflammatory activity by inhibiting inflammatory cytokines in adipose tissue and expression of tumor necrosis factor (TNF- α). This studies have shown that xanthorrhizol prevents changes in immune cells in adipose tissue so that it can downregulate inflammatory cytokine genes. Xanthorrhizol also has the activity of reducing the expression of the interleukin gene (IL-1 β) in muscle. Another study showed that xanthorrhizol can reduce serum IL-6 levels and increase serum transformation growth factor (TGF- β) in SLE patients with hypovitamin D.	[63]
<i>Phyllanthus niruri</i>	Phyllanthin	The docking results for 6W41 and 5R7Y proteins showed a score of -84.20kCal/mol and -90.65kCal/mol which indicated the ability of the ligand inhibitor. Binding to 6W41 inhibited viral infiltration, while binding to 5R7Y was predicted to inhibit viral translation and replication.	[64]
<i>Kaempferia galangal</i> and <i>Alpinia galanga</i>	Galangin	Molecular docking was performed on the selected protein targets, namely RBD-S (PDB ID:6LXT), PD-ACE2 (PDB ID: 6VW1), and the SARS-CoV-2 protease (PDB ID: 6LU7). Galangin, phenylpropane from galangal was proven to bind three receptors with lower energy than the reference compound used: -12.96Kcal/mol, -7.89Kcal/mol, and -7.60Kcal/mol, respectively. These findings indicate that the two compounds have better binding interactions and can inhibit initial viral infection to host cells.	[65]
<i>Curcuma longa</i>	Curcumin	The anti-inflammatory mechanism of curcumin is by inhibiting the production and release of proinflammatory cytokines, such as IL-1, IL-6, IL-8, TNF- α . Curcumin also showed good binding affinity for nucleocapsids ($\Delta G = -8.75$ kcal/mol, $K_i = 0.39\mu M$), nsp10 ($\Delta G = -7.85$ kcal/mol, $K_i = 1.77\mu M$) which was comparable to ivermectin with nucleocapsids. ($\Delta G = -7.11$ kcal/mol, $K_i = 6.17$ M) and nsp10 ($\Delta G = -9.82$ kcal/mol, $K_i = 63.20$ nm), and remdesivir with nucleocapsid ($\Delta G = -6.30$ kcal/mol, $K_i = 23.94$) and nsp10 ($\Delta G = -6.54$ kcal/mol, $K_i = 16.02$ M)	[66], [67]
<i>Centella asiatica</i>	Asiatic acid	Docking was carried out on the main protease SARS-CoV-2 (6LU7) with <i>Centella asiatica</i> . The results showed that the asiatic acid derivative AA9 had the best affinity for inhibiting the main protease SARS-CoV-2 with a binding free energy (ΔG) value	[68]

		of -9.90 kcal/mol, compared to favipiravir which had a G value of -4.58 kcal/mol. AA9 also interacts with the main protease of SARS-CoV-2 via hydrogen bonding with Gly143. This study demonstrated that asiatic acid and its derivatives have a higher binding affinity for the main protease SARS-CoV-2 compared to favipiravir.	
<i>Cymbopogon citratus</i>	Citronella and patchouli alcohol	Inhibition of citronella essential oil from <i>C. citratus</i> against ACE2 showed an affinity value of -4.8kCal/mol and Patchouli alcohol -6.9kCal/mol. Patchouli alcohol scores were higher than the control so that <i>C. citratus</i> essential oil had potential antiviral activity through its active compound patchouli alcohol. Predictably, patchouli alcohol can inhibit Asn 667 and Ser 701 APN by hydrogen bonding.	[69]
<i>Tamarindus indica</i>	Tripsin (Tamarin Tripsin Inhibitor/TTI)	Molecular docking studies have been carried out to model the binding interactions of various pro-inhibitors of 3CL and other proteases, such as TMPRSS2 (37,82-84). The pTTI-derived peptide also proved to be a strong candidate to block this protease because TTI is known to inhibit the serine protease co-operating with SARS-CoV-2 infection and characterize neutrophil activity in viral-induced lung injury. In addition, trypsin acid inhibitors (TTIs) exhibit several beneficial effects on the reduction of inflammatory markers (tumor necrosis factor [TNF- α], leptin) and biochemical parameters (fasting glycemia, triglycerides, and very low-density lipoprotein [VLDL]), in addition to improve pancreatic functional and mucosal integrity in obesity models. Thus, TTI can contribute to combating two severe overlapping problems with high costs and complex social implications, obesity and COVID-19.	
<i>Caesalpinia sappan</i>	Brazilein	Molecular docking was performed on the selected protein targets, namely RBD-S (PDB ID:6LXT), PD-ACE2 (PDB ID: 6VW1), and the SARS-CoV-2 protease (PDB ID: 6LU7). Brasilien was shown to be able to bind to three receptors with lower energy than the reference compound used, namely -10.52 Kcal/mol, -7.56 Kcal/mol, and -7.43Kcal/mol, respectively.	[65]
<i>Syzygium aromaticum</i>	Eugenol	The results of the molecular docking test showed that the compound eugenol was very potential as an inhibitor of Mpro COVID-19 with the Angiotensin Converting Enzyme 2 (ACE2) receptor. Eugenol is also reported to have a good affinity of -6.3 kcal/mol to inhibit the COVID-19 protease.	[70]
<i>Amomun compactum</i>		<i>No publication yet</i>	-
<i>Andragraphis paniculata</i>	Andrographolide	Docking analysis of the compound with the SARS-CoV-2 protease yielded a negative value for the free energy of the binding site -3.094357 KJ/mol, indicating a high affinity for the binding pocket. All binding conformations of the compound in the active binding pocket involve H-bond and salt bridge interactions. These compounds indeed bind proteases with 4 hydrogen bonds with 3 residues, namely Gly143, Cys145 and Glu166. Andrographolide successfully docked at the SARS-CoV-2 Mpro binding site. The computational approach also predicts this molecule to have good solubility, pharmacodynamic properties and target accuracy. This molecule also complies with Lipinski's rules, which makes it a promising compound for further biochemical and cell-based assays to explore its potential for use against COVID-19.	[71]
<i>Tinospora cordifolia</i>	Tinosporaside	Docking between viral protein and Tinosporaside ligand resulted in affinity scores for Spike protein (-5.7kCal/mol), Main Protease (-7.9kCal/mol), and RNA dependent RNA polymerase virus (-6.6kCal/mol). In the prediction tool, Swiss-ADME, a cheminformatics platform was used to predict the pharmacokinetics and potential drug contingencies from phytoconstituents indicating that there would be no adverse pharmacokinetic-pharmacodynamic herb-drug interactions with concomitant drug therapies.	[72]
<i>Phyllanthus urinaria</i>	Quercetin and quercitrin	In the molecular docking test, the compounds quercetin and quercitrin in meniran were able to bind to Mpro from SARS-CoV-2 (lowest binding energy for quercitrin -10.36 kcal/mol, -8.47 kcal/mol for quercetin). Mpro is an enzyme that plays an important role in viral replication, binding to this protein will make bioactive compounds an inhibitor and prevent viral replication.	[73]

5. SUGGESTIONS FOR FURTHER DEVELOPMENT

In future research, it is hoped that it will be able to analyze herbal-drug interactions so that the potential and threats to the use of herbs together with modern medicines can be identified. Multiherbal interactions also need to be analyzed simultaneously for all components of herbal medicine to determine the overall interaction. Prediction of the interaction of all polyherbals from herbal medicine to the target of COVID-19 virus treatment also needs to be done as a holistic analysis.

6. CONCLUSION

There are 9 types of herbs from 22 types of herbal plants that have been widely used by the community in COVID-19 therapy. These herbal ingredients are widely used to increase endurance and relieve symptoms caused by COVID-19. There were no negative or antagonistic interactions between the herbal ingredients in each herb, so it was safe to consume without losing the efficacy of each plant, but instead was able to increase the pharmacological activity of several compounds such as anti-inflammatory, antibacterial, antiviral, antioxidant, and hepatoprotective activities. The results of the *in silico* test on each component of the herbal medicine show that all types of herbal medicine contain herbal plants that have the potential to be COVID-19 drugs. Most of the inhibitory activity was found in Jamu Pahitan and Wedang Secang. In addition to being a candidate for COVID-19 treatment, all herbs can be used to prevent COVID-19 233w by increasing immunity and reducing risk factors for each symptom. However, no field evidence has been found regarding the use of herbal medicine as a single therapy or in combination with modern medicine for COVID-19 patients in Indonesia.

AUTHORS' CONTRIBUTIONS

All of authors have made substantial and intellectual contribution to the work article, and given approval for the final version of manuscript.

REFERENCES

[1] R. Lu et al., "Genomic characterisation and epidemiology of 2019 novel coronavirus: implications for virus origins and receptor binding," *Lancet*, vol. 395, no. 10224, pp. 565–

574, 2020, doi: 10.1016/S0140-6736(20)30251-8.

- [2] S. Lee et al., "Clinical Course and Molecular Viral Shedding among Asymptomatic and Symptomatic Patients with SARS-CoV-2 Infection in a Community Treatment Center in the Republic of Korea," *JAMA Intern. Med.*, vol. 180, no. 11, pp. 1447–1452, 2020, doi: 10.1001/jamainternmed.2020.3862.
- [3] "Coronavirus Disease (COVID-19) Situation Reports." https://www.who.int/emergencies/diseases/novel-coronavirus-2019/situation-reports/?gclid=CjwKCAjwpMOIBhBAEiwAy5M6YN03aY1G0pleRnFR2eBtjLiafUzdcGpJyYwyNZ5CQLcz5Z2GWbHPYRoCGR8QAvD_BwE (accessed Aug. 10, 2021).
- [4] "Peta Sebaran | Covid19.go.id." <https://covid19.go.id/peta-sebaran> (accessed Aug. 08, 2021).
- [5] M. Nicola, N. O. Neill, C. Sohrabi, M. Khan, M. Agha, and R. Agha, "Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Manfromn on the novel coronavirus COVID- 19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information," *Int. J. Surg.*, vol. 77, no. January, pp. 206–216, 2020.
- [6] D. W. Kurniawan and A. Ikhsanudin, "Potential of Jamu in Nanotechnology Perspective as an Alternative Treatment for Covid-19," *Pharm. Sci. Res.*, vol. 7, no. 3, pp. 123–131, 2020, doi: 10.7454/psr.v7i3.1082.
- [7] C. P. Skipper et al., "Hydroxychloroquine in Nonhospitalized Adults With Early COVID-19: A Randomized Trial," *Ann. Intern. Med.*, vol. 173, no. 8, pp. 623–631, 2020, doi: 10.7326/M20-4207.
- [8] J. H. Beigel et al., "Remdesivir for the Treatment of Covid-19 — Final Report," *N. Engl. J. Med.*, vol. 383, no. 19, pp. 1813–1826, 2020, doi: 10.1056/nejmoa2007764.
- [9] J. Alijotas-reig, E. Esteve-valverde, and C. Belizna, "Immunomodulatory therapy for the management of severe COVID-19. Beyond the anti-viral therapy: A comprehensive review Jaume," no. January, 2020.

- [10] L. Luo et al., "Analysis on herbal medicines utilized for treatment of COVID-19," *Acta Pharm. Sin. B*, vol. 10, no. 7, pp. 1192–1204, 2020, doi: 10.1016/j.apsb.2020.05.007.
- [11] J. Liu and S. Liu, "The management of coronavirus disease 2019 (COVID-19)," *J. Med. Virol.*, vol. 92, no. 9, pp. 1484–1490, 2020, doi: 10.1002/jmv.25965.
- [12] S. Susilawati and H. Hikmatulloh, "Bisnis Ukm Jamu Raden Sri Rastra Di Masa Pandemi Covid-19," *Swabumi*, vol. 9, no. 1, pp. 57–63, 2021, doi: 10.31294/swabumi.v9i1.10133.
- [13] Q. Qiu et al., "Potential Therapeutic Effect of Traditional Chinese Medicine on Coronavirus Disease 2019: A Review," *Front. Pharmacol.*, vol. 11, no. November, pp. 1–16, 2020, doi: 10.3389/fphar.2020.570893.
- [14] K. Kesehatan, "Eksplorasi Pengetahuan Lokal Etnomedisin and Tumbuhan Obat Berbasis Komunitas di Indonesia," 2017.
- [15] BAAND POM, *Informatorium Obat Modern Asli Indonesia (OMAI) di Masa Pandemi COVID-19*. 2020.
- [16] BAAND POM, *Pedoman Penggunaan Herbal and Suplemen Kesehatan dalam Menghadapi COVID-19 di Indonesia*. 2020.
- [17] M. Vijay, K. R. Reddy, M. R. K. Reddy, K. Gnananath, and B. S. Babu, "Standardization In Vitro Antioxidant Activity And Brine Shrimp Lethality Of A Polyherbal Formulation," *Int. J. Ayurveda Pharma Res.*, vol. 4, no. 4, pp. 18–21, 2016.
- [18] B. A. Behbahani, F. Falah, F. L. Arab, M. Vasiee, and F. T. Yazdi, "Antiproliferative Activities of Cinnamomum zeylanicum Bark Essential Oil," vol. 2020, 2020.
- [19] O. S. Enejoh, I. O. Ogunyemi, and M. S. Bala, "Ethnomedical Importance of Citrus Aurantifolia (Christm) Swingle," vol. 4, no. 8, pp. 1–6, 2015.
- [20] A. D. Rini Daud Supu and Jutti Levita, "Red Ginger (Zingiber Officinale Var . Rubrum): Its Chemical Constituents , Pharmacological Activities And," *Fitofarmaka J. Ilm. Farm.*, vol. 8, no. 1, pp. 25–31, 2018, doi: 10.33751/jf.v8i1.11768 Fitofarmaka.
- [21] D. Hartanti, B. A. Dhiani, S. L. Charisma, and R. Wahyuningrum, "The Potential Roles of Jamu for COVID-19: A Learn from the Traditional Chinese Medicine," *Pharm. Sci. Res.*, vol. 7, no. 4, pp. 12–22, 2020, doi: 10.7454/psr.v7i4.1083.
- [22] H. Harikrishnan, I. Jantan, A. Haque, and E. Kumolosasi, "Anti-inflammatory effects of Phyllanthus amarus Schum . & Thonn . through inhibition of NF- κ B , MAPK , and PI3K-Akt signaling pathways in LPS-induced human macrophages," pp. 1–13, 2018.
- [23] S. S. P. Adristy Ratna Kusumo, Farrel Yumna Wiyoga, Haekal Putra Peranda, Izzatidiva Khairunnisa, Raihan Ibadurrohman Suhandi, "Jamu Tradisional Indonesia: Tingkatkan Imunitas Tubuh Secara Alami Selama Pandemi," *J. Public Serv.*, vol. 4, no. 2, pp. 465–471, 2020.
- [24] P. N. Cahyawati and Warmadewa, "Efek Analgetik and Antiinflamantory Kaempferia Galanga (Kencur)," *J. Lingkungan. Pembang.*, vol. 4, no. 1, pp. 15–19, 2020.
- [25] A. R. Khairullah, T. I. Solikhah, A. Nur, M. Ansori, A. Fadholly, and S. Cashyer, "A Review of an Important Medicinal Plant : Alpinia galanga (L .) Willd.," vol. 11, no. 10, pp. 387–395, 2020.
- [26] S. Young, M. Ling, Y. Hun, Y. Kim, and S. Joon, "International Immunopharmacology Anti-in fl ammatory effects of aromatic-turmerone through blocking of NF- κ B , JNK , and p38 MAPK signaling pathways in amyloid β -stimulated microglia," *Int. Immunopharmacol.*, vol. 14, no. 1, pp. 13–20, 2012, doi: 10.1016/j.intimp.2012.06.003.
- [27] F. F. Ugm, No Title. .
- [28] W. Pompimon, J. Jomduang, U. Prawat, S. Mankhetkorn, P. O. Box, and M. District, "Anti-Phytophthora capsici Activities and Potential Use as Antifungal in Agriculture of Alpinia galanga Swartz , Curcuma longa Linn , Boesenbergia pandurata Schut and Chromolaena odorata : Bioactivities Guided Isolation of Active Ingredients Laboratory of ," vol. 4, no. 1, pp. 83–91, 2009.
- [29] E. S. Nikmah Nur Khusnawati, Suwijjiyo Pramono, "Pengaruh Ekstrak Etanolik 50% Herba Pegagan (Centella Asiatica (L.) Urban) Terhadap Peningkatan Proliferasi Sel Limfosit Mencit Jantan Galur Balb/C Yang Diinduksi

- Vaksin Hepatitis B,” *Tradit. Med. J.*, vol. 20, no. September, pp. 164–169, 2015.
- [30] B. Widaryanti, N. Khikmah, A. Analis, K. Manggala, A. Analis, and K. Manggala, “Pengaruh Rebusan Sereh (*Cymbopogon Citratus*) Terhadap Respon Stress Oksidatif Pada Tikus Putih (*Rattus norvegicus*) DIABETES,” no. 2007, 2012.
- [31] I. Wahyuningsih and L. Widiyastuti, “Pengolahan Empon-Empon Menjadi Minuman Kesehatan Berbasis Zero Waste Home Industry,” *J. Berdikasi*, vol. 7, no. 1, pp. 53–61, 2019, doi: <https://doi.org/10.18196/bdr.7157>.
- [32] C. Rini et al., “The Potency And Use Of *Tamarindus Indica* On Various,” vol. 3, pp. 40–54.
- [33] Elfahmi, H. J. Woerdenbag, and O. Kayser, “Jamu: Indonesian traditional herbal medicine towards rational phytopharmacological use,” *J. Herb. Med.*, vol. 4, no. 2, pp. 51–73, 2014, doi: 10.1016/j.hermed.2014.01.002.
- [34] G. Du, M. Xiao, S. Yu, M. Wang, Y. Xie, and S. Sang, “*Phyllanthus urinaria*: a potential phytopharmacological source of natural medicine,” vol. 11, no. 7, pp. 6509–6520, 2018.
- [35] K. Pandey, C. Shevkar, K. Bairwa, and A. S. Kate, *Pharmaceutical perspective on bioactives from *Alstonia scholaris*: ethnomedicinal knowledge , phytochemistry , clinical status , patent space , and future directions*, vol. 0123456789. Springer Netherlands, 2020.
- [36] M. Tian, X. Wu, Y. Hong, H. Wang, G. Deng, and Y. Zhou, “Comparison of Chemical Composition and Bioactivities of Essential Oils from Fresh and Dry Rhizomes of *Zingiber zerumbet* (L .) Smith,” vol. 2020, 2020.
- [37] W.-W. C. and B.-F. Lin*, “Isolation and identification of bioactive compounds in *Andrographis paniculata* (Chuanxinlian),” *Chao Lin Chinese Med.*, vol. 5, no. 17, pp. 1–15, 2010.
- [38] M. Silalahi, “BOTANI AND BIOAKTIVITAS PULAI (*Alstonia scholaris*),” no. July, 2019.
- [39] T. Hartady, R. L. Balia, M. Rizky, A. Adipurna, S. Jasni, and B. Pontjo, “Bioactivity of *Amomum Compactum Soland Ex Maton* (Java Cardamom) as a Natural Antibacterial,” vol. 11, no. 9, pp. 384–387, 2020.
- [40] Y. Niu, S. Wang, C. Li, J. Wang, Z. Liu, and W. Kang, “Effective Compounds From *Caesalpinia sappan* L . on the Tyrosinase In Vitro and In Vivo,” 2020, doi: 10.1177/1934578X20920055.
- [41] G. E. Batiha, L. M. Alkazmi, and L. G. Wasef, “*Syzygium aromaticum* L . (Myrtaceae): Traditional Uses , Bioactive Chemical Constituents , Pharmacological and Toxicological Activities,” 2020.
- [42] Y. D. R. Nutrisia A.S, Indri Kusuma Dewi, “Pengembangan Formula Weandg Secang As Minuman Kemasan Rendah Kalori,” *J. Terpadu Ilmu Kesehat.*, vol. 7, no. 1, pp. 87–95, 2018.
- [43] M. K. R. Indonesia, “Formularium Ramuan Obat Tradisional Indonesia,” 2017.
- [44] K. A. J. Widyanata et al., *PemanfaatanTanaman Obat Keluarga (TOGA) di Masa Pandemi COVID-19*. 2020.
- [45] Y. K. Dewi and B. A. Riyanfrom, “Potensi Tanaman Lokal as Tanaman Obat dalam Menghambat Penyebaran COVID-19,” *J. Pharmascience*, vol. 7, no. 2, p. 112, 2020, doi: 10.20527/jps.v7i2.8793.
- [46] H. Breitinger, “Drug Synergy – Mechanisms and Methods of Analysis.”
- [47] S. Parasuraman, G. S. Thing, and S. A. Dhanaraj, “Polyherbal formulation : Concept of ayurveda,” vol. 8, no. 16, 2014, doi: 10.4103/0973-7847.134229.
- [48] I. Batubara, P. Wulanfrom, L. K. Darusman, and E. Rohaeti, “Optimum mixture of temulawak (*Curcuma xanthorriza*) and meniran (*Phyllanthus niruri*) extract as antiacne Optimum Mixture of Temulawak (*Curcuma xanthorriza*) and Meniran (*Phyllanthus niruri*) Extract as Antiacne,” vol. 030002, no. June, 2020.
- [49] Amor Tresna Karyawati, “Aktivitas Antivirus Simian Retrovirus Serotype-2 (SRV-2) from Ekstrak Meniran and Temu Lawak,” *J. Penelit. Sains*, vol. 14, no. D, pp. 52–55, 2011.
- [50] K. B. Elizabeth Williamson, Samuel Driver, Ed., *Herbal Medicines Interactions*, 1st ed. 1 Lambeth High Street, London SE1 7JN, UK: Pharmaceutical Press, 2009.
- [51] A. Tarun Belwal, Harish C. Andola, Maria S. Atanassova, Bhasker Joshi, Renu Suyal, Shiny

- Thakur and I. D. B. and R. S. R. Bisht, Arvind Jantwal, "Gotu Kola (*Centella asiatica*)," in *Nonvitamin and Nonmineral Nutritional Supplements*, Elsevier Inc., 2019, pp. 265–275.
- [52] T. A. Ajith, U. Hema, and M. S. Aswathy, "Zingiber officinale Roscoe prevents acetaminophen-induced acute hepatotoxicity by enhancing hepatic antioxidant status," vol. 45, pp. 2267–2272, 2007, doi: 10.1016/j.fct.2007.06.001.
- [53] S. Gao et al., "Antimicrobial Activity of Lemongrass Essential Oil (*Cymbopogon flexuosus*) and Its Active Component Citral Against Dual-Species Biofilms of *Staphylococcus aureus* and *Candida* Species," vol. 10, no. December, pp. 1–14, 2020, doi: 10.3389/fcimb.2020.603858.
- [54] G. A. K. D. P. Sri Mulyani, Bambang Admadi Harsojuwono, "Potensi Minuman Kunyit Asam (*Curcuma Domestica* Val. - *Tamarindus Indica* L.) As Minuman Kaya Antioxidant," *Agritech*, vol. 34, no. 1, pp. 65–71, 2014.
- [55] G. Abiramasunfrom, K. R. Sumalatha, and M. Sreepriya, "Effects of *Tinospora cordifolia* (*Menispermaceae*) on the proliferation, osteogenic differentiation and mineralization of osteoblast model systems *in vitro*," *J. Ethnopharmacol.*, vol. 141, no. 1, pp. 474–480, 2012, doi: 10.1016/j.jep.2012.03.015.
- [56] Y. W. Sari Haryanti, Ika Yanti M. Sholikhah, "Efek Sinergis Kombinasi Ekstrak Etanolik Kayu Secang and Rimpang Lempuyang pada Sel Kanker Payudara MCF-7 The Synergistic Effect of Sappan Wood and Bitter Ginger Rhizome Ethanolic Extract in mekanisme efek sitotoksik kuat terhadap sel kanker seskuiterpen," *J. Kefarmasian Indones.*, vol. 9, no. 1, pp. 1–9, 2019, doi: 10.22435/jki.v9i1.324 Jurnal.
- [57] M. E. A. Sayan, A. Sindel, "Traditional and complementary treatments do have a role to play in global health, but probably not in emerging pandemics," *Adv. Integr. Med.*, vol. 7, no. January, pp. 19–21, 2020.
- [58] D. hai Zhang, K. lun Wu, X. Zhang, S. qiong Deng, and B. Peng, "In silico screening of Chinese herbal medicines with the potential to directly inhibit 2019 novel coronavirus," *J. Integr. Med.*, vol. 18, no. 2, pp. 152–158, 2020, doi: 10.1016/j.joim.2020.02.005.
- [59] Y. Yang, "Use of herbal drugs to treat COVID-19 should be with caution," *Lancet*, vol. 395, no. 10238, pp. 1689–1690, 2020, doi: 10.1016/S0140-6736(20)31143-0.
- [60] "Traditional Chinese medicine for COVID-19 treatment," 2020.
- [61] J. K. R. da Silva, P. L. B. Figueiredo, K. G. Byler, and W. N. Setzer, "Essential oils as antiviral agents. Potential of essential oils to treat sars-cov-2 infection: An in-silico investigation," *Int. J. Mol. Sci.*, vol. 21, no. 10, 2020, doi: 10.3390/ijms21103426.
- [62] L. Cheng et al., "Citrus fruits are rich in flavonoids for immunoregulation and potential targeting ACE2," Preprints, no. February, p. 2020020313, 2020, [Online]. Available: <https://www.preprints.org/manuscript/202002.0313/v1>.
- [63] R. V. Nugraha, H. Ridwansyah, M. Ghozali, A. F. Khairani, and N. Atik, "Traditional Herbal Medicine Candidates as Complementary Treatments for COVID-19: A Review of Their Mechanisms, Pros and Cons," *Evidence-based Complement. Altern. Med.*, vol. 2020, 2020, doi: 10.1155/2020/2560645.
- [64] A. F. H. and T. S. W. Honey Dzikri Marhaeny, Aty Widyawaruyanti, Tri Widianandi, "Phyllanthin and hypophyllanthin, the isolated compounds of *Phyllanthus niruri* inhibit protein receptor of corona virus (COVID-19) through in silico approach," *J. Basic Clin. Physiol. Pharmacol.*, vol. 4, no. 32, pp. 809–815, 2021, doi: 10.1515/jbcpp-2020-0473.
- [65] R. Y. Utomo, M. Ikawati, and E. Meiyanto, "Revealing the Potency of Citrus and Galangal Constituents to Halt SARS-CoV-2 Infection," vol. 2, no. March, pp. 1–8, 2020, doi: 10.20944/preprints202003.0214.v1.
- [66] Z. Liu and Y. Ying, "The Inhibitory Effect of Curcumin on Virus-Induced Cytokine Storm and Its Potential Use in the Associated Severe Pneumonia," *Front. Cell Dev. Biol.*, vol. 8, no. June, pp. 1–10, 2020, doi: 10.3389/fcell.2020.00479.

- [67] R. Suravajhala, A. Parashar, B. Malik, and A. V. Nagaraj, "Comparative Docking Studies on Curcumin with COVID-19 Proteins," vol. 10, no. May, 2020, doi: 10.20944/preprints202005.0439.v1.
- [68] I. Musfiroh, A. R. Azura, and D. Rahayu, "Prediction of Asiatic Acid Derivatives Affinity Against SARS-CoV-2 Main Protease Using Molecular Docking," *Pharm. Sci. Res.*, vol. 7, no. 4, pp. 57–64, 2020, doi: 10.7454/psr.v7i4.1086.
- [69] D. S. D. Yoni Rina Bintari, "In Silico Screening of Potential Essential Oil of *Mentha piperita* and *Cymbopogon citratus* Against Covid-19 by Targeting Angiotensin- Converting Enzyme 2 (ACE2) and Aminopeptidase (APN): Molecular Docking Approach," vol. 2, no. 6, p. 18063, 2020.
- [70] O. Sekiou, I. Bouziane, Z. Bouslama, and A. Djemel, "In-Silico Identification of Potent Inhibitors of COVID-19 Main Protease (Mpro) and Angiotensin Converting Enzyme 2 (ACE2) from Natural Products: Quercetin, Hispidulin, and Cirsimaritin Exhibited Better Potential Inhibition than Hydroxy-Chloroquine against," *ChemRxiv*, vol. 2, 2020, doi: 10.26434/chemrxiv.12181404.
- [71] S. K. Enmozhi, K. Raja, I. Sebastine, and J. Joseph, "Andrographolide as a potential inhibitor of SARS-CoV-2 main protease: an in silico approach," *J. Biomol. Struct. Dyn.*, vol. 0, no. 0, pp. 1–7, 2020, doi: 10.1080/07391102.2020.1760136.
- [72] S. Borse et al., "Ayurveda botanicals in COVID-19 management: An in silico multi-target approach," vol. 16, no. 6 June 2021. 2021.
- [73] R. S. Patel, A. G. Vanzara, N. R. Patel, A. M. Vasava, S. M. Patil, and K. S. Rajput, "In-silico Discovery of Fungal Metabolites Bergenin, Quercitrin and Dihydroartemisinin as Potential Inhibitors against Main Protease of SARS-CoV-2," *Coronaviruses*, vol. 01, pp. 1–23, 2020, doi: 10.2174/2666796701999201223163604.



Part of **SPRINGER NATURE**

[PROCEEDINGS](#) | [JOURNALS](#) | [BOOKS](#)

Search



Series: [Advances in Health Sciences Research](#)

Proceedings of the 2nd International Conference on Contemporary Science and Clinical Pharmacy 2021 (ICCSCP 2021)

[HOME](#)

[PREFACE](#)

[ARTICLES](#)

[AUTHORS](#)

[ORGANIZERS](#)

[PUBLISHING INFORMATION](#)



The International Conference on Contemporary Science and Clinical Pharmacy 2021 (ICCSCP) was organized by The Faculty of Pharmacy, Universitas Andalas. ICCSCP was first time held in 2018. This year, we organized this conference for the second time, albeit virtually due to the COVID-19 pandemic. The ICCSCP 2021 proceedings cover all areas of the pharmaceutical and clinical sciences. This proceeding provides a permanent record of what was presented.

Please click [here](#) for the conference website.

Atlantis Press

Atlantis Press – now part of Springer Nature – is a professional publisher of scientific, technical & medical (STM) proceedings, journals and books. We offer world-class services, fast turnaround times and personalised communication. The proceedings and journals on our platform are Open Access and generate millions of downloads every month.

For more information, please contact us at: contact@atlantis-press.com

- ▶ PROCEEDINGS
- ▶ JOURNALS
- ▶ BOOKS
- ▶ POLICIES
- ▶ MANAGE COOKIES/DO NOT SELL MY INFO
- ▶ ABOUT
- ▶ NEWS
- ▶ CONTACT
- ▶ SEARCH

[Home](#) [Privacy Policy](#) [Terms of use](#)



Copyright © 2006-2023 Atlantis Press – now part of Springer Nature



Proceedings of the 2nd International Conference on Contemporary Science and Clinical Pharmacy 2021 (ICCSCP 2021)

Steering Committee

Prof. Dr. Fatma Sri Wahyuni

Faculty of Pharmacy, Universitas Andalas, Indonesia

Chairperson

Prof. Dr. Dian Handayani

Faculty of Pharmacy, Universitas Andalas, Indonesia

Vice Chairman

Dr. Friardi Ismed

Faculty of Pharmacy, Universitas Andalas, Indonesia

Secretary

Dr. Nova Syafni

Faculty of Pharmacy, Universitas Andalas, Indonesia

Finance

HOME

Difa Permatasari, M. Farm

PREFACE

Faculty of Pharmacy, Universitas Andalas, Indonesia

ARTICLES

Scientific Committee

AUTHORS

Prof. Dr. Erizal Zaini

ORGANIZERS

Faculty of Pharmacy, Universitas Andalas, Indonesia

PUBLISHING INFORMATION

Prof. Dr. Armenia

Faculty of Pharmacy, Universitas Andalas, Indonesia

Technical Committee

Dr. Salman

Faculty of Pharmacy, Universitas Andalas, Indonesia

Dr. Najmiatul Fitria

Faculty of Pharmacy, Universitas Andalas, Indonesia

Advisory Board

Yaowaluck Hongkew, PhD

Bumrungrad International Hospital, Bangkok Thailand

Pieter de Boer, PhD

National Institute for Public Health and the Environment, Netherland

Nurmila Sari, M. Ag.Sc., PhD

Division Molecular Medicine, University of Shizuoka, Shizuoka, and
Regenerative Medicine, Jichi Medical University, Shimotsuke, Japan

Thang Nguyen, PhD

**Deputy Head Pharmacology and Clinical Pharmacy, Deputy Head Science
and Technology, Faculty of Pharmacy, Can Tho University, Vietnam)**

PREFACE

Roza Dianita, PhD

School of Pharmaceutical Sciences, Universiti Sains Malaysia, Malaysia

ORGANIZERS

Hilwan Yuda Teruna

PUBLISHING INFORMATION

Department of Chemistry, Universitas Riau, Indonesia

Prof. Dr. Yandi Syukri, S.Si., M.Si

Department of Pharmacy, Universitas Islam Indonesia, Indonesia

Editors

Prof. Dr. Erizal Zaini

Faculty of Pharmacy, Universitas Andalas, Indonesia

Dr. Friardi Ismed

Faculty of Pharmacy, Universitas Andalas, Indonesia

Dr. Rini Agustin

Faculty of Pharmacy, Universitas Andalas, Indonesia

Dr. Nova Syafni

Faculty of Pharmacy, Universitas Andalas, Indonesia

Lili Fitriani, M.Pharm, Sc

Faculty of Pharmacy, Universitas Andalas, Indonesia

Rahmi Yosmar, M.Farm

Faculty of Pharmacy, Universitas Andalas, Indonesia



Proceedings of the 2nd International Conference on Contemporary Science and Clinical Pharmacy 2021 (ICCSCP 2021)

Search

[+ Advanced search](#)

SEARCH

51 articles

Proceedings Article

Immunomodulator Effect Test of Sungkai Leaves (*Peronema canescens* Jack.) Ethanol Extract Using Carbon Clearance Method

Dwisari Dillasamola, Yufri Aldi, Hendra Kurniawan, Ilza Millenia Jalius

Sungkai leaf extract (*Peronema canescens* Jack.) has been used traditionally to cure various diseases for humans. This research aimed to determine the immunostimulant effect of sungkai leaf extract using the carbon clearance method, the total of leukocyte cells, and the percentage of leukocyte cell types....

[+ Article details](#)[+ Download article \(PDF\)](#)

HOME

Proceedings Article

PREFACE

Antihyperglycemic of Olive Leaf (*Olea Europaea* L.) Ethanol Extract in Obesity Diabetic Male Mice (*Mus musculus*)

ARTICLES

Nurhidayati Harun, Elin Herlina, R Siti Rahmah Kurnia

Olive leaf (*Olea europaea*) contains the primary polyphenols: oleuropein, hydroxytyrosol, luteolin 7-glucoside, apigenin 7-glucoside, and verbascoside. Empirically olive leaf is used for lowering blood sugar

because it has antioxidant properties. This study aimed to determine the antihyperglycemic activity...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

***In Vivo* Cytotoxic Activity and Acute Toxicity Test of Ethanol Extract from *Voacanga foetida* (Blume) Rolfe Leaves**

Adriani Susanty, Emrizal, Emma Susanti, Alfonita BR Saragih, Husni Muchtar, Dachriyanus

In vivo cytotoxic activity and acute toxicity test of ethanolic extract *Voacanga foetida* leave against male white mice was done. The in vivo cytotoxic activity used Micronucleus Assay Method while the acute toxicity used LD50 Test and Delay Toxicity Method. The micronucleus assay of *V. foetida* ethanol...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

Evaluation of Antiemetic Drugs Use in Chemotherapy Breast Cancer Patients at Dr. Pirngadi Medan Hospital

[HOME](#) [Eva Sartika Dasopang, Fenny Hasanah, Desy Natalia Siahaan, Jamilah](#)

[Nasution](#)

[PREFACE](#)

[Breast cancer is excessive cell proliferation that attacks the breast tissue.](#)

[ARTICLES](#)

[Breast cancer is a problem that requires serious attention, especially for women. Breast cancer is currently the first cause of death in women in the world. Nausea and vomiting caused by chemotherapy is one of the side](#)

[AUTHORS](#)
[ORGANIZERS](#)
[effects...](#)

[PUBLISHING INFORMATION](#)

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

The Cytotoxic Effect of Ethanolic Extract of *Dasun Tunggal* and Garlic (*Allium sativum*) on Raw 264.7 Cells

Irene Puspa Dewi, Fatma Sri Wahyuni, Yufri Aldi, Dachriyanus

Traditionally, dasun tunggal and garlic (*Allium sativum*) have been widely used as immunomodulators. Therefore, scientific research is needed to support this traditional use. The first step that needs to be done is to see the cytotoxic effect of the ethanol extract of dasun tunggal and garlic on raw 264.7...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

Correlation Between Fasting Blood Glucose and Hemoglobin Level in Women of Childbearing Age: A Base-case Analysis

Najmiatul Fitria, Fivi Melva Diana, Yelly Oktavia Sari

The childbearing age for women is the period when their reproductive organs are still active, so that is when women can give birth to children. Many factors can affect women's health conditions during this fertile age,

including blood sugar and hemoglobin levels. These two factors are very

HOME

important...

PREFACE

[+ Article details](#)

ARTICLES

[+ Download article \(PDF\)](#)

AUTHORS

ORGANIZERS

Proceedings Article

PUBLISHING INFORMATION

Formula Optimization of a Sunscreen Cream of Tomato's Purified Extract

Fianny Rezka Sjahjadi, Febriyenti, Henny Lucida

Tomatoes are rich in lycopene; a strong antioxidant and a potential sunscreen agent. The use of tomatoe 's extract is limited as functional foods, not much as cosmeceutical raw materials. This study aims to optimize the formulation of sunscreen cream of purified tomato extract and to evaluate the sunscreen...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

TLC-Bioautographic and LC-MS/MS Detection of Antimicrobial Compounds from Four Semipolar Extracts of *Cladonia* Species

Friardi Ismed, Widi Novella Desti, Nurwahidatul Arifa, Rustini Rustini, Deddi Prima Putra

To continue the study of Sumatran lichen, six species of the genus *Cladonia* were screened for phytochemicals and their antibacterial bioactivity. The procedure included sequential extraction (semi polar solvent), antibacterial activity tests using the agar diffusion method, and microdilution to determine...

[+ Article details](#)

[HOME](#) [Download article \(PDF\)](#)

[PREFACE](#)

Proceedings Article

[ARTICLES](#)

[Characterization and Antioxidant Activity of Nanoparticles Loaded Jackfruit Leaf Extract \(*Artocarpus heterophyllus*](#)

[AUTHORS](#)

[ORGANIZERS](#)

[Lamk.\)](#)

[Dina Permata Wijaya, Mardiyanto, Herlina, Vabiola Besti Delmoda](#)

Jackfruit leaves extract (*Artocarpus heterophyllus* Lamk.) which contained flavonoid as an antioxidant was unstable and sensitive to environmental conditions. Preparation of nanoparticles loaded jackfruit leaf extract aimed to increase the antioxidant activity through optimal delivery system and maintain...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

[Inventory, Morphological and Antioxidant Profile of the Sumatera Sidaguri \(*Sida* Spp.\) Plants](#)

[Ema Ratna Sari, Netty Suhatri, Friardi Ismed, Deddi Prima Putra](#)

Sidaguri plant is a wild and potential plant that is promoted as a good challenging medicinal plant in the future. However, information related to the diversity of these plants in Indonesia has not been widely exposed, and there are several species of this sidaguri found in Indonesia that are still unknown...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

[Patient Adherence to Rheumatoid Arthritis Medication at an](#)

HOME [Outpatient Clinic Public Hospital in Padang City, West](#)

[Sumatera](#)

PREFACE

[Dita Permatasari, Juni Fitrah, Vonny Novia, Yelly Oktavia Sari](#)

ARTICLES

[Rheumatoid arthritis \(RA\) is a chronic autoimmune disease that attacks the joints and spreads to other organs. Most patients suffer from chronic pain,](#)

[tissue damage, deformity and can cause risks of mortality. The disease](#)

[requires long term and regular treatment to achieve optimal therapeutic](#)

[results....](#)

ORGANIZERS

PUBLISHING INFORMATION

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

[Anti-inflammatory Activities of Gel Extract Marine Sponges \(*Axinella Carteri*\) to White Mice Male](#)

[Delladari Mayefis, Nur Kamilah Idzan, Yunisa Friscia Yusri](#)

Indonesia, especially the Kepulauan Riau, has very wide marine waters, and is rich in natural resources and marine biota diversity. One of the marine biota that has the potential to be developed is the marine sponge as an anti-inflammatory because it contains the flavonoid compound which is able to inhibit...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

[Formulation and Antibacterial Activity of Natural Disinfectant Combination of *Psidium guajava* and *Piper betle* Leaf Infusion Against *Staphylococcus aureus*](#)

[Nori Wirahmi, Camelia Dwi Putri Masrijal, Zul Amri, Ikhsan, Muhammad Ichsan Triyansyah](#)

[HOME](#) **Spraying disinfectant all over the body in the disinfection room is being used to prevent the spread of the SARS- CoV-2 as the cause of the Covid-19 disease. Due to chemical disinfectants have high toxicity to the body, a relatively safe natural disinfectant is needed. The purposes of this study were...**

[PREFACE](#)
[ARTICLES](#)
[AUTHORS](#)

[+ Article details](#)

[ORGANIZERS](#)

[+ Download article \(PDF\)](#)

[PUBLISHING INFORMATION](#)

Proceedings Article

Gastroprotective Effect of Propolis Against Male White Mice Gastric Ulcers Induced by Aspirin

Elsa Badriyya, Zultia Mandasari, Yufri Aldi, A Almahdy

A peptic ulcer is an inflammation of the stomach wall that can cause wounds in the gastric mucosa. Consumption of aspirin over therapeutic doses can cause injury to the stomach. Propolis is known to have an effect to protect stomach damage. This study aims to find out the effect of propolis on male white...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

Multicomponent Crystal of Fenofibric Acid- Saccharin: Characterization and Antihyperlipidemic Effectiveness

Salman Umar, Nedita Putri Bandaro, Deni Anggraini, Erizal Zaini

Fenofibric acid is an active form of fenofibrate which has an antihyperlipidemic effect. Fenofibric acid belongs to BCS class II, which has low solubility and high permeability. The aim of this study was to form multicomponent crystal of fenofibric acid and saccharin that can increase the solubility...

[+ Article details](#)

HOME

[+ Download article \(PDF\)](#)PREFACE

ARTICLES

[Proceedings Article](#)

AUTHORS

[Antihyperlipidemia Activity of Ethyl Acetate Fraction from](#)[Melinjo \(*Gnetum gnemon* Linn.\) Leaf in White Male Wistar](#)[Rats Induced by Propylthiouracil](#)PUBLISHING INFORMATION

Herlina, Ferlina Hayati, Dina Permata Wijaya, Rahma Belinda

The antihyperlipidemia activity of ethyl acetate fraction from Melinjo leaf (*Gnetum gnemon* L.) against white male wistar rats induced by propylthiouracil has been carried out. Wistar male rats were divided into six groups, which are a normal, a positive control, a negative control, and the other three...

[+ Article details](#)[+ Download article \(PDF\)](#)

Proceedings Article

[The Impact of Covid-19 Pandemic on Economy and Health in Bengkulu City](#)

Diana Laila Ramatillah, Stefanus Lukas, Mutiara Nanda

Early 2020, a new type of virus emerged, namely the coronavirus (SARS-CoV-2), and the disease is called Coronavirus Disease 2019 (Covid-19). Covid-19 appeared in Wuhan City, and on March 2th, 2020, it was officially announced that Indonesia had 2 cases of this virus. The Covid-19 cases in Indonesia have...

[+ Article details](#)[+ Download article \(PDF\)](#)

Proceedings Article

Differences of Clinical Disease Activity Index (CDAI) in Rheumatoid Arthritis Patients Towards the Use of Disease Modifying Anti Rheumatic Drugs (DMARD)

ARTICLES

Rahmi Yosmar, Sindi Suija, Yoneta Srangenge, A. Almahdy

AUTHORS

Rheumatoid arthritis is an autoimmune disease commonly associated with progressive disability and systemic complications. Rheumatoid arthritis is characterized by inflammation and synovial hyperplasia, production of autoantibodies, and destruction of bone and cartilage. One of the assessments of rheumatoid...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

Chemical Components and Antibacterial Activity of Essential Oil Extracted from *Citrus x aurantifolia* Peel

Elidahanum Husni, Sofia Ramadani, Suryati, Mesa Irna Suryani, Dachriyanus

Citrus species are among the plants that generate essential oils, which include a variety of chemical components with antibacterial properties. *Citrus x aurantiifolia* is one of the citrus types developed by a cross between lime (*Citrus aurantifolia*) and lemon (*Citrus hystrix*). The purpose of this study...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

Evaluation of Analgetic Use Rationality and Level of Patient Knowledge in Primary Healthcare Facility in Padang

Yoneta Srangenge, Sri Oktavia, Nur Hazizah

Worldwide, oral analgesics are the most widely used drugs with prevalence ranging from 7% up to 35% in various countries. Naturally, this is accompanied by a high level of rationality of use as well. This study aims to evaluate the rationale of administering analgesic drugs and evaluate the level of...

AUTHORS

[+ Article details](#)

ORGANIZERS

[+ Download article \(PDF\)](#)

PUBLISHING INFORMATION

Proceedings Article

The Episiotomy Effect of Topical Combination of Cinnamon Oil and Red Betel on Skin Wound Healing Mechanism

Emelda, Ratih Devi Alfiana, Nurul Kusumawardani, Yolanda, Sitarina Widyarini

The process of episiotomy surgery during childbirth results in injury and pain in the perineum. This study aims to evaluate the effect of the topical combination of cinnamon oil and red betel extract on the healing process of episiotomy incision model in Swiss mice. The research subjects were divided...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

In Vitro Effect of Tetraprenyltoluquinone on Normal Human Leukocyte Cell

Dira Hefni, Fatma Sri Wahyuni, Surya Dharma, Dachriyanus, Rielenda Fadhilah

Tetraprenyltoluquinone is a compound that was successfully isolated from the hexane fraction of stem bark *Garcinia cowa*, Roxb. In previous studies, tetraprenyltoluquinone is known to have anticancer and anti-inflammatory activity. The aim of the study is to investigate the impact of

tetraprenyltoluquinone...
HOME

 [Article details](#)

 [Download article \(PDF\)](#)
ARTICLES

AUTHORS

Proceedings Article

ORGANIZERS

**Pharmacist Counselling Can Change Adherence to Iron
Supplementation and Physical Activity Lifestyle of Anaemic
Pregnant Women in Yogyakarta**

PUBLISHING INFORMATION

Nurul Kusumawardani, Endang Darmawan, Sri Suprapti, Eliza Dwinta, Daru Estiningsih, Emelda

Pharmacists have a role in reducing maternal mortality in 2030 to less than 70 per 100.000 live births. One of the causes of maternal mortality is anemia in pregnancy. The study objective was to assess the determinant factor and effect of solution-focused brief counseling (SFBC) by pharmacists to decrease...

 [Article details](#)

 [Download article \(PDF\)](#)

HOME

Proceedings Article

PREFACE

**Effectiveness Test of Natural Detoxification Facial Wash Gel
Activated Charcoal Palm Shells Using Habatussaudah Scrub**

ARTICLES

Uji Estari, Faizar Farid

Natural detoxification can be obtained by using Habatussaudah which has the ability to increase the natural excretion of toxins from our body. To increase the ability of facial wash gel, activated charcoal palm shell in terms of absorption of toxins, it was further developed with the addition of Habatussaudah...

PUBLISHING INFORMATION

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

**Chemical Content Profile of Essential Oil from Kaffir Lime
(*Citrus hystrix* DC.) in Tanah Datar Regency and
Antibacterial Activity**

Elidahanum Husni, Utari Septiana Putri, Dachriyanus

Essential oil is one type of vegetable oil with many benefits, with the main characteristics of being volatile and having a distinctive aroma. In addition, the essential oil from kaffir lime has been reported to have various bioactivities: antioxidant, antibacterial, antileukemic, and antitussive. This...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

**Analgesic Effect of the Ethanol Extract of Sisik Naga Leaf
(*Drymoglossum* sp.) on White Male Mice (*Mus musculus*)**

Yessi Febriani, Muharni Saputri, Elim Simatupang

HOME

Pain is a sense of discomfort and accompanied by pain due to cell damage to the body. Pain can be overcome with the use of modern medicine and folk remedies. One of the plants that are suspected to be analgesics is the sisik naga leaf (*Drymoglossum* sp.). This test aims to find out the analgesic effect...

PREFACE

ARTICLES

AUTHORS

ORGANIZERS

[+ Article details](#)

PUBLISHING INFORMATION

[+ Download article \(PDF\)](#)

Proceedings Article

Mortality of Hemodialysis Patients with Comorbidities at RSUP Prof Dr R.D Kandou, Manado, North Sulawesi

Stefanus Lukas, Diana Laila Ramatillah, Apricia Latuharhary, Yufri Aldi, Fatma Sri Wahyuni

Background: Hemodialysis is a process in renal replacement function. Comorbid is one of the contributing factors for the high mortality among hemodialysis patients. Objective: To determine the correlation between comorbidities and duration of HD treatment and the effect of HD treatment duration on mortality...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

Sunscreen Activity Test and Fragmentation Analysis of the Active Compound Ethyl Acetate Fraction of Pidada Merah (*Sonneratia caseolaris* L.)

Eka Siswanto Syamsul, Salman, Meri Susanti, Dachriyanus

The leaves of the pidada merah (*Sonneratia caseolaris* L.) are traditionally used by the people of Borneo as a composition of chilly powder. They are

used by applying it to the face when doing outdoor activities exposure for a long period of time. This study aims to determine the sunscreen activity, and...

ARTICLES

[+ Article details](#)

[+ Download article \(PDF\)](#)

ORGANIZERS

Proceedings Article

Primer Design of SNP rs4506565 Transcription Factor 7 like 2 (*TCF7L2*) Gene to Detect Type-2 Diabetes Mellitus

Elsa Badriyya, Afifatul Achyar, Syamsurizal

Diabetes Mellitus is a chronic disease caused by a deficiency or insufficient insulin that increases the levels of blood glucose. Diabetes Mellitus causes several complications that can increase the numbers of mortality. SNP rs4506565 Transcription Factor 7 Like 2 (*TCF7L2*) gene reported to be strongly...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

The Formation and Characterization of Multicomponent Crystal Caffeic Acid-Tromethamine Using the Solvent Drop Grinding Technique

Adhitya Jessica, Amrizal, Uswatul Hasanah, Erizal Zaini, Lili Fitriani

Caffeic acid is a polyphenol compound with low solubility in water. This study aims to prepare and characterize multi-component crystals of caffeic acid, using tromethamine as a coformer, to enhance the dissolution rate without lessening its antioxidant effect. Multi-component crystals were prepared...

[+ Article details](#)

[HOME](#) [Download article \(PDF\)](#)

PREFACE

Proceedings Article

ARTICLES

Phytochemical Screening and Antibacterial Activity of Senduduk Leaves (*Melastoma malabathricum* L.)

AUTHORS

ORGANIZERS

Sri Hainil, Henny Rachdiati, Dwi Prawita

PUBLISHING INFORMATION

Melastoma malabathricum L. leaves is a traditional plant native to tropical Asia, subtropical, and Pacific islands. This study aimed to investigate the antibacterial activity of senduduk leaves (*Melastoma malabathricum* L.) against bacteria *Streptococcus pyogenes*, and *Klebsiella pneumonia*. Method of extraction...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

Anticancer and Antioxidant Activity of Asam Kandis (*Garcinia cowa* Roxb) Leaf Extract and Fraction

Medlyn Jhofi, Elidahannum Husni, Dachriyanus Hamidi

Breast cancer is the second-highest cancer case in the world. One of the causes of cancer is oxidative stress by free radicals. Antioxidants are free radical neutralizing compounds that can prevent cancer. Leaves extract of asam kandis (*Garcinia cowa* Roxb) has been shown to have breast cancer and antioxidant...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

Total Phenolic, Total Flavonoid Content, and α -Glucosidase Inhibitory Activity of *Centella asiatica* (L.) Urb.

HOME Minarti, Megawati, Sofa Fajriah

PREFACE Public awareness about the side effects of chemical drugs has increased in recent years. This has resulted in the rapid use of natural ingredients as medicine. Diabetes mellitus is one of the leading causes of death in the

ARTICLES world. A large number of people with diabetes and the high level of need for drugs...

AUTHORS ORGANIZERS

+ Article details
PUBLISHING INFORMATION

+ Download article (PDF)

Proceedings Article

Study of Jamu as Indonesian Herbal Medicine for Covid-19 Treatment

Marisca Evalina Gondokesumo, Krisyanti Budipramana, Silmi Qurrotu Aini

The spread of the Severe Acute Respiratory Syndrome (SARS-CoV-2) virus has caused the COVID-19 pandemic for more than a year. Daily cases in Indonesia keep increasing following the high mortality rate, entitling it as the epicentre country of Asia. The government advises patients with mild symptoms to...

+ Article details

+ Download article (PDF)

Proceedings Article

Purified Gambier (*Uncaria gambir* Roxb.) and Propolis Performance in Male White Mice (*Mus musculus* L.) Antibody Titer with Measles Vaccine

Lailaturrahmi, Rabani Ahmad, Dwisari Dillasamola, A. Almahdy

Gambier (*Uncaria gambir* Roxb.) and propolis showed a beneficial effect on infectious diseases. This study aims to examine the effect of gambier and propolis on the antibodies of white male mice given measles vaccine using

[HOME](#)
[the antibody titer method, number, and percentage of different types of leukocytes....](#)

[PREFACE](#)

[+ Article details](#)

[ARTICLES](#)

[+ Download article \(PDF\)](#)

[AUTHORS](#)

[ORGANIZERS](#)

[Proceedings Article](#)

[PUBLISHING INFORMATION](#)

[Adverse Events Following Immunization Report and Vaccine Effectiveness of Sinovac. An Interim Report](#)

Abraham Simatupang, Robert H Sirait, Forman Erwin Siagian, Yunita RMB Sitompul, Luana Natingkaseh, Sudung Nainggolan

Vaccination is a prevention of transmission and spread and in order to establish herd immunity against the Covid-19 pandemic. Indonesia chose the Sinovac vaccine, which uses weakened viruses. The objective of the study was to measure the Adverse Events Following Immunization (AEFI) 24-72 hours post-vaccination...

[+ Article details](#)

[+ Download article \(PDF\)](#)

[Proceedings Article](#)

[Correlation Between the Incidence of Drug-Related Problems \(DRPs\) and Clinical Outcomes in Type 2 Diabetes Mellitus Patients with Dyslipidemia at Central Hospital Dr. M. Djamil](#)

Fitri Rachmaini, Yelly Oktavia Sari, Yori Yuliandra, Dini Wariska

Effective management of diabetes therapy is a major challenge in the management of chronic conditions that are accompanied by comorbidities like dyslipidemia. The potential of unwanted events or Drug-Related Problems (DRPs) increases the emergence of comorbidities. This study aims to identify the correlation...

[+ Article details](#)

[HOME](#) [Download article \(PDF\)](#)

PREFACE

Proceedings Article

ARTICLES

Immunostimulant Effect of *Peronema canescens*. Jack Leaves Extract and Propolis in Male White Mice

AUTHORS

ORGANIZERS
Dwisari Dillasamola, Syofyan, Almahdy, Fitri Rachmaini, Mila Yultri,

Hansfani L. L. L. L. L. Elya Gustia, Skunda Diliarosta, Biomechy Oktomalioputri

Peronema canescens leaves are one of the natural ingredients that have the potential to increase the immune system. This study determine the immunostimulant effect which identify the effectiveness of phagocytosis in killing pathogens using the carbon clearance method. The parameters that used in this...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

The Immunomodulatory Activities of Alkaloid (Vf-1) Isolated from Stem Bark of Tampa Badak (*Voacanga Foetida* (Bl) Rolfe) on Raw 264.7 Cells

Zuriva Azwari, Dian Ayu Juwita, Adriani Susanty, Nurdina Putri, Dira Hefni, Dachriyanus, Fatma Sri Wahyuni

Voacanga foetida (Bl) Rolfe (Apocynaceae) locally called Tampa Badak is a widely distributed plant in Indonesia. Indonesian people use this plant as a traditional medicine to treat infections, pain, diarrhea, itching, and swelling. VF-1 is an alkaloid compound that has been previously isolated from the...

[+ Article details](#)

[+ Download article \(PDF\)](#)

HOME

Proceedings Article

PREFACE

Subacute Toxicity of Water Fraction of Africa Leaves

(*Vernonia amygdalina Del.*) on Blood Parameters in Male

White Mice

Dian Ayu Juwita, Helmi Arifin, Muhammad Yasin Abdullah, Dita

ORGANIZERS

Permatasari

PUBLISHING INFORMATION

African leaves (*Vernonia amygdalina Del.*) is one of the plants used as traditional medicine. Although natural, traditional medicinal ingredients are not necessarily safe for use in humans. In this study, a subacute toxicity test of African leaves water fraction was carried out on male white mice. This...

+ Article details

+ Download article (PDF)

Proceedings Article

The Immunostimulant Activities of the Gambir (*Uncaria gambir* Roxb) on Raw 264.7 Cell

Elsya Gustia, Yufri Aldi, Dira Hefni, Sefrianita Kamal, Dachriyanus, Fatma Sri Wahyuni

Gambir (*Uncaria gambir* Roxb) is a plant that is known to have various activities, one of which is immunostimulant. This study aims to determine whether gambir from Payakumbuh affects the viability and immunostimulant activity on RAW 264.7 cells induced by lipopolysaccharide (LPS). This study was conducted...

+ Article details

+ Download article (PDF)

Proceedings Article

Chemical Profiling and Antibacterial Activity of Javanese

Turmeric (*Curcuma xanthorriza*) Essential Oil on Selected Wound Pathogen

HOME
PREFACE

Suryati, Dachriyanus, Irwandi Jaswir, Faridah Yusof

ARTICLES

Essential oil is a volatile liquid with a pungent taste and aroma. The

AUTHORS

essential oil has been used since many years ago for many purpose

ORGANIZERS

including as antimicrobial agent. Javanese turmeric reported containing

essential oil. Therefore, the purpose of this study was to identify chemical

PUBLISHING INFORMATION
profiling of essential...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

Preliminary Study: Bioautography Screening on Edible Local Plants with α -Glucosidase Inhibitor

Nova Syafni, Nurwahidatul Arifa, Friardi Ismed, Deddi Prima Putra

Indonesia is ranked as one of the top ten countries with higher diabetes mellitus cases in 2019. In total, International Diabetes Federation recorded 10.7 million diabetes mellitus cases in 2019. Many researchers reported that diabetes mellitus type 2 could be controlled and managed with diet. Since...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

Molecular Dynamics, Prediction of Toxicity, and Interaction of the Active Compound *Caesalpinia sappan* on Essential Lipids *Klebsiella pneumoniae*

Purnawan Pontana Putra, Fithriani Armin, Nola Florida, Gio Vanny Yusuf, Netty Suharti

Pneumonia is the highest cause of death in the world. The presence of

COVID-19 can worsen the patient's condition. One of the causes of pneumonia is *Klebsiella pneumoniae*. These bacteria can be resistant to antibiotics. In this case, an alternative treatment is needed. *Caesalpinia sappan*, based on pre-clinical...

[Article details](#)

[Download article \(PDF\)](#)

PUBLISHING INFORMATION

Proceedings Article

In vitro Antioxidant Activity and Phytochemical Study of Arbuscular mycorrhizal Fungi Induced Red Ginger (*Zingiber officinale* var. *rubrum*)

Netty Suharti, Dachriyanus, Henny Lucida, Fatma Sri Wahyuni, Purnawan Pontana Putra

Ginger (*Zingiber officinale*), a member of the Zingiberaceae family, has been shown to have anti-inflammatory, antioxidant, anti-nausea/antiemetic, antibacterial, cytotoxic, and antidiabetic effects. Red ginger rhizome has been used as a spice, culinary flavoring, and herbal medicine. Our research shows...

[Article details](#)

[Download article \(PDF\)](#)

Proceedings Article

Chemical Contents Profile of Essential Oil from Calamansi (*Citrus microcarpa* Bunge) Peels and Leaves and Its Antibacterial Activities

Elidahanum Husni, Fitri Yeni, Dachriyanus

Citrus is a plant that has many health benefits and has been widely reported for its pharmacological effects. One of the citrus that has been reported to have antibacterial activity is calamansi (*Citrus microcarpa* Bunge). This study aims to determine the chemical content profile of the

HOME [essential oil...](#)

[Article details](#)

[Download article \(PDF\)](#)

AUTHORS

Proceedings Article

ORGANIZERS

[Membrane of Usnic Acid in Solid Dispersion and Effectiveness in Burn Healing](#)

PUBLISHING INFORMATION

Lili Fitriani, Bella Fista, Friardi Ismed, Erizal Zaini

Usnic acid, a yellow crystalline of secondary metabolite of *Usnea* sp., has been known for its tremendous pharmacological activities including antibacterial and anti-inflammatory activities. The aim of this study was to prepare membrane containing usnic acid and evaluate its effectiveness for burns healing....

[Article details](#)

[Download article \(PDF\)](#)

Proceedings Article

[Evaluation the Knowledge, Perception, and Attitude of Pharmacist Service Towards Pharmacovigilance](#)

Rahmi Yosmar, Nailil Khaira Zulni, Armenia

Pharmacists play an important role in detection, assessment, understanding, and prevention of adverse drug reactions (ADRs) or any other drug related problems. The purpose of this study was to determine the relationship between knowledge, perceptions, and attitudes of pharmacists in the field of service...

[Article details](#)

[Download article \(PDF\)](#)

HOME

Proceedings Article

PREFACE

Antibacterial Activity of Ethyl Acetate Extracts from Mangrove Plants *Rhizophora apiculata* and *Sonneratia alba* Associated Fungi

AUTHORS

R Dyana Sartika, Annisa Rahma, Salsabillah Amelano, Dian Handayani

ORGANIZERS

This study aims to explore the source of antibacterial compounds from mangrove endophytic fungi. Nine fungi were isolated from the leaves, barks, and roots of mangrove *Rhizophora apiculata* and *Sonneratia alba* from West Sumatera, Indonesia. The fungal isolate was cultivated in rice for 4-6 weeks, and...

PUBLISHING INFORMATION

+ Article details

+ Download article (PDF)

Proceedings Article

Various Chloride Salt Addition in Mesoporous Material (SBA-15) Synthesis and Potential as Carrier for Dissolution Enhancer

Uswatul Hasanah, Saleh Wikarsa, Sukmadjaja Asyarie

Nano pores in Santa Barbara Amorphous (SBA-15) has potential to be employed as a carrier to enhance solubility and dissolution. Addition of salt on surfactant solution enabling the SBA-15 synthesis process carried on

room temperature instead of 35-50°C. Atorvastatin Ca, which belongs to

Biopharmaceuticals...

We offer world-class services, fast turnaround times and personalised

communication. The proceedings and journals on our platform are Open

Access and generate millions of downloads every month.

For more information, please contact us at: contact@atlantispress.com

Proceedings Article

Stimulant Activity of Arabica Coffee Leaves (*Coffea arabica*)

PROCEEDINGS

ABOUT

HOME JOURNALS NEWS
 BOOKS CONTACT
 PREFACE SEARCH
 POLICIES

Lily and Glove Flowers (*Syzygium aromaticum* (L.) Merr. & P. M. Perry) Ethanol Extract Mixture in Male White Mice

Rahmad Abdillah, Helmi Arifin, Fitri Rachmainsi, Melda Guswita

Articles are some of the plants that are used as plants are known to have secondary metabolites that have stimulant activity. This study aims to determine the stimulant activity of a combination of coffee leaf extract (*Coffea arabica*...

MANAGE COOKIES/DO NOT SELL MY INFO

ARTICLE DETAILS

Download article (PDF)

Part of **SPRINGER NATURE**

[PROCEEDINGS](#) | [JOURNALS](#) | [BOOKS](#)

Search

Advances in Health Sciences Research

[SERIES HOME](#)

[VOLUMES](#)

[SEARCH](#)

[AIMS & SCOPE](#)

[PUBLISHING INFO](#)

[ABSTRACTING & INDEXING](#)

ISSN (Online):

2468-5739

ISSN (Print):

N/A

Series Editor(s):

vacant



[NEW CONTENT ALERTS](#)

The proceedings series *Advances in Health Sciences Research* aims to publish proceedings from conferences in fields of health sciences and health professions and those addressing the use of science and technology in the delivery of healthcare. [Read full Aims & Scope](#)

All proceedings in this series are **open access**, i.e. the articles published in them are immediately and permanently free to read, download, copy & distribute. Each volume is published under the **CC BY-NC 4.0** user license

which defines the permitted 3rd-party reuse of its articles. The online publication of each proceedings is sponsored by the conference organizers and hence no additional publication fees are required.

Should you wish to publish a proceedings in this series, please then submit your conference proposal via our online submission system

<http://www.atlantis-press.com/account>. Your proposal will be evaluated by a Series Editor(s) and/or a scientific evaluation committee consisting of senior researchers in the relevant field. This is to ensure the integrity and quality of the proceedings that we publish.

For any questions about this series, please **[contact our publishing team](#)**.

Latest Volumes

Most Downloaded

Most Cited

Upcoming Volumes

Proceedings of the International Conference on
Advances in Nano-Neuro-Bio-Quantum (ICAN 2023)

Editors: R. Somashekhar; Preenon Bagchi; Kirthi S. Jawalkar; G.
Dhanalakshmi; Richard Hill; Sanjay N. Harke
Volume 69, November 2023

Proceedings of the 19 th Otorhinolaryngology head and
neck surgery national congress (PERHATIKL 2022)

Editors: Yussy Afriani Dewi; Hamsu Kadriyan; Sinta Sari Ratunanda;
Mohamad Razif Mohamad Yunus; Sandeep Uppal; Pongsakorn
Tantilipikorn;
Volume 68, October 2023

Proceedings of the Second International Nursing Conference "Nursing Profession in the Current Era" (INC 2023)

Editors: Ahmad Rayan; Nidal Eshah; Ghada Abu Shosha; Latefa Dardas; Kristin Wainwright
Volume 67, October 2023




[View More Latest Volumes](#)

Atlantis Press

Atlantis Press – now part of Springer Nature – is a professional publisher of scientific, technical & medical (STM) proceedings, journals and books. We offer world-class services, fast turnaround times and personalised communication. The proceedings and journals on our platform are Open Access and generate millions of downloads every month.

For more information, please contact us at: contact@atlantis-press.com

- ▶ PROCEEDINGS
- ▶ JOURNALS
- ▶ BOOKS
- ▶ POLICIES
- ▶ MANAGE COOKIES/DO NOT SELL MY INFO
- ▶ ABOUT
- ▶ NEWS
- ▶ CONTACT
- ▶ SEARCH

Home [Privacy Policy](#) [Terms of use](#)   

Copyright © 2006-2023 Atlantis Press – now part of Springer Nature

DISCOVER
ISSN SERVICES



SEARCH
OPEN ACCESS RESOURCES



KEEPERS
REGISTRY



ISSN
INTERNATIONAL CENTER



Search



Identifiers

ISSN :2468-5747

Linking ISSN (ISSN-L):
2468-5747



Links

URL

Google

Bing

Yahoo

Koninklijke Bibliotheek



Key-title Advances in biological sciences research

Identifiers

ISSN :2468-5747
Linking ISSN (ISSN-L): 2468-5747

Resource information



Title proper: Advances in biological sciences research.

Other variant title: ABSR

Country: Netherlands

Medium: Online

Record information

My Tools

Share

Print

Display linked data

Discover all the features of the complete ISSN records

Access the full version of the ISSN Portal

