







Original Research

The competency of Indonesian pharmacy students in handling a self-medication request for a cough: a simulated patient study

Cecilia BRATA , Steven V. HALIM , Eko SETIAWAN , Bobby PRESLEY , Yosi I. WIBOWO ,
Carl R. SCHNEIDER 

Received (first version): 2-Jan-2021

Accepted: 11-Apr-2021

Published online: 28-Apr-2021

Abstract

Background: Cough is a common symptom for which people frequently present to community pharmacies. Previous articles from developing countries have shown that the provision of self-medication consultation for cough in community pharmacies were suboptimal, with knowledge deficiency being a contributing factor. However, little is known regarding the ability of pharmacy students in handling self-medication consultations in developing countries.

Objectives: To measure the competency of Indonesian pharmacy students in providing self-medication consultations for patients with chronic cough and to identify factors associated with the provision of appropriate advice.

Method: This study is a cross-sectional study. A simulated patient method using a product and a symptom-based request of chronic cough was used in students from a pharmacy school in Indonesia. The nature and amount of information gathered and advice provided by pharmacy students were noted and audio-recorded. A logistic regression analysis was performed to identify factors associated with the provision of appropriate advice.

Results: The information gathered by participating students was not comprehensive. The most common types of information gathered were related to the nature and duration of the cough. Information relating to accompanying symptoms, medications, and allergies was gathered in less than 60% of the participating students for both product and symptom based scenarios. The appropriate advice of direct medical referral was provided in 54% and 56% of the 183 participating students for the product and symptom-based request scenarios respectively. Asking about symptom duration and prior medical conditions were positively associated with the provision of appropriate advice in the symptom and product based requests respectively.

Conclusion: Student competency in self-medication consultation for chronic cough needs to be improved. Appropriate information-gathering is a predictor of appropriate advice. Further qualitative research identifying factors affecting students' competence in providing self-medication consultation is required, so that suitable interventions are developed and implemented.

Keywords

Cough; Antitussive Agents; Self Medication; Referral and Consultation; Students, Pharmacy; Pharmacists; Patient Simulation; Logistic Models; Multivariate Analysis; Indonesia

INTRODUCTION

Cough, a respiratory reflex to clear the upper airways, is a common condition that affects people worldwide.¹⁻³ Based on the duration, cough can be categorised as acute, sub-acute and chronic.⁴ Acute cough is defined as having a duration of less than three weeks and is commonly caused by upper respiratory tract infections and acute bronchitis.^{4,5} Sub-acute cough usually lasts between three to eight weeks

and is most commonly due to post infectious cough or exacerbation of underlying disease, such as upper airways cough syndrome (UACS), asthma and other pulmonary diseases.⁴ Meanwhile, chronic cough is defined as cough lasting longer than eight weeks and can be attributed to bacterial infection such as *Mycobacterium tuberculosis* infections, environmental exposure such as tobacco smoke and air pollutants, adverse drug reactions to certain medications such as Angiotensin Converting Enzyme (ACE) inhibitors or Angiotensin Receptor Blockers (ARBs), and underlying medical conditions, including asthma and other pulmonary diseases, gastro-oesophageal reflux disease, upper airway cough syndrome, and cardiac origins.^{4,6}

Many patients with cough often self-medicate to alleviate their symptoms, with community pharmacies as well-known sources for medicine supply that are accessible to most patients.⁷⁻¹⁰ Self-medication in this paper refers to the use of medicines by people on their own initiative, without prescription by a medical doctor.^{11,12} Surveys have shown that cough medicines are one of the highest selling over the counter (OTC) product categories in community pharmacies in many countries.^{13,14} This situation emphasises the importance of the role of community pharmacists.⁹ They are health care professionals who are well positioned in the community to provide professional advice for patients with cough.

Cecilia BRATA. PhD. Lecturer. Centre of Medicine Information and Pharmaceutical Care, Faculty of Pharmacy, University of Surabaya. Surabaya (Indonesia). cecilia.brata@gmail.com

Steven Victoria HALIM. Mpharm. Lecturer. Centre of Medicine Information and Pharmaceutical Care, Faculty of Pharmacy, University of Surabaya. Surabaya (Indonesia). stevencvictoria@staff.ubaya.ac.id

Eko SETIAWAN. MSc (Clin Pharm). Lecturer. Centre of Medicine Information and Pharmaceutical Care, Faculty of Pharmacy, University of Surabaya. Surabaya (Indonesia). ekosetiawan.ap@gmail.com

Bobby PRESLEY. Mpharm. Lecturer. Centre of Medicine Information and Pharmaceutical Care, Faculty of Pharmacy, University of Surabaya. Surabaya (Indonesia). bobbypresley@staff.ubaya.ac.id

Yosi Irawati WIBOWO. PhD. Senior Lecturer. Centre of Medicine Information and Pharmaceutical Care, Faculty of Pharmacy, University of Surabaya. Surabaya (Indonesia). yosi_wibowo@staff.ubaya.ac.id

Carl R. SCHNEIDER. PhD. Senior Lecturer. School of Pharmacy, Faculty of Medicine and Health, University of Sydney. Sydney (Australia). carl.schneider@sydney.edu.au




In general, the provision of pharmaceutical consultation in community pharmacies include two important steps, which are: patient assessment and the provision of advice.¹⁵⁻¹⁷ During patient assessment, pharmacists should gather relevant information, such as information on patient identity, symptoms (such as nature and duration of the symptoms, accompanying symptoms, etc.), action that has been taken to alleviate the symptoms, medical conditions, current medications used, and allergies.¹⁸ Then, pharmacists need to use their knowledge to analyse the information gathered to decide what best advice that can be provided to patients. This could be medical referral, recommended products along with the product information, provided non-pharmacological advice, or other advice that are relevant to patient symptoms.^{17,19}

Internationally, the provision of pharmaceutical consultations for patients with cough in community pharmacies is suboptimal, including Indonesia.²⁰⁻²² There are deficiencies in many aspects of pharmaceutical counselling for cough, including lack of history taking and inappropriate advice provision.²⁰⁻²² In developing countries, studies have identified several factors contributing to these deficiencies.²³⁻²⁵ In Indonesia, previous research has identified three main factors influencing pharmaceutical consultation performance: pharmacy staff-patient interaction, the organisational context of the pharmacy, and the external pharmacy environment (Figure 1).²³ These factors are complex and inter-related; for example the lack

of knowledge and competency of pharmacy staff to provide quality consultation are influenced by deficiencies in pharmacy education with respect to management and pharmaceutical consultation for self-medication.²³ Indonesian pharmacists also perceive that there is a problem with relevancy between what is taught in their pharmacy education and the knowledge required for practice, and therefore they felt that their knowledge is insufficient to provide optimal self-medication services.^{23,26} Similar findings have been reported in the literature from other developing countries, such as Brazil and Vietnam.^{24,27}

In Indonesia, to become a pharmacist, students are required to complete a four-year Bachelor of Pharmacy (BPharm) degree, followed by a one-year pre-registration training programme (Apothecary programme). Pharmacy curricula are regulated by the Ministry of Education and are typically based on the curriculum developed by the Indonesian Association of Higher Education of Pharmacy (Asosiasi Perguruan Tinggi Farmasi Indonesia, or APTFI).²⁸ While each pharmacy school has the freedom to design their own curriculum, APTFI suggests that the courses or units of study within the BPharm curriculum are designed to be predominantly theory based, with the addition of laboratory practice or simulation, whereas the apothecary program focuses on applied pharmacy practice.²⁸

The quality of Indonesian pharmacy education varies widely across institutions.²⁸ Therefore, from 2018 onwards, students in the apothecary programme are required to pass



Pharmacy staff–patient interactions:

- Patients’ response
 - Patients’ awareness of the role of and trust in pharmacy staff
 - Types of patients’ requests
 - Patients’ cultural characteristics
- The professionalism of pharmacy staff
 - Knowledge and competence
 - Ability to focus on the pharmacy job
 - Availability and implementation of standard operating procedure

The organisational context of the pharmacy

- Availability of trained staff
- Availability of time to provide consultations
- Staff affordability
- Availability of facilities to provide consultation – counselling space
- Dual employment – leading to potential conflicts of interest.

The external pharmacy environment

- The number of trained pharmacy staff in the research setting
- Relevance of pharmacy education to the needs of pharmacy practice**
- Support from the Indonesian Pharmacists Association
- A competitive business environment
- The policy environment

Figure 1. Factors related to the provision of self-medication consultation in community pharmacies in eastern Indonesia from Brata et al.²³

a national, computer-based test (CBT) examination before they can be registered as pharmacists.²⁹ In addition, students also need to take an objective structured clinical examination (OSCE) as a formative assessment.²⁹ The OSCE is intended to become a summative barrier examination; however, the transition is currently postponed as a result of the COVID pandemic.

Studies looking at the situation of Indonesian pharmacy education are limited. A study conducted at the University of Gadjah Mada, has evaluated their 422 apothecary students' performance using an OSCE.³⁰ The authors reported that approximately 85% of the 422 students were able to pass the minor ailment station.³⁰ However, the study aim was to measure students' experience and perception of the OSCE and not to measure the competence of students in responding to patients with minor illness.

This study is conducted to measure the competency of Indonesian pharmacy students in providing self-medication consultations for patients with a cough. In this study, the competency of pharmacy students in providing self-medication consultation is defined as a combination of students asking/gathering essential information such as information related to patient identity, symptoms (nature, duration, action taken to alleviate symptoms, associated symptoms, etc.), medical conditions, medications used, and allergies during patient assessment and students subsequently providing appropriate advice to the patient. Therefore, the specific objectives are to: (1) describe the type and amount of information gathered; (2) describe the type and amount of advice provided, (3) measure the overall proportion of students providing appropriate advice, and (4) identify factors associated with the provision of appropriate advice.

METHODS

Ethics approval

Ethics approval was sought from the Ethical Committee University of Surabaya, Indonesia (No 057/KE/XII/2018). To protect the identity of the research setting and student participants, any identifying information was not allowed to be published.

Research context

In this particular university, topics related to self-medication services are provided to third year BPharm students. The course (COMM PHARM 1) comprises a 4 hour session per week, consisting of a 1-hour lecture and 3-hours of laboratory practice for 14 weeks. In the laboratory practice component, students are expected to be able to respond to a simulated patient (i.e., tutors or students acting as a patient) requesting self-medication for several minor ailments such as painful conditions, respiratory problems, gastrointestinal tract problems, skin problems, and eye and ear problems. The COMM PHARM 1 course is intended to provide competency in the provision of advice when responding to minor ailment requests to the level that supervised provision of advice to the public is subsequently able to be conducted in the apothecary programme.

Participant recruitment

All third year pharmacy students in the research site who had completed COMM PHARM 1 were eligible for recruitment. In the first week of semester 6, the researchers announced and explained the study to all enrolled students in COMM PHARM 2. Announcement was made in the COMM PHARM 2 class because this is the subsequent class that needs to be enrolled after third year students completing COMM PHARM 1. Several students who had completed COMM PHARM 1 but did not enrol in COMM PHARM 2, identified via the school data register, were also contacted separately. Students were asked to participate in a simulated patient assessment at a time that suited them. Free cakes/pastry/snacks were provided after participation and an extra-curricular student point which can be printed on their non-academic transcript upon graduation was also offered. Participation was voluntary with students who agreed to participate providing informed consent.

Study instrument

This study is a cross-sectional study. A simulated patient method was used to present the scenario of cough due to worsening of asthma. The scenario was adapted from Schneider et.al and was presented as both a product and symptom-based request (Online appendix).²¹ The data collection form was adapted from previous literature and consisted of a checklist of participants' characteristics, the type of information gathered and the advice provided.^{15,17} The types of information gathered included important information that needs to be gathered in patients with cough as stated by Blenkinsopp *et al.*¹⁶ The information elements comprised: patient identity, symptoms (including nature of cough, duration of cough, and accompanying symptoms including danger symptoms), action taken, medical conditions, medications used, and allergies. The types of advice provided were categorised as medical referral, recommending products, recommending non-pharmacological advice or other advice (including advice on follow up).¹⁷ As per previous literature, the appropriateness of the advice for this scenario was defined as direct medical referral.²¹ In order to adjust to the local context, four Indonesian academics and practitioners in pharmacy practice assessed the scenario for content validity and reviewed the data collection form. No significant changes were made in the scenario and the data collection form after being revalidated.

Seven simulated patients, consisting of two pharmacists who were PhD students, three recent pharmacist graduates, and two drug information pharmacists were recruited and trained beforehand. The training consisted of explanation of the scenarios, how to fill the data collection form, and role plays. Role plays were repeated until the simulated patients were deemed able to accurately enact the scenarios and fill in the data collection form as evaluated by the principal researcher (CB). Details of the characteristics of the simulated patients as well as the training is presented in Online appendix and is informed by an article by Resende *et al.*³¹

Table 1. The comparison of demographic characteristics between participants and the total population

Demographic characteristics	Sample n=183 (%)	Population n=304 (%)
Age in years; mean (SD)	20 (0.95)	20 (0.85) ∞
Gender: female	159 (87%)	248 (82%)
Previous education		
High school	125 (68%)	198 (70%)∞
Pharmacy assistant school (equivalent to high school)^	58 (32%)	84 (30%)∞
Have any experience of working or training in community pharmacy		
Yes	54 (30%)	80 (28%)∞
No	129 (70%)	202 (72%)∞
Average GPA*; mean (SD)	2.8 (0.52)	2.7 (0.56)
Mark for COMM PHARM 1 course: #		
A	0	1 (0,3%)
AB	9 (5%)	10 (3%)
B	65 (36%)	93 (31%)
BC	65 (36%)	100 (33%)
C	33 (18%)	67 (22%)
D	11 (6%)	33 (11%)
E	0	0

∞Data is from 282 students
 *The maximum Grade Point Average (GPA) is 4.
 #The interpretation of the marking score were as follows: A = more than or equal to 81, AB = 73 to 81, B=66 to 72, BC= 60 to 65, C=55 to 59, D=40 to 54, E= less than 40. The maximum marking score is 100.

Data collection

One simulated patient was assigned to each student. During data collection, each participating student was presented with a product and a symptom-based request. For the product-based request, the simulated patients said “I want to buy woods® merah” (woods® merah is an Indonesian brand of cough medicines that contain dextromethorphan HBr and diphenhydramine HCl). For the symptom-based requests, the simulated patient started the conversation by saying “I want to buy medicines for cough, what do you recommend?”. No further information was provided unless it was requested by the students. The whole process was audio-recorded. The simulated patients also recorded in writing all the information asked and the advice provided by participating students. After completing each meeting, the simulated patients validated the written data in the data collection form with the audio recorder.

Data analysis

Descriptive statistics (i.e. frequencies) were used to summarise data related to the types of information gathered and types of advice provided. Univariate followed by multivariate logistic regression was used to identify factors associated with the participants’ provision of appropriate advice. Independent variables were derived from participant characteristics (i.e., gender, age, previous education, having working/training experience in community pharmacies, grade point average (GPA), and mark for COMM-PHARM 1 class), the number of questions asked, and the type of question asked (i.e., who is the patient, nature of cough, duration of cough, accompanying symptoms, action taken, medical conditions, medications used, allergies). The significant variables from the univariate analysis (p<0.05) were then entered into the multivariate model. A backward elimination method with a significance threshold of 0.05 was implemented to obtain the final, multivariate model. Odds ratios (OR) and 95% confidence intervals were calculated. IBM SPSS Statistics for Windows version 23 (Armonk, NY: IBM Corp) was used for the analysis.

RESULTS

Based on the faculty registration data, the population of third year students who had completed COMM-PHARM 1 in 2018 was 304. Of these, 183 agreed to participate in this study, resulting in a response rate of 60.2%. Details related to participants’ characteristics are presented in Table 1.

All students (100%) asked questions before providing advice. The average number of questions asked for product and symptom-based requests for cough due to worsening asthma scenarios were 5 and 6 with the standard deviation of 1.5 and 1.4 respectively. Information related to patient identity, and the nature and duration of cough were the most common types of information gathered in both scenarios (≥ 80% of the participating students). In contrast, accompanying symptoms, medications, and allergies were the least frequently gathered types of information; they were gathered in less than 60% of the participating students. Details related to the types and amount of information gathered can be seen in Figure 2 and Table 2.

The types and amount of advice provided by participants are outlined in Table 3. In the product-based request scenario, medical referral was advised by 99 students (54% of the 183 participants). Of these 99 students, 21 and 26 students also recommended cough medicines and non-pharmacological/follow-up advice, respectively. In the symptom-based request scenario, 102 students (56% of the 183 participants) recommended direct medical referral.

Table 2. The types of information gathered

The types of information gathered	Product-based request n=183 (%)	Symptom-based request n=183 (%)
Patient identity	147 (80%)	81 (81%)
Symptoms		
Nature of cough	158 (86%)	165 (90%)
Duration of cough	161 (88%)	167 (91%)
Accompanying symptoms	79 (43%)	68 (37%)
Action taken	125 (68%)	131 (72%)
Medications	91 (50%)	106 (58%)
Medical conditions	119 (65%)	137 (75%)
Allergies	93 (51%)	88 (48%)






Figure 2. The number of questions asked by students

Fifteen and 23 of these 102 students also recommended cough medicines and non-pharmacological/follow up advice, respectively.

Appropriate advice, which was defined as advice that contained direct medical referral, was provided in 99 (54%), and 102 (56%) of 183 participating students in the product-based requests and symptom-based requests, respectively.

From the univariate analysis, significant variables for providing appropriate advice comprised of asking about action taken and duration of cough in the product-based request scenario (Table 4). Meanwhile, significant variables for providing appropriate advice in the symptom-based request scenario consisted of: GPA, number of questions asked, asking about duration of cough, action taken, and other medical conditions (Table 5).

After inputting these significant variables into the multivariate analysis, we determined that the nature of the information gathering by students is associated with the provision of appropriate advice. In the product-based request scenario, students who asked about the duration of cough were more likely to provide appropriate advice than students who did not ask this question (OR 4.8, 95%CI 1.68-13.57). Meanwhile, in the symptom-based request scenario, students who asked about other medical conditions were more likely to provide appropriate advice

than students who did not (OR 2.1, 95%CI 1.05-4.19). Demographic characteristics of participating students (i.e. age, gender, previous education, have working/training experience in community pharmacies, GPA, mark for COMM PHARM 1 course) were not associated with the provision of appropriate advice.

DISCUSSION

This is the first study in Indonesia that examined the ability of pharmacy students in gathering information and providing appropriate advice for a simulated patient with chronic cough. All students participated in this study were novice learners, which describes that they just learnt about self-medication consultation during their undergraduate pharmacy curriculum.

This study indicates that the information-gathering process among students in this school, needs to be improved. The lecture notes and laboratory practice module of COMM-PHARM1 class indicated that topics related to information-gathering have been provided. However, this study showed that students did not gather appropriate information when encountering our simulated patients. Important information such as medical conditions, medications used, and accompanying symptoms that are needed to enable students to identify the root of the problems were asked in 40% to 75% of the encounters. This contrasts with the results from an Australian study, in which most of their novice students perform well during the information-gathering process.³² While it is argued that complete information is not always necessary to be gathered in order to provide appropriate advice, it is safer for novice students to gather complete information for assessing patients correctly and hence appropriate advice.³³

In this case, advice was considered appropriate when students recommended direct medical referral. However, in both scenarios, only about 55% of them were able to do so. This figure is about similar to findings from an Australian study, which stated that appropriate advice was provided in 52% of their novice students in the case of childhood fever.³² Other than incomplete information gathered as explained above, these participating students

Table 3. The types and amount of advice provided

The types of advice provided	Symptom based requests n =183 (%)	Product based requests n=183 (%)
(1) Direct medical referral	68 (37%)	61 (33%)
(1) Direct medical referral and (2) suggesting medicines*	11 (6%)	12 (7%)
(1) Direct medical referral, (2) suggesting medicines, and (3) suggesting non-pharmacological advice or other advice.**	5 (3%)	9 (5%)
(1) Direct medical referral and (3) suggesting non-pharmacological advice, or other advice	18 (10%)	17 (9%)
(2) Suggesting medicines.	29 (16%)	29 (16%)
(2) Suggesting medicines and (3) suggesting non-pharmacological advice, or other advice.	41 (22%)	47 (26%)
(3) Suggesting non-pharmacological advice or other advice.	1 (0%)	4 (2%)
(4) No advice suggested.***	10 (6%)	4 (2%)

*Medicines suggested was a variety of cough medicine brand names.

**Examples of non-pharmacological advice or other advice including: enough rest, eat healthy food, drink a lot of water, avoid strenuous exercise, avoid dusty or unhygienic places, drink herbal medicines such as ginger and honey, and advising on follow up (such as go to a doctor if there is no improvement after trying the suggested medicines, go back to the pharmacist if there is no improvement after trying the suggested medicines).

***No advice suggested meant that the students said that they did not know the answer or said that they did not know how to respond to the scenario in the simulated patient test.

Table 4. Factors associated with the provision of appropriate advice – product-based request scenario			
Factors		Univariate analysis	
		OR (95%CI)	p-value
Age		0.95 (0.696, 1.286)	0.723
Gender			0.994
Male		1.00 (0.424, 2.374)	
Female		Reference [^]	
Previous education			0.449
High school		1.27 (0.682, 2.376)	
Pharmacy assistant school		Reference	
Have any experience in working or training in community pharmacies			0.693
Yes		0.88 (0.465, 1.663)	
No		Reference	
GPA		1.42 (0.810, 2.480)	0.222
Mark for COMM PHARM 1 course			
AB		0.71 (0.118, 4.319)	0.654
B		0.86 (0.228, 3.224)	0.820
BC		0.49 (0.131, 1.873)	0.290
C		0.69 (0.168, 2.799)	0.599
D		Reference	
Number of questions asked		1.16 (0.955, 1.417)	0.132
Asking who is the patient			0.845
Yes		0.93 (0.447, 1.934)	
No		Reference	
Asking nature of cough			0.821
Yes		1.10 (0.474, 2.566)	
No		Reference	
Asking duration of cough			0.003*
Yes		4.77 (1.677, 13.566)	
No		Reference	
Asking accompanying symptoms			0.413
Yes		0.78 (0.435, 1.408)	
No		Reference	
Asking action taken			0.020*
Yes		0.47 (0.250, 0.887)	
No		Reference	
Asking medications			0.264
Yes		1.40 (0.778, 2.500)	
No		Reference	
Asking medical conditions			0.415
Yes		1.29 (0.701, 2.370)	
No		Reference	
Asking allergies			0.326
Yes		0.75 (0.417, 1.338)	
No		Reference	

*significant p value; [^]the term “reference” category is a category of comparison for the other categories

may also have limited knowledge of disease and therapy and inability to analyse the information gathered; and hence inappropriate advice.^{17,32} The syllabus in this pharmacy school shows that during semester 1 to 4, students only learn about basic sciences (human anatomy & physiology, pathophysiology, and pharmacology). Topics related to diseases and drug therapy, drug related problems, and patient counselling are provided in semester 5 and above. It could be that these students may not know that the case is cough due to worsening of asthma since they haven't learnt much about asthma, and therefore they did not know that direct medical referral was needed.

A potential factor to sub-optimal student performance is the teacher-student ratio. A review of the COMM PHARM1 classroom student list and tutor schedule identified that the teacher-student ratio was 1:20 during laboratory practices and 1:80 during lectures. Such ratios may be insufficient for students to receive adequate supervised practice and feedback during simulated role-plays. As the effectiveness of simulation learning relies on practice and

feedback for each learner, measures should be undertaken to optimised the teacher-student ratio.³⁴ Furthermore, with the current curriculum design, students are expected to be ready for supervised provision of minor ailment care to the public in the apothecary programme after completing the COMM PHARM1 course. This indicates that this one semester course carries a lot of weight in the curriculum. Redesigning the BPharm curriculum, such as by adopting the spiral curriculum in which key concepts are revisited and reinforced throughout the curriculum with increasing levels of complexity, could be more appropriate than a stand-alone course.^{35,36} Further qualitative research looking at the design of the current curriculum, students and teachers' perceptions, the teaching-learning process including the mode of delivery is required for designing interventions for the optimisation of student performance.

Simulated patient technique has been used widely in pharmacy education, including for training in consultation for self-medication requests.³⁴ Using simulated patient technique as an educational tool may improve training

Table 5. Factors associated with the provision of appropriate advice – symptom-based request scenario			
Factors		Univariate analysis	
		OR (95%CI)	p-value
Age		1.01 (0.742, 1.376)	0.949
Gender			
Male		1.13 (0.473-2.695)	0.784
Female		Reference [^]	
Previous education			
High school		1.56 (0.831, 2.909)	0.167
Pharmacy assistant school		Reference	
Have any experience in working or training in community pharmacies			
Yes		0.58 (0.307, 1.104)	0.098
No		Reference	
GPA		1.82 (1.029,3.217)	0.039*
Mark for COMM PHARM 1 course			
AB		1.40 (0.232, 8.464)	0.714
B		2.31 (0.616, 8.682)	0.214
BC		2.63 (0.698, 9.875)	0.153
C		2.10 (0.514, 8.573)	0.301
D		Reference	
Number of questions asked		1.26 (1.014, 1.553)	0.037*
Asking who is the patient			
Yes		1.24 (0.592, 2.594)	0.569
No		Reference	
Asking nature of cough			
Yes		1.33 (0.103, 1.034)	0.057
No		Reference	
Asking duration of cough			
Yes		3.05 (1.014, 9.167)	0.047*
No		Reference	
Asking accompanying symptoms			
Yes		1.110 (0.606, 2.032)	0.735
No		Reference	
Asking action taken			
Yes		2.14 (1.114, 4.108)	0.022*
No		Reference	
Asking medications			
Yes		1.19 (0.659, 2.148)	0.563
No		Reference	
Asking medical conditions			
Yes		2.18 (1.107, 4.310)	0.024*
No		Reference	
Asking allergies			
Yes		1.12 (0.627, 2.003)	0.701
No		Reference	

*significant p value, ^the term “reference” category is a category of comparison for the other categories

since it can reinforce knowledge, integrate between knowledge and practice, and improve professional skills and attitudes (e.g., communication skills, decision making, empathy, teamwork, social and cultural competency, and confidence).^{34,37} In addition as an educational tool, simulated patient technique can also be an assessment tool for evaluating the effectiveness of the training provided. Indeed, simulated patients in the form of OSCE is planned to be a compulsory element of the Indonesian national pharmacist registration examination.³⁰ Evidence suggests that examinations using simulated patient technique challenges students to think critically and indicates student performance in real clinical encounters.³⁷ Indonesian pharmacy schools, therefore, should use this technique in the teaching and evaluation of pharmacy students.

The multivariate analysis identified that asking a particular type of question (i.e., duration of cough and medical conditions) are factors positively associated with the provision of appropriate advice (Tables 4 and 5). This finding is in keeping with a previous Australian study in

pharmacy staff responding to a very similar scenario.²¹ It could be that by asking such questions, students were able to discern that the patient had a contraindication for self-medication and recommend medical referral. It is important to train students to be able to ask key questions according to the clinical situation to improve the appropriateness of advice for minor ailments. Further, as there is a current lack of formal guidance on information-gathering when responding to patients with self-medication requests in Indonesia, policy makers such as the Indonesian Pharmacists Association (IAI), the Ministry of Health, and the APTFI may need to develop and promote a national protocol to improve appropriate information-gathering and to optimise the provision of care.

Research evaluating the competence of pharmacy students providing self-medication consultation in Indonesia is limited. Kristina et.al reported that 85% of 422 apothecary students from the University of Gadjah Mada, passed the the minor ailment station in their OSCE assessment.³⁰ However, detailed aspects assessed in this minor illness

station were not provided in the article and the population used were different with this study's population (apothecary students vs third year students in this study); thus making direct comparison difficult. It could be hypothesized that final year/apothecary programme students may have superior outcomes compared to students at an earlier stage in their education, as found in an Australian study.³²

Despite the students in this study only being halfway through their pharmacist training, their performance was superior to Indonesian pharmacy staff, as reported in previous research.^{20,38} When responding to a simulated patient asking for a particular product for chronic cough, all of these students conducted information gathering and 54% of them were able to provide appropriate advice. In contrast, only 1% of pharmacy staff performed information gathering and none provided appropriate advice in the previous articles.^{20,38} Although the cases of chronic cough used between this study and the previous studies were not exactly the same, the essence and complexity of the cases are similar; and therefore we think they can be compared.

Better performance of these students compared to pharmacy staff as reported in the previous article may be due to several reasons. As stated in the theory of the Efficiency-Thoroughness Trade-Off (the ETTO principle), there is a balance or tradeoff between being efficient and being thorough.³⁹ If demands to productivity are high, thoroughness is usually reduced. On the other hand, if demands to safety are high, efficiency is reduced. Following this theory, it could be that pharmacy staff were being efficient due to demands of high productivity, whereas students were being thorough because they knew that they were being assessed and therefore they will try harder to show their best performance. Another possible reason of why Indonesian pharmacy staff performed more poorly than students might be that the person that typically handles patients with self-medication requests in pharmacies is a non-pharmacist, as has been reported in Indonesian and other developing countries' literature.⁴⁰⁻⁴² This being the case, measures should be taken to ensure that pharmacists are able to provide self-medication services when they are on duty. Further research on why students perform better than pharmacy staff is required.

Limitations

This study was only conducted in one pharmacy school in Indonesia. Because the quality of Indonesian pharmacy schools has been reported to vary, these findings may not be generalisable for all pharmacy students in Indonesia.²⁸ Therefore, further research using samples of different pharmacy schools are required to obtain a general picture of Indonesian pharmacy students' ability in self-medication consultation. Furthermore, the study was conducted in

novice students and therefore may not reflect the ability of pharmacy graduates. Future research on evaluating the ability of final year pharmacy students in handling self-medication consultation is also required. In addition, a Hawthorne effect may have been generated, because this is an overt method, in which participating students knew that they were being assessed. However, although these students have showed their best performance, only half of them were able to provide appropriate advice; and therefore there is still a need for improvement. Lastly, there was a possibility of selection bias in this study. However, the characteristics of the participating students were similar to the student cohort (Table 1) and therefore it might not significantly affect the findings.

CONCLUSIONS

The performance of novice students in responding appropriately to a simulated patient with a self-medication request for cough is sub-optimal. Appropriate information-gathering is a predictor of appropriate provision of advice. Measures to improve student performance, such as curriculum restructuring, optimising teacher-student ratio, and developing a protocol for appropriate information-gathering may be required. Further qualitative research to identify factors affecting student performance is required to develop strategies to improve student response to self-medication requests.

CONFLICT OF INTEREST

None.

FUNDING

This study was carried out with the grant from The University of Surabaya Institute of Research and Community Services.

AUTHOR ROLES (CRediT)

Conceptualization: CB, YIW, ES, SV.

Data curation: CB, SV.

Formal analysis: CB.

Funding acquisition: CB, SVH.

Investigation: CB, SVH, YIW, ES, BP.

Methodology: CB.

Project administration: CB, SVH.

Resources: SVH, BP.

Supervision: CB, CS.

Validation: CB, YIW, ES, SV.

Visualization: CS, CB.

Writing – original draft: CB.

Writing – review & editing: CS, CB.

References

1. Chung KF, Pavord ID. Prevalence, pathogenesis, and causes of chronic cough. *Lancet*. 2008;371(9621):1364-1374. [https://doi.org/10.1016/s0140-6736\(08\)60595-4](https://doi.org/10.1016/s0140-6736(08)60595-4)
2. Kauffmann F, Varraso R. The epidemiology of cough. *Pulm Pharmacol Ther*. 2011;24(3):289-294. <https://doi.org/10.1016/j.pupt.2010.10.012>
3. Tietze KJ. Cough. In: Krinsky DL, Ferreri SP, Hemstreet B, et al., editors. *Handbook of Nonprescription Drugs: An interactive approach to self-care*. Washington DC: APhA; 2015.



4. Irwin RS, French CL, Chang AB, Altman KW; CHEST Expert Cough Panel*. Classification of Cough as a Symptom in Adults and Management Algorithms: CHEST Guideline and Expert Panel Report. *Chest*. 2018;153(1):196-209. <https://doi.org/10.1016/j.chest.2017.10.016>
5. Holzinger F, Beck S, Dini L, Stöter C, Heintze C. The diagnosis and treatment of acute cough in adults. *Dtsch Arztebl Int*. 2014;111(20):356-363. <https://doi.org/10.3238/arztebl.2014.0356>
6. Perotin JM, Launois C, Dewolf M, et al. Managing patients with chronic cough: challenges and solutions. *Ther Clin Risk Manag*. 2018;14:1041-1051. <https://doi.org/10.2147/tcrm.s136036>
7. Limaye D, Limaye V, Krause G, Fortwengel G. A Systematic review of the literature to assess self medication practices. *Ann Med Health Sci Res*. 2017;7:1-15.
8. Ambizas EM. Acute cough—does anything help?. *US Pharm*. 2019;1:8-12.
9. Rutter P. Role of community pharmacists in patients' self-care and self-medication. *Integr Pharm Res Pract*. 2015;4:57-65. <https://doi.org/10.2147/iprp.s70403>
10. Darmanin Ellul R, Cordina M, Buhagiar A, Fenech A, Mifsud J. Health complaints and use of medicines among adolescents in Malta. *Pharm Pract (Granada)*. 2008;6(3):165-170. <https://doi.org/10.4321/s1886-36552008000300008>
11. Brandt N. Self-medication. In: Gellman MD, Turner JR, editors. *Encyclopedia of Behavioral Medicine*. New York: Springer; 2013.
12. International Pharmaceutical Federation. Joint statement by The International Pharmaceutical Federation and The World Self-Medication Industry: Responsible self-medication. <https://www.fip.org/file/1484> (accessed Feb 1, 2021).
13. The World Self-Medication Industry. Proceedings of WSMI 6th Asia Pacific Regional Conference. Beijing, 26 – 29 Oct 2004.
14. Connelly D. The OTC market in Britain in 2015. *Pharm J*. 2016;296:7887.
15. Schneider C. Provision of non-prescription medication by community pharmacists in Western Australia. University of Western Australia; 2010.
16. Blenkinsopp A, Duerden M, Blenkinsopp J. *Symptoms in the pharmacy: A guide to the management of common illnesses*. Oxford: Wiley-Blackwell; 2018.
17. Brata C. The provision of self-medication consultations in community pharmacies in an Eastern Indonesian setting. University of Western Australia; 2015.
18. Brata C, Gudka S, Schneider CR, Everett A, Fisher C, Clifford RM. A review of the information-gathering process for the provision of medicines for self-medication via community pharmacies in developing countries. *Res Social Adm Pharm*. 2013;9(4):370-383. <https://doi.org/10.1016/j.sapharm.2012.08.001>
19. Krinsky DL, Ferreri SP, Hemstreet B, et al, editors. *Handbook of nonprescription drugs: an interactive approach to self-care*. 18 ed. Washington DC: APhA; 2015.
20. Brata C, Schneider CR, Marjadi B, Clifford RM. The provision of advice by pharmacy staff in eastern Indonesian community pharmacies. *Pharm Pract (Granada)*. 2019;17(2):1452. <https://doi.org/10.18549/pharmpract.2019.2.1452>
21. Schneider CR, Everett AW, Geelhoed E, et al. Provision of primary care to patients with chronic cough in the community pharmacy setting. *Ann Pharmacother*. 2011;45(3):402-408. <https://doi.org/10.1345/aph.1p514>
22. Mináriková D, Fazekaš T, Minárik P, Jurišová E. Assessment of patient counselling on the common cold treatment at Slovak community pharmacies using mystery shopping. *Saudi Pharm J*. 2019;27(4):574-583. <https://doi.org/10.1016/j.jsps.2019.02.005>
23. Brata C, Fisher C, Marjadi B, Schneider CR, Clifford RM. Factors influencing the current practice of self-medication consultations in Eastern Indonesian community pharmacies: a qualitative study. *BMC Health Serv Res*. 2016;16:179. <https://doi.org/10.1186/s12913-016-1425-3>
24. Thang DX. An investigation of non-prescription medicine supply in community pharmacies in Hanoi, Vietnam. University of Nottingham; 2013.
25. Halila GC, Junior EH, Otuki MF, Correr CJ. The practice of OTC counseling by community pharmacists in Parana, Brazil. *Pharm Pract (Granada)*. 2015;13(4):597. <https://doi.org/10.18549/pharmpract.2015.04.597>
26. Hermansyah A, Sainsbury E, Krass I. Multiple policy approaches in improving community pharmacy practice: the case in Indonesia. *BMC Health Serv Res*. 2018;18(1):449. <https://doi.org/10.1186/s12913-018-3258-8>
27. Da Rocha CE, Bispo ML, Alcantara T, et al. What do Brazilian community pharmacists know about self-medication for minor illnesses? A pilot study in the northeast of Brazil. *J Appl Pharm Sci*. 2014;4(5):12-20. <https://doi.org/10.7324/JAPS.2014.40503>
28. Asosiasi Pendidikan Tinggi Farmasi Indonesia. [Academic manuscripts of graduate competency standards and pharmaceutical education curriculum standards]. <http://www.aptfi.or.id/dokumen/2016-01-01NAKKompetensi&KurAPTFI2013.pdf> (accessed Dec 26, 2020).
29. Fakultas Farmasi Universitas Indonesia. [The UI pharmacist followed UKAI OSCE 2019]. <https://farmasi.ui.ac.id/2019/07/apoteker-ui-mengikuti-ukai-osce-2019/> (accessed Mar 3, 2021).
30. Kristina SA, Gustriawanto N, Rokhman MR, Aditama H, Sari IP. Students' first experience with objective structured clinical examination in a pharmacy school in Indonesia. *J App Pharm Sci*. 2018;8(09):102-106. <https://doi.org/10.7324/JAPS.2018.8915>
31. Resende KA, Cavaco AM, Luna-Leite MD, et al. Training and standardization of simulated patients for multicentre studies in clinical pharmacy education. *Pharm Pract (Granada)*. 2020;18(4):2038. <https://doi.org/10.18549/pharmpract.2020.4.2038>
32. Wigmore BC, Collins JC, Schneider CR, Arias D, Moles RJ. Ability of Pharmacy Students, Pharmacists and Pharmacy Support Staff to Manage Childhood Fever via Simulation. *Am J Pharm Educ*. 2018;82(10):6445. <https://doi.org/10.5688/ajpe6445>

33. Newton GD, Divine H. Patient assessment and consultation. In: Krinsky DL FS, Hemstreet B, Hume AL, et al, editors. *Handbook of Non Prescription Drugs*. 18 ed. Washington: APhA; 2015.
34. Packeiser PB, Castro MS. Evaluation of simulated drug dispensing and patient counseling in the course of pharmaceutical improvement: 2009 to 2015. *Pharm Pract (Granada)*. 2020;18(4):1865. <https://doi.org/10.18549/pharmpract.2020.4.1865>
35. Harden RM. What is a spiral curriculum?. *Med Teach*. 1999;21(2):141-143. <https://doi.org/10.1080/01421599979752>
36. Setiadi AP, Wibowo Y, Brata C, Halim SV, Wardhani SA, Sunderland B. The role of pharmacists in community education to promote responsible self-medication in Indonesia: an application of the spiral educational model. *Int J Clin Pharm*. 2020;42(4):1088-1096. <https://doi.org/10.1007/s11096-020-01055-8>
37. Smithson J, Bellingan M, Glass B, Mills J. Standardized patients in pharmacy education: An integrative literature review. *Curr Pharm Teach Learn*. 2015;7(6):851-863. <https://doi.org/10.1016/j.cptl.2015.08.002>
38. Brata C, Marjadi B, Schneider CR, Murray K, Clifford RM. Information-gathering for self-medication via Eastern Indonesian community pharmacies: a cross-sectional study. *BMC Health Serv Res*. 2015;15:8. <https://doi.org/10.1186/s12913-014-0670-6>
39. Hollnagel E. *The ETTO Principle: efficiency-thoroughness trade-off*. London: Ashgate; 2009. ISBN 9780754676782.
40. Puspitasari HP, Faturrohman A, Hermansyah A. Do Indonesian community pharmacy workers respond to antibiotics requests appropriately?. *Trop Med Int Health*. 2011;16(7):840-846. <https://doi.org/10.1111/j.1365-3156.2011.02782.x>
41. Zawahir S, Lekamwasam S, Aslani P. Antibiotic dispensing practice in community pharmacies: A simulated client study. *Res Social Adm Pharm*. 2019;15(5):584-590. <https://doi.org/10.1016/j.sapharm.2018.07.019>
42. Amin M, Chewing B. Pharmacies without pharmacists: Absenteeism plagues pharmacies in developing countries. *Res Soc Adm Pharm*. 2017;14:400-401. <http://doi.org/10.1016/j.sapharm.2016.10.013>

Pharmacy Practice

HOME ABOUT ▾ CURRENT ARCHIVES AUTHORS FOR REVIEWERS


Box No.1 Content

Box No.2 Content

About the Journal

Pharmacy Practice is a full-text peer-reviewed journal with a scope on pharmacy practice. **Pharmacy Practice** is published quarterly.

[Read More](#)



Pharmacy Practice

[Current](#) [Archives](#) [Authors](#) [For reviewers](#) [About ▾](#)

[Home](#) / [About the Journal](#)

About Pharmacy Practice

Pharmacy Practice is a free available, open access, peer-reviewed journal with a scope on pharmacy practice. **Pharmacy Practice does not charge and will never charge any publication fee or article processing charge (APC) to the authors.** All the articles are free-access without delay (no embargo period). Pharmacy Practice is published using a continuous flow system with four issues quarterly closed. The current and future absence of any APCs is signed in the memorandum of understanding with the Center for Pharmacy Practice Innovation (CPPi) at the Virginia Commonwealth University (VCU) School of Pharmacy. **Pharmacy Practice** is the consequence of the efforts of a number of colleagues from different universities who believe in [collaborative publishing](#): nobody pays, nobody gets paid.

Although focusing on the practice of pharmacy, **Pharmacy Practice** covers a wide range of pharmacy activities, such as, pharmaceutical care, social pharmacy, pharmacy education, process and outcome research, clinical pharmacy, health promotion and education, health informatics, and pharmacoepidemiology, as well as many others.

Having an international scope is a major goal for **Pharmacy Practice**. See the [list of authors](#) since 2007.

Pharmacy Practice is being indexed and abstracted at [PubMed](#), [PubMed Central](#), Embase, [Scopus](#), Ebsco EJS, [Directory of Open Access Journals](#), Web of Science (ESCI), [Dimensions](#), Open J-gate, [ERIH Plus](#), [Scielo](#), [Dialnet](#), IBECS, [REDIB](#), [Latindex](#), [Redalyc](#).

For more information on **Pharmacy Practice** publication delay and peer-review process, visit our [reviewers](#) web page and the editorial [Pharm Pract \(Granada\) 2019;17\(1\):1502](#). For information about requirements for manuscripts, visit our [authors guidelines](#) web page.

Pharmacy Practice is a peer-reviewed journal. To make the peer review process work properly, it is necessary to count with a wide pool of prestigious peer reviewers, experts in different areas in journal's scope. **Pharmacy Practice** has created a peer reviewer selection process based on searching in PubMed for articles closely related to each manuscript awaiting evaluation (see a description of the peer [reviewer selection process](#)). We aim to ensure the expertise of each reviewer in the specific topic. Working as a peer-reviewer in **Pharmacy Practice** has no remuneration. See the [list of peer reviewers](#) who have collaborated in **Pharmacy Practice**.

Pharmacy Practice is funded through the corporate social responsibility funds of [Centro de Investigaciones y Publicaciones Farmaceuticas](#) and scientifically partnered with the [Center for Pharmacy Practice Innovation](#) at VCU.

Editorial & Advisory Boards

Pharmacy Practice Editorial Board

Fernando Fernandez-Llimos (editor)
University of Porto (Portugal)

David A. Holdford (associate editor)
Virginia Commonwealth University (USA)

Teresa M. Salgado (associate editor)
Virginia Commonwealth University (USA)

Shalom I. Benrimoj
University of Granada (Spain)

Karen B. Farris
University of Michigan (USA)

Martin Schulz
Freie Universitaet Berlin and ABDA – Federal Union of German Associations of Pharmacists (Germany)

Cassyano J. Correr
Federal University of Parana (Brazil)

Fernanda S. Tonin
Federal University of Parana (Brazil)

Joanna C. Moullin
Curtin University (Australia)

Jonathan Penm
University of Sydney (Australia)

Antoinette (Toni) B. Coe
University of Michigan (USA)

Meagen Rosenthal
University of Mississippi (USA)

Carrie M. Blanchard
University of North Carolina (USA)

Reidun L. Kjome
University of Bergen (Norway)

Sarah Dineen-Griffin
University of Technology Sydney (Australia)

Advisory Board


Claire Anderson, University of Nottingham (UK)
Marja Airaksinen, University of Helsinki (Finland)
Simon Bell, University of South Australia (Australia)
John Bucheit, Virginia Commonwealth University (USA)
Maria Cordina, University of Malta (Malta)
Lisa Guirguis, University of Alberta (Canada)
Souheil Hallit, Holy Spirit University of Kaslik (Lebanon)
Mohamed A. Hassali, University Sains Malaysia (Malaysia)
James Hoehns, University of Iowa (USA)
Carmel Hughes, Queen's University Belfast (UK)
Ulrich Jaehde, University of Bonn (Germany)
Rosemin Kassam, University of British Columbia (Canada)
David Nau, Pharmacy Quality Solutions (USA)
Lauren Pamulapati, Virginia Commonwealth University (USA)
Julie A. Patterson, Virginia Commonwealth University (USA)
Kimberly S. Plake, Purdue University (USA)
Bandana Saini, University of Sydney (Australia)
Anthony Serracino-Inglott, University of Malta (Malta)
Olayinka Shiyabola, University of Wisconsin-Madison (USA)
Mary Tully, University of Manchester (UK)

[Home](https://www.pharmacypractice.org/index.php/pp/index) (https://www.pharmacypractice.org/index.php/pp/index) / [Archives](https://www.pharmacypractice.org/index.php/pp/issue/archive) (https://www.pharmacypractice.org/index.php/pp/issue/archive)
/ Vol. 19 No. 2 (2021): Apr-Jun




ORIGINAL RESEARCH


[Feasibility of cardiovascular risk screening in Portuguese community pharmacies](https://www.pharmacypractice.org/index.php/pp/article/view/2255) (https://www.pharmacypractice.org/index.php/pp/article/view/2255)

Anabela A. Fonseca, Tacio M. Lima, Margarida Castel-Branco, Isabel V. Figueiredo
Abstract  1157 | pdf Downloads  555
 Page 2255





 [pdf](https://www.pharmacypractice.org/index.php/pp/article/view/2255/914) (https://www.pharmacypractice.org/index.php/pp/article/view/2255/914)


[An ounce of prevention is worth a pound of cure: considerations for pharmacists delivering the National Diabetes Prevention Program](https://www.pharmacypractice.org/index.php/pp/article/view/2426) (https://www.pharmacypractice.org/index.php/pp/article/view/2426)

Dave L. Dixon, Evan M. Sisson, Lauren G. Pamulapati, Rowan Spence, Teresa M. Salgado
Abstract  685 | pdf Downloads  277
 Page 2426




 [pdf](https://www.pharmacypractice.org/index.php/pp/article/view/2426/912) (https://www.pharmacypractice.org/index.php/pp/article/view/2426/912)


[The competency of Indonesian pharmacy students in handling a self-medication request for a cough: a simulated patient study](https://www.pharmacypractice.org/index.php/pp/article/view/2269) (https://www.pharmacypractice.org/index.php/pp/article/view/2269)

Cecilia Brata, Steven V. Halim, Eko Setiawan, Bobby Presley, Yosi I. Wibowo, Carl R. Schneider
Abstract  704 | pdf Downloads  407 online appendix Downloads  0
 Page 2269




 [pdf](https://www.pharmacypractice.org/index.php/pp/article/view/2269/907) (https://www.pharmacypractice.org/index.php/pp/article/view/2269/907)


[Improving outpatient medication counselling in hospital pharmacy settings: a behavioral analysis using the theoretical domains framework and behavior change wheel](https://www.pharmacypractice.org/index.php/pp/article/view/2271) (https://www.pharmacypractice.org/index.php/pp/article/view/2271)

Samirah N. Abdu-Aguye, Shafiu Mohammed, Nuhu M. Danjuma, Kamilu S. Labaran
Abstract  575 | pdf Downloads  340
 Page 2271




 [pdf](https://www.pharmacypractice.org/index.php/pp/article/view/2271/913) (https://www.pharmacypractice.org/index.php/pp/article/view/2271/913)


[Assessment of community pharmacists' communication and comfort levels when interacting with Deaf and hard of hearing patients](https://www.pharmacypractice.org/index.php/pp/article/view/2274) (https://www.pharmacypractice.org/index.php/pp/article/view/2274)

Elizabeth Y. Chong, Sabrina A. Jacob, Amutha Ramadas, Pei H. Goh, Uma D. Palanisamy
Abstract  911 | pdf Downloads  368
 Page 2274

 [pdf](https://www.pharmacypractice.org/index.php/pp/article/view/2274/911) (https://www.pharmacypractice.org/index.php/pp/article/view/2274/911)


[The role of primary care pharmacists in the response to the Zika epidemic](https://www.pharmacypractice.org/index.php/pp/article/view/2290) (https://www.pharmacypractice.org/index.php/pp/article/view/2290)

Cláudia B. Santos-Pinto, Claudia S. Osorio-de-Castro, Larissa M. Ferreira, Elaine S. Miranda
Abstract  476 | pdf Downloads  240
 Page 2290


 [pdf](https://www.pharmacypractice.org/index.php/pp/article/view/2290/924) (https://www.pharmacypractice.org/index.php/pp/article/view/2290/924)

[The Pharmacists' and pharmacy technicians' scopes of practice in the management of minor ailments at community pharmacies in Indonesia: a cross-sectional study](https://www.pharmacypractice.org/index.php/pp/article/view/2295) (https://www.pharmacypractice.org/index.php/pp/article/view/2295)

Vinci Mizranita, Tin F. Sim, Bruce Sunderland, Richard Parsons, Jeffery D. Hughes
Abstract  802 | pdf Downloads  314
 Page 2295



 [pdf](https://www.pharmacypractice.org/index.php/pp/article/view/2295/916) (https://www.pharmacypractice.org/index.php/pp/article/view/2295/916)


[The association between social support and medication adherence in patients with](https://www.pharmacypractice.org/index.php/pp/article/view/2300)

 [pdf](https://www.pharmacypractice.org/index.php/pp/article/view/2300/928) (https://www.pharmacypractice.org/index.php/pp/article/view/2300/928)

[hypertension: A systematic review](https://www.pharmacypractice.org/index.php/pp/article/view/2300) (https://www.pharmacypractice.org/index.php/pp/article/view/2300)



Wejdan Shahin, Gerard A. Kennedy, Ieva Stupans


Abstract  765 | pdf Downloads  309


 Page 2300

[Measuring depression and anxiety prevalence among Iraqi healthcare college students using hospital anxiety and depression scale](https://www.pharmacypractice.org/index.php/pp/article/view/2303) (https://www.pharmacypractice.org/index.php/pp/article/view/2303)

Sarmed H. Kathem, Ali A. Al Jumaili, Malak Noor-Aldeen , Noor Najah , Dema Ali Khalid



Abstract  651 | pdf Downloads  336


 Page 2303


 pdf (https://www.pharmacypractice.org/index.php/pp/article/view/2303/910)

[Public knowledge, beliefs, psychological responses, and behavioural changes during the outbreak of COVID-19 in the Middle East](https://www.pharmacypractice.org/index.php/pp/article/view/2306) (https://www.pharmacypractice.org/index.php/pp/article/view/2306)

Husam Abazid, Iman A. Basheti, Esraa E. Al-Jomaa, Ayham Abazid, Warda M. Kloub



Abstract  920 | pdf Downloads  329


 Page 2306


 pdf (https://www.pharmacypractice.org/index.php/pp/article/view/2306/915)

[Medication adherence and persistence of psoriatic arthritis patients treated with biological therapy in a specialty pharmacy in Brazil: a prospective observational study](https://www.pharmacypractice.org/index.php/pp/article/view/2312) (https://www.pharmacypractice.org/index.php/pp/article/view/2312)

Ana F. Souza, Michael R. da Silva, Jéssica B. dos Santos, Alessandra M. Almeida, Francisco A. Acurcio, Juliana Alvares-Teodoro




Abstract  730 | pdf Downloads  291


 Page 2312


 pdf (https://www.pharmacypractice.org/index.php/pp/article/view/2312/917)

[Descriptive assessment of graduates' perceptions of pharmacy-related competencies based on the Lebanese pharmacy core competencies framework](https://www.pharmacypractice.org/index.php/pp/article/view/2320) (https://www.pharmacypractice.org/index.php/pp/article/view/2320)

Rony M. Zeenny, Marwan Akel, Aline Hajj, Hala Sacre, Souheil Hallit, Pascale Salameh



Abstract  597 | pdf Downloads  208 online appendix Downloads  0


 Page 2320


 pdf (https://www.pharmacypractice.org/index.php/pp/article/view/2320/930)

[Why do pharmacists leave the profession? A mixed-method exploratory study](https://www.pharmacypractice.org/index.php/pp/article/view/2332) (https://www.pharmacypractice.org/index.php/pp/article/view/2332)

Trudi J. Aspden, Pushkar R. Silwal, Munyaradzi Marowa, Rhys Ponton





Abstract  677 | pdf Downloads  289


 Page 2332


 pdf (https://www.pharmacypractice.org/index.php/pp/article/view/2332/921)

[The Impact of Covid-19 containment on community pharmaceutical spending in Andalusia - Spain](https://www.pharmacypractice.org/index.php/pp/article/view/2346) (https://www.pharmacypractice.org/index.php/pp/article/view/2346)

Francisco Florido Alba, Nuria García-Agua, Angel Martín Reyes, Antonio Clavero Barranquero , Antonio J. Garcia Ruiz



Abstract  363 | pdf Downloads  200 online appendix 1 Downloads  0 online appendix 2 Downloads  0


 Page 2346


 pdf (https://www.pharmacypractice.org/index.php/pp/article/view/2346/925)

[Patient satisfaction with medication therapy adherence clinic services in a district hospital: a cross-sectional study](https://www.pharmacypractice.org/index.php/pp/article/view/2353) (https://www.pharmacypractice.org/index.php/pp/article/view/2353)


Yi C. Sim, Intan S. Mohd-Rosli, Boon T. Lau, Siew Y. Ng

Abstract  594 | pdf Downloads  346

 Page 2353




 pdf (https://www.pharmacypractice.org/index.php/pp/article/view/2353/920)

[Bridging health disparities: a national survey of ambulatory care pharmacists in](https://www.pharmacypractice.org/index.php/pp/article/view/2359)






 pdf (https://www.pharmacypractice.org/index.php/pp/article/view/2359/918)

underserved areas (<https://www.pharmacypractice.org/index.php/pp/article/view/2359>)





Morgan P. Stewart, Rhianna Fink, Emily Kosirog, Joseph J. Saseen

Abstract  420 | pdf Downloads  187 Page 2359**Patterns of medication errors involving pediatric population reported to the French Medication Error Guichet** (<https://www.pharmacypractice.org/index.php/pp/article/view/2360>)





Christine Azar, Delphine Allué, Marie B. Valnet-Rabier, Laurent Chouchana, Fanny Rocher, Dorothée Durand, Nathalie Grené-Lerouge, Nadine Saleh, Patrick Maison

Abstract  657 | pdf Downloads  282 online appendix Downloads  0 Page 2360 pdf (<https://www.pharmacypractice.org/index.php/pp/article/view/2360/922>)**INTERNATIONAL SERIES****Primary health care policy and vision for community pharmacy and pharmacists in Estonia** (<https://www.pharmacypractice.org/index.php/pp/article/view/2404>)




Kristiina Sepp, Anita Tuula, Veera Bobrova, Daisy Volmer

Abstract  1539 | pdf Downloads  306 Page 2404 pdf (<https://www.pharmacypractice.org/index.php/pp/article/view/2404/909>)**Community pharmacies and pharmacists in Brazil: A missed opportunity** (<https://www.pharmacypractice.org/index.php/pp/article/view/2467>)

Angelita C. Melo, Guilherme M. Trindade, Alessandra R. Freitas, Karina A. Resende, Tarcísio J. Palhano

Abstract  1168 | pdf Downloads  455 Page 2467 pdf (<https://www.pharmacypractice.org/index.php/pp/article/view/2467/929>)**REVIEW****Perspectives on the pharmacist's "product": a narrative review** (<https://www.pharmacypractice.org/index.php/pp/article/view/2430>)

David A. Holdford

Abstract  518 | pdf Downloads  303 Page 2430 pdf (<https://www.pharmacypractice.org/index.php/pp/article/view/2430/919>)Make a Submission (<https://www.pharmacypractice.org/index.php/pp/about/submissions>)**Browse**

Categories

- » [Pharm Pract \(Granada\) editorial policy](https://www.pharmacypractice.org/index.php/pp/catalog/category/EditPolicy) (<https://www.pharmacypractice.org/index.php/pp/catalog/category/EditPolicy>)
- » [Integration of community pharmacy in primary health care](https://www.pharmacypractice.org/index.php/pp/catalog/category/integration) (<https://www.pharmacypractice.org/index.php/pp/catalog/category/integration>)
- » [Simulated Patient Technique](https://www.pharmacypractice.org/index.php/pp/catalog/category/SimPatTech) (<https://www.pharmacypractice.org/index.php/pp/catalog/category/SimPatTech>)
- » [Medication Adherence](https://www.pharmacypractice.org/index.php/pp/catalog/category/medadh) (<https://www.pharmacypractice.org/index.php/pp/catalog/category/medadh>)

Pharmacy Practice | e-ISSN: 1886-3655 - p-ISSN: 1885-642X - CODEN: PPHRGO

Privacy Policy (<https://www.pharmacypractice.org/journal/index.php/pp/authors#a4>)

eISSN: 1886-3655 (Electronic)

ISSN: 1885-642X (Print)

OJS Hosting, Support, and Customization by: OpenJournalSystems.com (<https://openjournalssystems.com/ojs-hosting>)

Pharmacy Practice

COUNTRY	SUBJECT AREA AND CATEGORY	PUBLISHER	H-INDEX
Spain 	Health Professions Pharmacy Pharmacology, Toxicology and Pharmaceutics Pharmaceutical Science	Centro de Investigaciones y Publicaciones Farmaceuticas	22
PUBLICATION TYPE	ISSN	COVERAGE	INFORMATION
Journals	1885642X, 18863655	2006-2020	Homepage How to publish in this journal journal@pharmacypractice.org

[✕](#)

Call For Papers Elsevier Q4

Peer Reviewed Indexed Journal


Collaboration proposals are invited to provide a single platform for worldwide research.

turcomat.org

OPEN

SCOPE

Pharmacy Practice is a free full-text peer-reviewed journal with a scope on pharmacy practice. Pharmacy Practice is published quarterly. Pharmacy Practice does not charge and will never charge any publication fee or article processing charge (APC) to the authors. The current and future absence of any article processing charges (APCs) is signed in the MoU with the Center for Pharmacy Practice Innovation (CPPi) at Virginia Commonwealth University (VCU) School of Pharmacy. Pharmacy Practice is the consequence of the efforts of a number of colleagues from different Universities who believe in collaborative publishing: no one pays, no one receives. Although focusing on the practice of pharmacy, Pharmacy Practice covers a wide range of pharmacy activities, among them and not being comprehensive, clinical pharmacy, pharmaceutical care, social pharmacy, pharmacy education, process and outcome research, health promotion and education, health informatics, pharmacoepidemiology, etc.

 Join the conversation about this journal

[✕](#)

Impact Factor and Indexed

JNLS is a peer-reviewed journal publishing original research, review articles, and more

thieme.com

OPEN



Source details

Pharmacy Practice

Open Access ⓘ

Scopus coverage years: from 2006 to Present

Publisher: Centro de Investigaciones y Publicaciones Farmaceuticas

ISSN: 1885-642X E-ISSN: 1886-3655

Subject area: Health Professions: Pharmacy Pharmacology, Toxicology and Pharmaceutics: Pharmaceutical Science

Source type: Journal

CiteScore 2021

2.8 ⓘ

SJR 2021

0.409 ⓘ

SNIP 2021

0.994 ⓘ

[View all documents >](#)

[Set document alert](#)

[Save to source list](#)

[CiteScore](#) [CiteScore rank & trend](#) [Scopus content coverage](#)

CiteScore 2021 ⌵

$$2.8 = \frac{733 \text{ Citations } 2018 - 2021}{259 \text{ Documents } 2018 - 2021}$$

Calculated on 05 May, 2022

CiteScoreTracker 2022 ⓘ

$$3.0 = \frac{749 \text{ Citations to date}}{251 \text{ Documents to date}}$$

Last updated on 05 August, 2022 • Updated monthly

CiteScore rank 2021 ⓘ

Category	Rank	Percentile
Health Professions		
└ Pharmacy	#8/36	79th
Pharmacology, Toxicology and Pharmaceutics		
└ Pharmaceutical Science	#84/171	51st

[View CiteScore methodology >](#) [CiteScore FAQ >](#) [Add CiteScore to your site ↗](#)

About Scopus

[What is Scopus](#)

[Content coverage](#)

[Scopus blog](#)

[Scopus API](#)

[Privacy matters](#)

Language

[日本語版を表示する](#)

[查看简体中文版本](#)

[查看繁體中文版本](#)

[Просмотр версии на русском языке](#)

Customer Service

[Help](#)

[Tutorials](#)

[Contact us](#)

ELSEVIER

[Terms and conditions](#) ↗ [Privacy policy](#) ↗

Copyright © [Elsevier B.V](#) ↗. All rights reserved. Scopus® is a registered trademark of Elsevier B.V.

We use cookies to help provide and enhance our service and tailor content. By continuing, you agree to the [use of cookies](#) ↗.

 RELX