



Comparison of vitamin D status and physical activity related to obesity among tertiary education students

Amelia Lorensia^{1*}, Rivan Virlando Suryadinata², Irene Andriana Inu¹

¹Departement of Community Clinical Pharmaceutical, Faculty of Pharmacy, University of Surabaya (Universitas Surabaya). Jl. Raya Kalirungkut, Rungkut District, Surabaya City, East Java 60293, Indonesia.

²Departement of Public Health Science, Faculty of Medicine, University of Surabaya (Universitas Surabaya). Jl. Raya Kalirungkut, Rungkut District, Surabaya City, East Java 60293, Indonesia.

ARTICLE INFO

Received on: 30/09/2021
Accepted on: 27/12/2021
Available Online: 05/04/2022

Key words:

Physical activity, obesity, student, vitamin D status.

ABSTRACT

University students are young adults who are productive and have relatively high physical activity. Obesity is caused by imbalance between excessive food intake and low physical activity and is a risk factor for vitamin D deficiency. This research aimed to explore vitamin D status and physical activity in the obese and nonobese groups and the relationship between vitamin D status and physical activity. The research design used was cross-sectional using the International Physical Activity Questionnaire and vitamin D status questionnaires. The research location was the campus area of Surabaya and was carried out from March 2018 to January 2019. This study involved 143 respondents consisting of 71 people (49.65%) in the obese group and 72 people (50.34%) in the nonobese group. The results showed that there was a significant difference in physical activity ($p = 0.047$) and vitamin D status ($p < 0.05$) between the nonobese and obese groups. There was a significant relationship ($p < 0.05$) between vitamin D status and physical activity, although the relationship was low (correlation coefficient = 0.326). Therefore, increased physical activity can improve vitamin D status while still paying attention to other factors that affect vitamin D, such as diet and lifestyle. The condition of vitamin D also needs to be confirmed by measuring the 25(OH)D blood test.

INTRODUCTION

The World Health Organization (WHO) declared obesity a global epidemic (Jackson *et al.*, 2020; Jayawardena *et al.*, 2020). Overweight or obesity is a serious public health problem in Indonesia with a continuing increase in its prevalence (Rachmi *et al.*, 2017). Based on the results of the Basic Health Research in Indonesia (2018), it is stated that the prevalence of obesity in adults has increased since the three Riskesdas periods, i.e., by 10.5% (2007), 14.8% (2013), and 21.8% (2018) (Damayanti *et al.*, 2019). Obesity is caused by various factors including genetic, environmental, psychological, and health factors caused by the use of certain drugs and diseases, as well as factors in the development of adipose tissue (Fruh *et al.*, 2017; Panuganti *et al.*, 2021). However, the main cause of obesity is an imbalance between excessive food intake and low physical activity (Jehan *et al.*, 2020;

Romieu *et al.*, 2017). Prevention of obesity can be achieved by changing lifestyle (Salam *et al.*, 2020), such as increasing physical activity by exercising diligently and maintaining a healthy diet (Elliot and Hamlin, 2018; Koehler and Drenowatz, 2019).

High physical activity can be useful in the process of preventing obesity (Jakicic *et al.*, 2019; Kim *et al.*, 2017; Lorensia *et al.*, 2021); on the other hand, low physical activity results in an increased prevalence of the risk of noncommunicable diseases such as heart disease, diabetes, and cancer (Ahmed *et al.*, 2019; Saqib *et al.*, 2020). Physical activity is defined as movement of the body by the muscles and skeleton which requires energy expenditure. There are two categories, namely moderate (moderate) and vigorous (strong) physical activity (Füzéki and Banzer, 2018; Hamer and Stamatakis, 2018). Both these types of physical activity provide health benefits and also affect vitamin D status (Skalska *et al.*, 2019; Wiciński *et al.*, 2019). A person with vigorous and moderate activity has a higher vitamin D status than a person with low physical activity (Hall *et al.*, 2018; Orcees, 2019; Silva *et al.*, 2019). Vitamin D status is also seen by dietary intake of foods containing vitamin D and daily lifestyle habits

*Corresponding Author

Amelia Lorensia, Universitas Surabaya, Surabaya, Indonesia.
E-mail: amelia.lorensia@gmail.com

(Larson-Meyer *et al.*, 2019; Suryadinata *et al.*, 2020; Zadka *et al.*, 2018), such as the use of sunscreen, low intake of milk (Aji *et al.*, 2019), and avoiding sun exposure which causes a decrease in the synthesis of vitamin D by the skin resulting in vitamin D deficiency (Ramasamy, 2020; Suryadinata *et al.*, 2020; Webb *et al.*, 2018).

More than one billion people worldwide are deficient in vitamin D (Sizar *et al.*, 2021). About 30%–50% of children and adults in the United Arab Emirates, Australia, Turkey, India, and Lebanon are deficient in vitamin D (Zarooni *et al.*, 2019). The risk of vitamin D deficiency can occur in dark-skinned people, elderly patients, pregnant women, babies, people who do not get enough sun exposure, and obese individuals (Sizar *et al.*, 2021; Suryadinata *et al.*, 2021; Wacker and Holick, 2013). The body can get vitamin D by consuming milk, fish and fish oil, egg yolks, liver, and yeast (Maurya and Aggarwal, 2017; Polzonetti *et al.*, 2020), exposure to sunlight, and taking vitamin D supplements (Kift *et al.*, 2018; Lee *et al.*, 2020). Vitamin D functions in the formation and maintenance of bones, so vitamin D deficiency results in bone abnormalities, namely rickets in children and osteomalacia in adults (Michigami, 2018; Takeuchi, 2018; Umar *et al.*, 2018). Vitamin D deficiency also contributes to the development of cancer, especially breast, prostate, and colon cancer as well as heart disease, stroke, and autoimmune diseases (Charoenngam and Holick, 2020; Hassanalilou *et al.*, 2017; Wang *et al.*, 2017).

Early identification of vitamin D deficiency can be used as a preventive measure related to the risk of vitamin D deficiency (Aji *et al.*, 2019; Pilz *et al.*, 2019; Sizar *et al.*, 2021). The indicators used to determine the status of vitamin D include blood tests, dietary food intake containing vitamin D (Aji *et al.*, 2019; Lorensia *et al.*, 2020a; Pilz *et al.*, 2019), and questionnaires (Bolek-Berquist *et al.*, 2009; Larson-Meyer *et al.*, 2019). Evaluation of vitamin D status using a questionnaire is easier to carry out and more economical than blood serum examination. In addition, when compared to blood tests, clinical questionnaires can be used to identify long-term vitamin D status (Aji *et al.*, 2019; Annweiler *et al.*, 2017; Bolek-Berquist *et al.*, 2009). The questionnaire can be used to identify patients who are at high or low risk of experiencing vitamin D deficiency (Bolek-Berquist *et al.*, 2009). Obesity as a risk factor for vitamin D deficiency (Paschou *et al.*, 2019) needs to get prevention efforts as early as possible considering the dangers of vitamin D deficiency and low physical activity as causes of obesity which affect the development of chronic diseases and decreased quality of life (Kavadar *et al.*, 2015; Kim *et al.*, 2018; Oh *et al.*, 2017).

This study involved young adult respondents because adults were of productive age and relatively had high physical activity. However, nowadays, there are many young adults who actually experience vitamin D deficiency and rarely do physical activity (Lorensia *et al.*, 2019). This research was conducted on university students because education can affect a person's physical activity and lifestyle (Kotarska *et al.*, 2021). It was, therefore, necessary to identify vitamin D status and physical activity in obese and nonobese adults so that pharmacists can provide education to the public, both obese and nonobese adults, as a form of health workers' responsibility in early prevention efforts related to the risk of deficiency of vitamin D and other health problems.

METHODS

Research design

The design of this study was cross-sectional and was conducted on a group of students with nonobese and obese nutritional status in the campus area of the University of Surabaya. The research location chosen to conduct the research was the campus area in South Surabaya, Rungkut district, East Java, Indonesia, and the study was carried out from March 2018 to January 2019. This study was approved with Ethical Test No. 034/KE/I/2018 at Universitas Surabaya.

Research variable

The independent variable in this study was a group of obese and nonobese respondents. The dependent variables in this study were vitamin D status and physical activity. Vitamin D status can be defined as the presence of vitamin D in a certain amount in the body which was influenced by the intake of foods containing vitamin D, sun exposure, and consumption of vitamin D supplements. A person was said to be positive (+) (at risk) of vitamin D deficiency if the total answer score was ≥ 15 and negative (–) if the total answer score was < 15 . Vitamin D deficiency is a health problem related to vitamin D status. The risk of vitamin D deficiency in nonobese and obese respondents is calculated using the prevalence odds ratio equation.

Physical activity was defined as body movement by the muscles and skeleton which requires energy expenditure, including activities carried out while working, playing, doing household chores, traveling, and recreation. Physical activity in adults can be measured using the International Physical Activity Questionnaire (IPAQ) (Cleland *et al.*, 2018; Kavadar *et al.*, 2015; Sember *et al.*, 2020). This can be measured by asking seven questions related to daily activities. The data scale obtained from the measurement of physical activity was an ordinal data scale, where the results obtained from the patient questionnaire were categorized into low, medium, and high physical activity (Table 1).

Obesity was characterized by the accumulation of excess body fat tissue (De Lorenzo *et al.*, 2019). Determination of obesity status in individuals was based on the measurement of body mass index (BMI), a person's body weight (in kg) divided by the square of height (in m). The International Association for Study of Obesity Organization, International Obesity Task Force, and WHO determined the category of obesity for the Asian population if the BMI was ≥ 25 kg/m². Therefore, in this study, BMI points ≥ 25 kg/m² were used to determine the obesity category.

Population and sample

The population used in this study was all students at the University of Surabaya with a nutritional status of obesity or nonobesity. The sample used in this study was active students at the University of Surabaya with a nutritional status conditions of obesity or nonobesity who met the criteria, including 18–25-year-olds, those with no special diet (such as fasting or vegetarianism), those who did not have certain diseases like cardiovascular disorders such as angina, renal, and hepatic disorders (such as cirrhosis of the liver), and individuals who have no motoric problems. The sampling technique used was the purposive

Table 1. Guidelines for assessment of physical activity based on the IPAQ.

Type of physical activity	Description
Low physical activity	No reports of physical activity carried out Activities carried out not sufficient to meet the criteria for the type of moderate physical activity
Moderate physical activity (one of the three criteria)	Within 3 days or more doing high-intensity (vigorous) activity for 20 minutes per day Within 5 days or more doing moderate-intensity activity (moderate) or walking for 30 minutes Within 5 days or more doing a combination of walking, moderate-intensity activity, and vigorous activity of at least 600 MET-minute per week
High physical activity (one of the three criteria)	Performing vigorous activity for 3 days with a total MET accumulation of at least 1,500 MET-minute per week For 7 days or more doing a combination of walking and moderate-intensity or vigorous activity at a minimum of 3,000 MET-minute per week

Description: MET (multiples of the resting metabolic rate): the amount of energy released that is equivalent to 1 kcal/kg/hour (Cleland *et al.*, 2018; Kavadar *et al.*, 2015; Sember *et al.*, 2020).

sampling method, in which the sampling process was based on previously known characteristics or characteristics of the population and the consideration of the researcher.

Data collection methods and analysis

Measuring instruments used in the study were the vitamin D status questionnaire to identify vitamin D status and the IPAQ for physical activity data. Measurements of weight and height were carried out using a digital weight scale and a Microtoise stature meter, respectively. There were two types of questionnaires given, namely the vitamin D status questionnaire (Bolek-Berquist *et al.*, 2009; Cairncross *et al.*, 2017; Nabak *et al.*, 2014) and the IPAQ (Cleland *et al.*, 2018; Kavadar *et al.*, 2015; Sember *et al.*, 2020).

We conducted a trial or pilot study where data collection was preceded by testing the research instruments on 30 respondents. The questionnaire was validated using internal and external validity methods. Internal validity was enforced in a review step based on professional opinion judgments in the field of community pharmacy, while external validation was carried out by testing the questionnaire on the subject. Then, the results were analyzed using the Statistical Package for the Social Sciences (SPSS) version 24.0 computer program. The questionnaire was said to be valid if the value of $r_{\text{count}} > 0.361$. The questionnaire was rehabilitated using a single trial method technique with the alpha method (Cronbach), namely testing the reliability of the questionnaire once on respondents who have criteria such as research subjects, but these subjects are not used as samples. A reliability test can be carried out on valid questions. The calculation of the reliability coefficient uses the SPSS version 24.0 software. The questionnaire was declared reliable if the Cronbach's alpha value was >0.70 .

Comparative analysis of the risk of vitamin D deficiency was based on vitamin D status in the obese and nonobese groups of respondents using the prevalence odds ratio. Meanwhile, differences in vitamin D status and physical activity in the obese and nonobese respondent groups used the chi-square test. We then proceeded to test the relationship between vitamin D status and

physical activity in the groups of obese and nonobese respondents using a contingency coefficient.

RESULTS

The data obtained in this study were obtained through a vitamin D status questionnaire and the IPAQ. The data collection process uses instruments that are valid and reliable. Based on the data obtained, there were 143 respondents consisting of 2 groups, namely the obese group consisting of 71 people (49.65%) and the nonobese group consisting of 72 people (50.34%).

Validity and reliability of the IPAQ

The original IPAQ was available in English. The validation process was carried out by translating the questionnaire into Indonesian and then giving it to three professional judges in the field of community pharmacy. The validity was enforced in the analysis step based on the opinion of professional judges in the field of community pharmacy.

Validity and reliability of the vitamin D status questionnaire

The reliability of the questionnaire was determined based on Cronbach's alpha value. The questionnaire was declared reliable if Cronbach's alpha value was >0.60 . The Cronbach's alpha value of the vitamin D status questionnaire was 0.735, so the questionnaire was declared reliable. The questionnaire validation process used the SPSS version 24.0 program. The validation process was carried out on 30 respondents. Based on the Pearson product-moment r_{table} with a significance of 0.05 with a two-sided test and the amount of data (n) = 30, it was known that r_{table} value was 0.361 so that $r_{\text{count}} > 0.361$, and it can be concluded that the questionnaire was valid.

Respondent characteristics

In this study, the respondents included were students with a stratum 1 education level with an age range of 18–25 years. Table 2 explains the distribution of respondents' age, sex, and BMI.

The results of the answers to the two groups of respondents were classified into sections regarding sun exposure

Table 2. Characteristics of respondents.

Characteristics		Nonobese group (<i>n</i> = 72)		Obese group (<i>n</i> = 71)	
		Frequency	Percentage (%)	Frequency	Percentage (%)
Gender	Male	19	26.38	38	53.52
	Female	53	73.61	33	46.47
Age (years)	18	5	7.04	6	8.33
	19	14	19.72	7	0.72
	20	17	23.94	22	30.56
	21	20	28.17	14	19.44
	22	11	15.49	15	20.83
	23	4	5.63	4	5.56
	24			3	4.17
	25			1	1.39
BMI (kg/m ²)	Thin (<18.50)	12	16.66		
	Normal (18.50–22.90)	40	55.55		
	At risk (23.00–24.90)	20	27.77		
	Obese type I (25.00–29.90)			42	59.15
	Obese type II (≥30.00)			29	40.84
History of disease ^a	Respiratory tract disorders	3	4.16	6	8.45
	Digestive tract disorders	13	18.05	9	12.67
	Blood disease	5	6.94	3	4.22
	Infectious disease	7	9.72	5	7.04
	Others	9	12.50	4	5.63
	Does not have a history of disease	45	62.50	48	67.60
	Multivitamins	7	9.72	6	8.45
	Antibiotic	3	4.16	0	0.00
History of medicine ^a	Immune system booster	3	4.16	2	2.81
	Digestive tract disorders drugs	1	1.38	2	2.81
	Others	9	12.50	9	12.67
	Does not use drugs or supplements	52	72.22	52	73.23

^a The answer can be more than 1 in the category.

(Table 3), foods containing vitamin D (Table 4), and signs of vitamin D deficiency (Table 5).

The nonobese group had a more positive vitamin D status (≥ 15) (76.39%) than the obese group. At the same time, the level of activity in the nonobese group was dominated by moderate (55.55%) compared to the obese group, which was dominated by low activity levels (61.97%) (Table 6).

Based on the results of the analysis with the chi-square test to see the difference in vitamin D status between the nonobese and obese respondent groups, it was found that there was a significant difference (p -value < 0.05) in the vitamin D status of the nonobese and obese groups of respondents (Table 7).

Based on the results of the analysis with the chi-square test to see the difference in physical activity between the groups of nonobese and obese respondents, it was shown that there was significant difference between the two (p -value < 0.05) (Table 8).

Based on the contingency coefficient correlation test for vitamin D status and physical activity in the nonobese and obese respondent groups, the approximate significance value of 0.000 ($p = 0.000$) indicated that there was a significant relationship between

vitamin D status and physical activity and the relationship has a correlation value of 0.326, so it can be concluded that there is a significant relationship between physical activity and vitamin D status in the nonobese and obese respondent groups, but the relationship was low or weak (Table 9).

DISCUSSION

Respondents who were involved in this study were 143 people and consisted of male sex at 39.86% (57 people) and female sex at 60.13% (86 people). The group of respondents in the category of male obesity were as many as 53.52% (38 people) and female sex as many as 46.47% (33 people) and for respondents with nonobesity categories, male respondents as many as 26.38% (19 people) and women 73.61% (53 people). Gender can affect vitamin D status; the female sex has a low vitamin status compared to the male sex (Annweiler *et al.*, 2017; AlQuaiz *et al.*, 2018; Sanghera *et al.*, 2017) due to avoiding sun exposure and dress habits are one of the main factors (Kift *et al.*, 2018; Lee and Lee, 2020) such as wearing long-sleeved shirts, long skirts, and headscarves (Judistiani *et al.*, 2019; Pulungan *et al.*, 2021). Gender

Table 3. Distribution of respondents' answers regarding length of exposure and hours of sun exposure.

Component in question	Nonobese group (n = 72)		Obese group (n = 71)			
	Frequency	Percentage (%)	Frequency	Percentage (%)		
Duration of exposure and hours of sun exposure						
Duration of exposure (minute)	<15	20	27.77	21	29.57	
	>15	52	72.22	50	70.42	
Hours of exposure	07.00–09.00 am and 03.00–05.00 pm	35	48.61	41	57.74	
	10.00–11.00 am	37	51.38	30	42.25	
Use of protective equipment						
Use of protective equipment	Does not use protective equipment	5	6.94	15	21.12	
	Umbrella	8	11.11	8	11.26	
	Hat	7	9.72	8	11.26	
	Jacket	52	72.22	56	78.87	
	Sunblock/sun cream	48	66.66	40	56.33	
How often to use	Others (gloves, masks, etc.)	6	8.33	4	5.63	
	Always or every day	46	63.88	34	47.88	
	Sometimes	10	13.88	15	21.12	
	Rarely	11	15.27	17	23.94	
Applications use of protective equipment	Never	5	6.94	5	7.04	
	When leaving the house/boarding house	34	47.22	35	49.29	
	After 12 pm	28	38.88	25	35.21	
	When driving	9	12.50	10	14.08	
	Time at the beach	1	1.38	1	1.40	
Use of closed clothing and body parts that are protected from sun exposure						
Use of closed clothing	Not wearing closed clothes	30	41.66	44	61.97	
	Using closed clothes	42	58.33	27	38.02	
	Description of closed clothing	Using the hijab	12	16.66	3	4.22
		Jackets and trousers	17	23.61	14	19.71
Body part covered ^a	Long sleeves and trousers	13	18.05	10	14.08	
	Face	45	62.50	38	53.52	
	Hand	49	68.05	45	63.38	
	Sleeve	41	56.94	45	63.38	
	Foot	35	48.61	30	42.25	
	Back and shoulders	72	100.00	71	100.00	
Use of sunblock/sun cream						
Use of sunblock/sunscreen cream	Does not use sunblock/sun cream	24	33.33	31	43.66	
	Menggunakan sunblock/krim tabir surya	48	66.66	40	56.33	
	Always or every day	27	37.50	10	14.08	
How often to use sunblock/sun cream	Often	4	5.55	6	8.45	
	Sometimes	6	8.33	9	12.67	
	Rarely	11	15.27	15	21.12	
Duration of use of sunblock/sunscreen cream	1–3 hours	28	38.88	28	39.43	
	3–6 hours	8	11.11	4	5.63	
	>6 hours	12	16.66	8	11.26	
Repeated use of sunblock/sun cream in a day	No	62	86.11	67	94.36	
	Yes	<4 hours	1	1.38	1	1.40
		>4 hours	9	12.50	3	4.22
Use of cosmetics containing SPF	Yes	50	69.44	38	53.52	
	No	14	19.44	28	39.43	
	Does not know	8	11.11	5	7.04	

^aOne respondent can have more than one.

Table 4. Distribution of respondents' answers regarding foods containing vitamin D.

Component in question		Nonobese group (n = 72)		Obese group (n = 71)	
		Frequency	Percentage (%)	Frequency	Percentage (%)
Fish					
Consumption of fish in a week	Ate fish	53	73.61	47	66.19
	Did not eat fish	19	26.38	24	33.80
Type of fish consumed	Fish high in vitamin D (tuna, salmon, mackerel)	39	54.16	37	52.11
	Fish did not contain much vitamin D (salted fish, pindang, mujaer)	33	45.83	34	47.88
How the consumed fish was processed	Fried	48	66.66	50	70.42
	Burned	7	9.72	10	14.08
	Sushi	2	2.77	5	7.04
	Boiled	2	2.77	3	4.22
	Other	11	15.27	15	21.12
Where the food (fish) was acquired	Cook by myself	7	9.72	9	12.67
	Restaurant	27	37.50	40	56.33
	Catering	2	2.77	1	1.40
	Supermarket	20	27.77	16	22.53
Purpose of consuming fish	Love the taste	49	68.05	46	64.78
	Get health benefits	23	31.94	28	39.43
	Lose weight	3	4.16	6	8.45
	Price lower or more affordable	9	12.50	8	11.26
	Incidentally a fish dish available to eat	24	33.33	23	32.39
	Other	1	1.38	2	2.81
Egg and milk					
Consumption of milk (in the past 1 week)	Did not consume milk	20	27.77	25	35.21
	Consumed milk	52	72.22	46	64.78
Consumption of egg (in the past 1 week)	Did not consume egg	64	88.88	68	95.77
	Consumed egg	8	11.11	3	4.22
The portion of the egg consumed	Egg whites	9	12.50	1	1.40
	Egg yolk	0	0.00	1	1.40
	White and yellow	63	87.50	69	97.18
How the eggs consumed was processed	Fried	62	86.11	68	95.77
	Boiled	17	23.61	16	22.53
	Other	3	4.16	1	1.40
Behavior frequency of consumption of food and beverages containing vitamin D					
Egg	Often	18	25.00	29	40.84
	Rare	54	75.00	42	59.15
Fish	Often	5	6.94	5	7.04
	Rare	67	93.05	66	92.95
Milk	Often	19	26.38	21	29.57
	Rare	53	73.61	50	70.42
Behavior of consumption of fish oil and vitamin D supplements					
Consumption of fish oil	Did not consume fish oil	64	88.88	70	98.59
	Consumed fish oil	8	11.11	1	1.40
Consumption of supplements	Did not consume vitamin D supplements	52	72.22	52	73.23
	Consumed vitamin D supplements	2	2.77	0	0.00
	Containing vitamin D	18	25.00	0	0.00
	Not containing vitamin D	18	25.00	0	0.00

Table 5. Distribution of respondents' answers regarding signs of vitamin D deficiency.

Component in question	Nonobese group (n = 72)		Obese group (n = 71)	
	Frequency	Percentage (%)	Frequency	Percentage (%)
Muscle pain including lower back pain	27	37.50	30	42.25
Pain in the pelvis, back, and legs	23	31.94	28	39.43
Muscle weakness	8	11.11	10	14.08
Easy to get into a bad mood or get depressed easily	30	41.66	34	47.88
Low immunity, such as frequent colds in winter	23	31.94	26	36.61
Not at all experienced any of the conditions above	7	9.72	5	7.04

A respondent can experience more than one sign and symptom.

Table 6. Cross-tabulation of vitamin D status and physical activity in the nonobese and obese respondents.

Category		Nonobese group (n = 72)		Obese group (n = 71)	
		Frequency	Percentage (%)	Frequency	Percentage (%)
Vitamin D status (deficiency)	Negative	17	23.61	39	54.93
	Positive	55	76.39	32	45.07
Physical activity	Low physical activity	30	41.67	44	61.97
	Moderate physical activity	40	55.55	25	35.21
	High physical activity	2	2.78	2	2.82

Table 7. Risk analysis and differences test of vitamin D deficiency related to vitamin D status in nonobese and obese respondents.

Respondent group	Vitamin D status		Total	Odds ratio	Chi-square test
	Nondeficiency	Deficiency			
Nonobese (n = 72)	17	55	72	POR = 0.254 CI 95% (0.124–0.520)	p = 0.000 (p < 0.05)
Obese (n = 71)	39	32	71	Conclusion: The nonobese group had a 0.2-fold risk of vitamin D deficiency compared to the obese group	Conclusion: There were significant differences in vitamin D status in the obese and nonobese groups
Total	56	87	143		

Table 8. Risk analysis and differences test in physical activity in nonobese and obese respondents.

Respondent group	Physical activity			Total	Chi-square test
	Low	Moderate	High		
Nonobese (n = 72)	30	40	2	72	p = 0.047 (p < 0.05)
Obese (n = 71)	44	25	2	71	
Total	74	65	4	143	Conclusion: There was significant difference in physical activity in the obese and nonobese groups

Table 9. Results of analysis of the relationship between vitamin D status and physical activity.

Vitamin D status	Physical activity			Total	Correlation test
	Low	Moderate	High		
Deficiency	30	40	2	72	$(p = 0.000)$ KK = 0.326 Conclusion: There was a significant relationship with vitamin D status and physical activity, but the relationship was low or weak
Nondeficiency	44	25	2	71	
Total	74	65	4	143	

not only affects vitamin D status but also affects physical activity (Fernandes and Junior, 2017; Wiciński *et al.*, 2019). History of disease and use of drugs have a relationship with vitamin D status in the body. Certain pathological conditions, such as Crohn’s disease, cystic fibrosis, celiac disease, and surgical removal of parts of the stomach or intestines that play a role in fat absorption, can trigger vitamin D deficiency. Anticonvulsant drugs are also called antiepileptic drugs. The long duration of some antiepileptic drugs, such as phenobarbital, phenytoin, and carbamazepine, and the antimicrobial agent rifampin can cause osteomalacia, which is a condition caused by vitamin D deficiency. This is triggered by the induction of 1,25-dihydroxyvitamin D catabolism (Saket *et al.*, 2021). Men tend to be more active than women, because spend more time sitting and doing activities that smell of cosmetic, Women tend to be inactive compared to men. Women spend more time sitting and doing activities that smell of cosmetic while the male sex prefers to do sports so the activities carried out are higher than women (Sharifi *et al.*, 2016).

All respondents involved in this study were aged 18–25 years and were active students. This study uses young adult respondents because based on previous research young adults are known to be at risk of vitamin D deficiency which is influenced by habits such as consumption of vitamin D below the recommended limit per day, lack of outdoor physical activity, increased use of sunblock/sunscreen cream, and low milk intake (Bolek-Berquist *et al.*, 2009). In addition to young adults, the elderly are also one of the age groups that have a low risk of vitamin D status. Low vitamin D status can cause muscle weakness (Gunton and Girgis, 2018; Sizar *et al.*, 2021) and increase the risk of fractures in the elderly (Thanaplueitiwong *et al.*, 2020).

Obesity is a risk factor for vitamin D deficiency (Paschou *et al.*, 2019) based on the research by Kavadar *et al.*, (2015) that found a relationship between obesity and low vitamin D status, which has an impact on the risk of insulin resistance. The mechanism of vitamin D deficiency in obese patients is not precisely known, but several researchers have hypothesized that, related to the mechanism of vitamin D deficiency in obese patients, high fat accumulation results in a decrease in vitamin D bioavailability, where a lot of vitamin D is stored in fat cells in adipose tissue and increases vitamin D deficiency. Oxidative reactions, due to the fat-soluble nature of vitamin D, cause a decrease in the release of vitamin D from fat into the systemic circulation (Wimalawansa, 2019) in obese patients. One of the causes of obesity is excessive food intake, low physical activity (Romieu *et al.*, 2017), and lack of rest hours; besides that, there

is a relationship between age and workplace characteristics on the risk of becoming obese (Hruby *et al.*, 2016). Low physical activity is found in obese people (Kim *et al.*, 2017); therefore, increasing physical activity can be used as an appropriate step to prevent the risk of obesity and weight loss (Jakicic *et al.*, 2019).

Long exposure can affect the synthesis of vitamin D by the body; an exposure time of 15 minutes can synthesize vitamin D of 10,000–25,000 IU over a large area of the body surface and is characterized by a change in skin color to pink (Suryadinata *et al.*, 2021). In this study, the results showed that, in the two groups of respondents who were obese and nonobese, those who received longer sun exposure (>15 minutes) were the nonobese group of 36.36% (52 people) compared to the obese group of 36.36% (52 people). The most dominant group with less exposure to sunlight (<15 minutes) was the obese group at as much as 14.68% (21 people) and then the nonobese group at 13.98% (20 people). One of the factors that influence the duration of exposure to vitamin D synthesis is skin type (Wacker and Holick, 2013). Different skin types affect the length of time it takes the skin to synthesize vitamin D. Dark skin type (type IV skin) requires a long exposure time of six times longer than skin type I (Wacker and Holick, 2013). This is influenced by the pigment in the skin called melanin which functions as a natural sunscreen that can block the absorption of sunlight by the skin (Solano, 2020). Asian skin is in skin type IV or V; therefore, Asian people need a longer exposure time than other skin types. The recommended exposure time is more than 15 minutes or even up to 2 hours (Webb *et al.*, 2018). Exposure time is also an important factor in the process of vitamin D synthesis (Suryadinata *et al.*, 2021). It is known from the results of this study that the highest Ultraviolet B (UVB) intensity occurs from 11.00 am to 01.00 pm (Lorensia *et al.*, 2020b). UVB rays obtained from sunlight can help the synthesis of vitamin D in the skin (Wacker *et al.*, 2013; Lee and Lee, 2020).

One source of vitamin D is fish. The highest amount of vitamin D is found in fish liver (2–477 mg/kg), a maximum of 1,200 mg/kg and depending on the type of fish (Schmid and Walther, 2013). A total of 66.19% (47 people) of the obese respondent group, who consumed fish within the past week, consumed fish with vitamin D content of 52.11% (37 people), while in the nonobese respondent group of a total of 73.61% (53 people) who consumed fish as many as 54.16% (39 people) consumed fish with vitamin D content. The obese respondent group (7.04%) and the nonobese respondent group (6.94%) often consumed fish in the last week. The frequency of fish consumption can affect vitamin D status. In

Japan, fish is the main source for obtaining vitamin D (Nakamura *et al.*, 2002). The cooking process does not really affect the vitamin D content in food products because vitamin D is resistant to heat (Schmid and Walther, 2013), but the fish processing process also affects the amount of vitamin D.

Eggs are a source of vitamin D (Schmid and Walther, 2013). The obese respondents who consumed eggs were 95.77% (68 people), and the nonobese respondents who consumed eggs were 88.88% (64 people). The respondents from the obese group who often consumed eggs within 1 week were 40.84% (29 people), and the respondents from the nonobese group were 25% (18 people). Vitamin D in eggs is found in the yolk at as much as 20 IU of vitamin D3 or D2 per egg yolk (Schmid and Walther, 2013). In addition to eggs and fish, milk is also a fortification product that contains vitamin D (Lorensia *et al.*, 2020b; Schmid and Walther, 2013). Someone with low milk consumption can experience vitamin D deficiency (Bolek-Berquist *et al.*, 2009).

Based on the results of the analysis with the chi-square test, the *p*-value of 0.047 (*p*-value <0.05) can be concluded so it can be concluded that there is a significant difference in physical activity between the nonobese and obese groups of respondents. This is supported by evidence from previous studies which showed that there was a significant difference in physical activity in the normal group compared to the obese group, where the obese group had a longer sitting time than the nonobese group, and obesity had a significant relationship with low physical activity and physical function (Hong *et al.*, 2016).

In the analysis of the relationship between vitamin D status and physical activity using the contingency coefficient correlation test, we obtained an approximate significance value of 0.000 (*p*-value = 0.000), which indicates that there is a significant relationship between vitamin D status and physical activity and the relationship has a correlation value of 0.326. Based on the contingency coefficient table, the two variables have a low or weak relationship but are definitely related so it can be concluded that there is a relationship between vitamin D status and physical activity in the nonobese and obese respondent groups, but the relationship is low or weak. This is in accordance with previous research which showed that a person with low physical activity has a low vitamin D status (Aji *et al.*, 2019). In addition, high outdoor physical activity can increase vitamin D status (Lorensia *et al.*, 2021). Outdoor physical activity, which is in exposure to sunlight, is statistically significant in increasing vitamin D status (Lorensia *et al.*, 2020a). Therefore, physical activity and dietary intake of vitamin D should be increased in an effort to prevent vitamin D deficiency and other health problems such as metabolic risk factors (Lorensia *et al.*, 2020a). There are some limitations in this research, including the existence of variables that can affect the results of the study such as the uneven distribution of sexes between the two groups of respondents being compared, and there is no examination of serum 25(OH)D for determining vitamin D levels.

CONCLUSION

There is a significant difference in vitamin D status in the obese and nonobese groups of respondents, and there is a significant difference in physical activity in the obese and

nonobese groups of respondents. In addition, there is a significant relationship between vitamin D status and physical activity, but the relationship between the two is low or weak.

ACKNOWLEDGMENT

This research was funded by the Institute of Research and Community Service by Universitas Surabaya.

CONFLICTS OF INTEREST

The authors declare no conflicts of interest.

DATA AVAILABILITY

All data generated and analyzed are included within this research article.

PUBLISHER'S NOTE

This journal remains neutral with regard to jurisdictional claims in published institutional affiliation.

REFERENCES

- Ahmed SH, Meyer HE, Kjøllestad MK, Marjerrison N, Mdala I, Htet AS, Bjertness E, Madar AA. The prevalence of selected risk factors for non-communicable diseases in Hargeisa, Somaliland: a cross-sectional study. *BMC Public Health*, 2019; 19(1):878.
- Aji AS, Erwinda E, Yusrawati Y, Malik SG, Lipoeto NI. Vitamin D deficiency status and its related risk factors during early pregnancy: a cross-sectional study of pregnant Minangkabau women, Indonesia. *BMC Pregnancy Childbirth*, 2019; 19(1):183.
- AlQuaiz AM, Kazi A, Fouda M, Alyousefi N. Age and gender differences in the prevalence and correlates of vitamin D deficiency. *Arch Osteoporos*, 2018; 13(1):49.
- Annweiler C, Kabeshova A, Callens A, Paty ML, Duval GT, Holick MF. Self-administered vitamin D status predictor: older adults are able to use a self-questionnaire for evaluating their vitamin D status. *PLoS One*, 2017; 12(11):e0186578.
- Bolek-Berquist J, Elliott ME, Gangnon RE, Gemar D, Engelke J, Lawrence SJ, Hansen KE. Use of a questionnaire to assess vitamin D status in young adults. *Public Health Nutr*, 2009; 12(2):236–43.
- Cairncross CT, Stonehouse W, Conlon CA, Grant CC, McDonald B, Houghton LA, Eyles D, Camargo CA Jr, Coad J, von Hurst PR. Predictors of vitamin D status in New Zealand preschool children. *Matern Child Nutr*, 2017; 13(3):e12340.
- Charoenngam N, Holick MF. Immunologic effects of vitamin D on human health and disease. *Nutrients*, 2020; 12(7):2097.
- Cleland C, Ferguson S, Ellis G, Hunter RF. Validity of the International Physical Activity Questionnaire (IPAQ) for assessing moderate-to-vigorous physical activity and sedentary behaviour of older adults in the United Kingdom. *BMC Med Res Methodol*, 2018; 18(1):176.
- Damayanti RA, Sumarmi S, Mundiastuti L. The association between sleep duration with overweight and obesity among education staff in campus C of Universitas Airlangga. *Amerta Nutr*, 2019; 3(2):89–93.
- De Lorenzo A, Gratteri S, Gualtieri P, Cammarano A, Bertucci P, Di Renzo L. Why primary obesity is a disease? *J Transl Med*, 2019; 17(1):169.
- Elliott CA, Hamlin MJ. Combined diet and physical activity is better than diet or physical activity alone at improving health outcomes for patients in New Zealand's primary care intervention. *BMC Public Health*, 2018; 18(1):230.
- Fernandes MR, Junior DRBJ. Association between physical activity and vitamin D: a narrative literature review. *Rev Assoc Med Bras*, 2017; 63(6):550–6.
- Fruh SM. Obesity: risk factors, complications, and strategies for sustainable long-term weight management. *J Am Assoc Nurse Pract*, 2017; 29(S1):S3–14.

- Füzéki E, Banzer W. Physical activity recommendations for health and beyond in currently inactive populations. *Int J Environ Res Public Health*, 2018; 15(5):1042.
- Gunton JE, Girgis CM. Vitamin D and muscle. *Bone Rep*, 2018; 8:163–7.
- Hall JT, Ebeling M, Shary JR, Forestieri N, Wagner CL. The relationship between physical activity and vitamin D status in postpartum lactating and formula-feeding women. *J Steroid Biochem Mol Biol*, 2018; 177:261–5.
- Hamer M, Stamatakis E. Relative proportion of vigorous physical activity, total volume of moderate to vigorous activity, and body mass index in youth: the Millennium Cohort Study. *Int J Obes*, 2018; 42(6):1239–42.
- Hassanalilou T, Khalili L, Ghavamzadeh S, Shokri A, Payahoo L, Bishak YK. Role of vitamin D deficiency in systemic lupus erythematosus incidence and aggravation. *Auto Immun Highlights*, 2017; 9(1):1.
- Hong I, Coker-Bolt P, Anderson KR, Lee D, Velozo CA. Relationship between physical activity and overweight and obesity in children: findings from the 2012 National Health and Nutrition Examination Survey National Youth Fitness Survey. *Am J Occup Ther*, 2016; 70(5):7005180060p1–8.
- Hruby A, Manson JE, Qi L, Malik VS, Rimm EB, Sun Q, Willett WC, Hu FB. Determinants and consequences of obesity. *Am J Public Health*, 2016; 106(9):1656–62.
- Jackson SE, Llewellyn CH, Smith L. The obesity epidemic—nature via nurture: a narrative review of high-income countries. *SAGE Open Med*, 2020; 8:2050312120918265.
- Jakicic JM, Powell KE, Campbell WW, Dipietro L, Rate RR, Pescatello LS, Collins KA, Bloodgood B, Piercy KL. Physical activity and the prevention of weight gain in adults: a systematic review. *Med Sci Sports Exerc*, 2019; 51(6):1262–9.
- Jayawardena R, Jeyakumar DT, Misra A, Hills AP, Ranasinghe P. Obesity: a potential risk factor for infection and mortality in the current COVID-19 epidemic. *Diabetes Metab Syndr*, 2020; 14(6):2199–203.
- Jehan S, Zizi F, Pandi-Perumal SR, McFarlane SI, Jean-Louis G, Myers AK. Energy imbalance: obesity, associated comorbidities, prevention, management and public health implications. *Adv Obes Weight Manag Control*, 2020; 10(5):146–61.
- Judistiani RTD, Nirmala SA, Rahmawati M, Ghrahani R, Natalia YA, Sugianli AK, Indrati AR, Suwarsa O, Setiabudiawan B. Optimizing ultraviolet B radiation exposure to prevent vitamin D deficiency among pregnant women in the tropical zone: report from cohort study on vitamin D status and its impact during pregnancy in Indonesia. *BMC Pregnancy Childbirth*, 2019; 19(1):209.
- Kavadar G, Demircioğlu DT, Özgönel L, Emre TY. The relationship between vitamin D status, physical activity and insulin resistance in overweight and obese subjects. *Bosn J Basic Med Sci*, 2015; 15(2):62–6.
- Kift R, Rhodes LE, Farrar MD, Webb AR. Is sunlight exposure enough to avoid wintertime vitamin d deficiency in United Kingdom population groups? *Int J Environ Res Public Health*, 2018; 15(8):1624.
- Kim BY, Choi DH, Jung CH, Kang SK, Mok JO, Kim CH. Obesity and physical activity. *J Obes Metab Syndr*, 2017; 26(1):15–22.
- Kim SH, Oh JE, Song DW, Cho CY, Hong SH, Cho YJ, Yoo BW, Shin KS, Joe H, Shin HS, Son DY. The factors associated with vitamin D deficiency in community dwelling elderly in Korea. *Nutr Res Pract*, 2018; 12(5):387–95.
- Koehler K, Drenowatz C. Integrated role of nutrition and physical activity for lifelong health. *Nutrients*, 2019; 11(7):1437.
- Kotarska K, Nowak MA, Nowak L, Król P, Sochacki A, Sygit K, Sygit M. Physical activity and quality of life of university students, their parents, and grandparents in poland-selected determinants. *Int J Environ Res Public Health*, 2021; 18(8):3871.
- Larson-Meyer DE, Douglas CS, Thomas JJ, Johnson EC, Barcal JN, Heller JE, Hollis BW, Halliday TM. Validation of a vitamin D specific questionnaire to determine vitamin D status in athletes. *Nutrients*, 2019; 11(11):2732.
- Lee YM, Kim SA, Lee DH. Can current recommendations on sun exposure sufficiently increase serum vitamin d level? One-month randomized clinical trial. *J Korean Med Sci*, 2020; 35(8):e50.
- Lorensia A, Suryadinata RV, Saputra R. Physical activity and vitamin D level in asthma and non-asthma. *J Farmasi Indones*, 2019; 11(1):454–65.
- Lorensia A, Muntu CM, Suryadinata RV, Septiani R. Effect of lung function disorders and physical activity on smoking and non-smoking students. *J Prev Med Hyg*, 2021; 62(1):E89–96.
- Lorensia A, Wahyudi M, Yudiarto A, Kurnia SED. Effect of illness perception on improving asthma symptoms with omega-3 fish oil therapy: Pre-post design. *J Appl Pharm Sci*, 2020a; 10(6):62–71.
- Lorensia A, Suryadinata RV, Chandra NLMR. Profile of vitamin D status, physical activity, and lung health in construction workers (profil status vitamin D, Aktivitas Fisik, dan Kesehatan Paru pada Tukang Bangunan). *Community Med Public Health Indones J*, 2020b; 1(2):117–24.
- Maurya VK, Aggarwal M. Factors influencing the absorption of vitamin D in GIT: an overview. *J Food Sci Technol*, 2017; 54(12):3753–65.
- Michigami T. Rickets/osteomalacia. Consensus on Vitamin D deficiency and insufficiency in children. *Clin Calcium Jpn*, 2018; 28(10):1307–11.
- Nabak A, Johnson R, Keuler N, Hansen K. Can a questionnaire predict vitamin D status in postmenopausal women? *Public Health Nutr*, 2014; 17(4):739–46.
- Nakamura K, Nashimoto M, Okuda Y, Ota T, Yamamoto M. Fish as a major source of vitamin D in the Japanese diet. *Nutrition*, 2002; 18(5):415–6.
- Oh TR, Kim CS, Bae EH, Ma SK, Han SH, Sung SA, Lee K, Oh KH, Ahn C, Kim SW. Association between vitamin D deficiency and health-related quality of life in patients with chronic kidney disease from the KNOW-CKD study. *PLoS One*, 2017; 12(4):e0174282.
- Orces C. The association between body mass index and vitamin D supplement use among adults in the United States. *Cureus*, 2019; 11(9):e5721.
- Panuganti KK, Nguyen M, Kshirsagar RK. Obesity. *StatPearls*, Treasure Island, FL, 2021. [ONLINE] Available via <https://www.ncbi.nlm.nih.gov/books/NBK459357/> (Accessed 11 August 2021).
- Paschou SA, Kosmopoulos M, Nikas IP, Spartalis M, Kassi E, Goulis DG, Lambrinou I, Siasos G. The impact of obesity on the association between vitamin D deficiency and cardiovascular disease. *Nutrients*, 2019; 11(10):2458.
- Pilz S, Zittermann A, Trummer C, Theiler-Schwetz V, Lerchbaum E, Keppel MH, Grubler MR, Marz W, Pandis M. Vitamin D testing and treatment: a narrative review of current evidence. *Endocr Connect*, 2019; 8(2):R27–43.
- Polzonetti V, Pucciarelli S, Vincenzetti S, Polidori P. Dietary Intake of Vitamin D from dairy products reduces the risk of osteoporosis. *Nutrients*, 2020; 12(6):1743.
- Pulungan A, Soesanti F, Tridjaja B, Batubara J. Vitamin D insufficiency and its contributing factors in primary school-aged children in Indonesia, a sun-rich country. *Ann Pediatr Endocrinol Metab*, 2021; 26(2):92–8.
- Rachmi CN, Li M, Alison Baur L. Overweight and obesity in Indonesia: prevalence and risk factors—a literature review. *Public Health*, 2017; 147:20–9.
- Ramasamy I. Vitamin D metabolism and guidelines for vitamin D supplementation. *Clin Biochem Rev*, 2020; 41(3):103–26.
- Romieu I, Dossus L, Barquera S, Blottiere HM, Franks PW, Gunter M, Hwalla N, Hursting SD, Leitzmann M, Margetts B, Nishida C. Energy balance and obesity: what are the main drivers? *Cancer Causes Control*, 2017; 28(3):247–58.
- Saket S, Varasteh N, Halimi Asl AA, Saneifard H. How antiepileptics may change the serum level of vitamin D, calcium, and phosphorus in children with epilepsy. *Iran J Child Neurol*, 2021; 15(1):19–27.
- Salam RA, Padhani ZA, Das JK, Shaikh AY, Hoodbhoy Z, Jeelani SM, Lassi ZS, Bhutta ZA. Effects of lifestyle modification interventions to

prevent and manage child and adolescent obesity: a systematic review and meta-analysis. *Nutrients*, 2020; 12(8):2208.

Sanghera DK, Sapkota BR, Aston CE, Blackett PR. Vitamin D status, gender differences, and cardiometabolic health disparities. *Ann Nutr Metab*, 2017; 70(2):79–87.

Saqib ZA, Dai J, Menhas R, Mahmood S, Karim M, Sang X, Weng Y. Physical activity is a medicine for non-communicable diseases: a survey study regarding the perception of physical activity impact on health wellbeing. *Risk Manag Healthc Policy*, 2020; 13:2949–62.

Schmid A, Walther B. Natural vitamin D content in animal products. *Adv Nutr*, 2013; 4(4):453–62.

Sember V, Meh K, Sorić M, Starc G, Rocha P, Jurak G. Validity and reliability of international physical activity questionnaires for adults across EU countries: systematic review and meta-analysis. *Int J Environ Res Public Health*, 2020; 17(19):7161.

Sharifi N, Amani R, Hajiani E, Cheraghian B. Women may respond different from men to vitamin D supplementation regarding cardiometabolic biomarkers. *Exp Biol Med (Maywood)*, 2016; 241(8):830–8.

Silva ACMD, Cureau FV, de Oliveira CL, Giannini DT, Bloch KV, Kuschner MCC, Dutra ES, Schaan BD, de Carvalho KMB. Physical activity but not sedentary time is associated with vitamin D status in adolescents: study of cardiovascular risk in adolescents (ERICA). *Eur J Clin Nutr*, 2019; 73(3):432–40.

Sizar O, Khare S, Goyal A, Givler A. Vitamin D deficiency. *StatPearls*, Treasure Island, FL, 2021. [ONLINE] Available via <https://www.ncbi.nlm.nih.gov/books/NBK532266/> (Accessed 12 April 2021).

Skalska M, Nikolaidis PT, Knechtle B, Rosemann TJ, Radziminski L, Jastrzebska J, Kaczmarczyk M, Myśliwiec A, Dragos P, López-Sánchez GF, Jastrzębski Z. Vitamin D supplementation and physical activity of young soccer players during high-intensity training. *Nutrients*, 2019; 11(2):349.

Solano F. Photoprotection and skin pigmentation: melanin-related molecules and some other new agents obtained from natural sources. *Molecules*, 2020; 25(7):1537.

Suryadinata RV, Boengas S, Lorensia A. Effect of knowledge and attitude toward sun exposure related vitamin D to lung function. *Teikyo Med J*, 2021; 44(4):957–69.

Suryadinata RV, Wirjatmadi B, Andriani M, Lorensia A. Effect of age and weight on physical activity. *J Public Health Res*, 2020; 9(2):187–90.

Takeuchi Y. Rickets/osteomalacia. Treatment of vitamin D deficiency in adults. *Clin Calcium*, 2018; 28(10):1387–92.

Thanapluetiwong S, Chewcharat A, Takkavatakarn K, Praditpornsilpa K, Eiam-Ong S, Susantitaphong P. Vitamin D supplement on prevention of fall and fracture: a meta-analysis of randomized controlled trials. *Medicine (Baltimore)*, 2020; 99(34):e21506.

Umar M, Sastry KS, Chouchane AI. Role of vitamin D beyond the skeletal function: a review of the molecular and clinical studies. *Int J Mol Sci*, 2018; 19(6):1618.

Wacker M, Holick MF. Sunlight and vitamin D: a global perspective for health. *Dermatoendocrinol*, 2013; 5(1):51–108.

Wang H, Chen W, Li D, Yin X, Zhang X, Olsen N, Zheng SG. Vitamin D and chronic diseases. *Aging Dis*, 2017; 8(3):346–53.

Webb AR, Kazantzidis A, Kift RC, Farrar MD, Wilkinson J, Rhodes LE. Colour counts: sunlight and skin type as drivers of vitamin D deficiency at UK Latitudes. *Nutrients*, 2018; 10(4):457.

Wiciński M, Adamkiewicz D, Adamkiewicz M, Śniegocki M, Podhorecka M, Szycha P, Malinowski B. Impact of vitamin D on physical efficiency and exercise performance—a review. *Nutrients*, 2019; 11(11):2826.

Wimalawansa SJ. Vitamin D deficiency: effects on oxidative stress, epigenetics, gene regulation, and aging. *Biology (Basel)*, 2019; 8(2):30.

Zadka K, Pałkowska-Goździk E, Rosołowska-Huszcz D. The state of knowledge about nutrition sources of vitamin D, its role in the human body, and necessity of supplementation among parents in Central Poland. *Int J Environ Res Public Health*, 2018; 15(7):1489.

Zarooni AAR, Marzouqi FIA, Darmaki A, Prinsloo EAM, Nagelkerke N. Prevalence of vitamin D deficiency and associated comorbidities among Abu Dhabi Emirates population. *BMC Res Notes*, 2019; 12(503):1–6.

How to cite this article:

Lorensia A, Suryadinata RV, Inu IA. Comparison of vitamin D status and physical activity related to obesity among tertiary education students. *J Appl Pharm Sci*, 2022; 12(04):108–118.

About the Journal

On this page: [Aims & scope](#), [Journal policies](#), [Disclaimer](#)

AIMS & SCOPE

Journal of Applied Pharmaceutical Science (JAPS) is a monthly open-access journal dedicated to various disciplines of pharmaceutical and allied sciences. JAPS publishes manuscripts (Original research and review articles Mini-reviews, Short communication) on original work, either experimental or theoretical in the following areas;

- Pharmaceutics & Biopharmaceutics
- Novel & Targeted Drug Delivery
- Nanotechnology & Nanomedicine
- Pharmaceutical & Medicinal Chemistry
- Pharmacognosy & Ethnobotany
- Phytochemistry
- Pharmacology & Toxicology
- Pharmaceutical Biotechnology & Microbiology
- Pharmacy Practice & Hospital Pharmacy
- Pharmacovigilance
- Natural Product Research
- Drug Regulatory Affairs
- Analytical Chemistry

JOURNAL POLICIES

- [Open Access Policy](#)
- [Peer-review Policy](#)
- [Waiver Policy](#)
- [Copyright & Licensing policies](#)
- [Author Self-Archiving policies](#)
- [Publication ethics and malpractice statement](#)
- [Plagiarism Policy](#)
- [CrossMark Policy](#)
- [Digital Preservation](#)
- [Retraction policy](#)
- [Preprints policy](#)
- [Conflicts of Interest](#)
- [Complaints and appeals](#)
- [Misconduct and sanctions](#)

Open Access Policy

All the articles published in JAPS are freely available without any registration, subscription, or payment. Users are allowed to read, download, copy, distribute, print, search, or link to the full texts of the articles in this journal without asking prior permission from the publisher or the author, as long as the original authors are properly cited.

Furthermore, the journal also allows the authors to self-archive the final accepted version of the articles on any OAI-compliant institutional/subject-based repository.

This is in accordance with the **Budapest Open Access Initiative** definition, which states;

"By 'open access to this literature, we mean its free availability on the public internet, permitting any users to read, download, copy, distribute, print, search, or link to the full texts of these articles, crawl them for indexing, pass them as data to software, or use them for any other lawful purpose, without financial, legal, or technical barriers other than those inseparable from gaining access to the internet itself." [Read: [The Budapest Open Access Initiative](#)]

JAPS adopted a gold open-access model, which means that the published articles are funded by article processing charges paid at the time of acceptance. Author(s) retain the copyright of their work, with articles licensed to JAPS.

All the published articles are **licensed under a Creative Commons License (CC-BY)**, full details of which can be found [here](#).

Peer-review Policy

All the manuscripts submitted to JAPS will be subjected to the double-blinded peer-review process;

Initial Evaluation/Technical Review

CURRENT ISSUE

April, 2022

ARCHIVES

[READ MORE](#)

Follow us  

The Editor(s) first evaluates all the manuscripts. Manuscripts rejected at this stage are insufficiently original, have serious scientific flaws, have poor grammar or English language, or are outside the aims and scope of the journal, or not written as per the journal's guidelines. Those that meet the minimum criteria are forwarded to at least 2 experts for review.

Peer-review

- Generally, all the manuscripts are reviewed by a minimum of two suitable external experts in the respective subject area. The reports of all the reviewers will be considered while deciding on acceptance/revision or rejection of a manuscript, but the ultimate responsibility for acceptance or rejection lies with the Editor. Editor-In-Chief makes the final decision.
- EIC can ask one or more advisory board members for their suggestions upon a manuscript, before making the final decision.
- Associate editor and review editors provide administrative support to maintain the integrity of the peer-review process.
- In accordance with COPE recommendations on ethical editing for Editors, Editors will assign any submissions they cannot handle (e.g. if they are the author of an article submitted to their own journal) to a member of the Editorial Board or a guest editor.

Author's suggestion for the reviewer

The authors can suggest the peer-reviewers as per the following criteria:

- The suggested reviewers should have higher academic standing in the subject area of the submitted article and the same should be reflected in terms of their publications.
- The suggested reviewers should not be from the authors' own institute or from past affiliations.
- Authors must provide the **institutional/academic email addresses** of the suggested reviewers.
- Authors must provide the **web links to the academic profile pages** of the suggested reviewers.
- The journal reserve the right NOT to choose any reviewer from the suggested list.
- In case, authors challenge the editor's negative decision with suitable arguments, the manuscript can be sent to one more reviewer and the final decision will be made based upon his recommendations.

Note: Journal of Applied Pharmaceutical Science is a member of the World Association of Medical Editors. Please [click here](#) for more details about peer review and editorial policies.

Waiver Policy

JAPS may grant the partial/full waiver in article processing charges to the authors who are unable to pay the APCs due to financial restrictions. Only the authors from **low-income countries** can apply for the waiver. Please send your waiver requests to editor@japsonline.com with complete details.

Copyright and licensing policies

The journal allows the author(s) to hold the copyright of their work. That means the authors do not need to transfer the copyright of their work to the journal. However, the authors grant JAPS a license to publish the article and identify itself as the original publisher. All the articles published in JAPS are distributed under a creative commons license (CC-BY 4.0) (read detailed license information here: <https://creativecommons.org/licenses/by/4.0/>).

You are free to:

- Share** — copy and redistribute the material in any medium or format
- Adapt** — remix, transform, and build upon the material for any purpose, even commercially.

Under the following terms:

- Attribution** — You must give appropriate credit, provide a link to the license, and indicate if changes were made. You may do so in any reasonable manner, but not in any way that suggests the licensor endorses you or your use.
- No additional restrictions** — You may not apply legal terms or technological measures that legally restrict others from doing anything the license permits.

Author Self-Archiving Policy

JAPS is a [RoMEO green journal](#). The journal allows the authors to self-archive pre-print, post-print, and publisher's version of the articles in any OAI-compliant repository.

Plagiarism Policy

The authors should ensure the originality of their contents while preparing a manuscript draft. In case the authors have used the work and/or words of others this must be appropriately cited or quoted. All the articles submitted to JAPS shall be screened for plagiarism using iThenticate (online plagiarism detection software). In case, plagiarism is detected during the review/editorial process, such manuscript(s) will be rejected immediately. If the plagiarism is proven after publication, such manuscript(s) will be retracted from the journal and an appropriate announcement will be placed in this regard. We can also consider appropriate action against authors depending upon the seriousness of the case. It includes;

1. Debarring the authors from publication in the future.
2. We can bring such instances in notice of author's funding agencies, author's institutes (where they work) and to the original authors whose work has been plagiarized.

COPE flow charts (please follow the below link) shall also be considered while dealing with plagiarism complaints.

- [Suspected plagiarism in a submitted manuscript](#)
- [Suspected plagiarism in a published manuscript](#).

Retraction policy

JAPS follow the COPE Retraction Guidelines. (Link:<https://publicationethics.org/files/cope-retraction-guidelines-v2.pdf>)

Digital Preservation

All the contents published in JAPS are archived in [Portico](#) (an E-Journal Preservation Service), which provides permanent digital archiving for scholarly journals.

Preprints policy

- Preprints are defined as an author's version of a research manuscript prior to formal peer review at a journal, which is deposited on a public server. Journal encourages posting of preprints of primary research manuscripts on preprint servers, authors' or institutional websites. Preprints may be posted at any time during the peer-review process. Posting of preprints is not considered prior to publication and will not affect consideration at this journal.
- Authors should not assign copyright during the preprint process; authors should retain the copyright of their work when posting to a preprint server. Journal will consider for publication submissions that have previously been assigned CC-BY (-NC/-NC-ND) as preprints.
- Authors should disclose details of preprint posting, including DOI and licensing terms, upon submission of the manuscript to this journal. Once the preprint is published, it is the author's responsibility to ensure that the preprint record is updated with a publication reference, including the DOI and a URL link to the published version of the article on the journal website.

Conflicts of Interest

A conflict of interest defined as any relationship authors, reviewers or editors have, which interferes with the full and objective presentation, peer review, editorial decision-making, or publication of a manuscript. Conflicts of interest can be financial or non-financial, professional or personal, and can arise in relation to an organization or an individual.

Authors

When authors submit a manuscript of any type or format they are responsible for disclosing all relationships and activities that might bias or be seen to bias their work (e.g. employment, consulting fees, research contracts, stock ownership, patent licenses, honoraria, advisory affiliations, etc.). JAPS is committed to following [ICMJE recommendation on "Author Responsibilities—Conflicts of Interest"](#) in authors' conflict of interest issues. The ICMJE has developed a [Disclosure Form](#) to facilitate and standardize authors' disclosures, please [click this link to download the ICMJE COI form](#). Authors are required to upload the duly filled and signed form under the additional files section, during manuscript submission. They can also send it by email to editor@japsonline.com. Please follow the link to read the detailed ICMJE recommendations for [disclosure of financial and non-financial relationships and activities, and conflicts of interest](#).

Reviewers

Reviewers must disclose to journal editors any relationships or an activity that could bias their opinions of the manuscript and should recuse them from reviewing specific manuscripts if the potential for bias exists. Reviewers must not use knowledge of the work they're reviewing before its publication to further their own interests. Please use the review form provided at your reviewer login area to disclose the conflicts of interest.

Editors and Journal Staff

JAPS editors who make final decisions about manuscripts must have no personal, professional, or financial involvement in any of the issues they might judge. Other members of the editorial staff, if they participate in editorial decisions, must provide editors with a current description of their financial interests (as they might relate to editorial judgments) and recuse themselves from any decisions in which a conflict of interest exists. Editorial staff must not use information gained through working with manuscripts for private gain. When editors submit their own work to this journal, a colleague in the editorial office should manage the manuscript and the editor/author should recuse himself or herself from discussion and decisions about it.

Note: These policies are based upon the [ICMJE recommendations](#).

Additional Information

You may find the following useful resources to refer to for more information on Conflict of Interest policies, existing codes of practices and more general good practice in relation to journal publication ethics:

- View the [Guidelines published on good publication and the Code of Conduct by the Committee of Publication Ethics \(COPE\)](#).
- View the [guidelines published by the International Committee of Medical Journal Editors \(ICMJE\)](#).

Complaints and appeals

Authors & readers can submit their complaints/appeals to the editor or the publisher of the journal through email info@japsonline.com or through the complaint form provided on the contact us page [[Link](#)]. All the complaints will assign a unique complaint number by the publisher. Authors/readers can use complaint numbers for further follow-up.

If you do not receive a response to your complaint within 48 hrs or if you are not satisfied with the response please write to associate editor paraszee05@gmail.com or chief editor editor@japsonline.com. Please mention your unique complaint number in follow-up emails.

Disclaimer

The author(s) of each article appearing in this Journal is/are solely responsible for the content thereof; neither the Journal of Applied Pharmaceutical Science nor its editors or publisher nor anyone else involved in creating, producing, or distribution assumes any liability or responsibility for the accuracy, completeness, correctness or usefulness of any information provided in the Journal, nor shall they be liable for any direct, indirect, incidental, special, consequential or punitive damages arising out of

the use of the Journal of Applied Pharmaceutical Science. The journal remains neutral with regards to the jurisdictional claims in published institutional affiliations.

© 2011- 2022 Open Science Publishers LLP, unless otherwise stated.
By using this website, you agree to our [Terms and Conditions](#), [Privacy statement](#) and [Cookies](#) policy.



Editorial Board

CURRENT ISSUE

April, 2022

ARCHIVES

[READ MORE](#)Follow us  

Editor-in-Chief

Prof. Srinivas Mutalik

Coordinator, Centre for Drug Delivery Technologies,
Professor & Head, Department of Pharmaceutics, Manipal College of Pharmaceutical Sciences,
Manipal, India. Email: editor@japsonline.com [\[Website\]](#) [\[ORCID\]](#)

Interest: Novel drug delivery, Nanotechnology, pharmaceutical analysis.

Immediate Past Editor-in-chief

Dr. Pinaki Sengupta [\[Website\]](#) [\[View Profile\]](#)

Department of Pharmaceutical Analysis, National Institute of Pharmaceutical Education and Research - Ahmedabad,
Gandhinagar, India.

Interest: Pharmaceutical analysis, Spectroscopy, Bioanalysis, Pharmacokinetics, and bioequivalence.

Associate Editor

Paras Sharma [\[View Profile\]](#) [\[ORCID\]](#)

Department of Pharmacognosy, BVM College of Pharmacy, Gwalior, India.

Interest: Pharmacognosy, medicinal plants, antimicrobial resistance, pharmacy practice.

Advisory Board

Prof. Vinod Kumar Gupta [\[View Profile\]](#)

(Former Vice-Chancellor, Dr. Ram Manohar Lohia Awadh University, Faizabad, India)
Distinguished Professor, King Abdulaziz University, Jeddha, Saudi Arabia.

Interest: chemical synthesis, medicinal chemistry, pharmaceutical analysis.

Dr. K. Husnu Can Baser [\[View Profile\]](#)

Professor of Pharmacognosy, Near East University, Faculty of Pharmacy,
Head of the Department of Pharmacognosy, Nicosia, N. Cyprus, Turkey.

Interests: pharmacognosy, medicinal and aromatic plants, natural products, essential oils.

Prof. Guoyin Kai [\[View Profile\]](#)

Director, Laboratory of Medicinal Plant Biotechnology, College of Pharmaceutical Sciences
Zhejiang Chinese Medical University, Zhejiang Province, China.

Interests: biosynthesis of secondary metabolites, pharmacognosy, medicinal plants, plant biotechnology.

Prof. Imran Ali [\[View Profile\]](#)

Department of Chemistry, Jamia Millia Islamia, New Delhi, India.

Interests: nanotechnology, chemical synthesis, chromatographic & related separation techniques, analytical chemistry.

Prof. (Dr) Mainul Haque [\[View Profile\]](#)

The Unit of Pharmacology, Faculty of Medicine and Defence Health, Universiti Pertahanan,
National Defence University of Malaysia, Kem Perdana Sungai Besi, Kuala Lumpur, Malaysia.

Interests: pharmacology, clinical pharmacy, antimicrobial resistance, pharmacovigilance.

Roberto Pisano [\[View Profile\]](#)

Department of Applied Science and Technology, Politecnico di Torino, 24 corso Duca degli Abruzzi, Torino, Italy.

Interests: Pharmaceutical processing and formulation, Small molecules and biologics, Controlled Release.

Dr. Kirankumar Hullatti [\[View Profile\]](#)

Professor and Vice Principal, Cauvery College of Pharmacy, Mysore, India.

Interests: pharmacognosy, phytochemistry, medicinal and aromatic plants, natural products.

Prof. Saber A. Sakr [\[View Profile\]](#)

Faculty of Science, Menoufia University, Egypt.

Interests: pharmacology, toxicity studies, pharmaceutical biochemistry.

Prof. Shao Hong-Bo [[View profile](#)]

Qingdao University of Science & Technology, Qingdao, China.

Interests: pharmaceutical microbiology, biotechnology, nanotechnology.

Editorial Board

Dr. Subrahmanya Bhat K [[View Profile](#)]

Department of Chemistry, Manipal Institute of Technology, Manipal Academy of Higher Education, Manipal, India.

Interests: Organic Chemistry; Medicinal Chemistry; Chemical synthesis.

Prof. Roman Lesyk [[View Profile](#)]

Department of Pharmaceutical, Organic & Bioorganic Chemistry, Danylo Halytsky Lviv National Medical University, Lviv, Ukraine.

Interests: Pharmaceutical, medicinal and organic chemistry, drug design, synthesis of heterocyclic compounds as potential anticancer, anti-inflammatory, antiviral, antituberculosis and antimicrobial drugs.

Prof. Abdul Rohman [[View Profile](#)]

Department of Pharmaceutical Chemistry, Faculty of Pharmacy, Gadjah Mada University, Yogyakarta, Indonesia.

Interests: Pharmaceutical analysis, quality control, chemometrics, spectroscopy, chromatographic techniques.

Dr. Narendar Dudhipala [[View Profile](#)] [[Website](#)]

Department of Pharmaceutics and Drug Delivery, School of Pharmacy, University of Mississippi, USA.

Interests: Formulation development, preclinical evaluation of small molecules for ocular delivery and oral delivery, Tumor delivery, lipid nanoparticles, lyophilization.

Dr. Dinesh Kumar Mishra [[View Profile](#)]

Professor & Principal, Indore Institute of Pharmacy, Opposite IIM, Pithampur Road, Rau, Indore (M.P.), India.

Interests: Novel drug delivery systems, Nanomedicine, Vaccine, Transdermal Delivery, 3D Printing, Microneedles, Formulation and development.

Dr. Valery Dembitsky [[View Profile](#)]

Centre for Applied Research, Innovation & Entrepreneurship,

Lethbridge College, 3000 College Drive South Lethbridge, Canada.

Interests: Pharmaceutical Chemistry, Organic Synthesis, Liquid Chromatography, Analytical Chemistry, Cancer.

Dr. Eduardo Rocha [[View Profile](#)]

Professor, ICBAS-Institute of Biomedical Sciences Abel Salazar, University of Porto, 4050-313 Porto, Portugal.

Interests: Liver and Reproductive Toxicology, In vitro toxicology, cytotoxicity of bioactive extracts, metabolites derived from marine fungi and algae.

Dr. Pranav Kumar Prabhakar [[View Profile](#)] [[Website](#)]

Lovely Faculty of Applied Medical Sciences, Lovely Professional University, Phagwara, Punjab, India.

Interests: Combination therapy, secondary complications of diabetes, phytomedicine, cellular signaling.

Dr. Talha Bin Emran [[View Profile](#)]

Department of Pharmacy, BGC Trust University, Chittagong, Bangladesh.

Interests: Vaccinology, Applied Immunology, Phytomedicine, Natural Product Chemistry.

Dr. Bappaditya Chatterjee [[View Profile](#)] [[Website](#)]

Department of Pharmaceutics, SPPSPTM, SVKM's NMIMS, Mumbai, India.

Interests: Pharmaceutics, biopharmaceutics, Novel & Targeted Drug Delivery, Nanotechnology, amorphous dispersion, intranasal delivery systems.

Dr. Anoja Priyadarshani Attanayake [[View Profile](#)]

Department of Biochemistry, Faculty of Medicine, University of Ruhuna, Sri Lanka.

Interests: Bioactivity studies on medicinal plants, Clinical trials of herbal products, Nanonutraceuticals for diabetes, Preclinical studies on antidiabetic, nephroprotective agents.

Dr. Gurinder Singh [[View Profile](#)]

Formulation Scientist, Frankfurt am Main, Germany.

Interests: Nanocarriers, Resveratrol, Controlled delivery systems, DOE, in vitro/in vivo studies.

Dr. Uday Venkat Mateti [[View Profile](#)]

Dept. of Pharmacy Practice, NGSM Institute of Pharmaceutical Sciences,

Nitte (Deemed to be University), Mangaluru, Karnataka, India.

Interests: Patient Safety, Patient Reported Outcomes, Developing Pharmacoeconomics Modelling, Pharmaceutical care, Supportive Care, and Pharmacoepidemiology

Dr. Mrs. Karimunnisa S. Shaikh

Modern College of Pharmacy, Nigdi, Pune, India. [\[View Profile\]](#)

Interests: Polysaccharide-based drug delivery system, Nanosponge, Co-crystallization and co-amorphous systems, Nanotechnology, anticancer therapeutics, Nano drug delivery systems.

Prof. Alexander K. Nyarko [\[View Profile\]](#)

Professor of Pharmacology and Toxicology, Department of Pharmacology and Toxicology, School of Pharmacy, College of Health Sciences, University of Ghana, Legon, Ghana.

Interests: Natural products, pharmacology, biochemistry, medicinal plants.

Dr. Shazia Qasim Jamshed [\[View Profile\]](#)

Associate Professor Clinical Pharmacy and Practice, Faculty of Pharmacy, Universiti Sultan Zainal Abidin, Malaysia.

Interests: Clinical Pharmacy, Pharmacy Practice, Pharmacy education.

Prof. Oluwatoyin A. Odeku [\[Website\]](#) [\[View Profile\]](#)

Dean, Faculty of Pharmacy, University of Ibadan, Ibadan, Nigeria.

Interests: Pharmaceutics, biopharmaceutics, and pharmaceutical technology, Novel and targeted drug delivery, Nanotechnology, Excipients development.

Dr. Thirumal Kumar D [\[View Profile\]](#)

Assistant Research Director & Assistant Registrar (i/c), Meenakshi Academy of Higher Education and Research (Deemed to be University), Chennai, Tamil Nadu, India.

Interests: Bioinformatics, Structural biology, Drug Discovery, Drug Resistance, Molecular Docking.

Dr. Sameer Dhingra [\[View Profile\]](#) [\[Website\]](#)

Department of Pharmacy Practice, National Institute of Pharmaceutical Education and Research, Hajipur, India.

Interests: Pharmacy practice, clinical pharmacy, medication safety, rational use of drugs, antimicrobial stewardship, Pharmacovigilance.

Dr. Aysu YURDASIPER

Faculty of Pharmacy, Pharmaceutical Technology Department, Ege University, Izmir, Turkey.

Interests: Dermal delivery (topical, transdermal drug systems), controlled-release formulations, nanoparticles & microparticles for drug delivery, and nanomedicine in pulmonary delivery.

Dr. Wenyi Kang [\[View Profile\]](#)

Director, National R & D Center for Edible Fungus Processing Technology, Henan University, Kaifeng, China.

Dr. Ahmed Salih Sahib [\[View Profile\]](#)

Professor of Pharmacology and Toxicology,

Dean, College of Pharmacy, University of Kerbala, Karbala, Iraq.

Interests: Pharmacology, Toxicology, Biochemistry, antioxidants.

Dr. Dipankar Ghosh [\[View Profile\]](#)

Department of Biotechnology & Microbiology, JIS University, Agarpara, Kolkata, India.

Interests: Pharmaceutical Biotechnology, Antimicrobials, Microbial biosynthesis, antibiotics research.

Dr. Mohammad Javed Ansari [\[View Profile\]](#)

Department of Pharmaceutics, College of Pharmacy, Prince Sattam Bin Abdulaziz University, Al-Kharj, Saudi Arabia.

Interests: Pharmaceutics, nanopharmaceuticals, controlled-release formulations, microparticles for drug delivery, ocular delivery, bioavailability enhancement, pharmaceutical analysis.

Dr. Teerapol Srichana [\[View Profile\]](#)

Faculty of Pharmaceutical Sciences, Prince of Songkla University, Hat Yai, Songkla, Thailand.

Dr. Oluwafemi Omoniyi Oguntibeju [\[View Profile\]](#)

Department of Biomedical Sciences, Faculty of Health & Wellness Sciences, Cape Peninsula University of Technology, Bellville, South Africa.

Dr. Subrata Shaw

Broad Institute of MIT and Harvard, Cambridge, MA, USA. [\[View Profile\]](#)

Dr. U.S.Mahadeva Rao

Faculty of Medicine, Universiti Sultan Zainal Abidin, Malaysia. [\[View Profile\]](#)

Interests: Biochemistry, cancer, antioxidants, antidiabetic therapy.

Dr. Abdulhadi ALJAWISH

Polytech Lille, Institut Charles Violette, Université de Lille1, France. [\[View Profile\]](#)

Dr. Sai Prachetan Balguri

ORISE Fellow at U.S. FDA CDER/OPQ/OTR, New Hampshire Avenue Silver spring, MD, USA. [\[View Profile\]](#)

Dr. Farhad Shahsavar

Professor of Immunology, Lorestan University of Medical Sciences, Khorramabad, Iran. [\[View Profile\]](#)

Prof. Flavio Marques Lopes

UFG - School of Pharmacy, Goiânia, Brazil. [\[View Profile\]](#)

Dr. Elvis Adrian Fredrick Martis

Bombay College of Pharmacy, Kalina, Mumbai, India. [\[View Profile\]](#)

Note: All the above-mentioned editors and editorial board members are serving in their personal capacity and do not represent their affiliating institutions. Last Updated: 09/11/2021

© 2011-2022 Open Science Publishers LLP, unless otherwise stated.
By using this website, you agree to our [Terms and Conditions](#), [Privacy statement](#) and [Cookies](#) policy.




Current Issue

Volume: 12, Issue: 4, April, 2022




Rss Feed



Review Article: Effect of metformin treatment on inflammatory markers in type 2 diabetes mellitus—A systematic review and meta-analysis

Renuka Suvarna, Revathi P. Shenoy, M. Mukhyaprana Prabhu, Guruprasad Kalthur, Basavaraj S. Hadapad, Varashree Bolar Suryakanth

DOI: 10.7324/JAPS.2022.120401 Pages: 001-011

[\[Abstract\]](#) [\[Full Text PDF\]](#) [\[XML: Abstract + References\]](#)


Review Article: Chitosan and oral mucosal wound healing in dentistry—Journey from non-patented to patent introspection from 2010–2021

Meenakshi S, Raghunath N, Ramith Ramu


DOI: 10.7324/JAPS.2022.120402 Pages: 012-025

[\[Abstract\]](#) [\[Full Text PDF\]](#) [\[XML: Abstract + References\]](#)


Review Article: A review on the pharmacotherapy and case studies of hypercalcaemia and hyperoxaluria

Lika Ginanti Febriana, Sri Adi Sumiwi, Jutti Levita


DOI: 10.7324/JAPS.2022.120403 Pages: 026-036

[\[Abstract\]](#) [\[Full Text PDF\]](#) [\[XML: Abstract + References\]](#)


Review Article: Variations from SARS-CoV-2 to Omicron: A new threat knocking at World's door

Acharya Balkrishna, Akansha Rohela, Ashwani Kumar, Vedpriya Arya, Nazam Khan


DOI: 10.7324/JAPS.2022.120404 Pages: 037-043

[\[Abstract\]](#) [\[Full Text PDF\]](#) [\[XML: Abstract + References\]](#)


Research Article: Development of *Panax notoginseng* saponins-loaded orodispersible films: A potential approach to enhance delivery efficacy in older adults

Khanh Van Nguyen, Thu Kim Dang, Ha Thai Pham, Binh Thi Thanh Nguyen, Giang Thi Thu Vu, Hai Thanh Nguyen, Hiep Tuan Tran

DOI: 10.7324/JAPS.2022.120405 Pages: 044-053

[\[Abstract\]](#) [\[Full Text PDF\]](#) [\[XML: Abstract + References\]](#)


Research Article: Natural extracts derived from marine sponge-associated bacteria with diverse nonribosomal peptide synthetase genes have anticancer activities

Jepri Agung Priyanto, Widya Esti Purwaningtyas, Dewi Ferantika, Dudi Tohir, Rika Indri Astuti, Aris Tri Wahyudi

DOI: 10.7324/JAPS.2022.120406 Pages: 054-063

[\[Abstract\]](#) [\[Full Text PDF\]](#) [\[XML: Abstract + References\]](#)


Research Article: Assessing public knowledge of clinical and social issues of sickle cell disease: A study in the Kumasi metropolis of Ghana

Richmond Arthur, George Asumeng Koffuor

DOI: 10.7324/JAPS.2022.120407 Pages: 064-070

[\[Abstract\]](#) [\[Full Text PDF\]](#) [\[XML: Abstract + References\]](#)

Research Article: Effect of ethanolic extract from *Piper sarmentosum* on antihyperglycemic activity and complications in normal and streptozotocin-induced diabetic rats

Nittiya Suwannasom, Chutamas Thepmalee, Krissana Khoothiam, Chonthida Thephinlap


CURRENT ISSUE

April, 2022


ARCHIVES

[READ MORE](#)


Follow us

DOI: [10.7324/JAPS.2022.120408](https://doi.org/10.7324/JAPS.2022.120408) Pages: 071-079[[Abstract](#)] [[Full Text PDF](#)] [[XML: Abstract + References](#)]**Research Article: The effect of glycerin and polyethylene glycol 400 as humectant on stability and antibacterial activity of nanosilver biosynthetic peel-off mask**

Dian Eka Ermawati , Agung Putu Surya, Rini Setyawati, Sukma Uswatun Niswah

DOI: [10.7324/JAPS.2022.120409](https://doi.org/10.7324/JAPS.2022.120409) Pages: 080-089[[Abstract](#)] [[Full Text PDF](#)] [[XML: Abstract + References](#)]**Research Article: Phytochemical, antioxidant, and antimicrobial analysis of *Trichoderma asperellum* isolated from ascidian *Eudistoma* sp**


Deiske Adeliene Sumilat, Rosita Anggreini Junita Lintang, Suzanne Lidya Undap, Ahmad Akroman Adam, Trina Ekawati Tallei

DOI: [10.7324/JAPS.2022.120410](https://doi.org/10.7324/JAPS.2022.120410) Pages: 090-095[[Abstract](#)] [[Full Text PDF](#)] [[XML: Abstract + References](#)]**Research Article: Phytochemical screening, docking study, and *in vitro* assessment of cytotoxicity against MCF7 and 3T3 cell lines of Indonesian sea cucumber *Stichopus* sp.**


Teni Ernawati , Minarti Minarti, Galuh Widiyarti, Megawati Megawati, Faiza Maryani

DOI: [10.7324/JAPS.2022.120411](https://doi.org/10.7324/JAPS.2022.120411) Pages: 096-107[[Abstract](#)] [[Full Text PDF](#)] [[XML: Abstract + References](#)]**Research Article: Comparison of vitamin D status and physical activity related to obesity among tertiary education students**

Amelia Lorensia, Rivian Virlando Suryadinata, Irene Andriana Inu

DOI: [10.7324/JAPS.2022.120412](https://doi.org/10.7324/JAPS.2022.120412) Pages: 108-118[[Abstract](#)] [[Full Text PDF](#)] [[XML: Abstract + References](#)]**Research Article: A herbal combination attenuates muscle atrophy and cancer cachexia: A preclinical study**


Mohammed Mukhram Azeemuddin, Chamallamudi Mallikarjuna Rao, Mohamed Rafiq, Mallappa Onkaramurthy, Prachi Singh, Mirza Rizwan Baig, Uddagiri Venkanna Babu

DOI: [10.7324/JAPS.2022.120413](https://doi.org/10.7324/JAPS.2022.120413) Pages: 119-126[[Abstract](#)] [[Full Text PDF](#)] [[XML: Abstract + References](#)]**Research Article: Designing of next-generation dihydropyridine-based calcium channel blockers: An *in silico* study**


Sujoy Karmakar, Hriday Kumar Basak, Uttam Paswan, Arun Kumar Pramanik, Abhik Chatterjee

DOI: [10.7324/JAPS.2022.120414](https://doi.org/10.7324/JAPS.2022.120414) Pages: 127-135[[Abstract](#)] [[Full Text PDF](#)] [[XML: Abstract + References](#)]**Research Article: Acute toxicity evaluation and immunomodulatory potential of hydrodynamic cavitation extract of citrus peels**

Dyaningtyas Dewi Pamungkas Putri, Gergorius Gena Maran, Yuni Kusumastuti, Ratna Asmah Susidarti, Edy Meiyanto, Muthi Ikawati


DOI: [10.7324/JAPS.2022.120415](https://doi.org/10.7324/JAPS.2022.120415) Pages: 136-145[[Abstract](#)] [[Full Text PDF](#)] [[XML: Abstract + References](#)]**Research Article: Heat shock protein-70 expression in CSCs tumor-associated macrophages induced by *Typhonium flagelliforme* tuber extract**

Sugeng Ibrahim, Agung Putra, Nur Dina Amalina, Iqbal Pahlevi Adeputra Nasution

DOI: [10.7324/JAPS.2022.120416](https://doi.org/10.7324/JAPS.2022.120416) Pages: 146-152[[Abstract](#)] [[Full Text PDF](#)] [[XML: Abstract + References](#)]**Research Article: Imbalances of key parameters needed for energy-yielding processes in the serum of tramadol-treated male rats**

Husam Abazid , Sahar Jaffal, Mahmoud Abu-Samak

DOI: [10.7324/JAPS.2022.120417](https://doi.org/10.7324/JAPS.2022.120417) Pages: 153-158[[Abstract](#)] [[Full Text PDF](#)] [[XML: Abstract + References](#)]



Research Article: ADME studies of TUG-770 (a GPR-40 inhibitor agonist) for the treatment of type 2 diabetes using SwissADME predictor: *In silico* study

Khaldun M. Al Azzam, El-Sayed Negim, Hassan Y. Aboul-Enein

DOI: [10.7324/JAPS.2022.120418](https://doi.org/10.7324/JAPS.2022.120418) Pages: 159-169

[\[Abstract\]](#) [\[Full Text PDF\]](#) [\[XML: Abstract + References\]](#)




Mini Review: Ethnopharmacology, phytochemistry, and biological activity review of *Aleurites moluccana*

Aliefman Hakim, Jamaluddin Jamaluddin, Syarifah Wahidah Al Idrus, Abdul Wahab Jufri, Baiq Nila Sari Ningsih

DOI: [10.7324/JAPS.2022.120419](https://doi.org/10.7324/JAPS.2022.120419) Pages: 170-178

[\[Abstract\]](#) [\[Full Text PDF\]](#) [\[XML: Abstract + References\]](#)



Short Communication: Synthesis and *in vitro* cytotoxic activity of novel indazole analogues of curcumin against MCF-7, HeLa, WiDr, and vero cell lines

Hariyanti Hariyanti, Arry Yanuar, Kusmardi Kusmardi, Hayun Hayun

DOI: [10.7324/JAPS.2022.120420](https://doi.org/10.7324/JAPS.2022.120420) Pages: 179-184

[\[Abstract\]](#) [\[Full Text PDF\]](#) [\[XML: Abstract + References\]](#)

Letter to the Editor: Evacuation of yeast and fungal cells using the Sponge-Like Protocol: A new approach for fungal control

Amro Abd Al Fattah Amara, Nawal Abd El-Baky

DOI: [10.7324/JAPS.2022.120421](https://doi.org/10.7324/JAPS.2022.120421) Pages: 185-186

[\[Abstract\]](#) [\[Full Text PDF\]](#) [\[XML: Abstract + References\]](#)

© 2011-2022 Open Science Publishers LLP, unless otherwise stated.

By using this website, you agree to our [Terms and Conditions](#), [Privacy statement](#) and [Cookies](#) policy.



Comparison of vitamin D status and physical activity related to obesity among tertiary education students

by Amelia Lorensia

Submission date: 05-Apr-2022 09:10PM (UTC+0700)

Submission ID: 1802454828

File name: 1._isi.pdf (330.44K)

Word count: 8803

Character count: 43331



Comparison of vitamin D status and physical activity related to obesity among tertiary education students

Amelia Lorensia^{1*}, Rivan Virlando Suryadinata², Irene Andriana Inu¹

¹Departement of Community Clinical Pharmaceutical, Faculty of Pharmacy, University of Surabaya (Universitas Surabaya). Jl. Raya Kalirungut, Rungkut District, Surabaya City, East Java 60293, Indonesia.

²Departement of Public Health Science, Faculty of Medicine, University of Surabaya (Universitas Surabaya). Jl. Raya Kalirungut, Rungkut District, Surabaya City, East Java 60293, Indonesia.

5

ARTICLE INFO

Received on: 30/09/2021
 Accepted on: 27/12/2021
 Available Online: 05/04/2022

Key words:

Physical activity, obesity, student, vitamin D status.

ABSTRACT

University students are young adults who are productive and have relatively high physical activity. Obesity is caused by imbalance between excessive food intake and low physical activity and is a risk factor for vitamin D deficiency. This research aimed to explore vitamin D status and physical activity in the obese and nonobese group and the relationship between vitamin D status and physical activity. The research design used was cross-sectional using the International Physical Activity Questionnaire and vitamin D status questionnaires. The research location was the campus area of Surabaya and was carried out from March 2018 to January 2019. This study involved 143 respondents consisting of 53 people (49.65%) in the obese group and 72 people (50.34%) in the nonobese group. The results showed that there was a significant difference in physical activity ($p = 0.047$) and vitamin D status ($p < 0.05$) between the nonobese and obese groups. There was a significant relationship ($p < 0.05$) between vitamin D status and physical activity, although the relationship was low (correlation coefficient = 0.326). Therefore, increased physical activity can improve vitamin D status while still paying attention to other factors that affect vitamin D, such as diet and lifestyle. The condition of vitamin D also needs to be confirmed by measuring the 25(OH)D blood test.

52

INTRODUCTION

The World Health Organization (WHO) declared obesity a global epidemic (Jackson *et al.*, 2020; Jayawardena *et al.*, 2020). Overweight or obesity is a serious public health problem in Indonesia with a continuing increase in its prevalence (Rachmi *et al.*, 2017). Based on the results of Basic Health Research in Indonesia (2018), it is stated that the prevalence of obesity in adults has increased since the three Riskesdas periods, i.e., by 10.5% (2007), 14.8% (2013), and 21.8% (2018) (Damayanti *et al.*, 2019). Obesity is caused by various factors including genetic, environmental, psychological, and health factors caused by the use of certain drugs and diseases, as well as factors in the development of adipose tissue (Fruh *et al.*, 2017; Panuganti *et al.*, 2021). However, the main cause of obesity is an imbalance between excessive food intake and low physical activity (Jehan *et al.*, 2020;

Romieu *et al.*, 2017). Prevention of obesity can be achieved by changing lifestyle (Salam *et al.*, 2020), such as increasing physical activity by exercising diligently and maintaining a healthy diet (Elliot and Hamlin, 2018; Koehler and Drenowatz, 2019).

High physical activity can be useful in the process of preventing obesity (Leticia *et al.*, 2019; Kim *et al.*, 2017; Lorensia *et al.*, 2021); on the other hand, low physical activity results in an increased prevalence of the risk of noncommunicable diseases such as heart disease, diabetes, and cancer (Ahmed *et al.*, 2019; Saqib *et al.*, 2020). Physical activity is defined as movement of the body by the muscles and skeleton which requires energy expenditure. There are two categories, namely moderate (moderate) and vigorous (strong) physical activity (Füzéki and Banzer, 2018; Hamer and Stamatakis, 2018). Both these types of physical activity provide health benefits and also affect vitamin D status (Skalska *et al.*, 2019; Wiciński *et al.*, 2019). A person with vigorous and moderate activity has a higher vitamin D status than a person with low physical activity (Hall *et al.*, 2018; Ces, 2019; Silva *et al.*, 2019). Vitamin D status is also seen by dietary intake of foods containing vitamin D and daily lifestyle habits

47

*Corresponding Author

Amelia Lorensia, Universitas Surabaya, Surabaya, Indonesia.
 E-mail: amelia.lorensia@gmail.com

47

2

(Larson-Meyer *et al.*, 2019; Suryadinata *et al.*, 2020; Zadka *et al.*, 2018), such as the use of sunscreen, low intake of milk (Aji *et al.*, 2019), and avoiding sun exposure which causes a decrease in the synthesis of vitamin D by the skin resulting in vitamin D deficiency (Ramasamy, 2020; Suryadinata *et al.*, 2020; Webb *et al.*, 2018).

More than one billion people worldwide are deficient in vitamin D (Sizar *et al.*, 2021). About 30%–50% of children and adults in the United Arab Emirates, Australia, India, and Lebanon are deficient in vitamin D (Zarooni *et al.*, 2019). The risk of vitamin D deficiency can occur in dark-skinned people, elderly patients, pregnant women, babies, people who do not get enough sun exposure, and obese individuals (Sizar *et al.*, 2021; Suryadinata *et al.*, 2021; Wacker and Holick, 2013). The body can get vitamin D by consuming milk, fish and fish oil, egg yolks, liver, and yeast (Maurya and Aggarwal, 2017; Polzonetti *et al.*, 2020). Exposure to sunlight, and taking vitamin D supplements (Kift *et al.*, 2018; Lee *et al.*, 2020). Vitamin D functions in the formation and maintenance of bones, so vitamin D deficiency results in bone abnormalities, namely rickets in children and osteoporosis in adults (Michigami, 2018; Takeuchi, 2018; Umar *et al.*, 2018). Vitamin D deficiency also contributes to the development of cancer, especially breast, prostate, and colon cancer as well as heart disease, stroke, and autoimmune diseases (Charoenngam and Holick, 2020; Hassanailou *et al.*, 2017; Wang *et al.*, 2017).

Early identification of vitamin D deficiency can be used as a preventive measure related to the risk of vitamin D deficiency (Aji *et al.*, 2019; Pilz *et al.*, 2019; Sizar *et al.*, 2021). The indicators used to determine the status of vitamin D include blood tests, dietary food intake containing vitamin D (Aji *et al.*, 2019; Lorensia *et al.*, 2020a; Pilz *et al.*, 2019), and questionnaires (Lelek-Berquist *et al.*, 2009; Larson-Meyer *et al.*, 2019). Evaluation of vitamin D status using a questionnaire is easier to carry out and more economical than blood serum examination. In addition, when compared to blood tests, clinical questionnaires can be used to identify long-term vitamin D status (Aji *et al.*, 2019; Annweiler *et al.*, 2015; Bolek-Berquist *et al.*, 2009). The questionnaire can be used to identify patients who are at high risk of experiencing vitamin D deficiency (Bolek-Berquist *et al.*, 2009). Obesity as a risk factor for vitamin D deficiency (Paschou *et al.*, 2019) needs to get prevention efforts as early as possible considering the dangers of vitamin D deficiency and low physical activity as causes of obesity which affect the development of chronic diseases and decreased quality of life (Kavadar *et al.*, 2015; Kim *et al.*, 2018; Oh *et al.*, 2017).

This study involved young adult respondents because adults were of productive age and relatively had high physical activity. However, nowadays, there are many young adults who actually experience vitamin D deficiency and rarely do physical activity (Lorensia *et al.*, 2019). This research was conducted in university students because education can affect a person's physical activity and lifestyle (Karska *et al.*, 2021). It was, therefore, necessary to identify vitamin D status and physical activity in obese and nonobese adults so that pharmacists can provide education to the public, both obese and nonobese adults, as a form of health workers' responsibility in early prevention efforts related to the risk of deficiency of vitamin D and other health problems.

METHODS

Research design

The design of this study was cross-sectional and was conducted on a group of students with nonobese and obese nutritional status in the campus area of the University of Surabaya. The research location chosen to conduct the research was the campus area in South Surabaya, Rungkut district, East Java, Indonesia, and the study was carried out from March 2018 to January 2019. This study was approved with Ethical Test No. 034/KE/I/2018 at Universitas Surabaya.

Research variable

The independent variable in this study was a group of obese and nonobese respondents. The dependent variables in this study were vitamin D status and physical activity. Vitamin D status can be defined as the presence of vitamin D in a certain amount in the body which was influenced by the intake of foods containing vitamin D, sun exposure, and consumption of vitamin D supplements. A person was said to be positive (+) (at risk) of vitamin D deficiency if the total answer score was ≥ 15 and negative (-) if the total answer score was < 15 . Vitamin D deficiency is a health problem related to vitamin D status. The risk of vitamin D deficiency in nonobese and obese respondents is calculated using the prevalence odds ratio equation.

Physical activity was defined as body movement by the muscles and skeleton which requires energy expenditure, including activities carried out while working, playing, doing household chores, traveling, and recreation. Physical activity in adults can be measured using the International Physical Activity Questionnaire (IPAQ) (Cleland *et al.*, 2018; Kavadar *et al.*, 2015; Sembler *et al.*, 2020). This can be measured by asking seven questions related to daily activities. The data scale obtained from the measurement of physical activity was an ordinal data scale, where the results obtained from the patient questionnaire were categorized into low, medium, and high physical activity (Table 1).

Obesity was characterized by the accumulation of excess body fat tissue (De Lorenzo *et al.*, 2019). Determination of obesity status in individuals was based on the measurement of body mass index (BMI), a person's body weight (in kg) divided by the square of height (in m). The International Association for Study of Obesity Organization, International Obesity Task Force, and WHO determined the category of obesity for the Asian population if the BMI was ≥ 25 kg/m². Therefore, in this study, BMI points ≥ 25 kg/m² were used to determine the obesity category.

Population and sample

The population used in this study was all students at the University of Surabaya with a nutritional status of obesity or nonobesity. The sample used in this study was active students at the University of Surabaya with a nutritional status conditions of obesity or nonobesity who met the criteria, including 18–25-year-olds, those with no special diet (such as fasting or vegetarianism), those who did not have certain diseases like cardiovascular disorders such as angina, renal, and hepatic disorders (such as cirrhosis of the liver), and individuals who have no motoric problems. The sampling technique used was the purposive

Table 1. Guidelines for assessment of physical activity based on the IPAQ.

Type of physical activity	Description
	No reports of physical activity carried out
Low physical activity	Activities carried out not sufficient to meet the criteria for the type of moderate physical activity
Moderate physical activity (one of the three criteria)	Within 24 days or more doing high-intensity (vigorous) activity for 20 minutes per day Within 5 days or more doing moderate-intensity activity (moderate) or walking for 30 minutes
High physical activity (one of the three criteria)	Within 5 days or more doing a combination of walking, moderate-intensity activity, and vigorous activity of at least 600 MET-minute per week Performing vigorous activity for 3 days with a total MET accumulation of at least 1,500 MET-minute per week For 7 days or more doing a combination of walking and moderate-intensity or vigorous activity at a minimum of 3,000 MET-minute per week

Description: MET (multiples of the resting metabolic rate): the amount of energy released that is equivalent to 1 kcal/kg/hour (Cleland *et al.*, 2018; Kavadar *et al.*, 2015; Sember *et al.*, 2020).

sampling method, in which the sampling process was based on previously known characteristics or characteristics of the population and the consideration of the researcher.

Data collection methods and analysis

Measuring instruments used in the study were the vitamin D status questionnaire to identify vitamin D status and the IPAQ for physical activity data. Measurements of weight and height were carried out using a digital weight scale and a Microtoise stature meter, respectively. There were two types of questionnaires, namely the vitamin D status questionnaire (Bolek-Berquist *et al.*, 2009; Cairncross *et al.*, 2017; Nabak *et al.*, 2014) and the IPAQ (Cleland *et al.*, 2018; Kavadar *et al.*, 2015; Sember *et al.*, 2020).

We conducted a trial or pilot study where data collection was preceded by testing the research instruments on 30 respondents. The questionnaire was validated using internal and external validity methods. Internal validity was enforced in a review step based on professional opinion judgments in the field of community pharmacy, while external validation was carried out by testing the questionnaire on the subject. Then, the results were analyzed using the Statistical Package for the Social Sciences (SPSS) version 24.0 computer program. The questionnaire was said to be valid if the value of $r_{\text{count}} > 0.361$. The questionnaire was rehabilitated using a single trial method technique with the alpha method (Cronbach), namely testing the reliability of the questionnaire once on respondents who have criteria such as research subjects, but these subjects are not used as samples. A reliability test can be carried out on valid questions. The calculation of the reliability coefficient uses the SPSS version 24.0 software. The questionnaire was declared reliable if the Cronbach's alpha value was >0.70 .

Comparative analysis of the risk of vitamin D deficiency was based on vitamin D status in the obese and nonobese groups of respondents using the prevalence odds ratio. Meanwhile, differences in vitamin D status and physical activity in the obese and nonobese respondent groups used the chi-square test. We then proceeded to test the relationship between vitamin D status and

physical activity in the groups of obese and nonobese respondents using a contingency coefficient.

RESULTS

The data obtained in this study were obtained through a vitamin D status questionnaire and the IPAQ. The data collection process uses instruments that are valid and reliable. Based on the data obtained, there were 143 respondents consisting of 2 groups, namely the obese group consisting of 71 people (49.65%) and the nonobese group consisting of 72 people (50.34%).

Validity and reliability of the IPAQ

The original IPAQ was available in English. The validation process was carried out by translating the questionnaire into Indonesian and then giving it to three professional judges in the field of community pharmacy. The validity was enforced in the analysis step based on the opinion of professional judges in the field of community pharmacy.

Validity and reliability of the vitamin D status questionnaire

The reliability of the questionnaire was determined based on Cronbach's alpha value. The questionnaire was declared reliable if Cronbach's alpha value was >0.60 . The Cronbach's alpha value of the vitamin D status questionnaire was 0.735, so the questionnaire was declared reliable. The questionnaire validation process used the SPSS version 24.0 program. The validation process was carried out on 33 respondents. Based on the Pearson product-moment r_{table} with a significance of 0.05 with a two-sided test and the amount of data ($n = 30$), it was known that r_{table} value was 0.361 so that $r_{\text{count}} > 0.361$, and it can be concluded that the questionnaire was valid.

Respondent characteristics

In this study, the respondents included were students with a stratum 1 education level with an age range of 18–25 years. Table 2 explains the distribution of respondents' age, sex, and BMI.

The results of the answers to the two groups of respondents were classified into sections regarding sun exposure

Table 2. Characteristics of respondents.

Characteristics	Nonobese group (n = 72)		Obese group (n = 71)		
	Frequency	Percentage (%)	Frequency	Percentage (%)	
Gender	Male	19	26.38	38	53.52
	Female	53	73.61	33	46.47
Age (years)	18	5	7.04	6	8.33
	19	14	19.72	7	0.72
	20	17	23.94	22	30.56
	21	20	28.17	14	19.44
	22	11	15.49	15	20.83
	23	4	5.63	4	5.56
	24			3	4.17
	25			1	1.39
BMI (kg/m ²)	Thin (<18.50)	12	16.66		
	Normal (18.50–22.90)	40	55.55		
	At risk (23.00–24.90)	20	27.77		
	Obese type I (25.00–29.90)			42	59.15
	Obese type II (≥30.00)			29	40.84
	Respiratory tract disorders	3	4.16	6	8.45
History of disease ^a	Digestive tract disorders	13	18.05	9	12.67
	Blood disease	5	6.94	3	4.22
	Infectious disease	7	9.72	5	7.04
	Others	9	12.50	4	5.63
	Does not have a history of disease	45	62.50	48	67.60
History of medicine ^a	Multivitamins	7	9.72	6	8.45
	Antibiotic	3	4.16	0	0.00
	Immune system booster	3	4.16	2	2.81
	Digestive tract disorders drugs	1	1.38	2	2.81
	Others	9	12.50	9	12.67
Does not use drugs or supplements	52	72.22	52	73.23	

^a The answer can be more than 1 in the category.

(Table 3), foods containing vitamin D (Table 4), and signs of vitamin D deficiency (Table 5).

The nonobese group had a more positive vitamin D status (≥ 15) (76.39%) than the obese group. At the same time, the level of activity in the nonobese group was dominated by moderate (55.55%) compared to the obese group, which was dominated by low activity levels (61.97%) (Table 6).

Based on the results of the analysis with the chi-square test to see the difference in vitamin D status between the nonobese and obese respondent groups, it was found that there was a significant difference (p -value < 0.05) in the vitamin D status of the nonobese and obese groups of respondents (Table 7).

Based on the results of the analysis with the chi-square test to see the difference in physical activity between the nonobese and obese respondents, it was shown that there was a significant difference between the two (p -value < 0.05) (Table 8).

Based on the contingency coefficient correlation test for vitamin D status and physical activity in the nonobese and obese respondent groups, approximate significance value of 0.000 ($p = 0.000$) indicated that there was a significant relationship between

vitamin D status and physical activity and the relationship has a correlation value of 0.326, so it can be concluded that there is a significant relationship between physical activity and vitamin D status in the nonobese and obese respondent groups, but the relationship was low or weak (Table 9).

DISCUSSION

Respondents who were involved in this study were 143 people and consisted of male sex at 39.86% (57 people) and female sex at 60.13% (86 people). The group of respondents in the category of male obesity were as many as 53.52% (38 people) and female sex as many as 46.47% (33 people) and for respondents with nonobesity categories, male respondents as many as 26.38% (19 people) and women 73.61% (53 people). Gender can affect vitamin D status; the female sex as a low vitamin status compared to the male sex (Annweiler *et al.*, 2017; AlQuaiz *et al.*, 2018; Sanghera *et al.*, 2017) due to avoiding sun exposure and dress habits, one of the main factors (Kift *et al.*, 2018; Lee and Lee, 2020) such as wearing long-sleeved shirts, long skirts, and headscarves (Judistiani *et al.*, 2019; Pulungan *et al.*, 2021). Gender

Table 3. Distribution of respondents' answers regarding length of exposure and hours of sun exposure.

Component in question	Nonobese group (n = 72)		Obese group (n = 71)			
	Frequency	Percentage (%)	Frequency	Percentage (%)		
	Duration of exposure and hours of sun exposure					
Duration of exposure (minute)	<15	20	27.77	21	29.57	
	>15	52	72.22	50	70.42	
Hours of exposure	07.00–09.00 am and 03.00–05.00 pm	35	48.61	41	57.74	
	10.00–11.00 am	37	51.38	30	42.25	
Use of protective equipment						
Use of protective equipment	Does not use protective equipment	5	6.94	15	21.12	
	Umbrella	8	11.11	8	11.26	
	Using Hat	7	9.72	8	11.26	
	Using protective equipment ^a Jacket	52	72.22	56	78.87	
	Sunblock/sun cream	48	66.66	40	56.33	
	Others (gloves, masks, etc.)	6	8.33	4	5.63	
How often to use	Always or every day	46	63.88	34	47.88	
	Sometimes	10	13.88	15	21.12	
	Rarely	11	15.27	17	23.94	
Applications use of protective equipment	Never	5	6.94	5	7.04	
	When leaving the house/boarding house	34	47.22	35	49.29	
	After 12 pm	28	38.88	25	35.21	
	When driving	9	12.50	10	14.08	
	Time at the beach	1	1.38	1	1.40	
Use of closed clothing and body parts that are protected from sun exposure						
Use of closed clothing	Not wearing closed clothes	30	41.66	44	61.97	
	Using closed clothes	42	58.33	27	38.02	
	Description of closed clothing Using the hijab	12	16.66	3	4.22	
	Jackets and trousers	17	23.61	14	19.71	
Body part covered ^a	Long sleeves and trousers	13	18.05	10	14.08	
	Face	45	62.50	38	53.52	
	Hand	49	68.05	45	63.38	
	Sleeve	41	56.94	45	63.38	
	Feet	35	48.61	30	42.25	
	Back and shoulders	72	100.00	71	100.00	
Use of sunblock/sun cream						
Use of sunblock/sunscreen cream	Does not use sunblock/sun cream	24	33.33	31	43.66	
	Menggunakan sunblock/krim tabir surya	48	66.66	40	56.33	
How often to use sunblock/sun cream	Always or every day	27	37.50	10	14.08	
	Often	4	5.55	6	8.45	
	Sometimes	6	8.33	9	12.67	
	Rarely	11	15.27	15	21.12	
Duration of use of sunblock/sunscreen cream	1–3 hours	28	38.88	28	39.43	
	3–6 hours	8	11.11	4	5.63	
	>6 hours	12	16.66	8	11.26	
Repeated use of sunblock/sun cream in a day	No	62	86.11	67	94.36	
	Yes	<4 hours	1	1.38	1	1.40
		>4 hours	9	12.50	3	4.22
	Yes	50	69.44	38	53.52	
Use of cosmetics containing SPF	No	14	19.44	28	39.43	
	Does not know	8	11.11	5	7.04	

^a One respondent can have more than one.

Table 4. Distribution of respondents' answers regarding foods containing vitamin D.

Component in question		Nonobese group (n = 72)		Obese group (n = 71)	
		Frequency	Percentage (%)	Frequency	Percentage (%)
Fish					
Consumption of fish in a week	Ate fish	53	73.61	47	66.19
	Did not eat fish	19	26.38	24	33.80
Type of fish consumed	Fish high in vitamin D (tuna, salmon, mackerel)	39	54.16	37	52.11
	Fish did not contain much vitamin D (salted fish, pindang, mujaer)	33	45.83	34	47.88
How the consumed fish was processed	Fried	48	66.66	50	70.42
	Burned	7	9.72	10	14.08
	Sushi	2	2.77	5	7.04
	Boiled	2	2.77	3	4.22
	Other	11	15.27	15	21.12
Where the food (fish) was acquired	Cook by myself	7	9.72	9	12.67
	Restaurant	27	37.50	40	56.33
	Catering	2	2.77	1	1.40
	Supermarket	20	27.77	16	22.53
Purpose of consuming fish	Love the taste	49	68.05	46	64.78
	Get health benefits	23	31.94	28	39.43
	Lose weight	3	4.16	6	8.45
	Price lower or more affordable	9	12.50	8	11.26
	Incidentally a fish dish available to eat	24	33.33	23	32.39
	Other	1	1.38	2	2.81
Egg and milk					
Consumption of milk (in the past 1 week)	Did not consume milk	20	27.77	25	35.21
	Consumed milk	52	72.22	46	64.78
Consumption of egg (in the past 1 week)	Did not consume egg	64	88.88	68	95.77
	Consumed egg	8	11.11	3	4.22
The portion of the egg consumed	Egg whites	9	12.50	1	1.40
	Egg yolk	0	0.00	1	1.40
	White and yellow	63	87.50	69	97.18
How the eggs consumed was processed	Fried	62	86.11	68	95.77
	Boiled	17	23.61	16	22.53
	Other	3	4.16	1	1.40
Behavior frequency of consumption of food and beverages containing vitamin D					
Egg	Often	18	25.00	29	40.84
	Rare	54	75.00	42	59.15
Fish	Often	5	6.94	5	7.04
	Rare	67	93.05	66	92.95
Milk	Often	19	26.38	21	29.57
	Rare	53	73.61	50	70.42
Behavior of consumption of fish oil and vitamin D supplements					
Consumption of fish oil	Did not consume fish oil	64	88.88	70	98.59
	Consumed fish oil	8	11.11	1	1.40
Consumption of supplements	Did not consume vitamin D supplements	52	72.22	52	73.23
	Consumed vitamin D supplements	2	2.77	0	0.00
	Containing vitamin D Not containing vitamin D	18	25.00	0	0.00

Table 5. Distribution of respondents' answers regarding signs of vitamin D deficiency.

Component in question	Nonobese group (n = 72)		Obese group (n = 71)	
	Frequency	Percentage (%)	Frequency	Percentage (%)
Muscle pain including lower back pain	27	37.50	30	42.25
Pain in the pelvis, back, and legs	23	31.94	28	39.43
Muscle weakness	8	11.11	10	14.08
Easy to get into a bad mood or get depressed easily	30	41.66	34	47.88
Low immunity, such as frequent colds in winter	23	31.94	26	36.61
Not at all experienced any of the conditions above	7	9.72	5	7.04

A respondent can experience more than one sign and symptom.

Table 6. Cross-tabulation of vitamin D status and physical activity in the nonobese and obese respondents.

Category		Nonobese group (n = 72)		Obese group (n = 71)	
		Frequency	Percentage (%)	Frequency	Percentage (%)
Vitamin D status (deficiency)	Negative	17	23.61	39	54.93
	Positive	55	76.39	32	45.07
Physical activity	Low physical activity	30	41.67	44	61.97
	Moderate physical activity	40	55.55	25	35.21
	High physical activity	2	2.78	2	2.82

Table 7. Risk analysis and differences test of vitamin D deficiency related to vitamin D status in nonobese and obese respondents.

Respondent group	Vitamin D status		Total	Odds ratio	Chi-square test
	Nondeficiency	Deficiency			
Nonobese (n = 72)	17	55	72	POR = 0.254 CI 95% (0.124–0.520)	$p = 0.000$ ($p < 0.05$)
Obese (n = 71)	39	32	71	Conclusion: The nonobese group had a 0.2-fold risk of vitamin D deficiency compared to the obese group	Conclusion: There were significant differences in vitamin D status in the obese and nonobese groups
Total	56	87	143		

Table 8. Risk analysis and differences test in physical activity in nonobese and obese respondents.

Respondent group	Physical activity			Total	Chi-square test
	Low	Moderate	High		
Nonobese (n = 72)	30	40	2	72	$p = 0.047$ ($p < 0.05$)
Obese (n = 71)	44	25	2	71	
Total	74	65	4	143	Conclusion: There was significant difference in physical activity in the obese and nonobese groups

Table 9. Results of analysis of the relationship between vitamin D status and physical activity.

Vitamin D status	Physical activity			Total	Correlation test
	Low	Moderate	High		
Deficiency	30	40	2	72	Conclusion: There was a significant relationship with vitamin D status and physical activity, but the relationship was low or weak (p = 0.000) KK = 0.326
Nondeficiency	44	25	2	71	
Total	74	65	4	143	

not only affects vitamin D status but also affects physical activity (Fernandes and Junior, 2017; Wiciński *et al.*, 2019). History of disease and use of drugs have a relationship with vitamin D status in the body. Certain pathological conditions, such as Crohn's disease, cystic fibrosis, celiac disease, and surgical removal of parts of the stomach or intestines that play a role in fat absorption, can trigger vitamin D deficiency. Anticonvulsant drugs are also called antiepileptic drugs. The long duration of some antiepileptic drugs, such as phenobarbital, phenytoin, and carbamazepine, and the antimicrobial agent rifampin can cause osteomalacia, which is a condition caused by vitamin D deficiency. This is triggered by the induction of 1,25-dihydroxyvitamin D catabolism (Saket *et al.*, 2021). Men tend to be more active than women, because they spend more time sitting and doing activities that smell of cosmetic. Women tend to be inactive compared to men. Women spend more time sitting and doing activities that smell of cosmetic while the male sex prefers to do sports so the activities carried out are higher than women (Sharifi *et al.*, 2016).

All respondents involved in this study were aged 18–25 years and were active students. This study uses young adults as respondents because based on previous research young adults are known to be at risk of vitamin D deficiency which is influenced by habits such as consumption of vitamin D below the recommended limit per day, lack of outdoor physical activity, increased use of sunblock/sunscreen cream, and low milk intake (Bolek-Berquist *et al.*, 2009). In addition to young adults, the elderly are also one of the age groups that have a low risk of vitamin D status. Low vitamin D status can cause muscle weakness (Gunton and Girgis, 2018; Sizar *et al.*, 2021) and increase the risk of fractures in the elderly (Napluetiwong *et al.*, 2020).

Obesity is a risk factor for vitamin D deficiency (Paschou *et al.*, 2012) based on the research by Kavadar *et al.*, (2015) that found a relationship between obesity and low vitamin D status, which has an impact on the risk of insulin resistance. The mechanism of vitamin D deficiency in obese patients is not precisely known, but several researchers have hypothesized that, related to the mechanism of vitamin D deficiency in obese patients, high fat accumulation results in a decrease in vitamin D bioavailability, where a lot of vitamin D is stored in fat cells in adipose tissue and increases vitamin D deficiency. Oxidative reactions, due to the fat-soluble nature of vitamin D, cause a decrease in the release of vitamin D from fat into the systemic circulation (Wimalawansa, 2019) in obese patients. One of the causes of obesity is excessive food intake, low physical activity (Romieu *et al.*, 2017), and lack of rest hours; besides that, there

is a relationship between age and workplace characteristics on the risk of becoming obese (Hruby *et al.*, 2016). Low physical activity is found in obese people (Kim *et al.*, 2017); therefore, increasing physical activity can be used as an appropriate step to prevent the risk of obesity and weight loss (Jakicic *et al.*, 2019).

Long exposure can affect the synthesis of vitamin D by the body; an exposure time of 15 minutes can synthesize vitamin D of 10,000–25,000 IU over a large area of the body surface and is characterized by a change in skin color to pink (Suryadinata *et al.*, 2021). In this study, the results showed that, in the two groups of respondents who were obese and nonobese, those who received longer sun exposure (>15 minutes) were the nonobese group of 36.36% (52 people) compared to the obese group of 36.36% (52 people). The most dominant group with less exposure to sunlight (<15 minutes) was the obese group at as much as 14.68% (21 people) and then the nonobese group at 13.98% (20 people). One of the factors that influence the duration of exposure to vitamin D synthesis is skin type (Wacker and Holick, 2013). Different skin types affect the length of time it takes the skin to synthesize vitamin D. Dark skin type (type IV skin) requires a long exposure time of six times longer than skin type I (Wacker and Holick, 2013). This is influenced by the pigment in the skin called melanin which functions as a natural sunscreen that can block the absorption of sunlight by the skin (Solano, 2020). Asian skin is in skin type IV or V; therefore, Asian people need a longer exposure time than other skin types. The recommended exposure time is more than 15 minutes or even up to 2 hours (Webb *et al.*, 2018). Exposure time is also an important factor in the process of vitamin D synthesis (Suryadinata *et al.*, 2021). It is known from the results of this study that the highest Ultraviolet B (UVB) intensity occurs from 11.00 am to 01.00 pm (Lor *et al.*, 2020b). UVB rays obtained from sunlight can help the synthesis of vitamin D in the skin (Wacker *et al.*, 2013; Lee and Lee, 2000).

One source of vitamin D is fish. The highest amount of vitamin D is found in fish liver (2–477 mg/kg), a maximum of 1,200 mg/kg and depending on the type of fish (Schmid and Walther, 2013). A total of 66.19% (47 people) of the obese respondent group, who consumed fish within the past week, consumed fish with vitamin D content of 52.11% (37 people), while in the nonobese respondent group of a total of 73.61% (53 people) who consumed fish as many as 54.16% (39 people) consumed fish with vitamin D content. The obese respondent group (7.04%) and the nonobese respondent group (6.94%) often consumed fish in the last week. The frequency of fish consumption can affect vitamin D status. In

Japan, fish is the main source for obtaining vitamin D (Nakamura *et al.*, 2002). The cooking process does not really affect the vitamin D content in food products because vitamin D is resistant to heat (Schmid and Walther, 2013), but the fish processing process also affects the amount of vitamin D.

Eggs are a source of vitamin D (Schmid and Walther, 2013). The obese respondents who consumed eggs were 95.77% (68 people), and the nonobese respondents who consumed eggs were 88.88% (64 people). The respondents from the obese group who often consumed eggs within 1 week were 40.84% (29 people), and the respondents from the nonobese group were 25% (18 people). Vitamin D in eggs is found in the yolk at as much as 20 IU of vitamin D3 or D2 per egg yolk (Schmid and Walther, 2013). In addition to eggs and fish, milk is also a fortification product that contains vitamin D (Lorensia *et al.*, 2020b; Schmid and Walther, 2013). Someone with low milk consumption can experience vitamin D deficiency (Bolek-Berquist *et al.*, 2009).

Based on the results of the analysis with the chi-square test, the *p*-value of 0.047 (*p*-value <0.05) can be concluded so it can be concluded that there is a significant difference in physical activity between the nonobese and obese groups of respondents. This is supported by evidence from previous studies which showed that there was a significant difference in physical activity in the normal group compared to the obese group, where the obese group had a longer sitting time than the nonobese group, and obesity had a significant relationship with low physical activity and physical function (Ng *et al.*, 2016).

In the analysis of the relationship between vitamin D status and physical activity using the contingency coefficient correlation test, we obtained an approximate significance value of 0.000 (*p*-value = 0.00) which indicates that there is a significant relationship between vitamin D status and physical activity and the relationship has a correlation value of 0.326. Based on the contingency coefficient table, the two variables have a low or weak relationship but are definitely related so it can be concluded that there is a relationship between vitamin D status and physical activity in the nonobese and obese respondent groups, but the relationship is low or weak. This is in accordance with previous research which showed that a person with low physical activity has a low vitamin D status (Aji *et al.*, 2019). In addition, high outdoor physical activity can increase vitamin D status (Lorensia *et al.*, 2021). Outdoor physical activity, which is in exposure to sunlight, is statistically significant in increasing vitamin D status (Lorensia *et al.*, 2020a). Therefore, physical activity and dietary intake of vitamin D should be increased in an effort to prevent vitamin D deficiency and other health problems such as metabolic risk factors (Lorensia *et al.*, 2020a). There are some limitations in this research, including the existence of variables that can affect the results of the study such as the uneven distribution of sexes between two groups of respondents being compared, there is no examination of serum 25(OH)D for determining vitamin D levels.

CONCLUSION

There is a significant difference in vitamin D status in the obese and nonobese groups of respondents, and there is a significant difference in physical activity in the obese and

nonobese groups of respondents. In addition, there is a significant relationship between vitamin D status and physical activity, but the relationship between the two is low or weak.

ACKNOWLEDGMENT

This research was funded by the Institute of Research and Community Service by Universitas Surabaya.

CONFLICTS OF INTEREST

The authors declare no conflicts of interest.

DATA AVAILABILITY

All data generated and analyzed are included within this research article.

PUBLISHER'S NOTE

This journal remains neutral with regard to jurisdictional claims in published institutional affiliation.

REFERENCES

- Ahmed SH, Meyer HE, Kjøllestad MK, Marjerrison N, Mdala I, Htet AS, Bjertness E, Madar AA. The prevalence of selected risk factors for non-communicable diseases in Hargeisa, Somaliland: a cross-sectional study. *BMC Public Health*, 2019; 19(1):878.
- Aji AS, Erwinda E, Yusrawati Y, Malik SG, Lipoeto NI. Vitamin D deficiency status and its related risk factors during early pregnancy: a cross-sectional study of pregnant Minangkabau women, Indonesia. *BMC Pregnancy Childbirth*, 2019; 19(1):183.
- AlQuaiz AM, Kazi A, Fouda M, Alyousefi N. Age and gender differences in the prevalence and correlates of vitamin D deficiency. *Arch Osteoporos*, 2018; 13(1):49.
- Annweiler C, Kabeshova A, Callens A, Paty ML, Duval GT, Holick MF. Self-administered vitamin D status predictor: older adults are able to use a self-questionnaire for evaluating their vitamin D status. *PLoS One*, 2017; 12(11):e0186578.
- Bolek-Berquist J, Elliott ME, Gangnon RE, Gemar D, Engelke J, Lawrence SJ, Hansen KE. Use of a questionnaire to assess vitamin D status in young adults. *Public Health Nutr*, 2009; 12(2):236–43.
- Cairncross CT, Stonehouse W, Conlon CA, Grant CC, McDonald B, Houghton LA, Eyles D, Camargo CA Jr, Coad J, von Hurst PR. Predictors of vitamin D status in New Zealand preschool children. *Matern Child Nutr*, 2017; 13(3):e12340.
- Charoenngam N, Holick MF. Immunologic effects of vitamin D on human health and disease. *Nutrients*, 2020; 12(7):2097.
- Cleland C, Ferguson S, Ellis G, Hunter RF. Validity of the International Physical Activity Questionnaire (IPAQ) for assessing moderate-to-vigorous physical activity and sedentary behaviour of older adults in the United Kingdom. *BMC Med Res Methodol*, 2018; 18(1):176.
- Damayanti RA, Sumarmi S, Mundiastuti L. The association between sleep duration with overweight and obesity among education staff in campus C of Universitas Airlangga. *Amerta Nutr*, 2019; 3(2):89–93.
- De Lorenzo A, Gratterer S, Gualtieri P, Cammarano A, Bertucci P, Di Renzo L. Why primary obesity is a disease? *J Transl Med*, 2019; 17(1):169.
- Elliott CA, Hamlin MJ. Combined diet and physical activity is better than diet or physical activity alone at improving health outcomes for patients in New Zealand's primary care intervention. *BMC Public Health*, 2018; 18(1):230.
- Fernandes MR, Junior DRBJ. Association between physical activity and vitamin D: a narrative literature review. *Rev Assoc Med Bras*, 2017; 63(6):550–6.
- Fruh SM. Obesity: risk factors, complications, and strategies for sustainable long-term weight management. *J Am Assoc Nurse Pract*, 2017; 29(S1):S3–14.

- Füzéki E, Banzer W. Physical activity recommendations for health and beyond in currently inactive populations. *Int J Environ Res Public Health*, 2018; 15(5):1042.
- Gunton JE, Girgis CM. Vitamin D and muscle. *Bone Rep*, 2018; 8:163–7.
- Hall JT, Ebeling M, Shary JR, Forestieri N, Wagner CL. The relationship between physical activity and vitamin D status in postpartum lactating and formula-feeding women. *J Steroid Biochem Mol Biol*, 2018; 177:261–5.
- Hamer M, Stamatakis E. Relative proportion of vigorous physical activity, total volume of moderate to vigorous activity, and body mass index in youth: the Millennium Cohort Study. *Int J Obes*, 2018; 42(6):1239–42.
- Hassanalilou T, Khalili L, Ghavamzadeh S, Shokri A, Payahoo L, Bishak YK. Role of vitamin D deficiency in systemic lupus erythematosus incidence and aggravation. *Auto Immun Highlights*, 2017; 9(1):1.
- Hong I, Coker-Bolt P, Anderson KR, Lee D, Velozo CA. Relationship between physical activity and overweight and obesity in children: findings from the 2012 National Health and Nutrition Examination Survey National Youth Fitness Survey. *Am J Occup Ther*, 2016; 70(5):7005180060p1–8.
- Hruby A, Manson JE, Qi L, Malik VS, Rimm EB, Sun Q, Willett WC, Hu FB. Determinants and consequences of obesity. *Am J Public Health*, 2016; 106(9):1656–62.
- Jackson SE, Llewellyn CH, Smith L. The obesity epidemic—nature via nurture: a narrative review of high-income countries. *SAGE Open Med*, 2020; 8:2050312120918265.
- Jakicic JM, Powell KE, Campbell WW, Dipietro L, Rate RR, Pescatello LS, Collins KA, Bloodgood B, Piercy KL. Physical activity and the prevention of weight gain in adults: a systematic review. *Med Sci Sports Exerc*, 2019; 51(6):1262–9.
- Jayawardena R, Jayakumar DT, Misra A, Hills AP, Ranasinghe P. Obesity: a potential risk factor for infection and mortality in the current COVID-19 epidemic. *Diabetes Metab Syndr*, 2020; 14(6):2199–203.
- Jehan S, Zizi F, Pandi-Perumal SR, McFarlane SI, Jean-Louis G, Myers AK. Energy imbalance: obesity, associated comorbidities, prevention, management and public health implications. *Adv Obes Weight Manag Control*, 2020; 10(5):146–61.
- Judistiani RTD, Nirmala SA, Rahmawati M, Ghrahani R, Natalia YA, Sugianli AK, Indrati AR, Suwarsa O, Setiabudiawan B. Optimizing ultraviolet B radiation exposure to prevent vitamin D deficiency among pregnant women in the tropical zone: report from cohort study on vitamin D status and its impact during pregnancy in Indonesia. *BMC Pregnancy Childbirth*, 2019; 19(1):209.
- Kavadar G, Demircioğlu DT, Özgönenel L, Emre TY. The relationship between vitamin D status, physical activity and insulin resistance in overweight and obese subjects. *Bosn J Basic Med Sci*, 2015; 15(2):62–6.
- Kift R, Rhodes LE, Farrar MD, Webb AR. Is sunlight exposure enough to avoid wintertime vitamin d deficiency in United Kingdom population groups? *Int J Environ Res Public Health*, 2018; 15(8):1624.
- Kim BY, Choi DH, Jung CH, Kang SK, Mok JO, Kim CH. Obesity and physical activity. *J Obes Metab Syndr*, 2017; 26(1):15–22.
- Kim SH, Oh JE, Song DW, Cho CY, Hong SH, Cho YJ, Yoo BW, Shin KS, Joe H, Shin HS, Son DY. The factors associated with vitamin D deficiency in community dwelling elderly in Korea. *Nutr Res Pract*, 2018; 12(5):387–95.
- Koehler K, Drenowatz C. Integrated role of nutrition and physical activity for lifelong health. *Nutrients*, 2019; 11(7):1437.
- Kotarska K, Nowak MA, Nowak L, Król P, Sochacki A, Sygit K, Sygit M. Physical activity and quality of life of university students, their parents, and grandparents in poland-selected determinants. *Int J Environ Res Public Health*, 2021; 18(8):3871.
- Larson-Meyer DE, Douglas CS, Thomas JJ, Johnson EC, Barcal JN, Heller JE, Hollis BW, Halliday TM. Validation of a vitamin D specific questionnaire to determine vitamin D status in athletes. *Nutrients*, 2019; 11(11):2732.
- Lee YM, Kim SA, Lee DH. Can current recommendations on sun exposure sufficiently increase serum vitamin d level? One-month randomized clinical trial. *J Korean Med Sci*, 2020; 35(8):e50.
- Lorensia A, Suryadinata RV, Saputra R. Physical activity and vitamin D level in asthma and non-asthma. *J Farmasi Indones*, 2019; 11(1):454–65.
- Lorensia A, Muntu CM, Suryadinata RV, Septiani R. Effect of lung function disorders and physical activity on smoking and non-smoking students. *J Prev Med Hyg*, 2021; 62(1):E89–96.
- Lorensia A, Wahyudi M, Yudiarto A, Kurnia SED. Effect of illness perception on improving asthma symptoms with omega-3 fish oil therapy: Pre-post design. *J Appl Pharm Sci*, 2020a; 10(6):62–71.
- Lorensia A, Suryadinata RV, Chandra NLMR. Profile of vitamin D status, physical activity, and lung health in construction workers (profil status vitamin D, Aktivitas Fisik, dan Kesehatan Paru pada Tukang Bangunan). *Community Med Public Health Indones J*, 2020b; 1(2):117–24.
- Maurya VK, Aggarwal M. Factors influencing the absorption of vitamin D in GIT: an overview. *J Food Sci Technol*, 2017; 54(12):3753–65.
- Michigami T. Rickets/osteomalacia. Consensus on Vitamin D deficiency and insufficiency in children. *Clin Calcium Jpn*, 2018; 28(10):1307–11.
- Nabak A, Johnson R, Keuler N, Hansen K. Can a questionnaire predict vitamin D status in postmenopausal women? *Public Health Nutr*, 2014; 17(4):739–46.
- Nakamura K, Nashimoto M, Okuda Y, Ota T, Yamamoto M. Fish as a major source of vitamin D in the Japanese diet. *Nutrition*, 2002; 18(5):415–6.
- Oh TR, Kim CS, Bae EH, Ma SK, Han SH, Sung SA, Lee K, Oh KH, Ahn C, Kim SW. Association between vitamin D deficiency and health-related quality of life in patients with chronic kidney disease from the KNOW-CKD study. *PLoS One*, 2017; 12(4):e0174282.
- Orces C. The association between body mass index and vitamin D supplement use among adults in the United States. *Cureus*, 2019; 11(9):e5721.
- Panuganti KK, Nguyen M, Kshirsagar RK. Obesity. *StatPearls*, Treasure Island, FL, 2021. [ONLINE] Available via <https://www.ncbi.nlm.nih.gov/books/NBK459357/> (Accessed 11 August 2021).
- Paschou SA, Kosmopoulos M, Nikas IP, Spartalis M, Kassi E, Goulis DG, Lambrinoudaki I, Siasos G. The impact of obesity on the association between vitamin D deficiency and cardiovascular disease. *Nutrients*, 2019; 11(10):2458.
- Pilz S, Zittermann A, Trummer C, Theiler-Schwetz V, Lerchbaum E, Keppel MH, Grubler MR, Marz W, Pandis M. Vitamin D testing and treatment: a narrative review of current evidence. *Endocr Connect*, 2019; 8(2):R27–43.
- Polzonetti V, Pucciarelli S, Vincenzetti S, Polidori P. Dietary Intake of Vitamin D from dairy products reduces the risk of osteoporosis. *Nutrients*, 2020; 12(6):1743.
- Pulungan A, Soesanti F, Tridjaja B, Batubara J. Vitamin D insufficiency and its contributing factors in primary school-aged children in Indonesia, a sun-rich country. *Ann Pediatr Endocrinol Metab*, 2021; 26(2):92–8.
- Rachmi CN, Li M, Alison Baur L. Overweight and obesity in Indonesia: prevalence and risk factors—a literature review. *Public Health*, 2017; 147:20–9.
- Ramasamy I. Vitamin D metabolism and guidelines for vitamin D supplementation. *Clin Biochem Rev*, 2020; 41(3):103–26.
- Romieu I, Dossus L, Barquera S, Blottiere HM, Franks PW, Gunter M, Hwalla N, Hursting SD, Leitzmann M, Margets B, Nishida C. Energy balance and obesity: what are the main drivers? *Cancer Causes Control*, 2017; 28(3):247–58.
- Saket S, Varasteh N, Halimi Asl AA, Saneifard H. How antiepileptics may change the serum level of vitamin D, calcium, and phosphorus in children with epilepsy. *Iran J Child Neurol*, 2021; 15(1):19–27.
- Salam RA, Padhani ZA, Das JK, Shaikh AY, Hoodbhoy Z, Jeelani SM, Lassi ZS, Bhutta ZA. Effects of lifestyle modification interventions to

prevent and manage child and adolescent obesity: a systematic review and meta-analysis. *Nutrients*, 2020; 12(8):2208.

Sanghera DK, Sapkota BR, Aston CE, Blackett PR. Vitamin D status, gender differences, and cardiometabolic health disparities. *Ann Nutr Metab*, 2017; 70(2):79–87.

Saqib ZA, Dai J, Menhas R, Mahmood S, Karim M, Sang X, Weng Y. Physical activity is a medicine for non-communicable diseases: a survey study regarding the perception of physical activity impact on health wellbeing. *Risk Manag Healthc Policy*, 2020; 13:2949–62.

Schmid A, Walther B. Natural vitamin D content in animal products. *Adv Nutr*, 2013; 4(4):453–62.

Sember V, Meh K, Sorić M, Starc G, Rocha P, Jurak G. Validity and reliability of international physical activity questionnaires for adults across EU countries: systematic review and meta-analysis. *Int J Environ Res Public Health*, 2020; 17(19):7161.

Sharifi N, Amani R, Hajjani E, Cheraghian B. Women may respond different from men to vitamin D supplementation regarding cardiometabolic biomarkers. *Exp Biol Med (Maywood)*, 2016; 241(8):830–8.

Silva ACMD, Cureau FV, de Oliveira CL, Giannini DT, Bloch KV, Kuschnir MCC, Dutra ES, Schaan BD, de Carvalho KMB. Physical activity but not sedentary time is associated with vitamin D status in adolescents: study of cardiovascular risk in adolescents (ERICA). *Eur J Clin Nutr*, 2019; 73(3):432–40.

Sizar O, Khare S, Goyal A, Givler A. Vitamin D deficiency. StatPearls, Treasure Island, FL, 2021. [ONLINE] Available via <https://www.ncbi.nlm.nih.gov/books/NBK532266/> (Accessed 12 April 2021).

Skalska M, Nikolaidis PT, Knechtle B, Rosemann TJ, Radziminski L, Jastrzebska J, Kaczmarczyk M, Myśliwiec A, Dragos P, López-Sánchez GF, Jastrzebski Z. Vitamin D supplementation and physical activity of young soccer players during high-intensity training. *Nutrients*, 2019; 11(2):349.

Solano F. Photoprotection and skin pigmentation: melanin-related molecules and some other new agents obtained from natural sources. *Molecules*, 2020; 25(7):1537.

Suryadinata RV, Boengas S, Lorensia A. Effect of knowledge and attitude toward sun exposure related vitamin D to lung function. *Teikyō Med J*, 2021; 44(4):957–69.

Suryadinata RV, Wirjatmadi B, Andriani M, Lorensia A. Effect of age and weight on physical activity. *J Public Health Res*, 2020; 9(2):187–90.

Takeuchi Y. Rickets/osteomalacia. Treatment of vitamin D deficiency in adults. *Clin Calcium*, 2018; 28(10):1387–92.

Thanapluetiwong S, Chewcharat A, Takkavatakarn K, Praditpornsilpa K, Eiam-Ong S, Susantitaphong P. Vitamin D supplement on prevention of fall and fracture: a meta-analysis of randomized controlled trials. *Medicine (Baltimore)*, 2020; 99(34):e21506.

Umar M, Sastry KS, Chouchane AI. Role of vitamin D beyond the skeletal function: a review of the molecular and clinical studies. *Int J Mol Sci*, 2018; 19(6):1618.

Wacker M, Holick MF. Sunlight and vitamin D: a global perspective for health. *Dermatoendocrinol*, 2013; 5(1):51–108.

Wang H, Chen W, Li D, Yin X, Zhang X, Olsen N, Zheng SG. Vitamin D and chronic diseases. *Aging Dis*, 2017; 8(3):346–53.

Webb AR, Kazantzidis A, Kift RC, Farrar MD, Wilkinson J, Rhodes LE. Colour counts: sunlight and skin type as drivers of vitamin D deficiency at UK Latitudes. *Nutrients*, 2018; 10(4):457.

Wiciński M, Adamkiewicz D, Adamkiewicz M, Śniegocki M, Podhorecka M, Szycha P, Malinowski B. Impact of vitamin D on physical efficiency and exercise performance—a review. *Nutrients*, 2019; 11(11):2826.

Wimalawansa SJ. Vitamin D deficiency: effects on oxidative stress, epigenetics, gene regulation, and aging. *Biology (Basel)*, 2019; 8(2):30.

Zadka K, Pałkowska-Goździk E, Rosołowska-Huszcz D. The state of knowledge about nutrition sources of vitamin D, its role in the human body, and necessity of supplementation among parents in Central Poland. *Int J Environ Res Public Health*, 2018; 15(7):1489.

Zarooni AAR, Marzouqi FIA, Darmaki A, Prinsloo EAM, Nagelkerke N. Prevalence of vitamin D deficiency and associated comorbidities among Abu Dhabi Emirates population. *BMC Res Notes*, 2019; 12(503):1–6.

How to cite this article:

Lorensia A, Suryadinata RV, Inu IA. Comparison of vitamin D status and physical activity related to obesity among tertiary education students. *J Appl Pharm Sci*, 2022; 12(04):108–118.

Comparison of vitamin D status and physical activity related to obesity among tertiary education students

ORIGINALITY REPORT

24%

SIMILARITY INDEX

19%

INTERNET SOURCES

20%

PUBLICATIONS

8%

STUDENT PAPERS

PRIMARY SOURCES

- 1 oamjms.eu Internet Source 1%
- 2 Submitted to Surabaya University Student Paper 1%
- 3 Fatimawali Fatimawali, Kalalo Marko Jeremia, Simanjuntak Sibbantua Broolin, Heber Tri Andira et al. "Immunomodulatory potential of bioactive compounds of betel leaf extract targeting COVID-19 immunological human host proteins: An in silico study", Journal of Applied Pharmaceutical Science, 2022 Publication 1%
- 4 global-uploads.webflow.com Internet Source 1%
- 5 Basera Ishita A., Verma Preeti D., Bhatt Vijay P., Shah Mamta B.. "Determination of dioecy in Hippophae salicifolia by evaluating gallic acid content through a validated HPTLC method", Journal of Applied Pharmaceutical Science, 2022 1%

6	Vitamin D, 2010. Publication	1 %
7	napier-surface.worktribe.com Internet Source	1 %
8	strathprints.strath.ac.uk Internet Source	1 %
9	www.jpmmh.org Internet Source	1 %
10	nutritionj.biomedcentral.com Internet Source	1 %
11	www.moh.gov.sa Internet Source	1 %
12	Giuseppe Saggese, Francesco Vierucci, Flavia Prodam, Fabio Cardinale et al. "Vitamin D in pediatric age: consensus of the Italian Pediatric Society and the Italian Society of Preventive and Social Pediatrics, jointly with the Italian Federation of Pediatricians", Italian Journal of Pediatrics, 2018 Publication	<1 %
13	Handbook of Growth and Growth Monitoring in Health and Disease, 2012. Publication	<1 %
14	Evitha Latifah, Kusnandar Kusnandar, Yulia Lanti Retno Dewi. "Physical Activity, Sensory	<1 %

Threshold of Sweetness, and Waist-to-Height Ratio (WHtR) in Adolescents", Electronic Journal of General Medicine, 2020

Publication

15

"Handbook of Anthropometry", Springer Science and Business Media LLC, 2012

Publication

<1 %

16

Wibowo, Yosi, Bruce Sunderland, and Jeffery Hughes. "Pharmacist and physician perspectives on diabetes service delivery within community pharmacies in Indonesia: a qualitative study : Perspectives on pharmacy-based diabetes services", International Journal of Pharmacy Practice, 2015.

Publication

<1 %

17

www.alliedacademies.org

Internet Source

<1 %

18

www.duo.uio.no

Internet Source

<1 %

19

Peterson, C. A., A. K. Tosh, and A. M. Belenchia. "Vitamin D insufficiency and insulin resistance in obese adolescents", Therapeutic Advances in Endocrinology and Metabolism, 2014.

Publication

<1 %

20

Submitted to University of Ulster

Student Paper

<1 %

21 Stefanos Tyrovolas, Vassiliki Bountziouka, Natassa Papairakleous, Akis Zeimbekis et al. "Adherence to the Mediterranean diet is associated with lower prevalence of obesity among elderly people living in Mediterranean islands: the MEDIS study", International Journal of Food Sciences and Nutrition, 2009
Publication

22 digibug.ugr.es
Internet Source

23 Sajida Maryam, Saman Saba, Waseem Haider, Muhmmad Sohail Afzal et al. "Community-based social and demographic assessment of knowledge, attitudes, practices and medical conditions related to vitamin D deficiency in Gilgit Baltistan, Pakistan", Journal of Biosocial Science, 2021
Publication

24 Submitted to Mahidol University
Student Paper

25 [Www.japsonline.com](http://www.japsonline.com)
Internet Source

26 www.frontiersin.org
Internet Source

27 Submitted to Mississippi State Board for Community & Junior Colleges
Student Paper

28

hdl.handle.net

Internet Source

<1 %

29

link.springer.com

Internet Source

<1 %

30

repositorii.urindo.ac.id

Internet Source

<1 %

31

www.danonenutrindo.org

Internet Source

<1 %

32

www.valleyinternational.net

Internet Source

<1 %

33

icge.unespadang.ac.id

Internet Source

<1 %

34

Deepika Dhawan, Sheel Sharma. "Abdominal Obesity, Adipokines and Non-communicable Diseases", *The Journal of Steroid Biochemistry and Molecular Biology*, 2020

Publication

<1 %

35

Siobhan Cusack, Christian Mølgaard, Kim F. Michaelsen, Jette Jakobsen, Christel J.E. Lamberg-Allardt, Kevin D. Cashman. "Vitamin D and estrogen receptor- α genotype and indices of bone mass and bone turnover in Danish girls", *Journal of Bone and Mineral Metabolism*, 2006

Publication

<1 %

doaj.org

36

Internet Source

<1 %

37

jfcmonline.com

Internet Source

<1 %

38

Maša Hribar, Hristo Hristov, Matej Gregorič, Urška Blaznik et al. "Nutrihealth Study: Seasonal Variation in Vitamin D Status Among the Slovenian Adult and Elderly Population", *Nutrients*, 2020

Publication

<1 %

39

academic.oup.com

Internet Source

<1 %

40

e-journal.unair.ac.id

Internet Source

<1 %

41

www.scielo.br

Internet Source

<1 %

42

Submitted to Chester College of Higher Education

Student Paper

<1 %

43

Michael J. Berridge. "Vitamin D deficiency: infertility and neurodevelopmental diseases (attention deficit hyperactivity disorder, autism and schizophrenia)", *American Journal of Physiology - Cell Physiology*, 2017

Publication

<1 %

44

discovery.dundee.ac.uk

Internet Source

<1 %

45

ejournal.aibpm.org

Internet Source

<1 %

46

www.ijhcum.net

Internet Source

<1 %

47

"Effect of illness perception on improving asthma symptoms with omega-3 fish oil therapy: Pre-post design", Journal of Applied Pharmaceutical Science, 2020

Publication

<1 %

48

propertibazar.com

Internet Source

<1 %

49

worldnutrijournal.org

Internet Source

<1 %

50

Submitted to University of Leeds

Student Paper

<1 %

51

Submitted to University of South Florida

Student Paper

<1 %

52

docplayer.net

Internet Source

<1 %

53

www.science.gov

Internet Source

<1 %

54

Abd-Alla Howaida Ibrahim, Hassan Amal Zaki, Soltan Maha Mohamed, Abdelwahab Ahmed

<1 %

Baker, Hanna Atef Gobran. "Potential protein antiglycation, antiproliferation, and in silico study on the antidiabetic enzymes of bioactive metabolites from Adonis microcarpa DC and their ADMET properties", Journal of Applied Pharmaceutical Science, 2022

Publication

55

Submitted to Glasgow Caledonian University

Student Paper

<1 %

56

Submitted to Howard University

Student Paper

<1 %

57

Valerie A. Holmes, Maria S. Barnes, H. Denis Alexander, Peter McFaul, Julie M. W. Wallace.

"Vitamin D deficiency and insufficiency in pregnant women: a longitudinal study",

British Journal of Nutrition, 2009

Publication

<1 %

58

centaur.reading.ac.uk

Internet Source

<1 %

59

jabonline.a2hosted.com

Internet Source

<1 %

60

msjonline.org

Internet Source

<1 %

61

www.biomedcentral.com

Internet Source

<1 %

62

www.mjms.mk

<1 %

63

Ayesha Aslam, Ali Bahadar, Rabia Liaquat, Muhammad Saleem, Adeel Waqas, Mohammed Zwawi. "Algae as an attractive source for cosmetics to counter environmental stress", Science of The Total Environment, 2021

Publication

<1 %

64

Kazutoshi Nakamura. "Vitamin D insufficiency in Japanese populations: from the viewpoint of the prevention of osteoporosis", Journal of Bone and Mineral Metabolism, 2005

Publication

<1 %

65

Lubna I. Al Asoom. "Assessment of plasma level of 25(OH)D and its correlation with cardiorespiratory fitness in young females of Dammam City, KSA", Journal of Taibah University Medical Sciences, 2016

Publication

<1 %

66

Submitted to Manchester Metropolitan University

Student Paper

<1 %

67

www.cureus.com

Internet Source

<1 %

68

Ioannis Dimakopoulos, Emmanuella Magriplis, Anastasia-Vasiliki Mitsopoulou, Dimitra

<1 %

Karageorgou et al. "Association of serum vitamin D status with dietary intake and sun exposure in adults", Clinical Nutrition ESPEN, 2019

Publication

69

McDaid, David, Sassi, Franco, Merkur, Sherry. "EBOOK: Promoting Health, Preventing Disease: The Economic Case", EBOOK: Promoting Health, Preventing Disease: The Economic Case, 2015

Publication

70

Rasoul Abdollahzadeh, Mohammad Hossein Shushizadeh, Mina Barazandehrokh, Sepideh Choopani et al. "Association of Vitamin D receptor gene polymorphisms and clinical/severe outcomes of COVID-19 patients", Infection, Genetics and Evolution, 2021

Publication

71

Salah Gariballa, Iltaf Shah, Javed Yasin, Awad Alessa. "Vitamin D [25(OH)D] metabolites and epimers in obese subject: interaction and correlations with adverse metabolic health risk factors", The Journal of Steroid Biochemistry and Molecular Biology, 2021

Publication

72

Submitted to University of Wollongong

Student Paper

<1 %

<1 %

<1 %

<1 %

73	www.dovepress.com Internet Source	<1 %
74	biochemia-medica.com Internet Source	<1 %
75	bmcendocrdisord.biomedcentral.com Internet Source	<1 %
76	garuda.kemdikbud.go.id Internet Source	<1 %
77	jcspk.pk Internet Source	<1 %
78	masood abdukkareem abdukkrahman, Suad Yousif Alkass, Noor Isam Mohammed. "Free Vitamin D Status Among Apparently Healthy Adults Living in Duhok Governorate", Research Square Platform LLC, 2021 Publication	<1 %
79	mdpi-res.com Internet Source	<1 %
80	opendata.uni-halle.de Internet Source	<1 %
81	www.bjbms.org Internet Source	<1 %
82	www.thefreelibrary.com Internet Source	<1 %

83

www.vitamind-journal.it

Internet Source

<1 %

84

"Dietary reference values for vitamin D", EFSA Journal, 2016

Publication

<1 %

85

"Explaining Health Across the Sciences", Springer Science and Business Media LLC, 2020

Publication

<1 %

86

"Nutritional Influences on Bone Health", Springer Nature, 2013

Publication

<1 %

87

Cai, Fin ZJ, Sue Lester, Tim Lu, Helen Keen, Karyn Boundy, Susanna M Proudman, Anne Tonkin, and Maureen Rischmueller. "Mild autonomic dysfunction in primary Sjögren's syndrome: a controlled study", Arthritis Research & Therapy, 2008.

Publication

<1 %

88

G. Naureen, K. M. Sanders, L. Busija, D. Scott, K. Lim, J. Talevski, C. Connaughton, S. L. Brennan-Olsen. "Prediction models and questionnaires developed to predict vitamin D status in adults: a systematic review", Osteoporosis International, 2020

Publication

<1 %

89

N. Alonso, S. Zelzer, G. Eibinger, M. Herrmann. "Vitamin D Metabolites: Analytical Challenges and Clinical Relevance", *Calcified Tissue International*, 2022

Publication

<1 %

90

[bmcnutr.biomedcentral.com](https://www.bmcnutr.biomedcentral.com)

Internet Source

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On

Comparison of vitamin D status and physical activity related to obesity among tertiary education students

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8

PAGE 9

PAGE 10

PAGE 11
