



Detection of *Salmonella* spp., *Escherichia coli*, and *Listeria monocytogenes* in Tuna by Multiplex PCR

Kristina Natalia^{1,2}, Mariana Wahjudi¹ and Tjie Kok^{1*}

¹Faculty of Biotechnology, University of Surabaya, Jalan Raya Kali Rungkut, Surabaya 60293, East Java, Indonesia

²Regional Fish Quarantine and Inspection Agency, Surabaya I 61254, East Java, Indonesia

*Correspondence :
tjie_kok@staff.ubaya.ac.id

Received : 2022-06-10

Accepted : 2022-08-11

Keywords :
Escherichia coli, *Listeria monocytogenes*, Multiplex PCR (mPCR), *Salmonella* spp., Tuna

Abstract

Fishery products are types of commodity with various levels of development including the addition of food additives. They are very susceptible to pathogenic microbial contamination hence they can cause food poisoning and disease outbreaks in consumers. In this study, tuna fish spiked with *Salmonella* spp., *Escherichia coli*, and *Listeria monocytogenes* bacteria is examined for bacteria contamination. The examination of these pathogenic bacteria is carried out using multiplex PCR (mPCR), a modern technique developed to detect target DNAs of bacteria simultaneously. The results of mPCR are compared to those of biochemical tests for sensitivity and time-efficiency. Based on the results obtained, the mPCR method is ± 1250 -fold more sensitive and ± 80 -fold more time efficient than the biochemical test. Therefore, mPCR method could be recommended to be used as a tool for the detection of bacteria in fishery products.

INTRODUCTION

Fish is a source of protein that is generally the people's choice because it has a relatively cheap price. Fresh fish products are the main commodities of fishery products in Indonesia, one of which is tuna. Tuna is a kind of fish with a very high protein content (22.6-26.2 g/100 g) and fat (0.2-2.7 g/100 g). Tuna also contains minerals (calcium, phosphorus, iron, sodium), vitamin A (retinol), and vitamin Bs (thiamin, riboflavin, and niacin). In general, the edible portion of tuna (edible portion) ranges from 50-60% of the fish body (Alverson and Stansby, 1963).

Original fresh tuna products are usually sold in the local market and are usually purchased and then reprocessed into various processed fish, both raw and

cooked. Post-catching fresh fish products have several drawbacks due to the lack of quality control processes starting from catching, distribution and marketing, hence the quality of fresh fish products produced is very diverse.

Fresh tuna has a very high risk of contamination because metabolic processes in dead fish are uncontrolled. The catabolism process results in the formation of substrates for bacteria, and hence fresh fish are growth substrates for bacteria so raw fish products damage faster than processed fish products (Palawe *et al.*, 2016). The presence of bacterial contamination in fresh tuna is one of the quality parameters for checking. Based on data on the FDA's RAS website, in 2020, for tuna fish products in

Indonesia, there were 13 cases of *Salmonella* spp. contamination. In June 2021, there were 7 cases of *Salmonella* spp. contamination in tuna fish products and 1 case of *L. monocytogenes* contamination in shrimp.

Test for pathogenic bacteria (*Salmonella* spp., *Escherichia coli*, and *L. monocytogenes*) on tuna fish products that are used by most state laboratories in Indonesia is using the biochemical SNI method. The test takes approximately 10-12 days to obtain results. In addition, the equipment and media needed for the test vary a lot, and they render higher operational costs.

Considering that there are shortcomings in the biochemical method, the development of a new diagnostic method providing solutions for lower cost and more time efficiency is essential. The singleplex PCR method for the test of several food-borne pathogenic bacteria has been used with sensitive results (Patil-Joshi *et al.*, 2021). The quality of fishery products must fulfill export requirements before the delivery of products to destination countries such as Japan, the USA, the UK, Belgium, Canada, and some European countries. The ingestion of fish with high histamine levels is toxic, hence maximum permissible levels of < 200 or < 400 mg/kg of histamine are applied. The levels of pathogenic bacteria contamination are also regulated, for example, the level of *L. monocytogenes* contamination should not exceed 10² cfu/g (based on shelf life of fewer than 5 days) (Health Protection Agency Working Group, 2016).

In this study, a more time-efficient test is developed to detect the presence of the bacteria in fresh tuna fish products. Triwibowo *et al.* (2020) reported that mPCR offers a lower cost per sample than the biochemical method due to cheaper reagents and lower energy consumption for analysis. Hence mPCR is considered a promising new diagnostic method for the detection of food-borne pathogens. In this study, the sensitivity and time-efficiency of mPCR method are compared to those of

biochemical test for the detection of *Salmonella* spp., *E. coli*, and *L. monocytogenes* in fresh tuna fish products.

METHODOLOGY

Place and Time

This research was carried out in the Laboratory of Regional Fish Quarantine and Inspection Agency (Laboratorium Balai Karantina Ikan, Pengendalian Mutu dan Keamanan Hasil Perikanan – KIPM) Surabaya I for three months, January to March 2022.

Research Materials

The materials used in this study are *Salmonella* spp. test media according to SNI ISO 6579:2015: Bismut sulfite agar, Buffered peptone water, MKTTN, Xylose lysine deoxycholate agar, Triple sugar iron agar, Tryptone broth, Rappaport-Vassiliadis medium, Reagen covacs, ONPG, nutrient agar; *E. coli* test media according to SNI 2332.1-2015: Lauryl Tryptose broth, EC broth, Levine's eosin methylene blue, Tryptone broth, MR-VP broth, Simmon citrate broth, Plate count agar, Lactose broth, Covacs reaction, Methyl red indicator; *L. monocytogenes* test media according to SNI ISO 11290-2:2015: Fraser Broth, Fraser listeria ammonium iron III supplement, Listeria agar acc Ottaviani and agosti, Listeria agar selective-supplement, PALCAM agar acc van et al base, PALCAM listeria selective supplement, Rhamnose monohydrate, Xylose, Yeast extract, Enzymatic digest animal tissue) and Polymerase Chain Reaction (PCR) materials: Silica kit extraction, Nuclease free water, TAE buffer, primer, DNA ladder 20, dNA staining, agarose.

Meanwhile, the equipment used in this study were stomacher, incubator, water bath, autoclave, oven, laminar air flow, microwave, microcentrifuge, vortex, thermal cycler, electrophoresis, spin down, micro tube, and micro pipette.

Research Design

This study used a descriptive research design. The results of biochemical and PCR tests on tuna spiked with *Salmonella* spp., *E. coli*, and *L. monocytogenes* are presented in the Table and Figures based on data from Laboratory tests.

Work Procedure

This research was performed in several stages starting with checking pure cultures of *Salmonella* spp., *E. coli*, and *L. monocytogenes* bacteria. Detection of the pathogenic bacteria was carried out using biochemical and mPCR tests. For PCR, a singleplex PCR was initially performed to find the optimum temperature for each bacterium. After obtaining the optimum temperature of the three bacteria, mPCR test was performed using the optimum temperature obtained during the singleplex PCR. Tuna free from pathogenic bacteria (unspiked tuna) was used as a negative control for tuna contaminated with each of the three pathogenic bacteria (spiked tuna).

Bacterial Culture Preparation

Salmonella spp. ATCC 14028, *E. coli* ATCC 25922, and *L. monocytogenes* ATCC 7644 were respectively on NaCl TSA growing media. After each bacterium was grown on selective media, XLD agar,

LEMB agar, and ALOA agar, biochemical test. Meanwhile, to enrich *Salmonella* spp., *E. coli*, and *L. monocytogenes*, the corresponding isolate was cultured on non-selective Brain Heart Infusion Broth (BHIB) for 18-24 hours at 37 °C (*Salmonella* spp.), on Brain Heart Infusion Broth (BHIB) for 24 hours at 5 °C (*E. coli*), and on Listeria Enrichment Broth (LEB) for 42-48 hours at 27-30 °C (*L. monocytogenes*) in order to get stock (Sjahriani *et al.*, 2021). Then each culture was divided into two parts: part 1 for the colony test (encoded A for *Salmonella* spp., B for *E. coli*, and C for *L. monocytogenes*) (Table 1) and part 2 for the tuna sample test (Zhang *et al.*, 2015).

Tuna Sample Preparation

Tuna fish samples were crushed and divided into 4 parts. Part 1 was the unspiked tuna sample, encoded D. Part 2 was the tuna sample spiked with *Salmonella* spp. encoded E. Part 3 was tuna sample spiked with *E. coli* encoded F. Part 4 was a tuna sample spiked with *L. monocytogenes*, encoded G. In addition, 50 gr of each B, C, and D were taken and mixed to be sample encoded H (Table 1). Each sample weighing 25 g and 20 mg was taken for biochemical and PCR tests respectively. The sample was initially homogenized using a vortex and allowed to stand for 15 min-1 hour before the test.

Table 1. Sample coding.

Sample type	Code
Pure colony of <i>Salmonella</i> spp.	A
Pure colony of <i>Escherichia coli</i>	B
Pure colony of <i>Listeria monocytogenes</i>	C
Unspiked tuna sample	D
Tuna sample spiked with <i>Salmonella</i> spp.	E
Tuna sample spiked with <i>Escherichia coli</i>	F
Tuna sample spiked with <i>Listeria monocytogenes</i>	G
Mixture of samples E+F+G	H

Biochemical Test

Pure bacterial colonies and the tuna samples, both unspiked and spiked with target bacteria, were tested biochemically. Test for *Salmonella* spp. followed the SNI ISO 6579:2015, *E. coli* SNI 2332.1-2015,

and *L. monocytogenes* follows SNI ISO 11290-2:2015 procedure.

Bacterial DNA Isolation

Each isolate of the bacterial colony was performed using the silica extraction

method, following the standard procedure at the KIPM Biomolecular Lab. The isolate of unspiked and spiked tuna samples were also extracted using the same procedure. In addition, the extracted DNA from each

spiked tuna sample was taken, and mixed together to obtain a mixture containing DNAs of *Salmonella* spp., *E. coli*, and *L. monocytogenes*. The primers used for PCR were as follows in Table 2.

Table 2. Primer for PCR.

Gen Target (species)	Dimension (bp)	Label	Sequence (5' → 3')	Tm* (°C)	Reference
invA (<i>Salmonella</i> spp.)	275	Forward	AAT TAT CGC CAC GTT CGG GCA A	68	(Rahn <i>et al.</i> , 1992)
		Reverse	TCG CAC CGT CAA AGG AAC C CAT TAG TGG	63	
hlyA (<i>Listeria monocytogenes</i>)	730	Forward	AAA GAT GGA ATG	N/A	(Bilung <i>et al.</i> , 2018)
		Reverse	GTA TCC TCC AGA GTG ATC GA	N/A	
uidA (<i>E. coli</i>)	147	Forward	TGG TAA TTA CCG ACG AAA ACG GC	66	(Jefferson <i>et al.</i> , 1986)
		Reverse	ACG CGT GGT TAC AGT CTT	64	

*) Tm: primer melting temperature.

Optimization of PCR Condition

Optimization of PCR conditions was carried out by selecting a primer master mix according to the corresponding DNA

templates for optimizing the primer annealing temperature. The compositions of the mixtures and primers used in PCR were as follows (Table 3).

Table 3. Composition of mixture in PCR processes.

Single PCR	(μ l)	Multiplex PCR	(μ l)
Master mix	12.5 μ l	Master mix	12.5 μ l
Forward primer	1 μ l	Forward primer inv A	1 μ l
Reverse primer	1 μ l	Reverse primer inv A	1 μ l
Nuclease free	8.5 μ l	Forward primer uid A	1 μ l
Water		Reverse primer uid A	1 μ l
Template	2 μ l	Forward primer hly A	1 μ l
Total	25 μ l	Reverse primer hly A	1 μ l
		Nuclease free water	4.5 μ l
		DNA mixture	2 μ l
		Total	25 μ l

Note: Final concentration for each primer is 10 μ M.

Orientation processes of the singleplex PCR process were carried out using annealing temperatures ranging at 55.0-63.0 °C to find the optimum annealing temperature for each target DNA of bacteria. Next, mPCR was

performed at the annealing optimum temperature, i.e., 58.1 °C. Using optimum annealing temperature can increase PCR efficiency for unknown DNA (Wei *et al.*, 2007). The complete PCR stage and temperatures were as follows (Table 4):

Table 4. PCR stages and temperatures.

Stage	Temperature (°C)	Time (minutes)	
Pre-denaturation	95	3	} 35 cycles
Denaturation	95	0.5	
Annealing	58	0.5	
Elongation	72	0,5	
Final extension	72	5	

The PCR results were visualized by gel electrophoresis. A total of 2 g of agarose was dissolved in TAE buffer which has been two-fold diluted. The solution was heated using a microwave at 100°C for 4 minutes to ensure complete dissolution. Five µl of fluoro-safe DNA was then added and the mixture was allowed to stand for a while before being poured into molds and left to freeze. The frozen gel was placed in the placed in electrophoresis chamber and 1x TAE running buffer was poured into the chamber until the gel was submerged. The sample containing DNA and the marker were then put into the well of the gel. The electrophoresis process was run at 120 Volts for 35 min until completion. The gel was then put on a UV transilluminator to be visualized.

RESULTS AND DISCUSSION

Biochemical Test Results

In the biochemical test, there are three steps that must be carried out: pre-enrichment in broth media, followed by selective culture on agar media, and biochemical test. Pre-enrichment is an important step for the detection of pathogens from food or environmental product samples to suppress the growth of competitive, non-target bacteria, and also to avoid false negative results (Hoorfar and Baggesen, 1998). Based on the biochemical test results of *Salmonella* spp., *E. coli*, and *L. monocytogenes* it took approximately 5 days, 10 days, and 5 days, respectively. The biochemical test of unspiked tuna (sample D) for *Salmonella* spp., *E. coli*, and *L. monocytogenes* showed negative results (Figure 1).



Figure 1. Biochemical test results of sample D on (a) XLD media, (b) LEMB Media, (c) ALOA media.

For the *Salmonella* spp. there was no colony growing on the XLD media (Figure 1a). For *E. coli*, the LEMB media was clear (Figure 1b), indicating there is no H₂S produced. And for *L. monocytogenes*, there was no colony growing on ALOA media (Figure 1c).

The biochemical test for pure colonies (samples A, B, and C) gave positive results (Figure 2) and so did that for spiked tuna samples (samples E, F, G, and H) (Figures 3 and 4).




Figure 2. Biochemical test results of (a) sample A on XLD media, (b) sample B on LEMB media; (c) sample C on ALOA media.




Figure 3. Biochemical test results of (a) sample E on XLD media, (b) sample F on LEMB media, (c) sample G on ALOA media.




Figure 4. Biochemical test results of sample H on (a) XLD media, (b) LEMB media, (c) on ALOA media.

For pure colony of *Salmonella* spp. (Figures 3a and 4a), there were black colonies growing on XLD media indicating

that *Salmonella* spp. is producing H₂S. Further biochemical test results confirmed the presence of the bacteria (Table 5).

Table 5. *Salmonella* spp. biochemical test results.

Media Test	Appearance	Results		
		C+1	C+3	E
TSI agar	Yellow	no	no	No
Urea Agar	Pink	yes	yes	Yes
L-Lysine decarboxylation medium	Purple	yes	yes	Yes
Galactosidase Reaction	Yellow	no	no	No
Voges-Proskauer reaction	Red/Pink	no	no	No
Indo Production	Red ring	no	no	No
	Positive for	<i>Salmonella typh</i>	<i>Salmonella typh</i>	<i>Salmonella typh</i>

For a pure colony of *E. coli* and tuna samples spiked with *E. coli* initially, LTB test gave positive results, i.e., cloudy and gassy appearance. Then, some portions from the LTB tube were taken using an ose needle and inoculated into EC broth tubes containing Durham tubes, this also gave

positive results. Hence, they were further grown on LEMB media. There were black and metallic green colonizing appearing (Figures 2b, 3a, and 4b). Further biochemical test results for samples B, F, and H confirmed the presence of the bacteria (Table 6).

Table 6. *Escherichia coli* biochemical test results.

Criteria	Sightings	Results		
		B+1	B+3	E
LTB	Gas	yes	yes	yes
Indo	Red ring	No	no	no
Methyl red	Red	yes	yes	yes
Voges Proskauer	red/pink	No	no	no
Citric	Blue	yes	yes	yes
	Positive for	<i>Escherichia coli</i>	<i>Escherichia coli</i>	<i>Escherichia coli</i>

For the pure colony of *L. monocytogenes* test (Figs. 2c) and tuna samples spiked with *L. monocytogenes* (Figure 3c and 4c), there were greenish

blue colonies with or without a white halo growing on ALOA media. Further biochemical test results confirmed the presence of the bacteria (Table 7).

Table 7. *Listeria monocytogenes* test results.

Criteria	Appearance	Results		
		D+1	D+3	E
Microscopic test	Gram positive, thin, short rods or cocci-shaped bacilli	Gram positive, thin, short rods or cocci-shaped bacilli	Gram positive, thin, short rods or cocci-shaped bacilli	Gram positive, thin, short rods or cocci-shaped bacilli
Hemolysis	lines are narrowed, clean, and the hemolysis area is bright	Yes	yes	yes
L- Rhamnose	Yellow	Yes	yes	yes
D-Xylose	Yellow	No	no	no
	Positive for	<i>Listeria monocytogenes</i>	<i>Listeria monocytogenes</i>	<i>Listeria monocytogenes</i>

PCR Test

Prior to the mPCR, a singleplex PCR was carried out by initially performing the process at a temperature range of 55 to 63.0 °C to determine the optimum

annealing temperature for each target DNA of bacteria.

The electrophoresis visualization of singleplex PCR result for *Salmonella* spp. DNA was shown in Figure 5.




Figure 5. Visualization of singleplex PCR result for *Salmonella* spp. DNA at various temperatures: A. 63.0 °C; B. 62.4 °C; C. 61.4 °C; D. 59.9 °C; E. 58.1 °C; F. 56.5 °C; G. 55.6 °C; H. 55 °C; M is the marker.

The singleplex PCR for *Salmonella* spp. DNA at various temperatures gave a band at 275 bp on an electrophoresis gel.

The electrophoresis visualization of singleplex PCR results for *E. coli* DNA was shown in Figure 6.




Figure 6. Visualization of singleplex PCR result for *E. coli* DNA at various temperatures: A. 63.0 °C; B. 62.4 °C; C. 61.4 °C; D. 59.9 °C; E. 58.1 °C; F. 56.5 °C; G. 55.6 °C; H. 55 °C; M is the maker.

The Singleplex PCR for *E. coli* DNA at various temperatures gave a band at 147 bp on the electrophoresis gel. The

Electrophoresis visualization of singleplex PCR result for *L. monocytogenes* DNA was shown in Figure 7.




Figure 7. Electrophoresis visualization of singleplex PCR result DNA of *L. monocytogenes* at various temperatures: M is the marker, A. 63.0 °C; B. 62.4 °C; C. 61.4 °C; D. 59.9 °C; E. 58.1 °C; F. 56.5 °C; G. 55.6 °C; H. 55 °C.

The singleplex PCR for *L. monocytogenes* DNA at various temperatures gave a band at 730 bp on an electrophoresis gel.

three target DNAs of bacteria was 58.1 °C. Hence, mPCR was carried out at the temperature to examine the three target DNAs of positive controls (mixture of samples A, B, and C) and spiked tuna samples (mixture of samples E, F, and G) (Figure 8).

Based on the results of the singleplex PCR and confirmed by mPCR, the optimum annealing temperature for the




Figure 8. Visualization of mPCR results for the three target DNAs of positive controls and spiked tuna samples at 58.1 °C. M is the marker, A is target DNAs of positive controls, B is the target DNAs of spiked tuna samples.

The sensitivity of PCR to detect the presence of pathogenic bacteria in a food product, in this case, tuna product, depending on the sample matrix, pre-enrichment method, DNA extraction, and PCR condition (Law *et al.*, 2014). The specificity of mPCR result on electrophoresis gel showed the bands at respective bp of all the three target DNAs of bacteria (Figure 8), indicating the specificity of this method.

Based on this study, mPCR is more sensitive than a biochemical test: mPCR requires much less sample than a biochemical test. In the biochemical test, for every 225 mL of media ± 20 mg is needed. Hence, mPCR is $\pm(25 \times 1000)/20 = 1250$ -fold more sensitive than biochemical test. In addition, mPCR provides faster analysis than the biochemical test. In the biochemical test, it takes at least 5 days for *Salmonella* spp. analysis, 10 days for *E. coli* analysis, and 5 days for *L. monocytogenes* analysis. Meanwhile, in mPCR, it takes ± 3 hours to finish the analysis. Hence for the examination of *Salmonella* spp., *E. coli*, and *L. monocytogenes*, mPCR is $\pm(10 \times 24)/3 = 80$ -fold more time-efficient than the biochemical test.

Even though mPCR provides an advantage over the biochemical test, i.e., more sensitive and more efficient, this method has some weaknesses. First, if the extraction process is not good, then the isolate obtained is not pure and when the PCR is run, there is a possibility of obtaining false negative results. DNA is a material that is most likely covered by fat. Hence, DNA extraction is somewhat tricky and care is needed to obtain pure target DNAs from bacteria. Secondly, mPCR is unable to distinguish whether the DNAs are from living or not living bacteria (Dwivedi and Jaykus, 2011). Hence, an additional step is needed to confirm the presence of living bacterial DNAs rather than for quantification, excluding the Most Probable Number PCR (MPN PCR) test (Bonny *et al.*, 2018). Moreover, PCR cannot be applied to samples with a low PH.

Further researches need to be conducted to provide a more comprehensive study on the use of mPCR for simultaneous detection of pathogenic bacteria in fish products, for standardization of this method.

CONCLUSION

The biochemical test and mPCR can be applied to detect the presence of pathogenic bacteria in tuna samples. In this case, mPCR provides several advantages: it can detect *Salmonella* spp., *E. coli*, and *L. monocytogenes* simultaneously, and it is more sensitive and more time-efficient than a biochemical test. Hence mPCR could be recommended as a standard test for replacement of biochemical tests.

ACKNOWLEDGMENT

The authors would like to express deep gratitude to the Laboratory of Regional Fish Quarantine and Inspection Agency Surabaya I during the research.

REFERENCES

- Alverson, D.L. and Stansby, M.E., 1963. The spiny dogfish (*Squalus acanthias*) in the northeastern Pacific (No. 447). US Department of the Interior, Fish and Wildlife Service, Bureau of Commercial Fisheries. <https://spo.nmfs.noaa.gov/content/spiny-dogfish-squalus-acanthias-northeastern-pacific>.
- Bilung, M.L., Chai, L.S., Tahar, A.S., Ted, C.K. and Apun, K., 2018. Prevalence, genetic heterogeneity, and antibiotic resistance profile of *Listeria* spp. and *Listeria monocytogenes* at farm level: A highlight of ERIC-and BOX-PCR to reveal genetic diversity. *DioMed Research International*, 2018, 3067494. <https://doi.org/10.1155/2018/3067494>
- Bonny, S.Q., Hossain, M.M.A., Lin, T.K. and Ali, M.E., 2018. Multiplex MPN-PCR for the enumeration of three major Vibrios in raw fishes in Malaysia. *Food Control*, 90, pp.459-

465. <https://doi.org/10.1016/j.foodcont.2018.02.034>
- Dwivedi, H.P. and Jaykus, L.A., 2011. Detection of pathogens in foods: the current state-of-the-art and future directions. *Critical Reviews in Microbiology*, 27(1), pp.40-63. <https://doi.org/10.3109/1040841X.2010.506430>
- Health Protection Agency Working Group, 2016. Guidelines for assessing the microbiological safety of ready-to-eat foods placed on the market. https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/363146/Guidelines_for_assessing_the_microbiological_safety_of_ready-to-eat_foods_on_the_market
- Hoorfar, J. and Baggesen, D.L., 1998. Importance of pre-enrichment media for isolation of *Salmonella* spp. from swine and poultry. *FEMS Microbiology Letters*, 169(1), pp.125-130. <https://doi.org/10.1111/1/j.1574-6968.1998.tb13308.x>
- Jefferson, R.A., Burgess, S.M. and Hirsh, D., 1986. β -Glucuronidase from *E. coli* as a gene-fusion marker. *Proceedings of the National Academy of Sciences of the United States of America*, 83(22), pp.8447-8451. <https://doi.org/10.1073/pnas.83.2.8447>
- Law, J.W.F., Ab Mutalib, N.S., Chan, K.G. and Lee, L.H., 2014. Rapid methods for the detection of foodborne bacterial pathogens: principles, applications, advantages and limitations. *Frontiers in Microbiology*, 5, p.770. <https://doi.org/10.3389/fmicb.2014.00770>
- Palawe, J.F., Wodi, S.I.M. and Cayono, E., 2016. Analysis of Total Microbial Contamination in Some Fresh Fish in the Sangihe Islands Regency. *Jurnal Ilmiah Tindalung*, 2(1), pp.42-46. <https://doi.org/10.5281/jit.v2i1.80>
- Patil-Joshi, A., Rangaswamy, B.E. and Apte-Deshpande, A., 2021. Based PCR method development, validation and application for microbial detection. *Journal of Genetic Engineering and Biotechnology*, 19(1), pp.1-10. <https://doi.org/10.1186/s43141-020-00110-1>
- Rahn, K., De Grandis, S.A., Clarke, R.C., McEwen, S.A., Galán, J.E., Ginocchio, C., Curtiss III, R. and Gyles, C.L., 1992. Amplification of an *invA* gene sequence of *Salmonella typhimurium* by polymerase chain reaction as a specific method of detection of *Salmonella*. *Molecular and Cellular Probes*, 6(4), pp.271-279. [https://doi.org/10.1016/0890-8508\(92\)90002-f](https://doi.org/10.1016/0890-8508(92)90002-f)
- Sjahrani, T., Wasito, E.B. and Tyasningsih, W., 2021. Isolation and identification of *Escherichia coli* O157: H7 lytic bacteriophage from environment sewage. *International Journal of Food Science*, 2021, 7383121. <https://doi.org/10.1155/2021/7383121>
- SNI. 2332-11, 2017. Cara uji mikrobiologi - Dagian 11. Konfirmasi *Salmonella* spp. dengan teknik reaksi berantai polimerase (Polymerase Chain Reaction) konvensional pada produk perikanan. Badan Standardisasi Nasional, Jakarta.
- SNI. 2332.1, 2015. Cara uji mikrobiologi - Dagian 1. Penentuan koliform dan *Escherichia coli* pada produk perikanan. Badan Standardisasi Nasional, Jakarta.
- SNI. ISO 6579, 2015. Mikrobiologi bahan pangan dan pakan - Metode horizontal untuk deteksi *Salmonella* spp. Badan Standardisasi Nasional, Jakarta.
- SNI. ISO 11290—2, 2015. Mikrobiologi bahan pangan dan pakan - Metode horizontal untuk deteksi dan enumerasi *Listeria monocytogenes* - Dagian 2 : Metode enumerasi (ISO 11290-2:1998 dan Amd 1:2004, IDT.
- SNI. 2332.14, 2020. Cara Uji mikrobiologi - Dagian 14 : Deteksi *Listeria monocytogenes* pada produk

perikanan dengan metode polymerase chain reaction (PCR).

- Triwibowo, R., Rachmawati, N. and Dwiitno, 2020. Rapid and simultaneous detection of *Vibrio parahaemolyticus*, *Salmonella* spp. and *Escherichia coli* in fish by multiplex PCR. *Squalen Dulletin of Marine and Fisheries Postharvest and Diotechnology*, 15(2), pp.53-64. <http://dx.doi.org/10.15578/squalen.v15i2.444>
- Wei, L., Lanlan, L., Yun, N. and Xiaohong, L., 2007. Controllable nanocrystallization in Amorphous $Nd_5Fe_{85}B_6$ via combined application of severe plastic deformation and thermal annealing. *Applied Physics Letters*, 91(6), 062509. <https://doi.org/10.1063/1.2768023>
- Zhang, Z., Xiao, L., Lou, Y., Jin, M., Liao, C., Malakar, P.K., Pan, Y. and Zhao, Y., 2015. Development of a multiplex real-time PCR method for simultaneous detection of *Vibrio parahaemolyticus*, *Listeria monocytogenes* and *Salmonella* spp. in raw shrimp. *Food Control*, 51, pp.31-36. <https://doi.org/10.1016/j.foodcont.2014.11.007>



ISSN : 2301-7309
E-ISSN : 2528-0864

JOURNAL OF AQUACULTURE AND FISH HEALTH

VOLUME 11 NO. 3/SEPTEMBER 2022



FACULTY OF FISHERIES AND MARINE UNIVERSITAS AIRLANGGA

JAFH IS INDEXED BY :



Accredited by the Ministry of Research, Technology and Higher Education Number 23/E/KPT/2019



Editorial Team



Luthfiana Aprilianita Sari, S.Pi., M.Si

Editor in Chief

Department of Fish Health Management and Aquaculture, Faculty of Fisheries and Marine, Universitas Airlangga, Indonesia



-

-

57201778593



-



Prof. Felipe Polivanov Ottoni

Editorial Board

Laboratory of Systematics and Ecology of Aquatic Organisms (LASEOA), Center of Agrarian and Environmental Sciences (CCAA)

Universidade Federal do Maranhao, Brazil



-0000-0002-9390-0918

-

36619962800

-



Assoc Prof Nadirah binti Musa

Editorial Board

School of Fisheries and Aquaculture Sciences, Universiti Malaysia Terengganu, Kuala Terengganu, Terengganu, Malaysia Malaysia



-

-

23983138500



-



Prof. Marina Hassan

Editorial Board

Institute of Tropical Aquaculture, Universiti Malaysia Terengganu, 21030. Kuala Terengganu, Terengganu, Malaysia



-

-

36598787900



-



Siti Hudaidah, Ir., M.Sc

Editorial Board

Department of Aquaculture, Faculty of Agriculture, Universitas Lampung, Indonesia



-

-

57202393951



-



Prof. Mochammad Amin Alamsjah

Editorial Board

Department of Marine, Faculty of Fisheries and Marine, Universitas Airlangga, Indonesia



-

-

9635997700



-



Syifania Hanifah Samara, S.Pi., M.Sc

Managing Editor

Department of Fish Health Management and Aquaculture, Faculty of Fisheries and Marine, Universitas Airlangga, Indonesia



-

-

57207959121



-



Muhammad Browijoyo Santanumurti, S.Pi., M.Sc

Assistant Editor

Department of Fish Health Management and Aquaculture, Faculty of Fisheries and Marine, Universitas Airlangga, Indonesia



-

-

57201772053



-



Anita Erna Faricha, S. Ptk

Administrative

Faculty of Fisheries and Marine, Universitas Airlangga, Indonesia



-

-

-



-

Login

Username *

Password *

[Forgot your password?](#)

Keep me logged in

Login

[Register](#)

Accreditation Certificate

SERTIFIKAT



Vol. 11 No. 3 (2022): JAFH Vol. 11 No. 3 September 2022

Current Issue



Vol. 11 No. 3 (2022): JAFH Vol. 11 No. 3 September 2022

JAFH Vol. 11 No. 3 September 2022

Published: 2022-09-02

Articles

Effect of Spirulina (*Arthrospira platensis*) Powder on The Physico-chemical and Sensory Characterization of Dry Noodle

 Dinda Ersyah ⁽¹⁾, Abdul Aziz Jaziri ⁽²⁾, Dwi Setijawati ⁽³⁾

(1) Faculty of Fisheries and Marine Science, Universitas Brawijaya, Malang 65145, Indonesia ,

(2) 1 Faculty of Fisheries and Marine Science, Universitas Brawijaya, Malang 65145, Indonesia 2 Bioseafood Research Group, Faculty of Fisheries and Marine Science, Universitas Brawijaya, Malang 65145, Indonesia. 3 Halal Thoyyib Science Center, Universitas Brawijaya, Malang 65145, Indonesia ,


(3) Faculty of Fisheries and Marine Science, Universitas Brawijaya, Malang 65145, Indonesia

 277-288

 Abstract : 449

 pdf : 249

 pdf

 DOI : 10.20473/jafh.v11i3.20908

Effect of Feeding Fermented *Leucaena leucocephala* Seed Meal on Growth and Protein Content of *Osphronemus gouramy* Juvenile

 Putri Dista Ananda ⁽¹⁾, Safrida Safrida ⁽²⁾, Khairil Khairil ⁽³⁾, Hasanuddin Hasanuddin ⁽⁴⁾, Asiah M.D. ⁽⁵⁾

(1) Department of Biology Education, Faculty of Teacher Training and Education, Syiah Kuala University, Aceh 23111, Indonesia. ,


(2) Universitas Syiah Kuala ,

(3) Department of Biology Education, Faculty of Teacher Training and Education, Syiah Kuala University, Aceh 23111, Indonesia. ,

(4) Department of Biology Education, Faculty of Teacher Training and Education, Syiah Kuala University, Aceh 23111, Indonesia. ,


(5) Department of Biology Education, Faculty of Teacher Training and Education, Syiah Kuala University, Aceh 23111, Indonesia.

 289-297


 Abstract : 369

 pdf : 105

 pdf

 DOI : 10.20473/jafh.v11i3.25879

Morphometric and Meristic Analysis of *Rasbora* in East Java Province

 P Kumaladewi⁽¹⁾, Mufasirin⁽²⁾, Nunuk Dyah Retno Lastuti⁽³⁾, Mochammad Amin Alamsjah⁽⁴⁾, Win Darmanto⁽⁵⁾, Sapto Andriyono⁽⁶⁾

(1) Master Program of Fisheries Science, Faculty of Fisheries and Marine, Universitas Airlangga ,


(2) Faculty of Veterinary Medicine, Universitas Airlangga ,

(3) Faculty of Veterinary Medicine, Universitas Airlangga ,

(4) Department of Marine, Faculty of Fisheries and Marine, Universitas Airlangga ,

(5) Faculty of Science and Technology, Universitas Airlangga ,


(6) H-Indeks : 4, Scopus ID : 57193704905, Department of Marine Science, Faculty of Fisheries and Marine, Universitas Airlangga

 298-305


 Abstract : 242

 pdf : 150

 pdf

 DOI : 10.20473/jafh.v11i3.30956

Combination of *Nigella sativa* and *Phyllanthus niruri* as An Immunostimulant for The Prevention of White Spot Disease in *Litopenaeus vannamei*

 Wida Lesmanawati⁽¹⁾, Wasmen Manalu⁽²⁾, Min Rahminiwati⁽³⁾, Muhammad Agus Suprayudi⁽⁴⁾, Sri Nuryati⁽⁵⁾

(1) 1 Technology and Management of Applied Aquaculture Hatchery, College of Vocational Studies, IPB University, Jl. Kumbang 14, Bogor, West Java 16151 2 Department of Anatomy, Physiology and Pharmacology, Faculty of Animal Medicine, IPB University, Jl. Lingkar Kampus IPB Darmaga, Bogor, West Java 16680, Indonesia ,

(2) Department of Anatomy, Physiology and Pharmacology, Faculty of Animal Medicine, IPB University, Jl. Lingkar Kampus IPB Darmaga, Bogor, West Java 16680, Indonesia ,


(3) Department of Anatomy, Physiology and Pharmacology, Faculty of Animal Medicine, IPB University, Jl. Lingkar Kampus IPB Darmaga, Bogor, West Java 16680, Indonesia ,

(4) Department of Aquaculture, Faculty of Fisheries and Marine Science, IPB University, Jl. Lingkar Kampus IPB Darmaga, Bogor, West Java 16680, Indonesia ,


(5) Department of Aquaculture, Faculty of Fisheries and Marine Science, IPB University, Jl. Lingkar Kampus IPB Darmaga, Bogor, West Java 16680, Indonesia

 306-316


 Abstract : 252

 pdf : 121

 pdf

 DOI : 10.20473/jafh.v11i3.27470

Analysis of Nutritional Value and Consumer Behavior Diversification of Milkfish (*Chanos sp.*) Products From Aquaculture Products By Packaging Modification In MSME Sukolilo, Surabaya

 Patmawati Patmawati⁽¹⁾, Sri Subekti⁽²⁾, Wahyu Tjahjaningsih⁽³⁾, Ida Ayu Puspitasari⁽⁴⁾, Annisa Nur R. P.⁽⁵⁾, Mohamad Akmal A.H.⁽⁶⁾, Renyta Andini, W.⁽⁷⁾, Aliffiansyah Rizky E⁽⁸⁾

(1) Department of Marine, Faculty of Fisheries and Marine Universitas Airlangga, Jl. Mulyorejo, Surabaya 60115, East Java, Indonesia ,

(2) Department of Marine, Faculty of Fisheries and Marine Universitas Airlangga, Jl. Mulyorejo, Surabaya 60115, East Java, Indonesia ,

(3) Department of Marine, Faculty of Fisheries and Marine Universitas Airlangga, Jl. Mulyorejo, Surabaya 60115, East Java, Indonesia ,


(4) Study Programme of Fisheries Products Technology, Faculty of Fisheries and Marine, Universitas Airlangga, Jl. Mulyorejo, Surabaya 60115, East Java, Indonesia ,

(5) Study Programme of Fisheries Products Technology, Faculty of Fisheries and Marine, Universitas Airlangga, Jl. Mulyorejo, Surabaya 60115, East Java, Indonesia ,


(6) Study Programme of Fisheries Products Technology, Faculty of Fisheries and Marine, Universitas Airlangga, Jl. Mulyorejo, Surabaya 60115, East Java, Indonesia ,

(7) Study Programme of Fisheries Products Technology, Faculty of Fisheries and Marine, Universitas Airlangga, Jl. Mulyorejo, Surabaya 60115, East Java, Indonesia ,


(8) Study Programme of Fisheries Products Technology, Faculty of Fisheries and Marine, Universitas Airlangga, Jl. Mulyorejo, Surabaya 60115, East Java, Indonesia

 317-326

 Abstract : 339

 pdf : 106

 pdf

 DOI : 10.20473/jafh.v11i3.31382

Embryonic and Larval Development Stages of the African Catfish *Clarias gariepinus* (Burchell, 1822) (Teleostei, Clariidae) in The Ouargla, Algeria


 Ibrahim elkhail BEHMENE ⁽¹⁾, Benabdallah Bachir Bouiadjra ⁽²⁾, Miloud Attou ⁽³⁾, Abdelkader Homrani ⁽⁴⁾

(1) 1 Laboratory of Sciences and Technics of Animal Production (LSTAP), Abdelhamid Ibn Badis University, Mostaganem, Algeria 2 Department of Marine Science and Aquaculture, Abdelhamid Ibn Badis University, Mostaganem, Algeria ,

(2) 1 Laboratory of Sciences and Technics of Animal Production (LSTAP), Abdelhamid Ibn Badis University, Mostaganem, Algeria 2 Department of Marine Science and Aquaculture, Abdelhamid Ibn Badis University, Mostaganem, Algeria ,

(3) 1. Department of Marine Science and Aquaculture, Abdelhamid Ibn Badis University, Mostaganem, Algeria. 2. The experimental station fishing and inland aquaculture (mobile hatchery) located in Sidi bel abbes ,


(4) Laboratory of Sciences and Technics of Animal Production (LSTAP), Abdelhamid Ibn Badis University, Mostaganem, Algeria

 327-334


 Abstract : 203

 pdf : 115

 pdf

 DOI : 10.20473/jafh.v11i3.31871

Effect of Additional Aloe Vera (Aloe vera) on Artificial Feeds to Blood Cell Profile and Growth of Milkfish Seed (*Chanos chanos*)

 Linayati Linayati ⁽¹⁾, Nur Tri Jayanto ⁽²⁾, Tri Yusufi Mardiana ⁽³⁾, Muhammad Zulkham Yahya ⁽⁴⁾




(1) Aquaculture Department, Fisheries Faculty, Pekalongan University, Sriwijaya Street 3 Pekalongan, Central Java 51119, Indonesia ,

(2) Aquaculture Department, Fisheries Faculty, Pekalongan University, Sriwijaya Street 3 Pekalongan, Central Java 51119, Indonesia ,


(3) Aquaculture Department, Fisheries Faculty, Pekalongan University, Sriwijaya Street 3 Pekalongan, Central Java 51119, Indonesia ,

(4) Aquaculture Department, Fisheries Faculty, Pekalongan University, Sriwijaya Street 3 Pekalongan, Central Java 51119, Indonesia

Indonesia

 335-344 Abstract : 236 pdf : 79 pdf DOI : 10.20473/jafh.v11i3.32688

Vertical Infections and Prevalence of Viral Nervous Necrosis (VNN) in Milkfish (*Chanos chanos*)




 Sang Ayu Made Putri Suryani⁽¹⁾, Ni Made Darmadi⁽²⁾, Made Goura Premananda⁽³⁾, Gusti Ngurah Permana⁽⁴⁾

(1) Water Resources Management Study Program, Faculty of Agriculture, Warmadewa University, Jl. Terompong, Denpasar 80239, Bali, Indonesia ,

(2) Water Resources Management Study Program, Faculty of Agriculture, Warmadewa University, Jl. Terompong, Denpasar 80239, Bali, Indonesia ,

(3) Magister Programme of Environment Science, Udayana University. Bukit Campus, Jl. Raya Campus Unud Jimbaran, Badung Regency, 80361. Bali. Indonesia ,

(4) Center for Research and Development of Marine Cultivation, Buleleng

 345-351 Abstract : 179 pdf : 91 pdf DOI : 10.20473/jafh.v11i3.32798


Characterization of Salt Quality from *Artemia* (*Artemia franciscana*) Cultured Water with *Dunaliella salina* as Natural Feed

 Himna Sayyyidatul Islamiyah⁽¹⁾, Mochammad Amin Alamsjah⁽²⁾, Laksmi Sulmartiwi⁽³⁾


(1) Master Program of Fisheries and Marine Biotechnology, Universitas Airlangga, Jl. Mulyorejo, Surabaya 60115, East Java, Indonesia ,

(2) epartment of Marine, Faculty of Fisheries and Marine, Universitas Airlangga, Jl. Mulyorejo, Surabaya 60115, East Java, Indonesia ,

(3) epartment of Marine, Faculty of Fisheries and Marine, Universitas Airlangga, Jl. Mulyorejo, Surabaya 60115, East Java, Indonesia

 352-358 Abstract : 196 pdf : 115 pdf DOI : 10.20473/jafh.v11i3.34790

Determiation of The Best Concentration of Chitosan as a Recirculation Filter for Growth and Survival of Tilapia (*Oreochromis niloticus*)

 Miska Sanda Lembang⁽¹⁾, Widiawati Widiawati⁽²⁾

(1) Aquaculture Study Program, Faculty of Fisheries and Marine, University of Borneo Tarakan, Jalan Pantai Amal 1, Tarakan, North Kalimantan 77123, Indonesia ,

(2) Aquaculture Study Program, Faculty of Fisheries and Marine, University of Borneo Tarakan, Jalan Pantai Amal 1, Tarakan, North Kalimantan 77123, Indonesia

 359- Abstract : 192 pdf :



DOI : 10.20473/jafh.v11i3.34929

Solid Fermentation of Pelletized Fish Feeds Containing Black Soldier Fly (*Hermetia illucens*) Larvae Meal to Enhance Growth Performance of Catfish (*Clarias sp.*)

Cita Mahardika Hariyono ⁽¹⁾, Catur Sriherwanto ⁽²⁾, Harijono Harijono ⁽³⁾

(1) Department of Agricultural Product Technology, Faculty of Agricultural Technology, Brawijaya University, Veteran Street, Lowokwaru, Malang, 65145, Indonesia ,

(2) Applied Microbiology Research Centre, National Research and Innovation Agency (BRIN), Science and Technology Park, South Tangerang, Banten, 15314, Indonesia ,

(3) Department of Agricultural Product Technology, Faculty of Agricultural Technology, Brawijaya University, Veteran Street, Lowokwaru, Malang, 65145, Indonesia

367-379

Abstract : 227

pdf : 100



DOI : 10.20473/jafh.v11i3.34956

Vitamin C Improves the Efficacy of Oxytetracycline in Treatment of *Aeromonas hydrophila*-infected Juvenile Catfish (*Clarias gariepinus*)

Afisu Basiru ⁽¹⁾, Dr. Ajadi AA ⁽²⁾, Dr. Ahmed O. A ⁽³⁾, Dr. Suleiman K.O ⁽⁴⁾, Prof. Ameen S.A ⁽⁵⁾

(1) Department of Veterinary Physiology and Biochemistry, Faculty of Veterinary Medicine, University of Ilorin, Nigeria. ,

(2) Department of Veterinary Pathology, Faculty of Veterinary Medicine, University of Ilorin, Nigeria ,

(3) Department of Veterinary Microbiology, Faculty of Veterinary Medicine, University of Ilorin, Nigeria ,

(4) Department of Veterinary Physiology and Biochemistry, Faculty of Veterinary Medicine, University of Ilorin, Nigeria ,

(5) Department of Veterinary Medicine, Faculty of Veterinary Medicine, University of Ilorin, Nigeria

380-390

Abstract : 214

pdf : 78



DOI : 10.20473/jafh.v11i3.35153

Detection of *Salmonella* spp., *Escherichia coli*, and *Listeria monocytogenes* in Tuna by Multiplex PCR

Kristina Natalia ⁽¹⁾, Mariana Wahjudi ⁽²⁾, Tjie Kok ⁽³⁾

(1) Faculty of Biotechnology, University of Surabaya, Jalan Raya Kali Rungkut, Surabaya 60293, East Java, Indonesia 2 Regional Fish Quarantine and Inspection Agency, Surabaya I 61254, East Java, Indonesia ,

(2) Faculty of Biotechnology, University of Surabaya, Jalan Raya Kali Rungkut, Surabaya 60293, East Java, Indonesia ,

(3) Faculty of Biotechnology, University of Surabaya, Jalan Raya Kali Rungkut, Surabaya 60293, East Java, Indonesia

391-401


Abstract : 243

pdf : 109



DOI : 10.20473/jafh.v11i3.36509

Correlation of Density and Water Quality to Cortisol of Cantang Grouper Fish (*Epinephelus fuscoguttatus* X *Epinephelus lanceolatus*) in Floating Net Cages

 Laksmi Sulmartiwi ⁽¹⁾, Bintang Eka Nugraha Satia Aditama ⁽²⁾, Isnatul Umu Shohifah ⁽³⁾, Alfian Rahmadhani Satria Hani Saputra ⁽⁴⁾


(1) Department of Marine, Faculty of Fisheries and Marine, Universitas Airlangga, Jl. Mulyorejo, Surabaya 60115, East Java, Indonesia ,

(2) Study Program of Aquaculture, Faculty of Fisheries and Marine, Universitas Airlangga, Jl. Mulyorejo, Surabaya 60115, East Java, Indonesia ,

(3) Study Program of Aquaculture, Faculty of Fisheries and Marine, Universitas Airlangga, Jl. Mulyorejo, Surabaya 60115, East Java, Indonesia ,


(4) Study Program of Aquaculture, Faculty of Fisheries and Marine, Universitas Airlangga, Jl. Mulyorejo, Surabaya 60115, East Java, Indonesia

 402-411

 Abstract : 229

 pdf : 92



 DOI : 10.20473/jafh.v11i3.36661

Login

Username *

Password *

[Forgot your password?](#)

Keep me logged in

Login

Register

Accreditation Certificate



Focus and Scope

Publication Ethics

Article Processing Fee

Peer Reviewers Process

Peer Reviewers	Open Access Statement
Plagiarism	Copyright Notice
Contact	Article Withdrawal

Meet Our Editorial Team



Luthfiana Aprilianita Sari, S.Pi., M.Si
Editor Chief
Universitas Airlangga, Indonesia
57201778593



Syifania Hanifah Samara, S.Pi., M.Sc
Managing Editor
Universitas Airlangga, Indonesia
57207959121

[Read More](#)

Instruction For Author

Guide for authors	Online Submission
-----------------------------------	-----------------------------------

Article Template



Reference Manager



Grammar Editor



Back Issu JAFH



THIS JOURNAL IS SUPPORTED BY



Indexing






Current Issue

ATOM 1.0

RSS 2.0

RSS 1.0

Keywords



Stats



Address

Department of Aquaculture, Faculty of Fisheries and Marine, Universitas Airlangga Campus C UNAIR - Mulyorejo, Jl. Dharmahusada Permai No. 330, Mulyorejo, Surabaya, East Java 60115

Contact Info:

Phone: (031) 5911541

Fax: (031) 5911541

Email: luthfianaas@fpk.unair.ac.id



This work is licensed under a [Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License](https://creativecommons.org/licenses/by-nc-sa/4.0/).