

# Fingerprint and Multivariate Analysis of *Apium Graveolens* L. From Different Geographic with Spectroscopic ATR-FTIR

Wendy Nora Martian<sup>1</sup>, Dini Kesuma<sup>2,\*</sup>, Rima Via Angraini<sup>1</sup>

Wendy Nora Martian<sup>1</sup>, Dini Kesuma<sup>2,\*</sup>, Rima Via Angraini<sup>1</sup>

<sup>1</sup>Master of Industrial Pharmacy, Faculty of Pharmacy, University of Surabaya, Surabaya, East Java 60293, INDONESIA.

<sup>2</sup>Department of Pharmaceutical Chemistry, Faculty of Pharmacy, University of Surabaya, Surabaya, East Java 60293, INDONESIA.

## Correspondence

Dini Kesuma

Department of Pharmaceutical Chemistry, Faculty of Pharmacy, University of Surabaya, Surabaya, East Java 60293, INDONESIA.

E-mail: dinikesuma75@gmail.com

## History

- Submission Date: 27-10-2022;
- Review completed: 10-12-2022.
- Accepted Date: 10-12-2022.

DOI : 10.5530/pj.2022.14.205

Article Available online

<http://www.phcogj.com/v14/i6>

## Copyright

© 2022 Phcogj.Com. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International license.

## ABSTRACT

**Background:** *Apium graveolens* L. or widely known as celery is a plant that comes from the Apiaceae family. *Apium graveolens* L. is found in the Indonesian Herbal Pharmacopoeia as a nutritious plant. Plants that spread throughout Indonesia, both highlands and lowlands, contain 3-n-butylphthalide, D-limonene, Linalool, Luteolin, Apigenin and Apiin compounds. Differences in geographical location of growth resulted in differences in the content and concentration of *Apium graveolens* L. **Aim and Objectives:** This study was conducted to determine the differences of *Apium graveolens* L. based on its distribution in Indonesia. Indonesia does not yet have a standard for *Apium graveolens* L. **Material and Methods:** this research uses *Apium graveolens* provided by the Center for Research and Development of Traditional Medicinal and Medicinal Plants (B2P2TOOT) Tawangmangu. 10 samples from different regions in Indonesia will be read using the ATR-FTIR instrument in the finger print area as well as PCA (principal principal) multivariate analysis. component analysis) and HCA (Hierarchical Clustering Analysis). **Results:** The PCA results show the highest number of variants on PC 1 91,98% and PC 2 variants at 4,71%. In the quadrant 4 score plot, it was found that there was a closeness between the standard sample and the S2 sample (Materia Medika Batu). **Conclusion:** The result of this study is that there is a mutually supportive relationship from spectrum readings, sample location (temperature, soil moisture content, height (m a.s.l) with multivariate analysis (PCA, HCA results) when compared to the geographic location of each sample. **Key words:** *Apium graveolens* L, ATR-FTIR fingerprint area, Multivariate analysis.

## INTRODUCTION

Indonesia is identical with biological wealth which causes many plants to be used as alternative medicine, as well as making herbal medicine products widely circulated. Data from BPOM (food and drug regulatory agency) from 2017-2022, herbal medicine has a higher number than synthetic drugs.<sup>1,2</sup> The data is directly proportional to consumer demand. All herbal medicines are summarized in the Indonesian herbal pharmacopoeia. Starting from organoleptic to testing, everything is written in the Indonesian herbal pharmacopoeia.<sup>3</sup> This is done as a regulation and ensures the safety level of herbal medicine. *Apium graveolens* L. is one of the plants contained in the Indonesian Herbal Pharmacopoeia. This plant has properties as a blood pressure-lowering (anti-hypertensive) with several components in it, including 3-n-butylphthalide, D-limonene, Linalool, Luteolin, Apigenin and Apiin compounds.<sup>4-6</sup> Indonesian people very often use this plant as a vegetable in cooking and in the form of herbal medicine. In the Indonesian Herbal Pharmacopoeia mentioned *Apium graveolens* L. has flavonoids as much as 1.98% in the form of Apigenin. Differences in the chemical content of herbal medicines are influenced by several factors including the location of cultivation, harvest time, and differences in varieties of raw materials.<sup>7</sup> BPOM (regulatory agency in Indonesia) has issued many licenses for *Apium graveolens* L products, standardization is needed from the selection of raw materials to finished products.<sup>3</sup> This standardization requires QC (Quality Control) to improve the quality of herbal medicine products.

Starting from the evaluation of macroscopic and microscopic observations on specimens as well as chemical, biological and physicochemical analysis.<sup>8</sup> From the raw material, assessment of stability, efficacy, safety of the finished product, and the provision of product information to ensure proper use by consumers. The need for an instrument that provides fast, sensitive and accurate results is the ATR-FTIR spectroscopic.<sup>9-11</sup> Attenuated Total Reflectance - Fourier Transform Infrared (ATR - FTIR) is an instrument that helps to read the spectrum, and functional groups of a compound. FTIR spectrum contains multivariate data in the form of absorbance in the range of wave numbers 4000 - 650 cm<sup>-1</sup>.<sup>12</sup> Each sample has a different spectrum identity, the spectrum has dominant peaks that show the differences in each sample.<sup>8,12,13</sup> ATR-FTIR spectroscopy method combined with multivariate analysis is highly recommended for qualitative or quantitative QC tools for individual or mixed herbal medicine.<sup>14-16</sup> Multivariate analysis used is PCA (Principal component analysis) as could be used as classification and HCA (Hierarchical Cluster Analysis) relating the variables on a object to a discrete value of property one of wishes to predict, grouping, the herbal component from different origin.<sup>17,18</sup>

## MATERIALS AND METHODS

### Material

Standard of *Apium graveolens* L. came from the Center for Research and Development of Medicinal Plants and Traditional Medicines (B2P2TOOT) Tawangmangu and UPT Laboratory of Herbal

**Cite this article:** Martian WN, Kesuma D, Angraini RV. Fingerprint and Multivariate Analysis of *Apium Graveolens* L. From Different Geographic with Spectroscopic ATR-FTIR. Pharmacogn J. 2022;14(6)Suppl: 1022-1028.



**Figure 1:** Fresh *Apium graveolens* L.



**Figure 2:** Dry *Apium graveolens* L.



**Figure 3:** Powdered *Apium graveolens* L.

Materia Medika Batu. *Apium graveolens* L. samples were selected from 10 regions in Indonesia that have been determined.

Tools The tools used in this study include a blender, a sieve with a mesh size of 80, as well as checking the moisture content with a Moisturizer Analysis Hb43 Mettler-Toledo GmbH tool, using an Olympus Type x 23 microscope, ATR-FTIR (Attenuated Total Reflectance Fourier spectrophotometry Transform Infrared) Agilent carry 630, Origin 2018 multivariate analysis software.

### Methods sample preparation

Standard sample B2P2TOOT Tawangmangu was dried in an oven at a temperature of +/- 420C, while the other plant samples were air-dried in a room protected from direct sunlight. All plant samples were mashed with a blender and sieved with a mesh size of 80, and the moisture content was checked using Moisture Analysis Hb43 Mettler-Toledo GmbH.

### FTIR absorbance spectrum

In this study, a qualitative analysis will be carried out, selecting samples of celery plants taken from several islands in Indonesia with the aim of being able to describe the condition of celery plants scattered in Indonesia. The reading of all samples, both celery and capsules, used the Agilent carry 630 ATR-FTIR (Attenuated Total Reflectance Fourier Transform Infrared) spectrophotometric instrument using the Quality Assessment method in the Lab. Research Faculty of Pharmacy, University of Surabaya, Surabaya, East Java. Readings of all samples, both plant and commercial capsules, produced absorbance spectra, readings were carried out at a wavelength of  $4000\text{ cm}^{-1} - 650\text{ cm}^{-1}$  using a resolution of  $4\text{ cm}^{-1}$  which is commonly used for solid samples and 64 scans.<sup>11</sup> Tested on standard samples in accordance with the requirements in FHI (Indonesian Herbal Pharmacopoeia). Followed by qualification using polystyrene to ensure that the absorbance readings must be carried out before readings with the ATR-FTIR instrument. After that, the standard sample of celery was read 3 times for 3 days to see the results of absorption in the hope of having the same pattern. All samples of celery and capsules were read with the ATR-FTIR instrument to get the wave number and absorbance results. Selected finger print area at wave number  $1500\text{ cm}^{-1} - 650\text{ cm}^{-1}$  to get the pattern for each sample.<sup>19,20</sup> Absorbance results are continued for the calculation of %cv (coefficient variation) and SD (standard deviation). The requirements for the sample to be able to do multivariate test are to have %cv.

### Multivariate analysis

The multivariate method (MVA) is a statistical technique that allows simultaneous analysis of several variables. This helps to study relationships on a large scale as well as complex data sets. This study uses Origin 2018 Software as a tool for multivariate analysis. The multivariate analysis used is Principal Component Analysis (PCA) to explain the variance-covariance structure of a set of variables through linear combinations as well as a dimension reduction technique and Hierarchical Clustering Analysis (HCA) to classify samples in certain clusters so that they can see the closeness of each sample.

### RESULT

Physical characteristics of *Apium graveolens* L. This study used 10 samples of *Apium graveolens* spread across Indonesia. Initial observations in accordance with the provisions in the Indonesian



**Figure 4:** Physical observations.




Figure 5: Microscopic observations.




Figure 6: Absorbance spectrum (Finger Print) - standard replication in 3 days.




Figure 7: Absorbance spectrum standard overlay.




Figure 8: Absorbance spectrum standard.




Figure 9: Absorbance spectrum standard and 10 sample overlay.




Figure 10: PCA Plot Score (*Apium graveolens* L.).




Figure 11: HCA (*Apium graveolens* L.).

Table 1: Geographical location and climatic conditions of *Apium graveolens* L.

Code	Location	Geography position	Heigh	Rain fall	State of Water in Soil	Temperature (°C)
S1	Sumatera, Lampung Sedayu, semaka, Tanggamus, No. Determination : 074/497/102.7-A/2021	5°31'55"S 104°27'52"E	513 m a.s.l	200 - 300 mm	80 - 100%	27.563
S2	Jawa, Jawa Timur UPT lab. Matera Medika, Jl. Lahor no.87, Batu, No. Determination : 074/466/102.7-A/2021	7°55'04"S 112°31'10"E	895 m a.s.l	100 - 150 mm	80 - 100%	22.016
S3	Jawa, Jawa Timur Kebun Inspirasi Sultan Hidroponik, Sumber Pinang, Kecamatan Silo, Jember, No. Determination: 074/524/102.7-A/2021	8°14'42"S 113°50'38"E	241 m a.s.l	100 - 150 mm	-	27.44
S4	Kalimantan, Kalimantan Selatan Tariwin, Batu Mandi, Balangan, No. Determination : 074/520/102.7-A/2021	2°28'04"S 115°31'43"E	50 m a.s.l	200 - 300 mm	80 - 100%	26.75
S5	Sulawesi, Sulawesi Selatan Jalan Poros Loka, Bonto Marranu, Uluere, kab. Bantaeng, No. Determination : 074/521/102.7-A/2021	5°26'53"S 119°54'27"E	1.130 m a.s.l	100 - 150 mm	80 - 100%	26.77
S6	Bali, The Blooms garden, Cadikuning, Baturiti, Tabanan No. Determination : 074/522/102.7-A/2021	8°18'12"S 115°09'38"E	1.224 m a.s.l	100 - 150 mm	60 - 80%	27.86
S7	Lombok, NTB Mantang, Batukliang, lombok tengah No. Determination : 074/523/102.7-A/2021	8°36'37"S 116°19'21"E	366 m a.s.l	100 - 150 mm	80 - 100%	27.36
S8	Surabaya, Jawa timur Kampung Simolorejo, kel. Simomulyo, kec. Sukomanunggal, Surabaya No. Determination : 074/663/102.20-A/2022	7°15'24"S 112°42'22"E	4 m a.s.l	20 - 50 mm	20 - 40%	31
S9	Sampang, Madura Jl. Purba Gg: 2 No:11, Kec.Pamekasan, Kab. Pamekasan, Madura No. Determination : 074/664/102.20-A/2022	7°09'52"S 113°28'58"E	14 m a.s.l	20 - 50 mm	0 - 20%	32
S10	Nabire, Papua Tengah Asrama Denzipur 12, Jl. R.E. Martadinata, Sanoba, Distrik Nabire, Kabupaten Nabire, Papua tengah	3°20'22"S 135°32'26"E	24 m a.s.l	150 - 200 mm	60 - 80%	32
STD	Jawa, Jawa Tengah B2P2TOOT Jl. Raya lawu, tawangmangu, karanganyar No. Determinasi : YK.01.03/2/1175/2021	7°37'03"S 111°03'06"E	1.200 m a.s.l	100 - 150 mm	80 - 100%	22.81

**Table 2: Moisture content *Apium graveolens* L.**

No.	Sample	%MC
1	S1	2.7
2	S2	3.0
3	S3	2.7
4	S4	2.8
5	S5	2.7
6	S6	2.5
7	S7	2.5
8	S8	2.8
9	S9	2.6
10	S10	2.8
11	STD	2.8

Herbal Pharmacopoeia include physical observations, microscopic tests and testing of water content. From the shape of a single leaf, thin, brittle, jagged leaf, pinnate leaf bone, dark green in Figure 4 in accordance with the provisions in FHI. In addition to looking at the physical form of *Apium graveolens*, it was found macroscopically using an Olympus Type X 23 microscope with a magnification of 40x/0.6 in the form of petiole epidermis, upper epidermis with stomata, transport bundle with ladder type thickening and idioblasts (lower epidermis oil cells with stomata) in accordance with the provisions FHI in Figure 5. This shows the conformity of the sample with the provisions of FHI.

Moisture content (MC) can be measured with a %MC less than 10% as shown in Table 2. Sample S2 results. The reading is carried out at a wavelength of 1500 – 650  $\text{cm}^{-1}$  or what is often called the fingerprint area due to the large number of dominant peaks. From samples. The reading of the absorbance spectrum of the sample is carried out based on Lambert-beer's law which states that there is a linear relationship between absorbance and concentration.<sup>14</sup> From Figure 8. Shows (Materia medika Batu) has a high MC compared to the other samples, but all samples are included in the FHI provisions which state that the water content for celery plants is less than 10%.<sup>3</sup>

### ATR-FTIR absorbance spectrum

From *Apium graveolens* L. The reading of a standard sample of *Apium graveolens* (Tawangmangu) using the ATR-FTIR Agilent 630 Carry is shown in Figure 6. performed 3 times a day and for 3 consecutive days, in order to provide accurate, the dominant peak of *Apium graveolens* at wave numbers 883.3790  $\text{cm}^{-1}$  and 928.1071  $\text{cm}^{-1}$  which indicates a strong absorption of the functional group = C-H, at wave numbers 1017.3369  $\text{cm}^{-1}$ , 1075.3369  $\text{cm}^{-1}$ , 1196.4754  $\text{cm}^{-1}$ , 1244.9308  $\text{cm}^{-1}$ , 1319.4775  $\text{cm}^{-1}$  and 1375.3876  $\text{cm}^{-1}$  which indicate a strong absorption of the C – O functional group.

The results of standard readings with other samples were made in overlay view, to find out the similarities or differences. (Figure 9) shows the similarity between the standard and the S2 sample (Materia medika).

### PCA multivariate analysis (PCA)

The first multivariate analysis is principal component analysis (PCA) which reduces all data without eliminating existing data but is made in more representative results. In Figure 7. are the results of 10 samples and 1 standard sample? The PCA results show the highest number of variants on PC 1 91,98% and PC 2 variants at 4,71%. In the quadrant 4 score plot, it was found that there was a closeness between the standard sample and the S2 sample (Materia, Medika, Batu). It can be interpreted that the standard and S2 have the same chemical properties, as well as samples S8 and S9, S6 and S4, S7 and S5. This is supported by Table 1. Regarding the temperature and altitude of the location.

### HCA multivariate analysis (HCA)

(HCA) is a continuation of PCA, HCA is a dendrogram that describes groups in research, as well as to compare PCA results. In Figure 8 there are 2 large clusters namely STD, S2, S1, S5, S7, S3, S4, S6 and S8, S10, S9. HCA gives the same results as PCA, which means that HCA and PCA are chemometric analysis tests that support each other in differentiating samples qualitatively.

### DISCUSSION

The results of this study are that the ATR-FTIR instrument can provide precise qualitative testing to make it easier for researchers to differentiate each sample by reading the absorbance spectrum. The resulting spectrum results give dominant peaks which are signs on the test sample. each peak shows a wave number which can determine its functional group.

This replication is very important to provide accurate results in research. Indonesia does not have herbal plant standards, but the B2P2TOOT Tawangmangu government agency has celery plants (*Apium graveolens* L) and issued a letter of determination. *Apium graveolens* L from B2P2TOOT Tawangmangu is used as a standard, but replication testing is needed to ensure the correctness of the ATR-FTIR standard and instrument. On the first day, the standard *Apium graveolens* L was read 3 times, followed by the second and third days. the results of this replication reading are overlaid so that the similarities of each spectrum can be seen, if there are differences it is necessary to re-test to ensure the accuracy of the instrument.

replication of all samples was also read, after all samples were confirmed to be the same as the replication, then proceed to data analysis with Origin 2018 software. PCA and HCA multivariate data analysis was carried out. for qualitative testing, PCA and HCA analysis is enough, but in the future PLS-DA and SIMCA tests are needed.

In the PCA test, it was found that the standard score plot display has the same quadrant as the S2 sample which proves that the S2 sample from the government agency Materia Medika Batu has the same quality as the standard. The PCA results can be related to the geographic location in table 1. The PCA plot score results will be processed by HCA analysis. HCA makes it easier for researchers to differentiate and classify.

### CONCLUSION

The use of the ATR-FTIR instrument with a combination of multivariate analysis gives good results for qualitative research. It is necessary to optimize the standard (Tawangmangu) to ensure the accuracy of the instrument and the authenticity of the standard, bearing in mind that Indonesia does not issue herbal medicine plant standards. Spectrum readings based on the fingerprint area (1500 - 650 $\text{cm}^{-1}$ ) can determine the functional group of each peak. PCA results, HCA has a mutually supportive relationship to the geographic location of each sample related to temperature, rainfall, High, water content in the soil. Provide initial screening results to determine the quality and authenticity of *Apium graveolens* L.

### ACKNOWLEDGEMENT

This research was funded by the Institute of the Ministry of Research and Technology of the Republic of Indonesia with a letter of assignment no. 159/E5/P6.02.00.PT/2022 dated 10 May 2022.

### REFERENCES

1. <https://cekbpom.pom.go.id/> Indonesia: Badan Pengawas Obat dan Makanan. [updated 2022 october 12; cited 2022 November 9].
2. Kesuma D, Makayasa CHA, Suhud F, Azminah A, Yuniarta T, Sumartha IGA, et al. Structure Modification: Effect of Lipophilic,

- Electronic, And Steric Parameters of N-Benzoyl-N'-Phenylthiourea Compounds On Antiviral Activity of Covid-19 By in Silico. *Rasayan J Chem.* 2022;15(2):1445-9.
3. Farmakope Herbal Indonesia Edisi Ii 2017 Kementerian Kesehatan Republik Indonesia 615.1. 2017.
  4. Hedayati N, Naeini MB, Mohammadinejad A, Mohajeri SA. Beneficial effects of celery (*Apium graveolens*) on metabolic syndrome: A review of the existing evidences. *Phytotherapy Res.* 2019;33(12):3040-53.
  5. Jorge VG, Ángel JL, Adrián T, Francisco A, Anuar S, Samuel E. Vasorelaxant activity of extracts obtained from *Apium graveolens*: Possible source for vasorelaxant molecules isolation with potential antihypertensive effect. *Asian Pac J Trop Biomed.* 2013;3(10):776-9.
  6. Tashakori-Sabzevar F, Razavib BM, Imenshahidic M, Daneshmandia M, Fatehia H, Sarkarizi YE, *et al.* Evaluation of mechanism for antihypertensive and vasorelaxant effects of hexanic and hydroalcoholic extracts of celery seed in normotensive and hypertensive rats. *Revista Brasileira de Farmacognosia.* 2016;26(5):619-26.
  7. Ingallina C. Phytochemical and biological characterization of Italian "sedano bianco di Sperlonga" Protected Geographical Indication celery ecotype: A multimethodological approach. *Food Chem.* 2020;309:125649.
  8. Indrayanto G. Recent Development of Quality Control Methods for Herbal Derived Drug Preparations. *SAGE J.* 2018.
  9. Rytwo G, Zakai R, Wicklein B. The use of ATR-FTIR spectroscopy for quantification of adsorbed compounds. *J Spectroscopy.* 2015;2015.
  10. Wulandari L, Idroes R, Noviandy TR, Indrayanto G. Application of chemometrics using direct spectroscopic methods as a QC tool in pharmaceutical industry and their validation. *Profiles Drug Subst Excip Relat Methodol.* 2022;47:327-79.
  11. Canteri MHG, Renard CMGC, le Bourvellec C, Bureau S. ATR-FTIR spectroscopy to determine cell wall composition: Application on a large diversity of fruits and vegetables. *Carbohydr Polym.* 2019;212:186-96.
  12. Noviana E, Indrayanto G, Rohman A. Advances in Fingerprint Analysis for Standardization and Quality Control of Herbal Medicines. *Front Pharmacol.* 2022;13:853023.
  13. Ren J. Analytical strategies for the discovery and validation of quality-markers of traditional Chinese medicine. *Phytomedicine.* 2020;67:153165.
  14. Indrayanto G, Rohman A. The Use of FTIR Spectroscopic Combined with Multivariate Analysis in food composition Analysis. Springer Nature Singapore Pte Ltd.; 2020.
  15. Dubey RS, Saravanan S. Optical Study of Sol-Gel Spin-Coating TiO<sub>2</sub>/Si<sub>3</sub>N<sub>4</sub> Multilayers On Glass Substrates. *Rasayan J Chem.* 2022;15(4):2304-9.
  16. Syafri S, Jaswir I, Yusof F, Rohman A, Hamidi D. The Use of GC-MS and Ftir Spectroscopy Coupled with Multivariate Analysis for The Detection of Red Ginger Oil Adulteration. *Rasayan J Chem.* 2022;15(4):2231-6.
  17. Prikeržnik M, Srčič S. Multivariate analysis for optimization and validation of the industrial tablet-manufacturing process. *Drug Dev Ind Pharm.* 2021;47(1):61-71.
  18. Ferreira AP, Toba M. Multivariate analysis in the pharmaceutical industry: Enabling process understanding and improvement in the PAT and QbD era. *Pharm Dev Technol.* 2015;20(5):513-27.
  19. Easmin S. Rapid investigation of  $\alpha$ -glucosidase inhibitory activity of *Phaleria macrocarpa* extracts using FTIR-ATR based fingerprinting. *J Food Drug Anal.* 2017;25(2):306-15.
  20. Sam S, Alisha Md Isha Ali S, Muddukrishna BS, Vasantharaju SG. Chemical Fingerprinting of Piperine and Tannic Acid in an Ayurvedic Formulation of Dasanakanthi Churnam Using RP-HPLC. *Rasayan J Chem.* 2022;15(4):2218-25.

## ABOUT AUTHORS



DR. DINI KESUMA Is an Associate Professor in the Department of Pharmaceutical Chemistry, Faculty of Pharmacy, University of Surabaya, Indonesia. She has experience in medicinal chemistry and analytical chemistry. The main focus of her research and publication is cancer drug development in silico and in vitro.



WENDY NORA MARTIAN is currently a Master of Industrial Pharmacy student, Faculty of Pharmacy, University of Surabaya, Indonesia.



RIMA VIA ANGRAINI is currently a Master of Industrial Pharmacy student, Faculty of Pharmacy, University of Surabaya, Indonesia.

**Cite this article:** Martian WN, Kesuma D, Angraini RV. Fingerprint and Multivariate Analysis of *Apium Graveolens* L. From Different Geographic with Spectroscopic ATR-FTIR. *Pharmacogn J.* 2022;14(6)Suppl: 1022-1028.

[Home](#)

[About Journal](#) ▾

[Editorial Board](#)

[For Authors](#) ▾

[Contact Us](#)

# Pharmacognosy Journal

An Open Access, Peer Reviewed Journal in the field of  
Pharmacognosy



# Editorial Board (2020-21) | Pharmacognosy Journal

## Editors & Editorial Board Members (2021)

### **Dr.Djemli Samir**

Department of Biology , Applied Neuroendocrinology Laboratory  
Badji Mokhtar Annaba University  
Algeria

### **Dr. Raghava Naidu, Ph.D**

Department of Human Oncology,  
University of Wisconsin,  
1111, Highland Ave, Madison,  
Wisconsin 53705, USA

### **Dr.Karim Raafat**

Associate Professor of Pharmacognosy and Phytochemistry,  
Pharmaceutical Sciences Department,  
Faculty of Pharmacy,  
Beirut Arab University (BAU),  
Beirut 115020, Lebanon

### **Ourlad Alzeus Tantengco, MD-PhD Molecular Medicine**

College of Medicine, University of the Philippines Manila  
Pedro Gil Street, Ermita, Manila, Philippines, 1000

### **Janib Achmad**

Lecturer of Faculty of Fisheries and Marine Science,  
University of Khairun Ternate  
Kampus 2 JalanPertamina, KelurahanGambesi,  
Ternate Selatan

### **Muammar Fawwaz, Ph.D**

Department of Pharmaceutical Chemistry  
Faculty of Pharmacy  
Universitas Muslim Indonesia  
Makassar 90231, South Sulawesi, Indonesia

### **Hany Ezzat Khalil**

Associate Professor,  
College of Clinical Pharmacy,  
King Faisal University,  
KSA

### **Emad Yousif**

Department of Chemistry  
College of Science  
Al-Nahrain University  
Baghdad,Iraq

### **Sughosh Upasani**

R.C Patel Institute of pharmacy,  
Shirpur,Dist-Dhule,Maharashtra,  
India.

### **Gurusiddaiah suresh kumar**

Scientist  
Dept of biochemistry  
CSIR-CFTRI  
Mysore, Karnataka, INDIA

### **Arjun Patra**

Assistant Professor  
School of Pharmaceutical Sciences  
Guru Ghasidas Central University  
Koni, Bilaspur - 495 009  
Chattisgarh, India

### **Francis O. Atanu, Ph.D**

Department of Biochemistry  
Faculty of Natural Sciences  
Kogi State University

University of the West Indies  
St. Augustine, Trinidad & Tobago.

**Dr. Kunle Okaiyeto, PhD**

Applied and Environmental Microbiology Research Group (AEMREG)  
Department of Biochemistry and Microbiology  
University of Fort Hare  
Alice campus  
5700, Alice  
South Africa.

**Dr. Srisailam Keshetti, Ph.D**

Principal, University College of Pharmaceutical Sciences, Satavahana University  
Karimnagar 505001  
Telangana  
INDIA

**Dr. Gayathri M Rao**

Associate Professor  
Department of Biochemistry  
Kasturba Medical College, Mangaluru.

**Shuge Tian**

Experimental Teaching Demonstration Center of TCM in Xinjiang Medical University  
Department of traditional medicine, TCM  
Xinjiang Medical University  
Xinjiang CHINA 830054

**Dr. Ramachandra Setty Siddamsetty,**

Professor, Govt College of Pharmacy,  
Mission Road, Bengaluru, INDIA

**Dr. (Mrs.) Sayyada Khatoon**

HOD, Pharmacognosy Division  
CSIR-National Botanical Research Institute,  
Rana Pratap Marg, Post Box 436,  
Lucknow-226001 (U.P.) India

**Dr. A. Sajeli Begum**

Department of Pharmacy  
Birla Institute of Technology & Science  
Hyderabad, India

**Olga Silva**

Department of Pharmacological Sciences,  
Faculdade de Farmácia,  
Universidade de Lisboa, Portugal

**Xinwen Wang**

Department of Clinical Pharmacy  
University of Michigan  
USA

**Roman Lysiuk**

Department of Pharmacognosy and Botany,  
Danylo Halytsky Lviv National Medical University,  
Pekarska, 69., Lviv 79010, Ukraine

**Arif Nur Muhammad Ansori**

Universitas Airlangga  
Indonesia

# Pharmacognosy Journal (/)

An Open Access, Peer Reviewed Journal in the field of  
Pharmacognosy

Enter terms then hit Search...

[Articles In Press \(/inpress\)](#)[Current Issue \(/v14/i6\)](#)[Archives \(/archives\)](#)[RSS Feeds \(/rss.xml\)](#)[Submit Article \(https://www.phcogj.info\)](https://www.phcogj.info)

(<http://www.phcogj.info>)

HOME (/) / PHARMACOGNJ VOL 14, ISSUE 6 (SUPPL), NOV-DEC, 2022

## PharmacognJ Vol 14, Issue 6 (Suppl), Nov-Dec, 2022

[View \(/pharmacognj-vol-14-issue-6-suppl-nov-dec-2022\)](#)[What links here \(/node/1893/backlink\)](#)

Submitted by sys1 (/users/sys1) on Wed, 12/28/2022 - 14:46

**Medicinal Plants Adopted to Treat Children's Diseases by Traditional Pediatrics "Women Healers" In The Souss Massa Region (Agadir Idaoutanan, Inzegane Ait Meloul and Chtouka Ait Baha) Morocco (/article/1924)**

**Differences in interleukin-6 and interleukin-17 expression in covid-19 post-mortem lung tissue biopsy compared with noncovid- 19 (/article/1925)**

**Kaempferia galanga L. Extract Administration Attenuate Aquaporin-4 Expression in Traumatic Brain Injury: An Experimental Study in Rats (/article/1926)**

**Effect of Pomegranate Extract On N-Terminal Pro Brain Natriuretic Peptide and Asymmetric Dimethylarginine Levels in Children with Pulmonary Artery Hypertension in Acyanotic Congenital Heart Disease (/article/1927)**

**Evaluation of Experimental Cerebral Malaria of Curcumin and Kaempferol in Plasmodium**



**berghei ANKA-Infected Mice (/article/1928)**

**Molecular Docking Estrogen Receptor Alpha Antagonist and P53- MDM2 Inhibitor, ADMET Prediction of Alkaloid Compound from Mitragyna speciosa for Breast Cancer Therapy (/article/1929)**

**Left Atrial Myxoma Presented with an Obstructive Shock, Right Ventricle Dysfunction and Pulmonary Hypertension (/article/1930)**

**Risk Factors for Hepatotoxicity From L-Asparaginase Chemotherapy In Children With Acute Lymphoblastic Leukemia (/article/1931)**

**Phytochemical Approach Including Total Phenolic and Flavonoid Contents and Evaluation of in vitro ABTS Antioxidant Capacity and Lipoxygenase Inhibition of Anisoscadium lanatum (/article/1894)**

**Cichorioside a biocoumarin modulates lipid and glucose storage on 3T3-L1 cell lines: In vitro and in silico approach (/article/1895)**

**Relationship Histopathology Grading of Meningioma with the Use of Medroxyprogesterone Acetate (MPA) as A Hormonal Contraceptive (/article/1932)**

**Correlation Between the Suitability of Empirical and Definitive Therapies According to Culture Results with the Clinical Outcomes of Patients with Bacteremia Due to Carbapenem- Resistant Acinetobacter baumannii (CRAB) at Dr. Soetomo Tertiary Referral Hosp (/article/1933)**

**GC-MS Analysis of Volatiles Present in Papeea Capensis Extracts (/article/1934)**

**Phytochemical Analysis and Antioxidant Activity of Water Hyacinth Flowers (Eichhornia Crassipes) Extract (/article/1935)**

**Research on External Signs and Chemical Composition of Medicinal Plant Raw Material -Leaves of Ficus Elastica (/article/1936)**

**Antiparasitic effect of Psidium guajava on promastigotes and axenic amastigotes of Leishmania (/article/1937)**

**The Serum Formulation of Hati Tanah Tuber Ethanol Extract from Central Kalimantan (/article/1938)**

**Job fulfilment and its related variables among pharmacy certificate holders in Jordan: A Cross-sectional Study (/article/1939)**

**Chemical Profiling of Nonpolar Compounds of Onopordum Acanthium using GCMASS (/article/1940)**

**Antioxidant, Antimicrobial, and Antiplasmodial Activities of Sonchus arvensis L. Leaf Ethyl Acetate Fractions (/article/1941)**

**Senna Siamea Hexane Extract: Potent Antifungal Activity Against Candida albicans, Candida Krusei and Identification of Its Chemicals Content (/article/1942)**



DFT and Pharmacokinetic Study of Some Heterocyclic Aspirin Derivatives as The Cyclooxygenase Inhibitors: An In-Silico Approach (/article/1943)

Fingerprint and Multivariate Analysis of Apium Graveolens L. From Different Geographic with Spectroscopic ATR-FTIR (/article/1944)

Antibacterial Effect of Nigella sativa L. Seed from Indonesia (/article/1945)

Successful Intrapericardial Fibrinolysis on Acute Recurrent Purulent Pericarditis with Impending Cardiac Tamponade during Pandemic Situation: A Rare Case Report (/article/1946)

Bio-Evaluation, In-Vitro and In-Vivo Anti-Inflammatory Activity, Therapeutic Efficacy, and Genotoxicity of the Potentials of the Green Seaweed Valoniopsis Pachynema using Zebra Fish Larvae (Danio Rerio) as an Animal Model (/article/1947)

Aquaporine 4 Expression on End Feet Astrocyte Before and After Cerebrospinal Fluid Drainage of Hydrocephalus Mice Model (/article/1948)

Effects of Pomegranate Extract Supplementation (Punica granatum L.) on Clinical Manifestations of Pulmonary Arterial Hypertension in Children with Acyanotic Congenital Heart Disease (/article/1949)

The Role of Breast Milk on Reducing the Risk of Neonatal Sepsis in Preterm and Low Birth Weight Infants: A Systematic Review and Meta-Analysis (/article/1950)

Histopathological Perspectives of Multiple Organs in a Red- Footed Tortoise (Chelonoidis carbonaria) with Suspected Metabolic Bone Disease: A Case Report (/article/1951)

Medicinal Plants Adopted to Treat Children's Diseases by Traditional Pediatrics "Women Healers" In The Souss Massa Region (Agadir Idaoutanan, Inzegane Ait Meloul and Chtouka Ait Baha) Morocco › (/article/1924)

73 reads

## SHARE THIS ARTICLE

EMAIL	FACEBOOK	LINKEDIN	TWITTER	GOOGLE+	PINTEREST
(MAILTO:?SUBJECT=PHARMACOGNOSIS%20ISSUE%206%20%28SUPPL%29%2C%20NOV-DEC%2C%202022&BODY=HTTPS%3A%2F%2FWWW.PHCOGJ.COM%2FPHARMACOGNOSIS%20ISSUE-6-SUPPL-NOV-DEC-2022)	/SHARER.PHP?U=HTTPS%3A%2F%2FWWW.PHCOGJ.COM%2FPHARMACOGNOSIS%20ISSUE-6-SUPPL-NOV-DEC-2022)	/SHAREARTICLE?MINI=TRUE&URL=HTTPS%3A%2F%2FWWW.PHCOGJ.COM%2FPHARMACOGNOSIS%20ISSUE-6-SUPPL-NOV-DEC-2022&TITLE=PHARMACOGNOSIS%20ISSUE-6-SUPPL-NOV-DEC-2022	/TWEET?TEXT=PHARMACOGNOSIS%20ISSUE%206%20%28SUPPL%29%2C%20NOV-DEC%2C%202022&URL=HTTPS%3A%2F%2FWWW.PHCOGJ.COM%2FPHARMACOGNOSIS%20ISSUE-6-SUPPL-NOV-DEC-2022	/SHARE?URL=PHARMACOGNOSIS%20ISSUE%206%20%28SUPPL%29%2C%20NOV-DEC%2C%202022&HTTPS%3A%2F%2FWWW.PHCOGJ.COM%2FPHARMACOGNOSIS%20ISSUE-6-SUPPL-NOV-DEC-2022	PHARMACOGNOSIS%20ISSUE%206%20%28SUPPL%29%2C%20NOV-DEC%2C%202022

---

## About

Pharmacognosy Journal (Phcog J.) covers different topics in natural product drug discovery, and also publishes manuscripts that describe pharmacognostic investigations, evaluation reports, methods, techniques and applications of all forms of medicinal plant research

**Distinctions:** The most widely read, cited, and known Pharmacognosy journal and website is well browsed with all the articles published. More than 50,000 readers in nearly every country in the world each month

**ISSN :** 0975-3575 ; **Frequency :** Rapid at a time publication (6 issues/year)

**Indexed and Abstracted in :** SCOPUS, Scimago Journal Ranking, Chemical Abstracts, Excerpta Medica / EMBASE, Google Scholar, CABI Full Text, Index Copernicus, Ulrich's International Periodical Directory, ProQuest, Journalseek & Genamics, PhcogBase, EBSCOHost, Academic Search Complete, Open J-Gate, SciACCESS.

**Rapid publication:** Average time from submission to first decision is 30 days and from acceptance to In Press online publication is 45 days.

**Open Access Journal:** Pharmacognosy Journal is an open access journal, which allows authors to fund their article to be open access from publication.

## Submit your Next Article

- Online submission
- Highly indexed and abstracted
- 10 years of successful publishing
- Wider visibility though open access
- Higher impact with wider visibility
- Prompt review

## **Submit** your next article to Phcog J

and be a part of many successful authors.

**Create free account** (<http://phcogj.info>) / **Login** (<https://www.phcogj.com/submissions/index.php/phcogj/index>)



**Pharmacognosy Journal** and its contents are licensed under a Creative Commons Attribution-Non Commercial-No Derivs 4.0 License. Permissions beyond the scope of this license may be available with [editor@phcogj.com](mailto:editor@phcogj.com) (<mailto:editor@phcogj.com>)

[Home \(/\)](#)

[Advertise with us \(/\)](#)

[Privacy Statement \(/\)](#)



# Pharmacognosy Journal

COUNTRY	SUBJECT AREA AND CATEGORY	PUBLISHER	H-INDEX
<a href="#">India</a>  Universities and research institutions in India	<a href="#">Pharmacology, Toxicology and Pharmaceutics</a> <a href="#">Drug Discovery</a> <a href="#">Pharmacology</a>	<a href="#">EManuscript Technologies</a>	<h1>25</h1>
PUBLICATION TYPE	ISSN	COVERAGE	INFORMATION
Journals 	09753575	2009-2021	<a href="#">Homepage</a>  <a href="#">How to publish in this journal</a> <a href="mailto:editor@phcogj.com">editor@phcogj.com</a>

**SCOPE**

Pharmacognosy Journal (Phcog J.) covers different topics in natural product drug discovery, and also publishes manuscripts that describe pharmacognostic investigations, evaluation reports, methods, techniques and applications of all forms of medicinal plant research

 Join the conversation about this journal









 Quartiles

**FIND SIMILAR JOURNALS** 

<p>1 <b>Oriental Pharmacy and Experimental Medicine</b> USA 92%</p>	<p>2 <b>Pharmacognosy Research</b> IND 81%</p>	<p>3 <b>Indian Journal of Natural Products and Resources</b> IND 77%</p>	<p>4 <b>Journal of Herbs, Spices and Medicinal Plants</b> USA 76%</p>	<p>5 <b>Pharmacognosy Magazine</b> IND 76%</p>
---	--	--	---	--

options 





Pharmacognosy Journal

← Show this widget in your own website

Q3 Pharmacology  
best quartile

SJR 2021  
0.26

powered by scimagojr.com

Just copy the code below and paste within your html code:

<a href="https://www.scim."

SCImago Graphica

Explore, visually communicate and make sense of data with our new data visualization tool.

Metrics based on Scopus® data as of April 2022

E **edi utomo** 3 weeks ago

I want to get information about drug poison crimes, is there a journal about this?

reply

**Melanie Ortiz** 3 weeks ago

Dear Edi,

SCImago Team

# Source details

## Pharmacognosy Journal

Scopus coverage years: from 2009 to 2022

Publisher: Pharmacognosy Network Worldwide

ISSN: 0975-3575

Subject area: [Pharmacology, Toxicology and Pharmaceutics: Pharmacology](#)

[Pharmacology, Toxicology and Pharmaceutics: Drug Discovery](#)

Source type: Journal

[View all documents >](#)

[Set document alert](#)

[Save to source list](#)

CiteScore 2021

1.9



SJR 2021

0.258



SNIP 2021

0.718



[CiteScore](#) [CiteScore rank & trend](#) [Scopus content coverage](#)

CiteScore 2021

$$1.9 = \frac{1,760 \text{ Citations 2018 - 2021}}{946 \text{ Documents 2018 - 2021}}$$

Calculated on 05 May, 2022

CiteScoreTracker 2022

$$1.7 = \frac{1,479 \text{ Citations to date}}{847 \text{ Documents to date}}$$

Last updated on 06 December, 2022 • Updated monthly

### CiteScore rank 2021

Category	Rank	Percentile
Pharmacology, Toxicology and Pharmaceutics <a href="#">Pharmacology</a>	#219/303	27th
Pharmacology, Toxicology and Pharmaceutics <a href="#">Drug Discovery</a>	#116/154	25th

[View CiteScore methodology >](#) [CiteScore FAQ >](#) [Add CiteScore to your site](#)

## About Scopus

[What is Scopus](#)

[Content coverage](#)

[Scopus blog](#)

[Scopus API](#)

[Privacy matters](#)

## Language

[日本語版を表示する](#)

[查看简体中文版本](#)

[查看繁體中文版本](#)

[Просмотр версии на русском языке](#)

## Customer Service

[Help](#)

[Tutorials](#)

[Contact us](#)

---

## ELSEVIER

[Terms and conditions ↗](#) [Privacy policy ↗](#)

Copyright © Elsevier B.V. ↗. All rights reserved. Scopus® is a registered trademark of Elsevier B.V.

We use cookies to help provide and enhance our service and tailor content. By continuing, you agree to the use of cookies ↗.

