

DOI <http://dx.doi.org/10.36722/sst.v8i1.1494>

Antibacterial Activity of *Drosera* Sp. Ethanolic Extract Against *Staphylococcus Aureus* ATCC 25923

Alfian Hendra Krisnawan^{1*}, Ridho Islamie², Rahma Dhianitasari Wahyu Putri³, Ni Putu Citra Ratna Dewi³

¹Departement of Pharmaceutical Biology, Faculty of Pharmacy, University of Surabaya, Jl. Raya Kalirungkut, Surabaya, 60293

²Departement of Clinical Pharmacy and Community, Faculty of Pharmacy, University of Surabaya, Jl. Raya Kalirungkut, Surabaya, 60293

³Bachelor of Pharmacy Program, Faculty of Pharmacy, University of Surabaya, Jl. Raya Kalirungkut, Surabaya, 60293

*Corresponding author: alfian_hendra_k@staff.ubaya.ac.id

Abstract – *Drosera sp.* is an insectivorous plant with secondary metabolites, including naphthoquinones which can be used as antibacterial agents. *Staphylococcus aureus* is a bacterium that often causes nosocomial infections, such as ARI. This research aims to determine the antibacterial activity of four *Drosera* species, namely *D. burmannii*, *D. sessilifolia*, *D. capillaris*, and *D. filiformis*, against *Staphylococcus aureus* ATCC 25923. The agar diffusion method was employed in antibacterial testing by measuring the inhibition zone. The inhibition zone measured using a 7,000 ppm concentration of *Drosera sp.* ethanolic extract was 6.74 ± 0.21 mm, 9.21 ± 0.32 mm, 18.18 ± 0.60 mm, and 15.67 ± 0.37 mm for *D. burmannii*, *D. sessilifolia*, *D. capillaris*, and *D. filiformis*, respectively. The four *Drosera sp.* extracts were proven to have antibacterial potential against *Staphylococcus aureus* ATCC 25923. The largest value was produced by *D. capillaris* which was close to 19.73 ± 0.55 mm from the positive control (chloramphenicol).

Keywords – Antibacterial activity, *Drosera sp.*, *Staphylococcus aureus*

INTRODUCTION

Nosocomial infections initiate various diseases presented in almost all healthcare facilities, specifically in hospitals, both in Indonesia and worldwide. These are caused by pathogenic microorganisms, including *Staphylococcus aureus* which has developed resistance to certain antibiotics, including penicillin, ampicillin, and vancomycin [1], leading to treatment difficulty. *Staphylococcus aureus* causes an acute respiratory infection (ARI) [2], with symptoms such as fever, cough, and sore throat. ARI prevalence in Indonesia is 4.4% and the highest, 12.8%, occurs in children below five years [3].

Infectious diseases are generally managed with antibiotics, but in some cases of bacterial resistance, alternative treatments are needed, such as using herbs against gram-positive and gram-

negative bacteria. *Drosera* is a species of carnivorous herbal plant that can be used as an alternative, containing secondary metabolites applied traditionally as raw materials for pharmaceutical preparations [4]. In Indonesia, the extract has been used as a mixture of active ingredients in formulating cough medicines. *Drosera* contains phenols, flavonoids (Quercetin), anthocyanins, and naphthoquinones, namely plumbagin and 7-methyljuglone [5], [6], which are used to treat respiratory tract infections, such as bronchitis and tuberculosis [7], some of its pharmacological activities include antioxidant, anti-inflammatory, and anticancer effects [8].

The application of *D. rotundifolia*'s extract against gram-negative bacteria, such as *Escherichia coli*, *Salmonella enteritidis*, *Yersinia enterocolitica*, and gram-positive bacteria such as *Bacillus thuringiensis*, *Clostridium perfringens*, *Listeria*

monocytogenes, and *Staphylococcus aureus* [9], [10], the combination of *D. binata* extract with silver nanoparticles (AgNPs) increases the activity against *Staphylococcus aureus* [11], in this research, the antibacterial activity of *D. burmannii*, *D. sessilifolia*, *D. capillaris*, and *D. filiformis* in Indonesia was determined against *Staphylococcus aureus*. These four species have similarities, a tendency to grow in the southern hemisphere [12], and contain compounds including phenols, flavonoids, alkaloids, and anthraquinone with antioxidant effects [13], [14].

METHODS

This study is an in vitro experimental study using a Post-Test Only Control Group Design conducted at the Phytochemical and Microbiology Laboratory, Faculty of Pharmacy, University of Surabaya for 5 months.

Materials

The herbal plant samples used were four *Drosera* species, namely *D. burmannii*, *D. sessilifolia*, *D. capillaris*, and *D. filiformis*. The plants were grown on sphagnum-moss media in Surabaya-East of Java, Indonesia. The test bacterium employed was *Staphylococcus aureus* ATCC 25923 which was grown at the Faculty of Pharmacy, Surabaya University.

Preparation of *Drosera sp.* Herb Extract

Dry *Simplicia* powder of *Drosera sp.* was extracted by maceration method using 96% ethanol as solvent. Maceration was carried out for 24 hours with 6 hours of immersion and stirring. The filtrate was separated from the residue which was macerated again twice. This was combined with the second filtrate and processed into a concentrated extract, then the yield of each sample was determined [15].

Preparation of Test Solution

Approximately 70 mg of the concentrated extract was weighed and dissolved in 10 ml of 10% DMSO. Therefore, a concentration of 7,000 ppm was obtained and graded from 3,000 ppm to 6,000 ppm. The 10% DMSO was applied as negative control while the positive was chloramphenicol. *Staphylococcus aureus* ATCC was cultured on Nutrient Agar slanted (NA) media and incubated at 37°C for 24 hours. This was suspended in liquid media with the addition of 3 mL of 0.9% NaCl and homogenized, then the turbidity of the bacterial

suspension was measured using a 0.5 McFarland densitometer [16].

Antibacterial Activity Test

A petri dish containing liquid agar and 100 L of *Staphylococcus aureus* suspension was allowed to solidify. Afterward, wells were made using a cork bore with a 6 mm diameter and each was filled with *Drosera sp.* at five different concentrations, negative control, and positive control. The treatment was incubated at 37°C for 24 hours and then a clear inhibition zone was marked on the agar medium to indicate the absence of bacterial growth [17].

RESULTS AND DISCUSSION

Drosera sp. Herb Extract

The maceration method used 96% ethanol solvent to extract the active compound content of *Drosera sp.* herb powder. This allows the obtained compounds including plumbagin, anthraquinone, and flavonoid to be undamaged and more stable than the extraction method conducted with high temperatures which can affect antibacterial activity [18], in this research, the ethanol extract of *D. spatulata* var. *bakoensis* produced better antibacterial activity than aqueous and methanol extracts [7], the yield obtained from the four *Drosera sp.* ranged between 23 - 66%, meaning many compounds were extracted from the herb with ethanol solvent. It is also a *Simplicia*'s non-specific parameter that can be used as quality control of a traditional medicinal raw material [19], and the data are presented in Table 1.

Table 1. The yield of *Drosera sp.* Extract

| Drosera Species | Yield (%) |
|------------------------|------------------|
| <i>D. burmannii</i> | 24.51 |
| <i>D. sessilifolia</i> | 23.73 |
| <i>D. capillaris</i> | 37.30 |
| <i>D. filiformis</i> | 66.81 |

Antibacterial activity of *Drosera sp.*

Antibacterial activity was determined through the ability of *Drosera sp.* to hinder bacterial growth on solid NA media which was characterized by an inhibition zone. The four species at concentrations of 3,000, 4,000, 5,000, 6,000, and 7,000 ppm produced inhibition zones against *Staphylococcus aureus*, and the results are presented in Table 2.

Table 2. Determination of Inhibition Zone of *Drosera sp.* Extract against *Staphylococcus aureus*

| Spesies <i>Drosera</i> | Diameter of Inhibition Zone (mm) | | | | | Chloramphenicol | DMSO 10% |
|------------------------|----------------------------------|--------------|--------------|--------------|--------------|-----------------|----------|
| | a | b | c | d | e | | |
| <i>D.burmanniei</i> | 0 | 0 | 7.32 ± 0.5 | 7.81 ± 0.3 | 8.19 ± 0.16 | 19.64 ± 1.25 | 0 |
| <i>D. sessilifolia</i> | 0 | 0 | 8.02 ± 0.69 | 8.82 ± 0.33 | 9.13 ± 0.49 | 20.71 ± 0.9 | 0 |
| <i>D. capillaris</i> | 9.90 ± 0.31 | 11.02 ± 0.38 | 12.24 ± 0.31 | 14.21 ± 0.4 | 17.70 ± 0.61 | 19.37 ± 1.09 | 0 |
| <i>D. filiformis</i> | 8.97 ± 1.2 | 9.68 ± 0.3 | 10.58 ± 0.29 | 14.04 ± 0.37 | 16.78 ± 0.39 | 20.30 ± 0.65 | 0 |

Description: a) 3,000 ppm, b) 4,000 ppm, c) 5,000 ppm, d) 6,000 ppm, e) 7,000 ppm

Based on the antibacterial test, the concentration of *Drosera sp.* used ranged from 3,000 to 7,000 ppm. *D. burmannii* and *D. sessilifolia* extracts did not produce inhibition zones at concentrations of 3,000 ppm and 4,000 ppm, but they began at 5,000 ppm – 7,000 ppm. Meanwhile, *D. capillaris* and *D. filiformis* extracts produced inhibition zones from concentrations of 3,000 to 7,000 ppm. Greater extract concentration generates a higher inhibition zone. From the four samples, 7,000 ppm led to a high inhibition zone as demonstrated in Fig. 1. Similarly, Rita et al. (2018) explained that elevations in both the inhibition zone and extract concentration were directly proportional [20].

Also, the content of active compounds in the extract and antibacterial activity increased concurrently. In the comparison of antibacterial activity between samples at a 7,000 ppm concentration, the *D. capillaris* extract produced the largest inhibition zone diameter of 17.70 mm±0.61 mm. This was followed by *D. Filiformis*, *D. Burmannii*, and *D. sessilifolia* with values of

16.78 mm ± 0.39 mm, 8.19 mm ± 0.16 mm, and 9.13 mm ± 0.49 mm, respectively. The chloramphenicol positive control had 19.37–20.71 mm, while two *Drosera sp.* which had values close to this range were retested simultaneously and they produced inhibition zones as presented in Table 3.

Table 3. Inhibition Zone Test for 7000 ppm concentration of *Drosera sp.* against *Staphylococcus aureus*

| <i>Drosera</i> Species | Diameter of Inhibition Zone (mm) |
|------------------------|----------------------------------|
| <i>D. burmannii</i> | 6.74 ± 0.21 |
| <i>D. sessilifolia</i> | 9.21 ± 0.32 |
| <i>D. capillaris</i> | 18.18 ± 0.60 |
| <i>D. fliformis</i> | 15.67 ± 0.37 |
| Chloramphenicol | 19.73 ± 0.55 |





Figure 1. Inhibition zone of *Drosera sp.* extract against *Staphylococcus aureus*Figure 2. Inhibition zone of *Drosera sp.* extract at 7000 ppm against *Staphylococcus aureus*

Another species, *D. magnifica*, has antibacterial activity that is synergistic with antibiotics and can suppress bacterial virulence factors in *in vitro* tests [21], the compound in *Drosera sp.* which acts as an antibacterial against gram-positive bacteria is plumbagin from the naphthoquinone group [11], other compounds such as alkaloids, anthraquinones, and flavonoids also have antimicrobial effects against microbes causing respiratory tract infections [7], the mechanism of antibacterial action is by inhibiting nucleic acid synthesis, cytoplasmic membrane function, energy metabolism, and porins in cell membranes, as well as changing membrane permeability [22].

D. capillaris and *D. filiformis* extracts at 7,000 ppm had antibacterial activity close to the positive control, namely chloramphenicol with an inhibition zone of 19.73±0.55 mm as demonstrated in Fig. 2. Natural ingredients can be classified as strong antibacterials once their inhibition zone value is equal to or better than the positive control. Compounds with an inhibition area diameter of 10-20 mm or more are claimed to contain strong antibacterial potential [23], the extracts of *D. capillaris*, *D. filiformis*, *D. burmannii*, and *D. sessilifolia* can have antibacterial potential at higher concentrations. However, their toxicity effect needs to be considered, because it increases alongside the concentration. In 2019, Lee did not reported death

and toxicity in mice but obtained an LD₅₀ value of 4,800 mg/kg by using single (300-1,200 mg/kg) and repeated doses (150-600 mg/kg) of *D. rotundifolia* for 4 weeks [24].

CONCLUSION

The 96% ethanol extract of four *Drosera sp.* produced antibacterial activity against *Staphylococcus aureus* ATCC 25923 as indicated by the inhibition zones, hence they are potential antibacterial agents. *D. burmannii* and *D. sessilifolia* as well as *D. capillaris* and *D. filiformis* produced an inhibition zone at concentrations of 5000ppm and 3000 ppm, respectively. At a 7000 ppm concentration, the inhibition zones of each sp. extract were 6.74±0.21 mm, 9.21±0.32 mm, and 15.67±0.37 mm for *D. burmannii*, *D. sessilifolia*, and *D. filiformis*, respectively. Meanwhile, *D. capillaris* had the largest inhibition zone, which was 18.18±0.60 mm, and the activity closest to 19.73±0.55 mm from the positive control (*chloramphenicol*).

ACKNOWLEDGMENT

The authors are grateful to the Faculty of Pharmacy, University of Surabaya, for providing

support and laboratory facilities in carrying out this research which was independently funded.

REFERENCES

- [1] M. A. Akindolire, O. O. Babalola, and C. N. Ateba, "Detection of antibiotic resistant Staphylococcus aureus from milk: A public health implication," *Int J Environ Res Public Health*, vol. 12, no. 9, pp. 10254–10275, Aug. 2015, doi: 10.3390/ijerph120910254.
- [2] A. Fusvita and A. Umar, "Identifikasi Bakteri Pernafasan Penyebab Infeksi Saluran Pernafasan (ISPA) Pada Usia Balita di Rumah Sakit Bahteramas," *Jurnal Analisis kesehatan Kendari*, vol. 1, no. 1, pp. 40–46, 2016.
- [3] Ministry of Health of the Republic of Indonesia, "Laporan Nasional Riskesdas 2018," Jakarta, 2018.
- [4] P. A. Egan and F. van der Kooy, "Review - Phytochemistry of the Carnivorous Sundew Genus Drosera (Droseraceae) - Future Perspectives and Ethnopharmacological Relevance," *Chem Biodivers*, vol. 10, pp. 1774–1790, 2013, doi: 10.1002/cbdv.201200359.
- [5] B. Banyai, C. Bäcker, C. Reich, and U. Lindequist, "The production of 7-methyljuglone, plumbagin and quercetin in wild and cultivated Drosera rotundifolia and Drosera intermedia," *Mires and Peat*, vol. 18, 2016, doi: 10.19189/MaP.2016.OMB.228.
- [6] R. Ravee, F. I. M. Salleh, and H. H. Goh, "Discovery of digestive enzymes in carnivorous plants with focus on proteases," *PeerJ*, 2018, doi: 10.7717/peerj.4914.
- [7] G. Dileep Kumar, C. D. S. L. N. Tulasi, and G. Ramakrishnaiah, "Phytochemical screening and evaluation of in vitro antimicrobial activity of drosera spatulata var. Bakoensis - An indigenous carnivorous plant against respiratory tract infectious microbes," *Asian Journal of Pharmaceutical and Clinical Research*, vol. 9, no. 6, pp. 274–283, Nov. 2016, doi: 10.22159/ajpcr.2016.v9i6.14382.
- [8] R. Goswami, T. Sinha, and K. Ghosh, "Drosera Sp: A Critical Review on Phytochemical and Ethnomedicinal Aspect," *International Journal of Pharmacy and Biological Sciences-IJPBS TM*, no. 1, p. 9, 2019, doi: 10.21276/ijpbs.2019.9.1.76.
- [9] D. Ďurechová, M. Kačániová, M. Terentjeva, J. Petrová, L. Hleba, and I. Kata, "Antibacterial Activity of Drosera Rotundifolia L. Against Gram-Positive and Gram-Negative Bacteria," *Journal of microbiology, biotechnology and food sciences*, vol. 5, no. special 1, pp. 20–22, Feb. 2016, doi: 10.15414/jmbfs.2016.5.special1.20-22.
- [10] M. Kačániová *et al.*, "Antimicrobial Activity of Drosera rotundifolia L," *Scientific Papers: Animal Science and Biotechnologies*, vol. 47, no. 2, pp. 366–369, 2014.
- [11] M. Krychowiak *et al.*, "Combination of Silver Nanoparticles and Drosera binata Extract as a Possible Alternative for Antibiotic Treatment of Burn Wound Infections Caused by Resistant Staphylococcus aureus," *PLoS One*, vol. 9, no. 12, Dec. 2014, doi: 10.1371/journal.pone.0115727.
- [12] D. K. Biswal, S. Yanthan, R. Konhar, M. Debnath, S. Kumaria, and P. Tandon, "Phylogeny and biogeography of the carnivorous plant family Droseraceae with representative Drosera species from Northeast India," *F1000Res*, vol. 6, no. 1454, Aug. 2017, doi: 10.12688/f1000research.12049.1.
- [13] N. B. Ghate, D. Chaudhuri, A. Das, S. Panja, and N. Mandal, "An Antioxidant Extract of the Insectivorous Plant Drosera burmanii Vahl. Alleviates Iron Induced Oxidative Stress and Hepatic Injury in Mice," *PLoS One*, vol. 10, no. 5, 2015.
- [14] D. M. Septy, T. Saptawati, and F. A. Rachma, "The Anatomical Characteristic, Phytochemical Screening and Antioxidant Activity Test of Drosera capillaris Herbs Using the FRAP Method," in *Prosiding Seminar Nasional UNIMUS*, 2021, pp. 1272–1280.
- [15] Susanty and F. Bachmid, "Perbandingan Metode Ekstraksi Maserasi dan Refluks Terhadap Kadar Fenolik dari Ekstrak Tongkol Jagung (Zea mays L.)," *JURNAL KONVERSI*, vol. 5, no. 2, pp. 87–93, Oct. 2016, doi: 10.24853/konversi.5.2.87-92.
- [16] E. Gayathiri, B. Bharathi, and K. Priya, "Study of the Enumeration of Twelve Clinical Important Bacterial Populations at 0.5 McFarland Standard," *International*

- Journal of Creative Research Thoughts (IJCRT)*, vol. 6, no. 2, pp. 880–893, 2018.
- [17] H. Manurung, R. Agung Nugroho, and E. Marina, “Phytochemical Screening and Antibacterial Activity of Leaves Extract Balangla (*Litsea cubeba* (Lour) Pers.) from Malinau, East Borneo,” in *Proceeding of 5th International Seminar on New Paradigm and Innovation on Natural Sciences and Its Application (5th ISNPINSA)*, 2015, pp. 5–8.
- [18] C. Onyebuchi and D. Kavaz, “Effect of extraction temperature and solvent type on the bioactive potential of *Ocimum gratissimum* L. extracts,” *Sci Rep*, vol. 10, no. 1, Dec. 2020, doi: 10.1038/s41598-020-78847-5.
- [19] Ministry of Health of the Republic of Indonesia, “Farmakope Herbal Indonesia edisi II,” Jakarta, 2017.
- [20] W. S. Rita, I. M. D. Swantara, I. A. R. A. Asih, and N. K. Sinarsih, “Antibacterial Activity of *Samanea saman* Leaf Ethanol Extract Against *Escherichia coli* and *Staphylococcus aureus* and its Total Flavonoid And Phenolic Contents,” *Jurnal Kimia*, vol. 12, no. 2, pp. 121–127, 2018.
- [21] P. A. Singh and N. Bajwa, “Future Perspectives of *Drosera Magnifica*: The First Ever Plant Discovered on The Social Media,” *Journal of Drug Delivery and Therapeutics*, vol. 7, no. 4, pp. 12–19, Jul. 2017, doi: 10.22270/jddt.v7i4.1474.
- [22] Y. Xie, W. Yang, F. Tang, X. Chen, and L. Ren, “Antibacterial Activities of Flavonoids: Structure-Activity Relationship and Mechanism,” *Curr Med Chem*, vol. 22, no. 1, pp. 132–149, Sep. 2014, doi: 10.2174/0929867321666140916113443.
- [23] A. Trisia, R. Philyria, and A. N. Toemon, “Uji Aktivitas Antibakteri Ekstrak Etanol Daun Kalanduyung (*Guazuma ulmifolia* Lam.) Terhadap Pertumbuhan *Staphylococcus aureus* dengan Metode Difusi Cakram (Kirby-Bauer),” *Anterior Jurnal*, vol. 17, no. 2, pp. 136–143, 2018.
- [24] Y. E. Lee, J. H. Lee, Y. K. Lee, B. H. Lee, J. S. Kim, and H.-J. Lee, “A Study on Single-dose Toxicity and Repeated-Dose Toxicity of *Drosera Rotundifolia* L. Pharmacopuncture in Rodent Models,” *Korean Journal of Acupuncture*, vol. 36, no. 2, pp. 127–138, Jun. 2019, doi: 10.14406/acu.2019.013.

Jurnal AL-AZHAR INDONESIA

SERI SAINS DAN TEKNOLOGI

*SK Menteri Riset dan Teknologi/Badan Riset dan Inovasi Nasional
No. 158/E/KPT/2021*

Nutritional Knowledge of Prospective Brides In Indonesia: A Systematic Review
Mukhlidah Hanun Siregar

Kandungan Mineral dan Senyawa Fenolik Pada Kulat Basi (*Termitomyces sp.*) Asal Kabupaten Kapuas Hulu
Henny Sulistiany, Eka Trisianawati, Siti Suhartati

Sistem Identifikasi Titik Kritis Halal Menggunakan Algoritma *Forward Chaining*
Alexander Moya Hin, Adhi Kusnadi, Marlinda Vasty Overbeek, Oqke Prawira, Yaman Khaeruzzaman, Syarief Gerald Prasetya

Perencanaan Rute Distribusi Produk Pupuk NPK *Blending* Dengan Menggunakan Algoritma *Differential Evolution* di PT. XYZ
Qouman Sholihin, Dwi Sukma Donoriyanto

Anti bacterial Activity of *Drosera Sp.* Ethanolic Extract Against *Staphylococcus Aureus ATCC 25923*
Alfian Hendra Krisnawan, Ridho Islamie, Rahma Dhianitasari Wahyu Putri, Ni Putu Citra Ratna Dewi

Penerapan *Lean Tools Change Acceleration Process Model* dan *Spaghetti Diagram* Proses Produksi Sofa di PT. XYZ
Wan Habibi Rahman Barus, Iphov Kumala Sriwana

Karakteristik Mutu Sensori dan Kimia Liang Teh Pontianak Kaya Antioksidan Pada Berbagai Suhu Pengeringan
Devi Viola Florentina, Yohana Sutiknyawati Kusuma Dewi, Suko Priyono

Perilaku Gajah Sumatra (*Elephas maximus sumatranus*) Betina pada Kandang Dalam di Taman Margasatwa Ragunan, Jakarta
Nasya Aqila Firmanza, Luthfirda Sjahfirdi

Desain Model Kendali Mrac-pd Untuk Pengaturan Level Deaerator Pada PT Pertamina Ru II Dumai
Surya Andy Wananda, Putut Son Maria, Aulia Ullah, Jufrizel

Analisis Potensi Sampah di Tempat Pemrosesan Akhir Toisapu Sebagai Bahan Baku Pembangkit Listrik Tenaga Sampah di Kota Ambon
Heni Mutmainnah, Fajar Akbar Pandiangan, Akbar Khasemi Hamzah



Jurnal AL-AZHAR INDONESIA SERI SAINS DAN TEKNOLOGI

Volume 8, No. 1, Januari 2023

ISSN Cetak: 2087-9725

ISSN Online: 2355-8059

Akreditasi berdasarkan Surat Keputusan Direktur Jenderal Pendidikan Tinggi, Riset dan Teknologi
Nomor 158/E/KPT/2021, SINTA 3

Jurnal AL-AZHAR INDONESIA SERI SAINS DAN TEKNOLOGI merupakan jurnal ilmiah yang memuat makalah dan artikel hasil penelitian atau kajian teoritis dalam bidang Sains dan Teknologi.

Kata Al-Azhar berakar dari z-h-r. Verba zahara bermakna *to shine, give light, be radiant; to glow, gleam, glare, shine; to blossom, be in bloom* (Wehr, 1974: 384). Dengan demikian, kata Al-Azhar berarti sesuatu yang bersinar, yang memberi cahaya, yang berseri-seri atau yang mekar

Jurnal ini diterbitkan 3 (tiga) dalam setahun mulai tahun 2022 yaitu pada bulan Januari, Mei, dan September. Panduan penulisan disertakan pada halaman belakang jurnal

PIMPINAN REDAKSI

Dr. Dewi Elfidasari, S.Si., M.Si

DEWAN PENYUNTING

Dwi Astharini, ST., M.Sc

Ir. Winangsari Pradani, MT

Ema Komalasari, STP., M.Si

Amalina Ratih Puspa, S.P., M.Si

Dr. Ir. Syarif Hidayat, M.Eng.Sc., MM.

Dr. rer. nat. Yunus Effendi, S.Pd., M.Si., M.Sc.

SEKRETARIS REDAKSI

Kasyfil Azis, S.Pd

Nur Yunita Sari, A.Md

Selma Intan Praditya Sari Himawan, ST

MITRA BESTARI

Prof. Dr. Ir. Riri Fitri Sari, MM., M.Sc (UI)

Prof. Dr. drh. Retno D. Soejoedono MS (IPB)

Dr. Dasumiati, M. Si (UIN Syarif Hidayatullah Jakarta)

Dr. Iphov Kumala Sriwana, S.T, M. Si (Universitas Telkom)

Prof. Dr. Ir. Dadan Umar Daihani, DEA (Universitas Trisakti)

Irmawati, S. Kom, MMSI (Sekolah Tinggi Teknologi Bandung)

Prof. Dr. Ir. Andi Adriansyah, M. Eng (Universitas Mercubuana)

ALAMAT REDAKSI

Lembaga Penelitian dan Pengabdian Masyarakat

Universitas Al Azhar Indonesia, Komplek Masjid Agung Al Azhar,

Jl. Sisingamangaraja, Jakarta 12110, Telp: (021) 727 92753, Fax: (021) 724 4767,

email: lp2m@uai.ac.id/Website: <http://lp2m.uai.ac.id>

PENERBIT: LPPM Universitas Al Azhar Indonesia

DAFTAR ISI


DAFTAR ISI

i

| | |
|--|---------|
| Nutritional Knowledge of Prospective Brides In Indonesia: A Systematic Review <i>Mukhlidah Hanun Siregar</i> | 1 - 8 |
| Kandungan Mineral dan Senyawa Fenolik Pada Kulat Basi (<i>Termitomyces sp.</i>) Asal Kabupaten Kapuas Hulu <i>Henny Sulistiany, Eka Trisianawati, Siti Suhartati</i> | 9 - 17 |
| Sistem Identifikasi Titik Kritis Halal Menggunakan Algoritma <i>Forward Chaining</i> <i>Alexander Moya Hin, Adhi Kusnadi, Marlinda Vasty Overbeek, Oqke Prawira, Yaman Khaeruzzaman, Syarief Gerald Prasetya</i> | 18 - 23 |
| Perencanaan Rute Distribusi Produk Pupuk NPK <i>Blending</i> Dengan Menggunakan Algoritma <i>Differential Evolution</i> di PT. XYZ <i>Qouman Sholihin, Dwi Sukma Donoriyanto</i> | 24 - 32 |
| Antibacterial Activity of <i>Drosera Sp. Ethanolic</i> Extract Against <i>Staphylococcus Aureus ATCC 25923</i> <i>Alfian Hendra Krisnawan, Ridho Islamie, Rahma Dhianitasari Wahyu Putri, Ni Putu Citra Ratna Dewi</i> | 33 - 38 |
| Penerapan <i>Lean Tools Change Acceleration Process Model</i> dan <i>Spaghetti Diagram</i> Pada Proses Produksi Sofa di PT. XYZ <i>Wan Habibi Rahman Barus, Iphov Kumala Sriwana</i> | 39 - 48 |
| Karakteristik Mutu Sensori dan Kimia Liang Teh Pontianak Kaya Antioksidan Pada Berbagai Suhu Pengeringan <i>Devi Viola Florentina, Yohana Sutiknyawati Kusuma Dewi, Suko Priyono</i> | 49 - 57 |
| Perilaku Gajah Sumatra (<i>Elephas maximus sumatranus</i>) Betina pada Kandang Dalam di Taman Margasatwa Ragunan, Jakarta <i>Nasya Aqila Firmanza, Luthfiralda Sjahfirdi</i> | 58 - 66 |
| Desain Model Kendali Mrac-pd Untuk Pengaturan Level Deaerator Pada PT Pertamina Ru II Dumaiy <i>Surya Andy Wananda, Putut Son Maria, Aulia Ullah, Jufrizel</i> | 67 - 76 |
| Analisis Potensi Sampah di Tempat Pemrosesan Akhir Toisapu Sebagai Bahan Baku Pembangkit Listrik Tenaga Sampah di Kota Ambon <i>Heni Mutmainnah, Fajar Akbar Pandiangan, Akbar Khasemi Hamzah</i> | 77 - 85 |

Petunjuk Penulisan Artikel Jurnal AL-AZHAR INDONESIA SERI SAINS DAN TEKNOLOGI

Template Penulisan Jurnal AL AZHAR INDONESIA SERI SAINS DAN TEKNOLOGI



JURNAL AL-AZHAR INDONESIA SERI SAINS DAN TEKNOLOGI

[LPPM UNIVERSITAS AL AZHAR INDONESIA](#)

✳ P-ISSN : 20879725 <> E-ISSN : 23558059



1.5

Impact Factor



1370

Google Citations



Sinta 3

Current Accreditation

[Google Scholar](#)

[Garuda](#)

[Website](#)

[Editor URL](#)

History Accreditation

2019

2020

2021

2022

2023

2024

[Garuda](#)

[Google Scholar](#)

[Kandungan Mineral dan Senyawa Fenolik pada Kulit Basi \(Termitomyces sp.\) Asal Kabupaten Kapuas Hulu](#)

Universitas Al Azhar Indonesia [JURNAL AL-AZHAR INDONESIA SERI SAINS DAN TEKNOLOGI Vol 8, No 1 \(2023\): Januari 2023 9 - 17](#)

[2023](#) [DOI: 10.36722/sst.v8i1.1352](#) [Accred : Sinta 3](#)

[Desain Model Kendali MRAC-PD untuk Pengaturan Level Deaerator Pada PT Pertamina RU II Dumai](#)

Universitas Al Azhar Indonesia [JURNAL AL-AZHAR INDONESIA SERI SAINS DAN TEKNOLOGI Vol 8, No 1 \(2023\): Januari 2023 67 - 76](#)

[2023](#) [DOI: 10.36722/sst.v8i1.1761](#) [Accred : Sinta 3](#)

[Sistem Identifikasi Titik Kritis Halal Menggunakan Algoritma Forward Chaining](#)

Universitas Al Azhar Indonesia [JURNAL AL-AZHAR INDONESIA SERI SAINS DAN TEKNOLOGI Vol 8, No 1 \(2023\): Januari 2023 18 - 23](#)

[2023](#) [DOI: 10.36722/sst.v8i1.1285](#) [Accred : Sinta 3](#)




[Analisis Potensi Sampah Di TPA Toisapu Sebagai Bahan Baku Pembangkit Listrik Tenaga Sampah \(PLTSa\) Di Kota Ambon](#)

Universitas Al Azhar Indonesia [JURNAL AL-AZHAR INDONESIA SERI SAINS DAN TEKNOLOGI Vol 8, No 1 \(2023\): Januari 2023 77 - 85](#)

[2023](#) [DOI: 10.36722/sst.v8i1.1384](#) [Accred : Sinta 3](#)




[Perencanaan Rute Distribusi Produk Pupuk NPK Blending Dengan Menggunakan Algoritma Differential Evolution Di PT. XYZ](#)

Universitas Al Azhar Indonesia  [JURNAL AL-AZHAR INDONESIA SERI SAINS DAN TEKNOLOGI Vol 8, No 1 \(2023\): Januari 2023 24 - 32](#)
📅 2023  [DOI: 10.36722/sst.v8i1.1354](#)  [Accred : Sinta 3](#)

[Perilaku Gajah Sumatra \(*Elephas maximus sumatranus*\) Betina pada Kandang Dalam di Taman Margasatwa Ragunan, Jakarta](#)
Universitas Al Azhar Indonesia  [JURNAL AL-AZHAR INDONESIA SERI SAINS DAN TEKNOLOGI Vol 8, No 1 \(2023\): Januari 2023 58 - 66](#)
📅 2023  [DOI: 10.36722/sst.v8i1.1671](#)  [Accred : Sinta 3](#)

[Antibacterial Activity of *Drosera* Sp. Ethanolic Extract Against *Staphylococcus Aureus* ATCC 25923](#)
Universitas Al Azhar Indonesia  [JURNAL AL-AZHAR INDONESIA SERI SAINS DAN TEKNOLOGI Vol 8, No 1 \(2023\): Januari 2023 33 - 38](#)
📅 2023  [DOI: 10.36722/sst.v8i1.1494](#)  [Accred : Sinta 3](#)

[Nutritional Knowledge of Prospective Brides in Indonesia: A Systematic Review](#)
Universitas Al Azhar Indonesia  [JURNAL AL-AZHAR INDONESIA SERI SAINS DAN TEKNOLOGI Vol 8, No 1 \(2023\): Januari 2023 1 - 8](#)
📅 2023  [DOI: 10.36722/sst.v8i1.1251](#)  [Accred : Sinta 3](#)

[Penerapan Lean Tools Change Acceleration Process \(CAP\) Model dan Spaghetti Diagram Pada Proses Produksi Sofa di PT. XYZ](#)
Universitas Al Azhar Indonesia  [JURNAL AL-AZHAR INDONESIA SERI SAINS DAN TEKNOLOGI Vol 8, No 1 \(2023\): Januari 2023 39 - 48](#)
📅 2023  [DOI: 10.36722/sst.v8i1.1474](#)  [Accred : Sinta 3](#)

[Karakteristik Mutu Sensori Dan Kimia Liang Teh Pontianak Kaya Antioksidan Pada Berbagai Suhu Pengeringan](#)
Universitas Al Azhar Indonesia  [JURNAL AL-AZHAR INDONESIA SERI SAINS DAN TEKNOLOGI Vol 8, No 1 \(2023\): Januari 2023 49 - 57](#)
📅 2023  [DOI: 10.36722/sst.v8i1.1432](#)  [Accred : Sinta 3](#)

[View more ...](#)