

# HLS

International Journal of Health Services



## Healthcare in Low-resource Settings

2013 | Volume 2, Issue 1

www.hlsjournal.org

## Article

# Profil of omega-3 food intake and its association with socioeconomic status in smoker on online motorcycle drivers

Amelia Lorensia,<sup>1</sup> Rivan Virlando Suryadinata<sup>2</sup>

<sup>1</sup>Department of Clinical Pharmacy-Community, Faculty of Pharmacy, Universitas Surabaya, Jl. Raya Kalirungkut, Surabaya, Indonesia; <sup>2</sup>Department of Public Health, Faculty of Medicine, Universitas Surabaya, Jl. Raya Kalirungkut, Surabaya, Surabaya, Indonesia

## Abstract

**Introduction:** Omega-3 fatty acids were found to be effective in reducing inflammation and free radicals caused by air pollution (cigarette smoke), as well as improving lung function. Furthermore, economic conditions can influence a person's food consumption habits. The type of food consumed will certainly affect the amount of omega-3 absorbed by the body, and the quality of food consumed is influenced by socioeconomic status (SES). Therefore, the purpose of this study was to determine omega-3 intake and its relationship with the socioeconomic status of online motorcycle drivers.

**Design and Methods:** This was an observational study with a cross-sectional method of measuring omega-3 consumption using a 24-hour food recall questionnaire and assessing the mean of omega-3 intake after 3 days. The was conducted from May-August 2020, in Kali Rungkut, Surabaya City, and the subjects were online motorcycle drivers. A Chi-square test was used to assess the relationship between omega-3 intake and SES (income and education).

**Results:** There were 49 respondents in this study with most of them having a classification of omega-3 intake at the deficit level (57.14%), and none belonging to the good and more level. There was a significant difference in omega-3 intake on income ( $p=0.000$ ) and education ( $p=0.000$ ).

**Conclusions:** Income and education must be prioritized to improve a healthy diet that includes omega-3.

## Introduction

Despite the long distance, the community's high mobility necessitates quick movement from one location to another.<sup>1</sup> The existence of online motorcycle drivers is one of the newest transportation media that overcomes congestion. It is an online application-based transportation development supported by communication technology through smartphones, motorcycle services, and communication technology.<sup>2</sup> The high risk of online motorcycle

drivers being exposed to vehicle air pollution increases the prevalence of chronic lung diseases and this is supported by smoking habits. Previous studies show that smoking is significantly associated with the incidence of COPD (chronic obstructive pulmonary disease).<sup>3</sup> Therefore, smokers have greater respiratory problems and are less physically active than non-smokers.<sup>4</sup>

Smoking is a problem because it causes numerous losses in terms of social, moral, economic, health, death, and a decrease in human resources.<sup>5</sup> Tobacco content in cigarettes can affect almost all organs of the human body. Hence, smoking is linked to a variety of non-communicable diseases such as cancer, heart disease, chronic respiratory disease, and diabetes.

In Indonesia, smoking is responsible for a high proportion of morbidity and mortality. Furthermore, the percentage of Indonesian men who smoke has dramatically increased from 56.2% in 2000 to 76.2% in 2015.<sup>6</sup> Another effect is the loss of human resources as a result of smoking-related morbidity and mortality.<sup>7</sup> This contributes to poverty by diverting household spending away from basic needs such as food and shelter to cigarettes. Since the tobacco in cigarettes is very addictive, this behavior is difficult to overcome. Additionally, the economic costs of smoking are high because of the healthcare costs associated with smoking-related diseases.<sup>8,9</sup> Attempts to quit smoking often experience obstacles, such as low self-confidence and self-esteem among smokers, and depression is a significant factor in smoking cessation failure. There are several drug therapies for smoking cessation, such as nicotine replacement therapy which is the first line of smoking cessation treatment that should be used with caution in patients with arrhythmias and myocardial infarction. Nicotine can cause adverse cardiovascular effects by increasing myocardial workload through increased heart rate and blood pressure. It can also cause coronary artery constriction, resulting in cardiac ischemia.<sup>10,11,12</sup>

Nutrients, which are included in the COVID-19 prevention treatment, are particularly effective in maintaining health and improving lung function.<sup>13</sup> Omega-3 is effective in reducing inflammation and free radicals caused by air pollution and cigarette smoke and improves lung function.<sup>14,15</sup> Therefore, a person's food intake can affect the health of the lung function,<sup>16</sup> and

### Significance for public health

Due to continuous exposure to pollution, online motorcycle drivers, including 49 respondents, are at a high risk of chronic lung function decline. The health of lung function can be affected by the dietary intake of omega-3, while the quality of food can affect the amount of omega-3 in the body. Therefore, this study examines the relationship between socioeconomic status and omega-3 in 49 online motorcycle drivers' diets.

amount of omega-3 in the body. By causing oxidative stress, cigarette smoke can enhance PUFA (Polyunsaturated Fatty Acid) lipid peroxidation. Low PUFA concentrations will in turn affect neurotransmission in the central nervous system, such as dopaminergic increases nicotine addiction, thereby inhibiting efforts to quit smoking.<sup>17,18</sup>

According to lower levels of DHA (docosahexaenoic acid) and EPA (eicosapentaenoic acid), smokers consume less fish rich in omega-3 fatty acids than non-smokers. The importance of omega-3 intake is that it can normalize the dopaminergic system and reduce addiction, hence reducing the desire to smoke significantly. The anti-inflammatory effect of omega-3 is due to the content of EPA which is a competitive substrate with arachidonic. Therefore, it has the potential to reduce inflammation and bronchoconstriction in the respiratory tract. These anti-inflammatory effects can aid improve lung function and influence smoking patterns.<sup>18</sup> Cigarettes contain high concentrations of free radicals and oxidants which cause systemic oxidative stress and lipid peroxidation, hence affecting the levels of omega-3 PUFAs, specifically EPA and DHA. According to the theory, cigarette smoke induces oxidative stress by peroxidation of polyunsaturated fatty acids which lowers the levels of omega-3. Hence, it can affect nerve transmission and cause hypofunction in the mesocortical system associated with the dependence mechanism, and increased desire to smoke. Omega-3 deficiency, particularly EPA and DHA, has also been associated with poor psychological health and the inability to cope with stress.<sup>19,20</sup> The effectiveness data of omega-3 fish oil revealed a relationship between illness perception and chronic respiratory disease. Although the influence of illness perception was significant, the length of time that respiratory symptoms were experienced was not.<sup>21</sup>

Epidemiological studies reveal that omega 3 has a protective effect against cardiovascular disease, myocardial or cerebral infarction, hypertension, and hyperlipidemia. Additionally, it is beneficial against chronic inflammatory diseases including COPD, asthma, rheumatoid arthritis, and inflammation of the gut.<sup>14,22,23</sup> Higher omega-3 intake and fish consumption are associated with better lung function in smokers and ex-smokers, with EPA and DHA acting as antioxidants and reducing oxidative stress.<sup>17,18</sup> Epidemiological studies reveal that EPA and DHA are the main omega-3 PUFAs found in fish, fish oil, and fish oil supplements. Furthermore, omega-3 are easily obtained from foods such as salmon, sardines, seafood, and others.<sup>20,24,25</sup> According to the recommended nutritional adequacy rate for Indonesians, the nutritional adequacy rate for omega-3 is 1.6 g/day for men 18 years.<sup>26</sup>

Given the beneficial cardiovascular effects of omega-3s, it is important to understand the sociocultural factors that influence adequate intake to improve dietary quality in minorities.<sup>27</sup> According to the results of omega-3 intake, all respondents consumed less omega-3 containing foods (<1,600 mg), with the average total intake of foods containing omega-3 being 226.47 mg. The most consumed omega-3s were eggs (average: 19.46 mg/day) and chicken (average: 10.74 mg/day), both of which contain low omega-3. Meanwhile, based on data collection per day using the 24-hour food recall, the average daily consumption of respondents was only 249.05 mg. Therefore, further analysis of the processing pattern of the food consumed and the factors that influence their diet is necessary.<sup>26</sup>

Omega-3 cannot only be obtained by taking supplements, but also from the foods we consume on the daily basis.<sup>21</sup> Economic circumstances can have an impact on a person's food consumption habits and the type of food consumed will certainly affect the omega-3 content absorbed by the body.<sup>28</sup> Also, the quality of food consumed is influenced by socioeconomic status (SES). Several

studies linked an individual's socioeconomic status to health, with positive associations of SES (income, education, occupation), fruit, vegetable intake, and nutrition,<sup>28,29</sup> but negative associations with chronic disease.<sup>30</sup> For example, income, and educational attainment are recognized as fundamental factors influencing heart disease rates in minority and low-income populations.<sup>31</sup>

A 24-hour food recall can be used to determine the pattern of omega-3 intake.<sup>32,33</sup> Using this method, individuals are asked to describe the food and drinks consumed for three days. The 24-hour food recall method gathers information about the food consumed at a certain time. In addition, this can be performed at all survey locations such as the community, and hospitals or installations by housewives. The data obtained is in the form of protein intake, carbohydrates, sugar, water, salt, and calcium in calorie units.<sup>34,35</sup> Therefore, this study aims to determine omega-3 intake and its relationship to the socioeconomic status of online motorcycle drivers.

## Design and Methods

A retrospective cohort study was used as its design, and the measurement of omega-3 intake was performed using a 24-hour food recall. The subjects were online motorcycle drivers and this was performed in Kali Rungkut, Surabaya City from May to August 2020. Furthermore, ethical permission was received from the University of Surabaya Ethics Committee Number 016-OL/KE.VII/2020.

The independent variables were omega-3 food intake and socioeconomic status. This was all performed around the Rungkut area in Surabaya, East Java, Indonesia, which was conducted from May to August 2020. Online motorcycle drivers provided public transportation in the form of motorcycles that could be ordered over the internet or through mobile phone applications. They not only deliver and pick up customers but also foods and goods.

The 24-hour food recall method was used and interviews were conducted with the help of a food photo book to aid in the estimation of the household size. An analysis of the adequacy of values for adults was performed in the Rungkut area, Surabaya City on online motorcycle drivers. Subsequently, the value was adjusted to the Nutrition Adequacy Rate (RDA),<sup>34,35</sup> which was 75 g protein/day while omega-3 was 1600 mg/day.<sup>21</sup>

The participants were online motorcycle drivers in the Surabaya City area. Therefore, the target population that met the inclusion criteria were: i) Aged 18 years; ii) an active smoker smoking 100 cigarettes in his life until now, or respondents smoking 1 cigarette/day; iii) Worked on weekdays regularly (min  $\pm$ 7 hours/day); iv) Having no eating disorders and gastrointestinal diseases such as Gastroesophageal Reflux Disease (GERD), gastritis and dyspepsia. The purposive sampling technique was used, while the 24-hour recall method was an interview guide measuring food consumption in the preparation of interview questions. This method was performed three times but was not successive, namely twice on weekdays and once on holidays, because the scheme could illustrate the variability of calorie and nutrient intake. Respondents were interviewed about food and drinks consumed in the past 24-hours, including portion sizes with the help of a photograph of household sizes, such as a spoon, plate, glass, or another size commonly used daily as stated in Food Photo Book by the Ministry of Health of the Republic of Indonesia in 2014.<sup>36</sup> Then, the results were synchronized to the average intake per day. Primary data was obtained directly from study subjects through direct dialogue (interviews) and presented descriptively. The classification of omega-3 intake was divided into 5, namely: deficit

(<70% of the value of the minimum measure of intake); less (70-80% of the value of a minimum measure of intake); sufficient (80-100% of the value of a minimum measure of intake); good (100-110% of the minimum size measure of intake); and more (>110% of the minimum intake measure value).<sup>37</sup> Respondents were interviewed about food and drinks consumed in the past 24 hours, including portion sizes with the help of a photograph of household sizes, such as a spoon, plate, glass, or other sizes commonly used daily as stated in Food Photo Book. The results were then synchronized to an average intake per day. Table 1 shows the List of Food Intake Composition Containing Omega-3.<sup>38,39,40</sup> The data in this study was obtained from subjects through direct dialogue (interviews).

Subsequently, the obtained data were processed using one of the Nutrisurvey programs, which was software designed for non-commercial use. This was useful for analyzing food nutrients from a menu or a consumption survey. After collecting the calorie intake, the data was entered into SPSS version 24, and the chi-square test was used to assess the relationship between omega-3 intake and SES (income and education).

## Results and Discussions

### Characteristics of respondents

This study made use of 49 respondents working as online motorcycle drivers in the Rungkut area. During the collection of respondents, 5 drivers refused to be involved in the study because of a call to order from a passenger. The prevalence of smoking in Indonesia is very high, specifically in men ranging from children, adolescents, to adults. Furthermore, 12 of the respondents were 40-44 years (24.49%), and this is not in accordance with Tomioka et al.<sup>41</sup> that stated that people aged 45-64 years tend to smoke more than those aged <45 years or >65 years. Table 2 described the characteristics of the respondents. Most of the smoking severity was light (48.98%) with an income mostly around <1 million rupiahs (61.22%). In terms of education level, most of them were high school seniors (73.47%).

A total of 49 respondents' data from online motorcycle drivers were analyzed descriptively. Table 3 described the average (212.7 mg/day), minimum (0 mg/day), maximum (1,069.43 mg/day) and standard deviation (264.1 mg/day) of the consumption of omega-3. These fatty acids were one of the nutrients needed by the body. However, the body does not produce it naturally, and therefore needs to get it elsewhere. Eicosanoids, which are chemicals that provide signals to the cardiovascular, pulmonary, endocrine, and immunological systems, are formed by omega-3 fatty acids. According to the data obtained, consumption of omega-3 per day was not sufficient, both on workdays and holidays. This is because it did not meet the recommended nutritional adequacy rate (RDA) of 1600 mg/day. Furthermore, the minimum average consumption of omega-3 was 0 mg/day, the maximum average was 1069.43 mg/day and the standard deviation (SD) was 277.18 mg/day. These

results were influenced by the level of income obtained. Due to the corona pandemic and ignorance of foods containing omega-3, some respondents did not consume foods containing omega-3 at all, resulting in an average result was 0. Meanwhile, other respondents obtained omega-3 through eggs, chicken, catfish, and anchovies.

### Eggs

Respondents consumed chicken eggs, but not those containing omega-3. Omega-3 chicken eggs are produced from laying hens fed a diet containing omega-3 supplements for three consecutive

**Table 1. List of Food Intake Composition Containing Omega-3.**<sup>37, 38,39</sup>

| Food      | Gram | Omega-3 Content (mg/serving) |
|-----------|------|------------------------------|
| Anchovy   | 57   | 1,165                        |
| Catfish   | 85   | 151                          |
| Shells    | 85   | 241                          |
| Crab      | 85   | 351                          |
| Lobster   | 100  | 84                           |
| Salmon    | 85   | 1,825                        |
| Tuna      | 85   | 228                          |
| Sardines  | 57   | 556                          |
| Shrimp    | 85   | 267                          |
| Egg       | 63   | 22                           |
| Beef      | 100  | 22                           |
| Chicken   | 100  | 19                           |
| Goat meat | 100  | 18                           |

**Table 2. Characteristics of respondents.**<sup>9</sup>

| Characteristics of respondents | Frequency (n=49) | Percentage (%) |
|--------------------------------|------------------|----------------|
| Age (years)                    |                  |                |
| 18-25                          | 9                | 18.40          |
| 26-35                          | 17               | 34.70          |
| 36-45                          | 17               | 34.70          |
| 46-55                          | 4                | 8.20           |
| 56-65                          | 2                | 4.10           |
| Smoking severity               |                  |                |
| Light                          | 24               | 48.99          |
| Moderate                       | 23               | 46.93          |
| Severe                         | 2                | 4.08           |
| Income (IDR)                   |                  |                |
| <1 million                     | 30               | 61.22          |
| 1-3 million                    | 19               | 38.78          |
| Education Degree               |                  |                |
| Elementary school              | 3                | 6.10           |
| Junior high school             | 2                | 4.10           |
| Senior high school             | 36               | 73.50          |
| Bachelor degree                | 8                | 16.30          |

**Table 3. Omega-3 intake profile based on 24-hour food recall.**

|         | 1st Meeting (mg/day) | 2nd Meeting (mg/day) | 3rd Meeting (mg/day) | Average (mg/day) |
|---------|----------------------|----------------------|----------------------|------------------|
| Average | 275.0                | 152.7                |                      | 210.5 212.7      |
| Minimum | 0                    | 0                    |                      | 0 0              |
| Maximum | 2105.6               | 1062.1               |                      | 2107.26 1,069.1  |
| SD      | 581.0                | 279.2                |                      | 393.1 264.1      |



weeks. Farmers in Indonesia use waste oil that is waste from the fish processing industry. Consumption of enriched hen eggs containing a mixture of omega-3 PUFA (ALA+EPA+DHA), causes changes in the microvascular reactivity, blood pressure, and triglyceride level in healthy subjects that are associated with cardiovascular benefits. This suggested that daily consumption of omega-3 PUFA-enriched eggs in healthy individuals may potentially contribute to cardiovascular risk factor attenuation and disease prevention.<sup>42</sup>

### Chicken

The two types of chicken frequently consumed are local chicken (boras/kampung) and broiler chicken. Since chicken consumption has steadily increased in recent decades, meat enriched with long-chain polyunsaturated fatty acids (LC-PUFA) has become an important source of nutrients to humans. The meat of broilers fed diets with fish oil and either flaxseed or rapeseed for two weeks before the slaughter is believed to be "high in omega-3 fatty acids." A 100 g portion of such breast or thigh meat would provide on average 33% and 15.5%, respectively, of the recommended daily intake of EPA and DHA for humans. Therefore, enriched chicken meat was a superior source of LC-PUFA than lean fish meat.<sup>43</sup>

### Catfish

Catfish can serve as a potential source of essential fatty acids to human nutrition particularly in Nigeria with the growing popularity of catfish consumption.<sup>44</sup> *Pangasius micronemus* (Black pangasius sp.) and *Pangasius nasutus* (Fruit pangasius sp.) are two species of silver catfish. Both *Pangasius* sp. are good supplies of omega-3 and omega-6. Moreover, *P. micronemus* from Sg. Kancong was the best choice because it was high in EPA and DHA.<sup>45</sup> Most of the respondents' classification of omega-3 intake was at the deficit level (57.14%), and none belonged to the good and more level. Classification of omega-3 intake was divided into 5, namely, deficit (<70% of the value of the minimum measure of

omega-3 intake); less (70-80% of the value of a minimum measure of omega-3 intake); sufficient (80-100% of the value of a minimum measure of omega-3 intake); good (100-110% of the minimum size measure of omega-3 intake); and more (>110% of the minimum omega-3 intake measure value) (Table 4). Prolonged omega-3 deficiency can affect neurodevelopment and cause neurological and visual disturbances. In addition, this could lead to immune, memory, and mental disorders. By increasing neurogenesis and neurotransmitters in the brain, DHA in omega-3 can promote memory and cognitive development in children. It also plays a role in increasing the activity of the prefrontal part of the brain used for thinking and behaving.<sup>46</sup>

In this study, Socioeconomic Status (SES) consisted of Income (IDR) and education degree. Table 5 showed that there was a significant difference in omega-3 intake based on income ( $p=0.000$ ) and education ( $p=0.000$ ). Also, the amount of EPA and DHA consumed daily varied greatly depending on income. Much less is known about the potential role of other demographic factors on n-3 LCPUFA intakes, such as ethnicity, income, and education level. However, according to a recent analysis of NHANES 2011–2014, n-3 LCPUFA intake varies by ethnicity in the US, with non-Hispanic Asians consuming significantly more EPA and DHA than Hispanics, non-Hispanic Whites, and non-Hispanic Blacks.<sup>47</sup> There is a relationship between the price of omega-3 supplements with EPA and DHA content, as observed from relative percentage. Product A1 (with the lowest price), has a relatively lower percentage, compared to brand A2 (with the highest price), with a relatively higher grade.<sup>48</sup>

Participants with higher school education had a significantly higher dietary intake of EPA and combined EPA+DHA than those with lower school education.<sup>47</sup> Other studies revealed the association of SES with omega-3 intake. Although socioeconomic status may play a role in the relationship between these factors and maternal omega-3 DHA status.<sup>49</sup> After adjusting for energy intake, education was significantly correlated with EPA + DHA intakes, and acculturation was significantly correlated with Total n-3, ALA, and EPA + DHA intakes. Foods sources of EPA + DHA eaten by at least 50% of participants were chicken, shrimp, tuna, and eggs.<sup>29</sup>

The data collected using the 3 x 24-hour food recall method was the limitation of this study because it mainly relies on the respondent's memory. To overcome this bias, a description of the food that might be consumed was provided. Once the respondent could not still remember, a 3x24 hour food recall data processing is issued using the Nutri Survey program. This has a weakness of not covering all types of food consumed by the respondent, but can be analyzed with the program to enable manual calculation. Additionally, the process of serving food can also affect the levels of omega-3 in food which was not considered in this study.

**Table 4. Classification of Omega-3 Intake.**

| Classification of Omega-3 intake | Frequency (n=49) | Percentage (%) |
|----------------------------------|------------------|----------------|
| Deficit (<70%)                   | 28               | 57.15          |
| Less (70-80%)                    | 16               | 32.65          |
| Sufficient (80-100%)             | 5                | 10.20          |
| Good (100-110%)                  | -                | -              |
| More (>110%)                     | -                | -              |

**Table 5. Cross-tabulation of your Omega-3 food intake its association with socioeconomic status.**

| Socioeconomic Status | Classification of Omega-3 intake |      |            | Total | P value (Chi Square Test) |
|----------------------|----------------------------------|------|------------|-------|---------------------------|
|                      | Deficit                          | Less | Sufficient |       |                           |
| Income (IDR)         |                                  |      |            |       |                           |
| <1 million           | 21                               | 9    |            | 30    | 0.000                     |
| 1-3 million          | 7                                | 7    | 5          | 19    |                           |
| Education Degree     |                                  |      |            |       |                           |
| Elementary school    | 3                                | -    | -          | 3     | 0.000                     |
| Junior high school   | 2                                | -    | -          | 2     |                           |
| Senior high school   | 23                               | 12   | 1          | 36    |                           |
| Bachelor degree      |                                  | 4    | 4          | 8     |                           |

## Conclusions

The choice of omega-3 foods was influenced by Socioeconomic Position (SES), which includes Income (IDR) and education level. Therefore, the higher the income and education level, the better the intake of omega-3 by online motorcycle drivers.

Correspondence: Amelia Lorensia, Faculty of Pharmacy, Universitas Surabaya (UBAYA), Jl. Raya Kalirungkut 60293, Surabaya, Indonesia, Tel.: +62312981110, Fax:+62312981111, E-mail: amelia.lorensia@staff.ubaya.ac.id

Key words: Omega-3; recall 24 hours; income; education; motorcycle driver online.

Acknowledgment: Gratitude goes to the Faculty of Pharmacy Universitas Surabaya, Surabaya, Indonesia, for their kind support and motivation during this study.

Contributions: In this study, contributions were made equally. RVS contributed to data processing, while AL reviewed the final product.

Conflict of interest: The authors declare no conflicts of interest.

Funding: This study was funded by the Faculty of Pharmacy Universitas Surabaya and Ministry of Research, Technology and Higher Education of the Republic of Indonesia as well as supported by the Institute of Research and Community Service Universitas Surabaya.

Clinical Trials: This study has been approved by the Health Research Ethics Committee of Universitas Surabaya.

Availability of data and materials: All data generated or analyzed during this study are included in this published article.

Informed consent: Written informed consent was obtained from a legally authorized representative(s) for anonymized patient information to be published in this article.

Conference presentation: Part of this paper was presented at the 2<sup>nd</sup> International Nursing and Health Sciences Symposium that took place at the Faculty of Medicine, Universitas Brawijaya, Malang, Indonesia.

Received for publication: 4 December 2021.

Accepted for publication: 12 May 2022.

This work is licensed under a Creative Commons Attribution 4.0 License (by-nc 4.0).

©Copyright: the Author(s), 2023

Licensee PAGEPress, Italy

Healthcare in Low-resource Settings 2023; 11(s1):11164

doi:10.4081/hls.2023.11164

*Publisher's note: All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.*

## References

- Verlinghieri E, Schwanen T. Transport and mobility justice: Evolving discussions. *J Transp Geogr* 2020;87:102798.
- Irawan MZ, Rizki M, Joewono TB, et al. Exploring the intention of out-of-home activities participation during new normal conditions in Indonesian cities. *Transp Res Interdiscip Perspect* 2020;8:100237.
- Ramadhan MAH, Hartono B. Incidence of Chronic Obstructive Pulmonary Disease (COPD) in Online Ojek Drivers in Bogor City and Depok City in 2018 (A Case Study of Air Pollution). *Jurnal Nasional Kesehatan Lingkungan Global* 2020;1(1):76-85.
- Lorensia A, Muntu CM, Suryadinata RV, et al. Effect of lung function disorders and physical activity on smoking and non-smoking students. *J Prev Med Hyg* 2021;62:E89-96.
- West R. Tobacco smoking: Health impact, prevalence, correlates and interventions. *Psychol Health* 2017;32:1018-36.
- Holipah H, Sulistomo HW, Maharani A. Tobacco smoking and risk of all-cause mortality in Indonesia. *PLoS One* 2020;15:e0242558.
- Balatif R. Cigarettes and Its Effects on Health. *Scripta Score Scientific Medical Journal* 2020;2:44-52.
- Baumeister RF. Addiction, cigarette smoking, and voluntary control of action: Do cigarette smokers lose their free will?. *Addict Behav Rep* 2017;5:67-84.
- González-Roz A, Jackson J, Murphy C, et al. Behavioral economic tobacco demand in relation to cigarette consumption and nicotine dependence: a meta-analysis of cross-sectional relationships. *Addiction* 2019;114:1926-40.
- Joly B, Perriot J, d'Athis P, et al. Success rates in smoking cessation: Psychological preparation plays a critical role and interacts with other factors such as psychoactive substances. *PLoS One* 2017;12:e0184800.
- Nagano T, Katsurada M, Yasuda Y, et al. Current pharmacologic treatments for smoking cessation and new agents undergoing clinical trials. *Ther Adv Respir Dis* 2019;13:1753466619875925.
- Hersi M, Traversy G, Thombs BD, et al. Effectiveness of stop smoking interventions among adults: protocol for an overview of systematic reviews and an updated systematic review. *Syst Rev* 2019;8:28.
- Aman F, Masood S. How Nutrition can help to fight against COVID-19 Pandemic. *Pak J Med Sci* 2020;36:S121-S123.
- Lorensia A, Wahyudi M, Yudianto A, et al. Effect of illness perception on improving asthma symptoms with omega-3 fish oil therapy: Pre-post design. *J Appl Pharmaceut Sci* 2020;10:62-71.
- Lorensia A, Wahyudi M, Mayzika NA. Effectiveness of Fish Oil Containing Omega-3 In Improving Symptoms And Lung Function In Asthma Outpatient In Surabaya, Indonesia. *Int J Pharmaceut Quality Assurance* 2018;9:260-6.
- Lorensia A, Suryadinata RV, Sidabutar BCM. Effect Analysis of Protein Intake of Pedicab Driver in Surabaya. *J Trop Pharm Chem* 2021;5:188-93.
- Wiest EF, Walsh-Wilcox MT, Walker MK. Omega-3 Polyunsaturated Fatty Acids Protect Against Cigarette Smoke-Induced Oxidative Stress and Vascular Dysfunction. *Toxicol Sci* 2017;156:300-310.
- Sadeghi-Ardekani K, Haghghi M, Zarrin R. Effects of omega-3 fatty acid supplementation on cigarette craving and oxidative stress index in heavy-smoker males: A double-blind, randomized, placebo-controlled clinical trial. *J Psychopharmacol* 2018 Sep;32:995-1002.
- Scaglia N, Chatkin J, Chapman KR, et al. The relationship between omega-3 and smoking habit: a cross-sectional study. *Lipids Health Dis* 2016;15:61.
- Harris WS, Tintle NL, Imamura F, et al. Blood n-3 fatty acid

- levels and total and cause-specific mortality from 17 prospective studies. *Nat Commun* 2021;12:2329.
21. Gammone MA, Riccioni G, Parrinello G, et al. Omega-3 Polyunsaturated Fatty Acids: Benefits and Endpoints in Sport. *Nutrients* 2018;11:46.
  22. Calder PC. Omega-3 fatty acids and inflammatory processes: from molecules to man. *Biochem Soc Trans* 2017;45:1105-15.
  23. Buoite Stella A, Gortan Cappellari G, Barazzoni R, et al. Update on the Impact of Omega 3 Fatty Acids on Inflammation, Insulin Resistance and Sarcopenia: A Review. *Int J Mol Sci* 2018;19:218.
  24. Hu C, Yang M, Zhu X, et al. Effects of Omega-3 Fatty Acids on Markers of Inflammation in Patients With Chronic Kidney Disease: A Controversial Issue. *Ther Apher Dial* 2018;22:124-32.
  25. Weylandt KH, Serini S, Chen YQ, et al. Omega-3 Polyunsaturated Fatty Acids: The Way Forward in Times of Mixed Evidence. *Biomed Res Int* 2015;2015:143109.
  26. Zibaenezhad MJ, Ghavipisheh M, Attar A, et al. Comparison of the effect of omega-3 supplements and fresh fish on lipid profile: a randomized, open-labeled trial. *Nutr Diabetes* 2017;7:1.
  27. Lorensia A, Suryadinata RV. Assessment of Omega-3 Fatty Acid Food Intakes in Online motorcycle drivers. *Teikyo Med J* 2021;44:881-92.
  28. Muhammad A, D'Souza A, Meade B, et al. How income and food prices influence global dietary intakes by age and sex: evidence from 164 countries. *BMJ Glob Health* 2017;2:e000184.
  29. Lora KR, Lewis NM, Eskridge KM, et al. Correlation of omega-3 fatty acids intakes with acculturation and socioeconomic status in midwestern Latinas. *J Immigr Minor Health* 2011;13:111-8.
  30. Pechey R, Monsivais P. Socioeconomic inequalities in the healthiness of food choices: Exploring the contributions of food expenditures. *Prev Med* 2016;88:203-9.
  31. Viego V, Temporelli K. Socioeconomic Status and Self-Reported Chronic Diseases Among Argentina's Adult Population: Results Based on Multivariate Probability Models. *J Public Health Res* 2017;6:883.
  32. Singh GK, Daus GP, Allender M, et al. Social Determinants of Health in the United States: Addressing Major Health Inequality Trends for the Nation, 1935-2016. *Int J MCH AIDS* 2017;6:139-64.
  33. Roach LA, Russell KG, Lambert K, et al. Polyunsaturated fatty acid food frequency questionnaire validation in people with end stage renal disease on dialysis. *Nutr Diet* 2020;77:131-138.
  34. Ansari MR, Agustina R, Khusun H, et al. Development and evaluation of a semiquantitative food frequency questionnaire for estimating omega-3 and omega-6 fatty acid intakes in Indonesian children. *Asia Pac J Clin Nutr* 2016;25:S20-S29.
  35. Dao MC, Subar AF, Warthon-Medina M, et al. Dietary assessment toolkits: an overview. *Public Health Nutr* 2019;22:404-18.
  36. Badan Penelitian dan Pengembangan Kesehatan Kementerian Kesehatan Indonesia. *Buku Foto Makanan oleh Tim Survei Konsumsi Makanan Individu*. Jakarta: Badan Penelitian dan Pengembangan Kesehatan; 2014.
  37. Badan Pusat Statistik (BPS). *Konsumsi kalori dan protein penduduk Indonesia dan provinsi*. Jakarta: Badan Pusat Statistik; 2020.
  38. Bellows L, Clifford J, Niebaum K, et al. Omega-3 Fatty Acids. *CSU Extension. Fact Sheet* 2015;9:382.
  39. Tur J, Bibiloni M, Sureda A, et al. Dietary sources of omega 3 fatty acids: Public health risks and benefits. *Br J Nutr* 2012;107:S23-52.
  40. Schwalfenberg G. Omega-3 fatty acids: their beneficial role in cardiovascular health. *Can Fam Physician* 2006;52:734-40.
  41. Tomioka K, Shima M, Saeki K. Association between heaviness of cigarette smoking and serious psychological distress is stronger in women than in men: a nationally representative cross-sectional survey in Japan. *Harm Reduct J* 2021;18:27.
  42. Stupin A, Rasic L, Matic A, et al. Omega-3 polyunsaturated fatty acids-enriched hen eggs consumption enhances microvascular reactivity in young healthy individuals. *Appl Physiol Nutr Metab* 2018;43:988-95.
  43. Konieczka P, Czuderna M, Smulikowska S. The enrichment of chicken meat with omega-3 fatty acids by dietary fish oil or its mixture with rapeseed or flaxseed—Effect of feeding duration: Dietary fish oil, flaxseed, and rapeseed and n-3 enriched broiler meat. *Animal Feed Science and Technology* 2017;223:42-52.
  44. Mustapha RA, Bolajoko O, Akinola OO. Omega-3 and Omega-6 Fatty Acids Potential of Smoked and Boiled Catfish (*Clarias Gariepinus*). *Curr Res Nutr Food Sci* 2014;2: 94-7.
  45. Hashim RB, Jamil EF, Zulkiply FH, et al. Fatty acid compositions of silver catfish, *Pangasius sp.* farmed in several rivers of Pahang, Malaysia. *J Oleo Sci* 2015;64:205-9.
  46. Lauritzen L, Brambilla P, Mazzocchi A, et al. DHA Effects in Brain Development and Function. *Nutrients* 2016;8:6.
  47. Cave C, Hein N, Smith LM, et al. Omega-3 Long-Chain Polyunsaturated Fatty Acids Intake by Ethnicity, Income, and Education Level in the United States: NHANES 2003-2014. *Nutrients* 2020;12:2045.
  48. Lorensia A, Budiono R, Suryadinata RV, et al. Quantitative determination of EPA and DHA in fish oil capsules for cardiovascular disease therapy in Indonesia by GC-MS. *J Public Health Res* 2021;10:2159.
  49. Wilson NA, Mantzioris E, Middleton PF, et al. Influence of sociodemographic, lifestyle and genetic characteristics on maternal DHA and other polyunsaturated fatty acid status in pregnancy: A systematic review. *Prostaglandins Leukot Essent Fatty Acids* 2020;152:102037.



The power of the Web of Science™ on your mobile device, wherever inspiration strikes.

Dismiss

Learn More

### Already have a manuscript?

Use our Manuscript Matcher to find the best relevant journals!

Find a Match

### Filters

Clear All

Web of Science Coverage

Open Access

Category

Country / Region

Language

Frequency

Journal Citation Reports

## Refine Your Search Results

2281-7824

Search

Sort By: Title (A-Z)

### Search Results

Found 1 results (Page 1)

Share These Results

### Journals Relevant To "2281-7824"

#### HEALTHCARE IN LOW-RESOURCE SETTINGS

OPEN ACCESS

Publisher: PAGEPRESS PUBL , MEDITGROUP, VIA G BELLI, 4, PAVIA, Italy, 27100

ISSN / eISSN: 2281-7824

Web of Science Core Collection: Emerging Sources Citation Index

Share This Journal

View profile page

\* Requires free login.

1 - 1 of 1



Items per page: 10



**Editorial Disclaimer:** As an independent organization, Clarivate does not become involved in and is not responsible for the editorial management of any journal or the business practices of any publisher. Publishers are accountable for their journal performance and compliance with ethical publishing standards. The views and opinions expressed in any journal are those of the author(s) and do not necessarily reflect the views or opinions of Clarivate. Clarivate remains neutral in relation to territorial disputes, and allows journals, publishers, institutes and authors to specify their address and affiliation details including territory.

Criteria for selection of newly submitted titles and re-evaluation of existing titles in the Web of Science are determined by the Web of Science Editors in their sole discretion. If a publisher's editorial policy or business practices negatively impact the quality of a journal, or its role in the surrounding literature of the subject, the Web of Science Editors may decline to include the journal in any Clarivate product or service. The Web of Science Editors, in their sole discretion, may remove titles from coverage at any point if the titles fail to maintain our standard of quality, do not comply with ethical standards, or otherwise do not meet the criteria determined by the Web of Science Editors. If a journal is deselected or removed from coverage, the journal will cease to be indexed in the Web of Science from a date determined by the Web of Science Editors in their sole discretion – articles published after that date will not be indexed. The Web of Science Editors' decision on all matters relating to journal coverage will be final.

Clarivate.™ Accelerating innovation.

© 2023 Clarivate

[Copyright Notice](#)

[Terms of Use](#)

[Privacy Notice](#)

[Cookie Policy](#)

[Manage cookie preferences](#)

[Help Center](#)

Follow us:



# Vol. 11 No. s1 (2023): 2nd International Nursing and Health Sciences Symposium

Published: 30-12-2022



## ORIGINAL ARTICLES



### The effects of TeamSTEPPS implementation by nurses on situation monitoring in hospital

Ahsan Ahsan, Imeldha Monitasari, Evi Harwiati Ningrum, Ike Nesdia Rahmawati, Linda Wieke Noviyanti, Kuswantoro Rusca Putra

<https://doi.org/10.4081/hls.2023.11163>  0  0  0  0

 3  PDF: 0

 PDF



### Profil of omega-3 food intake and its association with socioeconomic status in smoker on online motorcycle drivers

Amelia Lorensia, Rivan Virlando Suryadinata

<https://doi.org/10.4081/hls.2023.11164>  0  0  0  0

 1  PDF: 0

 PDF



### $\beta$ -(1,3)-D-glucan from *Pleurotus ostreatus* correlates with lower plasma IL-6, IL-1 $\beta$ , HOMA-IR, and higher pancreatic beta cell count in High-Fat and High-Fructose Diet (HFFD) rats

Alma Maghfrotun Innayah, Elvira Nur Sa'idah Hariani, Husnul Khotimah, Inggita Kusumastuty, Ema Pristi Yunita, Dian Handayani

<https://doi.org/10.4081/hls.2023.11165>  0  0  0  0

 2  PDF: 0

 PDF

### The effect of community of practice in improving the role of clinical instructor in patient safety



### implementation by prelicensure nursing stude

Dyah Wiji Puspita Sari, Hanny Handiyani, Kuntarti Kuntarti, Yuni Su Arief

<https://doi.org/10.4081/hls.2023.11166> 0 0 0 0

2 PDF: 0

PDF



### Identification of new cases of tuberculosis during the COVID-19 pandemic using model strategic management

Eppy Setiyowati, Umi Hanik, Ni Njoman Juliasih, Priyo Susilo

<https://doi.org/10.4081/hls.2023.11167> 0 0 0 0

2 PDF: 0

PDF



### The effects of a 'COVID Nurse Assistant' application on patient satisfaction in COVID isolation rooms

Evi Harwiati Ningrum, Annisa Wuri Kartika, Ahmad Hasyim Wibisono, Ike Nesda Rahmawati, Linda Wieke Noviyanti, Ahsan Ahsan, Kuswanto Rusca Putra, Ungky Agus Setyawan, Lusita Titik Andayani, Ririn Widayanti, Arif Jati Purnanto, Gatot Subroto, Nurul Laili, Judith Anderson

<https://doi.org/10.4081/hls.2023.11168> 0 0 0 0

2 PDF: 0

PDF



### The three month's dietary brown rice intervention has not significantly decreased levels of CRP, TNF- $\alpha$ , and IL6 of type 2 diabetes mellitus patients

Fajar Ari Nugroho, Inggita Kusumastuty, Anggun Rindang Cempaka, Atifa Nafia Hasantie Latif, Dian Handayani

<https://doi.org/10.4081/hls.2023.11172> 0 0 0 0

2 PDF: 0

PDF



## Correlation of leg pain responses with ankle-brachial index and peripheral sensory responses in foot of type 2 diabetes mellitus

Heri Kristianto, Tina Handayani Nasution, Endah Panca Lidya Fatma, Efris Kartika Sari, Ahmad Hasyim Wibisono, Haryadi Kurniawan, Firdausy Ratna, Ica Cristiningtyas, Endang Listyowati

<https://doi.org/10.4081/hls.2023.11173> 0 0 0 0

3 PDF: 0

PDF



## The use of web-based digital media to enhance admission orientation for patients in the hospital

Ike Nesdia Rahmawati, Imeldha Monitasari, Debby Hamsa Putri, Delfira Arizda, Febry Pricila, Fitri Rosyidawati, Bagus Lanang Sejati

<https://doi.org/10.4081/hls.2023.11174> 0 0 0 0

3 PDF: 0

PDF



## The ethical principles and caring behavior of Indonesian nurses

Ilkafah Ilkafah, Anestasia Pangestu Mei Tyas, Rini Rachmawaty

<https://doi.org/10.4081/hls.2023.11176> 0 0 0 0

2 PDF: 0

PDF



## A cross-sectional study of the knowledge, skills, and 6 rights on medication administration by nurses at emergency department

Linda Wieke Noviyanti, Andri Junianto, Ahsan Ahsan

<https://doi.org/10.4081/hls.2023.11177> 0 0 0 0

2 PDF: 0

PDF





### The efficacy of Raspberry Pi-based automatic voice message education on knowledge level and prevention behavior of high-risk population

Mifetika Lukitasari, Wahyu Indah Windarti, Endah Panca Lydia Fatma, Tony Suharsono, Dwi Adi Nugroho

<https://doi.org/10.4081/hls.2023.11178> 📄 0 ✅ 0 🔄 0 ❓ 0

👁️ 2 📄 PDF: 0

PDF



### Patient's knowledge, gender, and physical activity level as the predictors of self-care in heart failure patients

Mifetika Lukitasari, Ulfia Fitriani Nafista

<https://doi.org/10.4081/hls.2023.11179> 📄 0 ✅ 0 🔄 0 ❓ 0

👁️ 2 📄 PDF: 0

PDF



### The correlation between family role and anemia prevention in pregnancy

Mira Triharini, Ni Ketut Alit Armini, Sekar Ayu Pitaloka

<https://doi.org/10.4081/hls.2023.11180> 📄 0 ✅ 0 🔄 0 ❓ 0

👁️ 2 📄 PDF: 0

PDF



### Relationship between postural stability and fall risk in young adult after lower limb muscle fatigue

Mohd Khairuddin Mohd Safee, Noor Azuan Abu Osman

<https://doi.org/10.4081/hls.2023.11182> 📄 0 ✅ 0 🔄 0 ❓ 0

👁️ 2 📄 PDF: 0

PDF



### Knowledge and peer support for increase Menstrual Hygiene Management (MHM) in adolescents

Ni Ketut Alit Armini, Arik Setyani, Aria Aulia Nastiti, Mira Triharini

 <https://doi.org/10.4081/hls.2023.11193>

 0  0  0  0

 1  PDF: 0

 PDF



### Family health task implementation and the health status of diabetes mellitus patients: A correlational study

Niko Dima Kristianingrum Kristianingrum, Delfira Arizda, Setyoadi Setyoadi, Yati Sri Hayati, Michael Leo Parchman

 <https://doi.org/10.4081/hls.2023.11195>

 0  0  0  0

 1  PDF: 0

 PDF



### Physical activity, exercise habits, and body mass index of adults

Ratna Candra Dewi, Bambang Wirjatmadi

 <https://doi.org/10.4081/hls.2023.11197>

 0  0  0  0

 1  PDF: 0

 PDF



### Correlation between family support with work productivity of people with severe mental disorders (PSMD) in Bantur Community Health Center, Malang, Indonesia

Renny Nova, Heni Dwi Windarwati, Erik Meidianto

 <https://doi.org/10.4081/hls.2023.11207>

 0  0  0  0

 1  PDF: 0

 PDF



### A societal adaptation model as a novel approach toward the recovery of people with schizophrenia

Retno Lestari, Ah Yusuf, Febri Endra Budi Setyawan, Ahsan Ahsan, Rachmat Hargono

 <https://doi.org/10.4081/hls.2023.11208>

 0  0  0  0

 1  PDF: 0

 PDF



### Mother's knowledge as a dominant factor for the success of exclusive breastfeeding in Indonesia

Rinik Eko Kapti, Yuni Sufyanti Arief, Nurona Azizah

 <https://doi.org/10.4081/hls.2023.11209>

 0  0  0  0

 1  PDF: 0

 PDF



### *Brassica oleracea* var. italica extract reducing free radicals and inflammation initiated by an exposure to cigarette smoke

Rivan Virlando Suryadinata, Dwi Martha Nur Aditya, Viera Permatasari Wiana, Diana Rahman

 <https://doi.org/10.4081/hls.2023.11210>

 0  0  0  0

 1  PDF: 0

 PDF



### The effects of exclusive breastfeeding booklets on mothers' knowledge in providing exclusive breastfeeding in Mataram City, Indonesia

Septa Katmawanti, Farah Paramita, Agung Kurniawan, Siti Khuzaimah A Sharoni, Rosuzeita Fauzi, Intan Gumilang Pratiwi, Dea Aflah Samah, Yassinda Thasia Audina, Oktavia Sri Wahyuni, Meyralda Dara Adisa, Alinda Rahmani

 <https://doi.org/10.4081/hls.2023.11211>

 0  0  0  0

 2  PDF: 0

 PDF



### The correlation between family support and health status in patients with diabetes mellitus

Setyoadi Setyoadi, Ah Yusuf, Niko Dima Kristianingrum, Yati Sri Hayati, Linda Wieke Noviyanti, Nurmalia Filda Syafiky

<https://doi.org/10.4081/hls.2023.11212>

0 0 0 0

1 PDF: 0

PDF



### Experience of nurses using the basic nursing science in performing nursing care for patients at RSUD Dr. Soedomo Trenggalek, Indonesia

Rizki Bahtiyar Ardyansah, Shila Wisnasari, Titin Andri Wihastuti, Dina Dewi Sartika Lestari Ismail

<https://doi.org/10.4081/hls.2023.11213>

0 0 0 0

1 PDF: 0

PDF



### The effectiveness of nursing psychoeducation toward family burden and quality life on caregiver of people with schizophrenia in the community

Suharsono Suharsono, Noor Faidah, Moh Hanafi

<https://doi.org/10.4081/hls.2023.11215>

0 0 0 0

1 PDF: 0

PDF



### The implementation of code blue by nurses as first responders in outpatient and inpatient rooms at Malang Indonesia Hospital

Tony Suharsono, Sunarmi Sunarmi, Nur Ida, Bella Nove Khirria, Nazla Asrin, Ikhda Ulya

<https://doi.org/10.4081/hls.2023.11217>

0 0 0 0

1 PDF: 0

PDF

## REVIEWS

### Strategies to improving patient safety in hospitals

Mohamad Taji, Kuswanto Rusca Putra, Dina Dewi Sartika Lestari Ismail





<https://doi.org/10.4081/hls.2023.11181> 0 0 0 0

2 PDF: 1

PDF



### A systematic review on telenursing as a solution in improving the treatment compliance of tuberculosis patients in the COVID-19 pandemic

Ni Nyoman Elfiyunai, Nursalam Nursalam, Tintin Sukartini, Ferry Efendi

<https://doi.org/10.4081/hls.2023.11194> 0 0 0 0

1 PDF: 0

PDF



### The effectiveness of self-care management in treating heart failure: A scoping review

Alfrina Hany, Ratih Arum Vatmasari

<https://doi.org/10.4081/hls.2023.11196> 0 0 0 0

1 PDF: 0 Supplementary Materials: 0

PDF

SUPPLEMENTARY MATERIALS



### The application of the public health model in pregnancy coaching for preventing high-risk pregnancy

Sri Wahyuni, Apriliani Yulianti Wuriningsih, Nursalam Nursalam, Yuni Sufyanti Arief

<https://doi.org/10.4081/hls.2023.11214> 0 0 0 0

1 PDF: 0 Supplementary Materials: 0

PDF

SUPPLEMENTARY MATERIALS

FOR AUTHORS

**SUBMIT YOUR PAPER**

Guide for Authors

Benefits for Authors

---

How to write a scientific paper

---

How to write a Review article

---

Article Processing Charge

---

#### FOR REVIEWERS

Benefits for Reviewers

---

How to review

---

Thanks to Reviewers

---

#### INDEXING

DOAJ

---

ESCI

---

#### MOST READ LAST MONTH

Efficacy and safety of Camosunate for the treatment of uncomplicated malaria in the University of Benin Teaching Hospital, Benin City, Nigeria

👁 80

---

Color coding: a tool to enhance the quality of health care in low resource settings

👁 60


---

Knowledge, attitude and practice of contraceptive use among female students of Dilla secondary and preparatory school, Dilla town, South Ethiopia, 2014

👁 57

---

#### KEYWORDS



## Healthcare in Low-resource Settings

*is an Open Access, peer-reviewed journal published by PAGEPress®, Pavia, Italy. All credits and honors to PKP for their OJS.*

eISSN: 2281-7824

About

About The Journal

Editorial Board

Info Submission

Submissions

Guidelines For Authors

Our App Is Free For Anyone!





## About the Journal

### Focus, Scope and Objectives

**Healthcare in Low-resource Settings** is an open access, peer-reviewed journal which aims to publish high quality, outcome-based research conducted in or applicable to low-resource settings. Potential topics of interest are also: comparison of different medical procedures in terms of their effects on the healthcare resources; education of health professionals in rural areas; strategies to formulate effective health policies in those areas, and guidelines targeted specifically to them; advances in healthcare resource management. **Healthcare in Low-resource Settings** publishes *research articles, reviews* (narrative, systematic and meta-analysis), *case reports, debate articles, short reports, letters to the Editor, and study protocols*. The Journal also publishes thematic issues focusing on a single topic within the scope of the journal. Contextually relevant announcements, book reviews and abstracts from scientific meetings may also be hosted. Every article published in the Journal will be peer-reviewed by experts in the field and decided on by members of the editorial board.

### Peer review process

Peer review is applied to all research articles and the majority of other article types published in PAGEPress journals. This usually entails at least two independent, expert peer reviewers.

### Policy

All submissions to PAGEPress journals are first checked for completeness (criteria for desk reject are available in the [Guide for Authors](#)) before being sent to an Editor, who decides whether they are suitable for peer review. If an Editor is on the author list or has a competing interest in a particular manuscript, another member of the Editorial Board will be assigned to oversee peer review. When making a decision, Editors will consider the peer-reviewed reports, but they will not be bound by the opinions or recommendations contained in them. A single peer reviewer's or the Editor's concern may result in the manuscript being rejected. Peer review reports are sent to authors along with the editorial decision on their manuscript.

PAGEPress journals use one of two types of peer review:

- **Single-blind:** the reviewers know the names of the authors, but the authors do not know who reviewed their manuscript.
- **Double-blind:** the reviewers do not know the authors' names, and the authors do not know who reviewed their manuscript.



## Selection

Peer reviewer selection is critical to the publication process. A variety of factors influence it, including expertise, reputation, specific recommendations, conflicts of interest, and previous performance. All of these qualities are highly desirable: speed, thoroughness, sound reasoning, and collegiality.

When two independent peer reviewers cannot be obtained, the Editor may act as a second reviewer or make a decision based on only one report. If acting as a second reviewer, the editor must have sufficient knowledge in the area and sign the review to ensure transparency in the peer review process.

Before accepting an invitation to review a manuscript, potential peer reviewers should notify the Editor of any potential conflicts of interest. Editors' and peer reviewers' communications contain confidential information that should not be shared with third parties.

PAGEPress journals are committed to making editorial decisions and publishing them as soon as possible, and we believe that an efficient editorial process benefits both our authors and the research community as a whole. As a result, we ask reviewers to respond within the agreed-upon time frame. If reviewers anticipate a delay, we ask that they notify us so that we can keep the authors updated and, if necessary, find alternative solutions.

## Diversity and equity

PAGEPress is committed to diversity, equity, and inclusion, and strives for demographic diversity among peer reviewers. When inviting peer reviewers, editors are bly encouraged to consider geographical regions, gender identities, racial/ethnic groups, and other groups.

## Misconduct

False or misleading information, such as identity theft and suggesting fake peer-reviewers, will result in the manuscript being rejected, further investigation in accordance with PAGEPress' misconduct policy ([https://www.pagepress.org/site/plagiarism\\_misconduct](https://www.pagepress.org/site/plagiarism_misconduct)), and notification to the authors' institutions/employers. PAGEPress journals follow the Committee on Publication Ethics (COPE) [guidance](#) about peer reviewer fraud/falsification.

## Guidance

The primary goal of peer review is to provide the Editor with the information needed to make a fair, evidence-based decision that adheres to the journal's editorial criteria. Review reports should also assist authors in revising their paper so that it can be accepted for publication. Reports that include a recommendation to reject the paper should explain the major flaws in the research; this will help the authors prepare their manuscript for possible re-submission (if allowed by the Editor) or submission to a different journal.

A quick guide to reviewing is available [here](#)

**Before submitting a report, reviewers should ask themselves the following questions:**

- **How would you react if you received this report?**
- **Do you find the tone offensive?**
- **Is it polite and professional?**
- **Do the authors or their competitors receive any unnecessary personal or antagonistic remarks?**

**Please keep in mind that any offensive language in your report may be removed by the Editor.**

## **Publication Frequency**

All papers are published as soon as they have been accepted, by adding them to the "current" volume's *Table of Contents*.

## **Article Processing Charge**

Open Access publishing does have its costs. However, as a promotional activity, all articles accepted for publication in **Healthcare in Low-resource Settings** will be published completely exempt from any charge.

## **Open Access Policy**

This journal provides immediate open access to its content on the principle that making research freely available to the public supports a greater global exchange of knowledge.

## **Archiving**

This journal utilizes the [PKP Preservation Network](#), the [Global LOCKSS Network](#) and [Portico](#) to create a distributed archiving system among participating libraries and permits those libraries to create permanent archives of the journal for purposes of preservation and restoration.

## **Pre- and post-prints**

PAGEPress allows and encourages authors to deposit both their pre- and post-prints in Open-Access institutional archives or repositories. The primary benefit of pre- and post-print self-archiving is reaching a larger audience which enhances the visibility and impact of your research.

### **FOR AUTHORS**

#### **SUBMIT YOUR PAPER**

---

[Guide for Authors](#)

---

[Benefits for Authors](#)

---

[How to write a scientific paper](#)

---

[How to write a Review article](#)

---

Article Processing Charge

---

**FOR REVIEWERS**

Benefits for Reviewers

---

How to review

---

Thanks to Reviewers

---

**INDEXING**

DOAJ

---

ESCI

---

**MOST READ LAST MONTH**

Efficacy and safety of Camosunate for the treatment of uncomplicated malaria in the University of Benin Teaching Hospital, Benin City, Nigeria

 80

---

Color coding: a tool to enhance the quality of health care in low resource settings

 60

---

Knowledge, attitude and practice of contraceptive use among female students of Dilla secondary and preparatory school, Dilla town, South Ethiopia, 2014

 57

---

**KEYWORDS**



## Healthcare in Low-resource Settings

*is an Open Access, peer-reviewed journal published by PAGEPress®, Pavia, Italy. All credits and honors to PKP for their OJS.*

eISSN: 2281-7824

About

About The Journal

Editorial Board

Info Submission

Submissions

Guidelines For Authors

Our App Is Free For Anyone!





[HOME](#) / Editorial Board

## Editorial Board

### Editor-in-Chief

***Luigi Barberini***

University of Cagliari and University Hospital of Cagliari, Italy | [Curriculum Vitae](#)

### Editorial Board

*Saurav Basu*

Indian Institute of Public Health (Public Health Foundation of India), New Delhi, India

*Mauro Carta*

Department of Medical Sciences and Public Health, University of Cagliari, Italy

*Vittorio Colizzi*

University of Rome Tor Vergata, Italy and Faculty of Science & Technology, Evangelic University of Cameroon

*Claudia Fattuoni*

Department of Chemical and Geological Sciences, University of Cagliari, Italy

*Simona Gurzu*

George Emil Palade University of Medicine, Pharmacy, Sciences and Technology, Targu-Mures, Romania (UMFST)

*Uzzam Ahmed Khawaja*

Division of Pulmonary Medicine and Critical Care, Medicare Cardiac and General Hospital (MCGH), Karachi, Pakistan

*Mohsen Khosravi*

Department of Psychiatry and Clinical Psychology, Zahedan University of Medical Sciences, Zahedan, Iran

*Roy Rillera Marzo*

Department Chair of Public Health, International Medical School, Management and Science University, Malaysia

*Tolbert Nyenswah*

s Hopkins University Bloomberg School of Public Health, United States

*Salvatore Sardo*

Dipartimento di Scienze Mediche e Sanità Pubblica, Università degli Studi di Cagliari, Italy

## FOR AUTHORS

### SUBMIT YOUR PAPER

---

[Guide for Authors](#)

---

[Benefits for Authors](#)

---

[How to write a scientific paper](#)

---

[How to write a Review article](#)

---

[Article Processing Charge](#)

---

## FOR REVIEWERS

[Benefits for Reviewers](#)

---

[How to review](#)

---

[Thanks to Reviewers](#)

---

## INDEXING

[DOAJ](#)

---

[ESCI](#)

---

## MOST READ LAST MONTH

Efficacy and safety of Camosunate for the treatment of uncomplicated malaria in the University of Benin Teaching Hospital, Benin City, Nigeria

 80

---


Color coding: a tool to enhance the quality of health care in low resource settings

60

Knowledge, attitude and practice of contraceptive use among female students of Dilla secondary and preparatory school, Dilla town, South Ethiopia, 2014

57

KEYWORDS



### Healthcare in Low-resource Settings

is an Open Access, peer-reviewed journal published by PAGEPress®, Pavia, Italy. All credits and honors to PKP for their OJS.

eISSN: 2281-7824

About

About The Journal

Editorial Board

Info Submission

Submissions

Guidelines For Authors

Our App Is Free For Anyone!





# Profil of omega-3 food intake and its association with socioeconomic status in smoker on online motorcycle drivers

*by Amelia Lorensia*

---

**Submission date:** 10-Feb-2023 10:11PM (UTC+0700)

**Submission ID:** 2010918981

**File name:** hls-11-s1-11164.pdf (2.58M)

**Word count:** 4609

**Character count:** 29528

## Article

## Profil of omega-3 food intake and its association with socioeconomic status in smoker on online motorcycle drivers

Amelia Lorensia,<sup>1</sup> Rivan Virlando Suryadinata<sup>2</sup>

<sup>1</sup>Department of Clinical Pharmacy-Community, Faculty of Pharmacy, Universitas Surabaya, Jl. Raya Kalirungkut, Surabaya, Indonesia; <sup>2</sup>Department of Public Health, Faculty of Medicine, Universitas Surabaya, Jl. Raya Kalirungkut, Surabaya, Surabaya, Indonesia

### Abstract

**Introduction:** Omega-3 fatty acids were found to be effective in reducing inflammation and free radicals caused by air pollution (cigarette smoke), as well as improving lung function. Furthermore, economic conditions can influence a person's food consumption habits. The type of food consumed will certainly affect the amount of omega-3 absorbed by the body, and the quality of food consumed is influenced by socioeconomic status (SES). Therefore, the purpose of this study was to determine omega-3 intake and its relationship with the socioeconomic status of online motorcycle drivers.

**Design and Methods:** This was an observational study with a cross-sectional method of measuring omega-3 consumption using a 24-hour food recall questionnaire and assessing the mean of omega-3 intake after 3 days. The was conducted from May-August 2020, in Kali Rungkut, Surabaya City, and the subjects were online motorcycle drivers. A Chi-square test was used to assess the relationship between omega-3 intake and SES (income and education).

**Results:** There were 49 respondents in this study with most of them having a classification of omega-3 intake at the deficit level (57.14%), and none belonging to the good and more level. There was a significant difference in omega-3 intake on income ( $p=0.000$ ) and education ( $p=0.000$ ).

**Conclusions:** Income and education must be prioritized to improve a healthy diet that includes omega-3.

### Introduction

Despite the long distance, the community's high mobility necessitates quick movement from one location to another.<sup>1</sup> The existence of online motorcycle drivers is one of the newest transportation media that overcomes congestion. It is an online application-based transportation development supported by communication technology through smartphones, motorcycle services, and communication technology.<sup>2</sup> The high risk of online motorcycle

drivers being exposed to vehicle air pollution increases the prevalence of chronic lung diseases and this is supported by smoking habits. Previous studies show that smoking is significantly associated with the incidence of COPD (chronic obstructive pulmonary disease).<sup>3</sup> Therefore, smokers have greater respiratory problems and are less physically active than non-smokers.<sup>4</sup>

Smoking is a problem because it causes numerous losses in terms of social, moral, economic, health, death, and a decrease in human resources.<sup>5</sup> Tobacco content in cigarettes can affect almost all organs of the human body. Hence, smoking is linked to a variety of non-communicable diseases such as cancer, heart disease, chronic respiratory disease, and diabetes.

In Indonesia, smoking is responsible for a high proportion of morbidity and mortality. Furthermore, the percentage of Indonesian men who smoke has dramatically increased from 56.2% in 2000 to 76.2% in 2015.<sup>6</sup> Another effect is the loss of human resources as a result of smoking-related morbidity and mortality.<sup>7</sup> This contributes to poverty by diverting household spending away from basic needs such as food and shelter to cigarettes. Since the tobacco in cigarettes is very addictive, this behavior is difficult to overcome. Additionally, the economic costs of smoking are high because of the healthcare costs associated with smoking-related diseases.<sup>8,9</sup> Attempts to quit smoking often experience obstacles, such as low self-confidence and self-esteem among smokers, and depression is a significant factor in smoking cessation failure. There are several drug therapies for smoking cessation, such as nicotine replacement therapy which is the first line of smoking cessation treatment that should be used with caution in patients with arrhythmias and myocardial infarction. Nicotine can cause adverse cardiovascular effects by increasing myocardial workload through increased heart rate and blood pressure. It can also cause coronary artery constriction, resulting in cardiac ischemia.<sup>10,11,12</sup>

Nutrients, which are included in the COVID-19 prevention treatment, are particularly effective in maintaining health and improving lung function.<sup>13</sup> Omega-3 is effective in reducing inflammation and free radicals caused by air pollution and cigarette smoke and improves lung function.<sup>14,15</sup> Therefore, a person's food intake can affect the health of the lung function,<sup>16</sup> and

### Significance for public health

Due to continuous exposure to pollution, online motorcycle drivers, including 49 respondents, are at a high risk of chronic lung function decline. The health of lung function can be affected by the dietary intake of omega-3, while the quality of food can affect the amount of omega-3 in the body. Therefore, this study examines the relationship between socioeconomic status and omega-3 in 49 online motorcycle drivers' diets.

amount of omega-3 in the body. By causing oxidative stress, cigarette smoke can enhance PUFA (Polyunsaturated Fatty Acid) lipid peroxidation. Low PUFA concentrations will in turn affect neurotransmission in the central nervous system, such as dopaminergic increases nicotine addiction, thereby inhibiting efforts to quit smoking.<sup>17,18</sup>

According to lower levels of DHA (docosahexaenoic acid) and EPA (eicosapentaenoic acid), smokers consume less fish rich in omega-3 fatty acids than non-smokers. The importance of omega-3 intake is that it can normalize the dopaminergic system and reduce addiction, hence reducing the desire to smoke significantly. The anti-inflammatory effect of omega-3 is due to the content of EPA which is a competitive substrate with arachidonic. Therefore, it has the potential to reduce inflammation and bronchoconstriction in the respiratory tract. These anti-inflammatory effects can aid improve lung function and influence smoking patterns.<sup>18</sup> Cigarettes contain high concentrations of free radicals and oxidants which cause systemic oxidative stress and lipid peroxidation, hence affecting the levels of omega-3 PUFAs, specifically EPA and DHA. According to the theory, cigarette smoke induces oxidative stress by peroxidation of polyunsaturated fatty acids which lowers the levels of omega-3. Hence, it can affect nerve transmission and cause hypofunction in the mesocortical system associated with the dependence mechanism, and increased desire to smoke. Omega-3 deficiency, particularly EPA and DHA, has also been associated with poor psychological health and the inability to cope with stress.<sup>19,20</sup> The effectiveness data of omega-3 fish oil revealed a relationship between illness perception and chronic respiratory disease. Although the influence of illness perception was significant, the length of time that respiratory symptoms were experienced was not.<sup>21</sup>

Epidemiological studies reveal that omega 3 has a protective effect against cardiovascular disease, myocardial or cerebral infarction, hypertension, and hyperlipidemia. Additionally, it is beneficial against chronic inflammatory diseases including COPD, asthma, rheumatoid arthritis, and inflammation of the gut.<sup>14,22,23</sup> Higher omega-3 intake and fish consumption are associated with better lung function in smokers and ex-smokers, with EPA and DHA acting as antioxidants and reducing oxidative stress.<sup>17,18</sup> Epidemiological studies reveal that EPA and DHA are the main omega-3 PUFAs found in fish, fish oil, and fish oil supplements. Furthermore, omega-3 are easily obtained from foods such as salmon, sardines, seafood, and others.<sup>20,24,25</sup> According to the recommended nutritional adequacy rate for Indonesians, the nutritional adequacy rate for omega-3 is 1.6 g/day for men 18 years.<sup>26</sup>

Given the beneficial cardiovascular effects of omega-3s, it is important to understand the sociocultural factors that influence adequate intake to improve dietary quality in minorities.<sup>27</sup> According to the results of omega-3 intake, all respondents consumed less omega-3 containing foods (<1,600 mg), with the average total intake of foods containing omega-3 being 226.47 mg. The most consumed omega-3s were eggs (average: 19.46 mg/day) and chicken (average: 10.74 mg/day), both of which contain low omega-3. Meanwhile, based on data collection per day using the 24-hour food recall, the average daily consumption of respondents was only 249.05 mg. Therefore, further analysis of the processing pattern of the food consumed and the factors that influence their diet is necessary.<sup>26</sup>

Omega-3 cannot only be obtained by taking supplements, but also from the foods we consume on the daily basis.<sup>21</sup> Economic circumstances can have an impact on a person's food consumption habits and the type of food consumed will certainly affect the omega-3 content absorbed by the body.<sup>28</sup> Also, the quality of food consumed is influenced by socioeconomic status (SES). Several

studies linked an individual's socioeconomic status to health, with positive associations of SES (income, education, occupation), fruit, vegetable intake, and nutrition,<sup>28,29</sup> but negative associations with chronic disease.<sup>30</sup> For example, income, and educational attainment are recognized as fundamental factors influencing heart disease rates in minority and low-income populations.<sup>31</sup>

A 24-hour food recall can be used to determine the pattern of omega-3 intake.<sup>32,33</sup> Using this method, individuals are asked to describe the food and drinks consumed for three days. The 24-hour food recall method gathers information about the food consumed at a certain time. In addition, this can be performed at all survey locations such as the community, and hospitals or installations by housewives. The data obtained is in the form of protein intake, carbohydrates, sugar, water, salt, and calcium in calorie units.<sup>34,35</sup> Therefore, this study aims to determine omega-3 intake and its relationship to the socioeconomic status of online motorcycle drivers.

## Design and Methods

A retrospective cohort study was used as its design, and the measurement of omega-3 intake was performed using a 24-hour food recall. The subjects were online motorcycle drivers and this was performed in Kali Rungkut, Surabaya City from May to August 2020. Furthermore, ethical permission was received from the University of Surabaya Ethics Committee Number 016-OL/KE.VII/2020.

The independent variables were omega-3 food intake and socioeconomic status. This was all performed around the Rungkut area in Surabaya, East Java, Indonesia, which was conducted from May to August 2020. Online motorcycle drivers provided public transportation in the form of motorcycles that could be ordered over the internet or through mobile phone applications. They not only deliver and pick up customers but also foods and goods.

The 24-hour food recall method was used and interviews were conducted with the help of a food photo book to aid in the estimation of the household size. An analysis of the adequacy of values for adults was performed in the Rungkut area, Surabaya City on online motorcycle drivers. Subsequently, the value was adjusted to the Nutrition Adequacy Rate (RDA),<sup>34,35</sup> which was 75 g protein/day while omega-3 was 1600 mg/day.<sup>21</sup>

The participants were online motorcycle drivers in the Surabaya City area. Therefore, the target population that met the inclusion criteria were: i) Aged 18 years; ii) an active smoker smoking 100 cigarettes in his life until now, or respondents smoking 1 cigarette/day; iii) Worked on weekdays regularly (min  $\pm$ 7 hours/day); iv) Having no eating disorders and gastrointestinal diseases such as Gastroesophageal Reflux Disease (GERD), gastritis and dyspepsia. The purposive sampling technique was used, while the 24-hour recall method was an interview guide measuring food consumption in the preparation of interview questions. This method was performed three times but was not successive, namely twice on workdays and once on holidays, because the scheme could illustrate the variability of calorie and nutrient intake. Respondents were interviewed about food and drinks consumed in the past 24-hours, including portion sizes with the help of a photograph of household sizes, such as a spoon, plate, glass, or another size commonly used daily as stated in Food Photo Book by the Ministry of Health of the Republic of Indonesia in 2014.<sup>36</sup> Then, the results were synchronized to the average intake per day. Primary data was obtained directly from study subjects through direct dialogue (interviews) and presented descriptively. The classification of omega-3 intake was divided into 5, namely: deficit



(<70% of the value of the minimum measure of intake); less (70-80% of the value of a minimum measure of intake); sufficient (80-100% of the value of a minimum measure of intake); good (100-110% of the minimum size measure of intake); and more (>110% of the minimum intake measure value).<sup>37</sup> Respondents were interviewed about food and drinks consumed in the past 24 hours, including portion sizes with the help of a photograph of household sizes, such as a spoon, plate, glass, or other sizes commonly used daily as stated in Food Photo Book. The results were then synchronized to an average intake per day. Table 1 shows the List of Food Intake Composition Containing Omega-3.<sup>38,39,40</sup> The data in this study was obtained from subjects through direct dialogue (interviews).

Subsequently, the obtained data were processed using one of the Nutrisurvey programs, which was software designed for non-commercial use. This was useful for analyzing food nutrients from a menu or a consumption survey. After collecting the calorie intake, the data was entered into SPSS version 24, and the chi-square test was used to assess the relationship between omega-3 intake and SES (income and education).

## Results and Discussions

### Characteristics of respondents

This study made use of 49 respondents working as online motorcycle drivers in the Rungkut area. During the collection of respondents, 5 drivers refused to be involved in the study because of a call to order from a passenger. The prevalence of smoking in Indonesia is very high, specifically in men ranging from children, adolescents, to adults. Furthermore, 12 of the respondents were 40-44 years (24.49%), and this is not in accordance with Tomioka et al.<sup>41</sup> that stated that people aged 45-64 years tend to smoke more than those aged <45 years or >65 years. Table 2 described the characteristics of the respondents. Most of the smoking severity was light (48.98%) with an income mostly around <1 million rupiahs (61.22%). In terms of education level, most of them were high school seniors (73.47%).

A total of 49 respondents' data from online motorcycle drivers were analyzed descriptively. Table 3 described the average (212.7 mg/day), minimum (0 mg/day), maximum (1,069.43 mg/day) and standard deviation (264.1 mg/day) of the consumption of omega-3. These fatty acids were one of the nutrients needed by the body. However, the body does not produce it naturally, and therefore needs to get it elsewhere. Eicosanoids, which are chemicals that provide signals to the cardiovascular, pulmonary, endocrine, and immunological systems, are formed by omega-3 fatty acids. According to the data obtained, consumption of omega-3 per day was not sufficient, both on workdays and holidays. This is because it did not meet the recommended nutritional adequacy rate (RDA) of 1600 mg/day. Furthermore, the minimum average consumption of omega-3 was 0 mg/day, the maximum average was 1069.43 mg/day and the standard deviation (SD) was 277.18 mg/day. These

results were influenced by the level of income obtained. Due to the corona pandemic and ignorance of foods containing omega-3, some respondents did not consume foods containing omega-3 at all, resulting in an average result was 0. Meanwhile, other respondents obtained omega-3 through eggs, chicken, catfish, and anchovies.

### Eggs

Respondents consumed chicken eggs, but not those containing omega-3. Omega-3 chicken eggs are produced from laying hens fed a diet containing omega-3 supplements for three consecutive

**Table 1. List of Food Intake Composition Containing Omega-3.**<sup>37, 38,39</sup>

| Food      | Gram | Omega-3 Content (mg/serving) |
|-----------|------|------------------------------|
| Anchovy   | 57   | 1,165                        |
| Catfish   | 85   | 151                          |
| Shells    | 85   | 241                          |
| Crab      | 85   | 351                          |
| Lobster   | 100  | 84                           |
| Salmon    | 85   | 1,825                        |
| Tuna      | 85   | 228                          |
| Sardines  | 57   | 556                          |
| Shrimp    | 85   | 267                          |
| Egg       | 63   | 22                           |
| Beef      | 100  | 22                           |
| Chicken   | 100  | 19                           |
| Goat meat | 100  | 18                           |

**Table 2. Characteristics of respondents.<sup>9</sup>**

| Characteristics of respondents | Frequency (n=49) | Percentage (%) |
|--------------------------------|------------------|----------------|
| Age (years)                    |                  |                |
| 18-25                          | 9                | 18.40          |
| 26-35                          | 17               | 34.70          |
| 36-45                          | 17               | 34.70          |
| 46-55                          | 4                | 8.20           |
| 56-65                          | 2                | 4.10           |
| Smoking severity               |                  |                |
| Light                          | 24               | 48.99          |
| Moderate                       | 23               | 46.93          |
| Severe                         | 2                | 4.08           |
| Income (IDR)                   |                  |                |
| <1 million                     | 30               | 61.22          |
| 1-3 million                    | 19               | 38.78          |
| Education Degree               |                  |                |
| Elementary school              | 3                | 6.10           |
| Junior high school             | 2                | 4.10           |
| Senior high school             | 36               | 73.50          |
| Bachelor degree                | 8                | 16.30          |

**Table 3. Omega-3 intake profile based on 24-hour food recall.**

|         | 1st Meeting (mg/day) | 2nd Meeting (mg/day) | 3rd Meeting (mg/day) | Average (mg/day) |         |
|---------|----------------------|----------------------|----------------------|------------------|---------|
| Average | 275.0                | 152.7                |                      | 210.5            | 212.7   |
| Minimum | 0                    | 0                    |                      | 0                | 0       |
| Maximum | 2105.6               | 1062.1               |                      | 2107.26          | 1,069.1 |
| SD      | 581.0                | 279.2                |                      | 393.1            | 264.1   |

weeks. Farmers in Indonesia use waste oil that is waste from the fish processing industry. Consumption of enriched hen eggs containing a mixture of omega-3 PUFA (ALA+EPA+DHA), causes changes in the microvascular reactivity, blood pressure, and triglyceride level in healthy subjects that are associated with cardiovascular benefits. This suggested that daily consumption of omega-3 PUFA-enriched eggs in healthy individuals may potentially contribute to cardiovascular risk factor attenuation and disease prevention.<sup>42</sup>

### Chicken

The two types of chicken frequently consumed are local chicken (boras/kampung) and broiler chicken. Since chicken consumption has steadily increased in recent decades, meat enriched with long-chain polyunsaturated fatty acids (LC-PUFA) has become an important source of nutrients to humans. The meat of broilers fed diets with fish oil and either flaxseed or rapeseed for two weeks before the slaughter is believed to be "high in omega-3 fatty acids." A 100 g portion of such breast or thigh meat would provide on average 33% and 15.5%, respectively, of the recommended daily intake of EPA and DHA for humans. Therefore, enriched chicken meat was a superior source of LC-PUFA than lean fish meat.<sup>43</sup>

### Catfish

Catfish can serve as a potential source of essential fatty acids to human nutrition particularly in Nigeria with the growing popularity of catfish consumption.<sup>44</sup> *Pangasius micronemus* (*Black pangasius* sp.) and *Pangasius nasutus* (*Fruit pangasius* sp.) are two species of silver catfish. Both *Pangasius* sp. are good supplies of omega-3 and omega-6. Moreover, *P. micronemus* from Sg. Kancong was the best choice because it was high in EPA and DHA.<sup>45</sup> Most of the respondents' classification of omega-3 intake was at the deficit level (57.14%), and none belonged to the good and more level. Classification of omega-3 intake was divided into 5, namely, deficit (<70% of the value of the minimum measure of

omega-3 intake); less (70-80% of the value of a minimum measure of omega-3 intake); sufficient (80-100% of the value of a minimum measure of omega-3 intake); good (100-110% of the minimum size measure of omega-3 intake); and more (>110% of the minimum omega-3 intake measure value) (Table 4). Prolonged omega-3 deficiency can affect neurodevelopment and cause neurological and visual disturbances. In addition, this could lead to immune, memory, and mental disorders. By increasing neurogenesis and neurotransmitters in the brain, DHA in omega-3 can promote memory and cognitive development in children. It also plays a role in increasing the activity of the prefrontal part of the brain used for thinking and behaving.<sup>46</sup>

In this study, Socioeconomic Status (SES) consisted of Income (IDR) and education degree. Table 5 showed that there was a significant difference in omega-3 intake based on income ( $p=0.000$ ) and education ( $p=0.000$ ). Also, the amount of EPA and DHA consumed daily varied greatly depending on income. Much less is known about the potential role of other demographic factors on n-3 LCPUFA intakes, such as ethnicity, income, and education level. However, according to a recent analysis of NHANES 2011–2014, n-3 LCPUFA intake varies by ethnicity in the US, with non-Hispanic Asians consuming significantly more EPA and DHA than Hispanics, non-Hispanic Whites, and non-Hispanic Blacks.<sup>47</sup> There is a relationship between the price of omega-3 supplements with EPA and DHA content, as observed from relative percentage. Product A1 (with the lowest price), has a relatively lower percentage, compared to brand A2 (with the highest price), with a relatively higher grade.<sup>48</sup>

Participants with higher school education had a significantly higher dietary intake of EPA and combined EPA+DHA than those with lower school education.<sup>47</sup> Other studies revealed the association of SES with omega-3 intake. Although socioeconomic status may play a role in the relationship between these factors and maternal omega-3 DHA status.<sup>49</sup> After adjusting for energy intake, education was significantly correlated with EPA + DHA intakes, and acculturation was significantly correlated with Total n-3, ALA, and EPA + DHA intakes. Foods sources of EPA + DHA eaten by at least 50% of participants were chicken, shrimp, tuna, and eggs.<sup>29</sup>

The data collected using the 3 x 24-hour food recall method was the limitation of this study because it mainly relies on the respondent's memory. To overcome this bias, a description of the food that might be consumed was provided. Once the respondent could not still remember, a 3x24 hour food recall data processing is issued using the Nutri Survey program. This has a weakness of not covering all types of food consumed by the respondent, but can be analyzed with the program to enable manual calculation. Additionally, the process of serving food can also affect the levels of omega-3 in food which was not considered in this study.

**Table 4. Classification of Omega-3 Intake.**

| Classification of Omega-3 intake | Frequency (n=49) | Percentage (%) |
|----------------------------------|------------------|----------------|
| Deficit (<70%)                   | 28               | 57.15          |
| Less (70-80%)                    | 16               | 32.65          |
| Sufficient (80-100%)             | 5                | 10.20          |
| Good (100-110%)                  | -                | -              |
| More (>110%)                     | -                | -              |

**Table 5. Cross-tabulation of your Omega-3 food intake its association with socioeconomic status.**

| Socioeconomic Status | Classification of Omega-3 intake |      |            | Total | P value (Chi Square Test) |
|----------------------|----------------------------------|------|------------|-------|---------------------------|
|                      | Deficit                          | Less | Sufficient |       |                           |
| Income (IDR)         |                                  |      |            |       |                           |
| <1 million           | 21                               | 9    |            | 30    | 0.000                     |
| 1-3 million          | 7                                | 7    | 5          | 19    |                           |
| Education Degree     |                                  |      |            |       |                           |
| Elementary school    | 3                                | -    | -          | 3     | 0.000                     |
| Junior high school   | 2                                | -    | -          | 2     |                           |
| Senior high school   | 23                               | 12   | 1          | 36    |                           |
| Bachelor degree      |                                  | 4    | 4          | 8     |                           |

## Conclusions

The choice of omega-3 foods was influenced by Socioeconomic Position (SES), which includes Income (IDR) and education level. Therefore, the higher the income and education level, the better the intake of omega-3 by online motorcycle drivers.

Correspondence: Amelia Lorensia, Faculty of Pharmacy, Universitas Surabaya (UBAYA), Jl. Raya Kalirungkut 60293, Surabaya, Indonesia, Tel.: +62312981110, Fax: +62312981111, E-mail: amelia.lorensia@staff.ubaya.ac.id

Key words: Omega-3; recall 24 hours; income; education; motorcycle driver online.

Acknowledgment: Gratitude goes to the Faculty of Pharmacy Universitas Surabaya, Surabaya, Indonesia, for their kind support and motivation during this study.

Contributions: In this study, contributions were made equally. RVS contributed to data processing, while AL reviewed the final product.

Conflict of interest: The authors declare no conflicts of interest.

Funding: This study was funded by the Faculty of Pharmacy Universitas Surabaya and Ministry of Research, Technology and Higher Education of the Republic of Indonesia as well as supported by the Institute of Research and Community Service Universitas Surabaya.

Clinical Trials: This study has been approved by the Health Research Ethics Committee of Universitas Surabaya.

Availability of data and materials: All data generated or analyzed during this study are included in this published article.

Informed consent: Written informed consent was obtained from a legally authorized representative(s) for anonymized patient information to be published in this article.

Conference presentation: Part of this paper was presented at the 2<sup>nd</sup> International Nursing and Health Sciences Symposium that took place at the Faculty of Medicine, Universitas Brawijaya, Malang, Indonesia.

Received for publication: 4 December 2021.

Accepted for publication: 12 May 2022.

This work is licensed under a Creative Commons Attribution 4.0 License (by-nc 4.0).

© Copyright: the Author(s), 2023

Licensee PAGEPress, Italy

Healthcare in Low-resource Settings 2023; 11(s1):11164

doi:10.4081/hls.2023.11164

**Publisher's note:** All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.

## References

- Verlinghieri E, Schwanen T. Transport and mobility justice: Evolving discussions. *J Transp Geogr* 2020;87:102798.

- Irawan MZ, Rizki M, Joewono TB, et al. Exploring the intention of out-of-home activities participation during new normal conditions in Indonesian cities. *Transp Res Interdiscip Perspect* 2020;8:100237.
- Ramadhan MAH, Hartono B. Incidence of Chronic Obstructive Pulmonary Disease (COPD) in Online Ojek Drivers in Bogor City and Depok City in 2018 (A Case Study of Air Pollution). *Jurnal Nasional Kesehatan Lingkungan Global* 2020;1(1):76-85.
- Lorensia A, Muntu CM, Suryadinata RV, et al. Effect of lung function disorders and physical activity on smoking and non-smoking students. *J Prev Med Hyg* 2021;62:E89-96.
- West R. Tobacco smoking: Health impact, prevalence, correlates and interventions. *Psychol Health* 2017;32:1018-36.
- Holipah H, Sulistomo HW, Maharani A. Tobacco smoking and risk of all-cause mortality in Indonesia. *PLoS One* 2020;15:e0242558.
- Balatif R. Cigarettes and Its Effects on Health. *Scripta Score Scientific Medical Journal* 2020;2:44-52.
- Baumeister RF. Addiction, cigarette smoking, and voluntary control of action: Do cigarette smokers lose their free will?. *Addict Behav Rep* 2017;5:67-84.
- González-Roz A, Jackson J, Murphy C, et al. Behavioral economic tobacco demand in relation to cigarette consumption and nicotine dependence: a meta-analysis of cross-sectional relationships. *Addiction* 2019;114:1926-40.
- Joly B, Perriot J, d'Athis P, et al. Success rates in smoking cessation: Psychological preparation plays a critical role and interacts with other factors such as psychoactive substances. *PLoS One* 2017;12:e0184800.
- Nagano T, Katsurada M, Yasuda Y, et al. Current pharmacologic treatments for smoking cessation and new agents undergoing clinical trials. *Ther Adv Respir Dis* 2019;13:1753466619875925.
- Hersi M, Traversy G, Thombs BD, et al. Effectiveness of stop smoking interventions among adults: protocol for an overview of systematic reviews and an updated systematic review. *Syst Rev* 2019;8:28.
- Aman F, Masood S. How Nutrition can help to fight against COVID-19 Pandemic. *Pak J Med Sci* 2020;36:S121-S123.
- Lorensia A, Wahyudi M, Yudianto A, et al. Effect of illness perception on improving asthma symptoms with omega-3 fish oil therapy: Pre-post design. *J Appl Pharmaceut Sci* 2020;10:62-71.
- Lorensia A, Wahyudi M, Mayzika NA. Effectiveness of Fish Oil Containing Omega-3 In Improving Symptoms And Lung Function In Asthma Outpatient In Surabaya, Indonesia. *Int J Pharmaceut Quality Assurance* 2018;9:260-6.
- Lorensia A, Suryadinata RV, Sidabutar BCM. Effect Analysis of Protein Intake of Pedicab Driver in Surabaya. *J Trop Pharm Chem* 2021;5:188-93.
- Wiest EF, Walsh-Wilcox MT, Walker MK. Omega-3 Polyunsaturated Fatty Acids Protect Against Cigarette Smoke-Induced Oxidative Stress and Vascular Dysfunction. *Toxicol Sci* 2017;156:300-310.
- Sadeghi-Ardekani K, Haghghi M, Zarrin R. Effects of omega-3 fatty acid supplementation on cigarette craving and oxidative stress index in heavy-smoker males: A double-blind, randomized, placebo-controlled clinical trial. *J Psychopharmacol* 2018 Sep;32:995-1002.
- Scaglia N, Chatkin J, Chapman KR, et al. The relationship between omega-3 and smoking habit: a cross-sectional study. *Lipids Health Dis* 2016;15:61.
- Harris WS, Tintle NL, Imamura F, et al. Blood n-3 fatty acid



- levels and total and cause-specific mortality from 17 prospective studies. *Nat Commun* 2021;12:2329.
21. Gammone MA, Riccioni G, Parrinello G, et al. Omega-3 Polyunsaturated Fatty Acids: Benefits and Endpoints in Sport. *Nutrients* 2018;11:46.
  22. Calder PC. Omega-3 fatty acids and inflammatory processes: from molecules to man. *Biochem Soc Trans* 2017;45:1105-15.
  23. Buoitte Stella A, Gortan Cappellari G, Barazzoni R, et al. Update on the Impact of Omega 3 Fatty Acids on Inflammation, Insulin Resistance and Sarcopenia: A Review. *Int J Mol Sci* 2018;19:218.
  24. Hu C, Yang M, Zhu X, et al. Effects of Omega-3 Fatty Acids on Markers of Inflammation in Patients With Chronic Kidney Disease: A Controversial Issue. *Ther Apher Dial* 2018;22:124-32.
  25. Weylandt KH, Serini S, Chen YQ, et al. Omega-3 Polyunsaturated Fatty Acids: The Way Forward in Times of Mixed Evidence. *Biomed Res Int* 2015;2015:143109.
  26. Zibaenezhad MJ, Ghavipisheh M, Attar A, et al. Comparison of the effect of omega-3 supplements and fresh fish on lipid profile: a randomized, open-labeled trial. *Nutr Diabetes* 2017;7:1.
  27. Lorensia A, Suryadinata RV. Assessment of Omega-3 Fatty Acid Food Intakes in Online motorcycle drivers. *Teikyo Med J* 2021;44:881-92.
  28. Muhammad A, D'Souza A, Meade B, et al. How income and food prices influence global dietary intakes by age and sex: evidence from 164 countries. *BMJ Glob Health* 2017;2:e000184.
  29. Lora KR, Lewis NM, Eskridge KM, et al. Correlation of omega-3 fatty acids intakes with acculturation and socioeconomic status in midwestern Latinas. *J Immigr Minor Health* 2011;13:111-8.
  30. Pechey R, Monsivais P. Socioeconomic inequalities in the healthiness of food choices: Exploring the contributions of food expenditures. *Prev Med* 2016;88:203-9.
  31. Viego V, Temporelli K. Socioeconomic Status and Self-Reported Chronic Diseases Among Argentina's Adult Population: Results Based on Multivariate Probability Models. *J Public Health Res* 2017;6:883.
  32. Singh GK, Daus GP, Allender M, et al. Social Determinants of Health in the United States: Addressing Major Health Inequality Trends for the Nation, 1935-2016. *Int J MCH AIDS* 2017;6:139-64.
  33. Roach LA, Russell KG, Lambert K, et al. Polyunsaturated fatty acid food frequency questionnaire validation in people with end stage renal disease on dialysis. *Nutr Diet* 2020;77:131-138.
  34. Ansari MR, Agustina R, Khusun H, et al. Development and evaluation of a semiquantitative food frequency questionnaire for estimating omega-3 and omega-6 fatty acid intakes in Indonesian children. *Asia Pac J Clin Nutr* 2016;25:S20-S29.
  35. Dao MC, Subar AF, Warthon-Medina M, et al. Dietary assessment toolkits: an overview. *Public Health Nutr* 2019;22:404-18.
  36. Badan Penelitian dan Pengembangan Kesehatan Kementerian Kesehatan Indonesia. *Buku Foto Makanan oleh Tim Survei Konsumsi Makanan Individu*. Jakarta: Badan Penelitian dan Pengembangan Kesehatan; 2014.
  37. Badan Pusat Statistik (BPS). *Konsumsi kalori dan protein penduduk Indonesia dan provinsi*. Jakarta: Badan Pusat Statistik; 2020.
  38. Bellows L, Clifford J, Niebaum K, et al. Omega-3 Fatty Acids. *CSU Extension. Fact Sheet* 2015;9:382.
  39. Tur J, Bibiloni M, Sureda A, et al. Dietary sources of omega 3 fatty acids: Public health risks and benefits. *Br J Nutr* 2012;107:S23-52.
  40. Schwalfenberg G. Omega-3 fatty acids: their beneficial role in cardiovascular health. *Can Fam Physician* 2006;52:734-40.
  41. Tomioka K, Shima M, Saeki K. Association between heaviness of cigarette smoking and serious psychological distress is stronger in women than in men: a nationally representative cross-sectional survey in Japan. *Harm Reduct J* 2021;18:27.
  42. Stupin A, Rasic L, Matic A, et al. Omega-3 polyunsaturated fatty acids-enriched hen eggs consumption enhances microvascular reactivity in young healthy individuals. *Appl Physiol Nutr Metab* 2018;43:988-95.
  43. Konieczka P, Czaudema M, Smulikowska S. The enrichment of chicken meat with omega-3 fatty acids by dietary fish oil or its mixture with rapeseed or flaxseed—Effect of feeding duration: Dietary fish oil, flaxseed, and rapeseed and n-3 enriched broiler meat. *Animal Feed Science and Technology* 2017;223:42-52.
  44. Mustapha RA, Bolajoko O, Akinola OO. Omega-3 and Omega-6 Fatty Acids Potential of Smoked and Boiled Catfish (*Clarias Gariepinus*). *Curr Res Nutr Food Sci* 2014;2: 94-7.
  45. Hashim RB, Jamil EF, Zulkipli FH, et al. Fatty acid compositions of silver catfish, *Pangasius sp.* farmed in several rivers of Pahang, Malaysia. *J Oleo Sci* 2015;64:205-9.
  46. Lauritzen L, Brambilla P, Mazzocchi A, et al. DHA Effects in Brain Development and Function. *Nutrients* 2016;8:6.
  47. Cave C, Hein N, Smith LM, et al. Omega-3 Long-Chain Polyunsaturated Fatty Acids Intake by Ethnicity, Income, and Education Level in the United States: NHANES 2003-2014. *Nutrients* 2020;12:2045.
  48. Lorensia A, Budiono R, Suryadinata RV, et al. Quantitative determination of EPA and DHA in fish oil capsules for cardiovascular disease therapy in Indonesia by GC-MS. *J Public Health Res* 2021;10:2159.
  49. Wilson NA, Mantzioris E, Middleton PF, et al. Influence of sociodemographic, lifestyle and genetic characteristics on maternal DHA and other polyunsaturated fatty acid status in pregnancy: A systematic review. *Prostaglandins Leukot Essent Fatty Acids* 2020;152:102037.



# Profil of omega-3 food intake and its association with socioeconomic status in smoker on online motorcycle drivers

## ORIGINALITY REPORT

**21** %  
SIMILARITY INDEX

**17** %  
INTERNET SOURCES

**12** %  
PUBLICATIONS

**5** %  
STUDENT PAPERS

## PRIMARY SOURCES

**1** Submitted to Surabaya University  
Student Paper 4%

**2** [www.researchgate.net](http://www.researchgate.net)  
Internet Source 4%

**3** [jtpc.farmasi.unmul.ac.id](http://jtpc.farmasi.unmul.ac.id)  
Internet Source 2%

**4** [pagepress.org](http://pagepress.org)  
Internet Source 1%

**5** Ridzwan B. Hashim, Elyna Fatinie Jamil, Farah Hanis Zulkipli, Jamaluddin Mohd Daud. "Fatty Acid Compositions of Silver Catfish, *Pangasius* sp. Farmed in Several Rivers of Pahang, Malaysia", *Journal of Oleo Science*, 2015  
Publication 1%

**6** [worldwidescience.org](http://worldwidescience.org)  
Internet Source 1%

**7** [jphres.org](http://jphres.org)  
Internet Source 1%

|    |  |      |
|----|--|------|
| 8  | "Integrative and Functional Medical Nutrition Therapy", Springer Science and Business Media LLC, 2020<br>Publication   | 1 %  |
| 9  | Caleb Cave, Nicholas Hein, Lynette M. Smith, Ann Anderson-Berry et al. "Omega-3 Long-Chain Polyunsaturated Fatty Acids Intake by Ethnicity, Income, and Education Level in the United States: NHANES 2003–2014", Nutrients, 2020<br>Publication                      | 1 %  |
| 10 | <a href="http://www.w-i-c.org">www.w-i-c.org</a><br>Internet Source  | <1 % |
| 11 | Muhamad Rizki, Jeanly Syahputri, Prawira Fajarindra Belgiawan, Muhammad Zudhy Irawan. "Electrifying Tourist Mobility in Bali, Indonesia: Setting the Target and Estimating the CO2 Reduction Based on Stated Choice Experiment", Sustainability, 2021<br>Publication | <1 % |
| 12 | <a href="http://www.pagepress.org">www.pagepress.org</a><br>Internet Source  | <1 % |
| 13 | <a href="http://lipidworld.biomedcentral.com">lipidworld.biomedcentral.com</a><br>Internet Source  | <1 % |
| 14 | <a href="http://bmcmmedresmethodol.biomedcentral.com">bmcmmedresmethodol.biomedcentral.com</a><br>Internet Source  | <1 % |

|    |   |      |
|----|---|------|
| 15 | <a href="http://coek.info">coek.info</a><br>Internet Source   | <1 % |
| 16 | <a href="http://www.scielo.org.mx">www.scielo.org.mx</a><br>Internet Source   | <1 % |
| 17 | <a href="http://www.aio.co.id">www.aio.co.id</a><br>Internet Source   | <1 % |
| 18 | Kiana Sadeghi-Ardekani, Mahmonir Haghghi, Rasoul Zarrin. "Effects of omega-3 fatty acid supplementation on cigarette craving and oxidative stress index in heavy-smoker males: A double-blind, randomized, placebo-controlled clinical trial", <i>Journal of Psychopharmacology</i> , 2018<br>Publication | <1 % |
| 19 | Patricia Budihartanti Liman, Rina Agustina, Ratna Djuwita, Jahja Umar et al. "Dietary and Plasma Carboxymethyl Lysine and Tumor Necrosis Factor- $\alpha$ as Mediators of Body Mass Index and Waist Circumference among Women in Indonesia", <i>Nutrients</i> , 2019<br>Publication                       | <1 % |
| 20 | <a href="http://webhosting.ubaya.ac.id">webhosting.ubaya.ac.id</a><br>Internet Source   | <1 % |
| 21 | <a href="http://www.medrxiv.org">www.medrxiv.org</a><br>Internet Source   | <1 % |

22

N.A. Wilson, E. Mantzioris, P.F. Middleton, B.S. Muhlhausler. "Influence of sociodemographic, lifestyle and genetic characteristics on maternal DHA and other polyunsaturated fatty acid status in pregnancy: A systematic review", Prostaglandins, Leukotrienes and Essential Fatty Acids, 2020

Publication

<1 %

23

[japsonline.com](http://japsonline.com)

Internet Source

<1 %

24

[www.science.gov](http://www.science.gov)

Internet Source

<1 %

25

P. Konieczka, M. Czauderna, S. Smulikowska. "The enrichment of chicken meat with omega-3 fatty acids by dietary fish oil or its mixture with rapeseed or flaxseed—Effect of feeding duration", Animal Feed Science and Technology, 2017

Publication

<1 %

Exclude quotes  On

Exclude matches  Off

Exclude bibliography  On

# Profil of omega-3 food intake and its association with socioeconomic status in smoker on online motorcycle drivers

---

GRADEMARK REPORT

---

FINAL GRADE

**/0**

GENERAL COMMENTS

**Instructor**

---

PAGE 1

---

PAGE 2

---

PAGE 3

---

PAGE 4

---

PAGE 5

---

PAGE 6

---