

# Brassica oleracea var. italica extract reducing free radicals and inflammation initiated by an exposure to cigarette smoke

*by Rivan Virlando Suryadinata*

---

**Submission date:** 16-Feb-2023 06:11AM (UTC+0700)

**Submission ID:** 2015171926

**File name:** hls-11-s1-11210.pdf (2.34M)

**Word count:** 3197

**Character count:** 20792

## Article

## ***Brassica oleracea* var. *italica* extract reducing free radicals and inflammation initiated by an exposure to cigarette smoke**

Rivan Virlando Suryadinata, Dwi Martha Nur Aditya, Viera Permatasari Wiana, Diana Rahman

Faculty of Medicine, Universitas Surabaya (UBAYA), Surabaya, Indonesia

### Abstract

**Introduction:** Herbal extracts are often administered to cigarette smokers to prevent excessive free radicals. These include *Brassica oleracea* var. *italica*, known to contain high antioxidant flavonoids and selenium micronutrients. Therefore, this study aims to determine the efficacy of *Brassica oleracea* var extract. *italica* in reducing the free radicals and inflammation present in experimental animals exposed to cigarette smoke.

**Design and Methods:** This research was conducted based on an experimental method using a randomized controlled trial (RCT) for 21 days. The animals used were divided into six groups (negative control, positive control, and four treatments). Particularly, the positive control and treatment groups were exposed to cigarette smoke for 2 minutes, twice a day, at 50 PPM CO levels. The treatment groups were administered the extract at different doses (0.5 ml; 0.75 ml; 1 ml; 1.25 ml), before assessing the blood level of malondialdehyde and C-Reactive Protein.

**Result:** The results showed the tendency for exposure to smoke to increase the number of free radicals and stimulate inflammation responses in the body ( $P < 0,05$ ). In addition, a strong correlation between variables was established ( $p = 0,000$ ;  $r = 0,713$ ).

**Conclusions:** Broccoli extracts (*Brassica oleracea* L. var. *italica*) administration has the potential to cause a decline in the two aspects, including free radicals and inflammation responses resulting from exposure to cigarette smoke.

### Introduction

Tobacco cigarette is one of the products consumed through smoking. In addition, there has been an increase in popularity despite the negative health effects. Numerous countries have explored various means to stop the use,<sup>1</sup> but these efforts have not yielded significant results. Based on previous reports, one of the serious potential problems of every smoker is the elevated risk of disease infections, which is implicated in around 6 million deaths, worldwide. Despite the practice of abstinence for several years, smokers are known to have a sustained higher risk of infection with earlier onset than non-smokers. These are strongly associated with 10 years of shorter life expectancy.<sup>2</sup> The total number of

smokers worldwide is estimated to have reached 1 billion people and consist predominantly of men.<sup>3</sup> In addition, the prevalence in developed countries, including America is relatively high, at about 15% of the total population, and consists more of people in the productive age groups.<sup>4</sup> The number in developing countries, including Armenia, Laos and Indonesia, is comparably more, reaching over 50% of the total population.<sup>5</sup>

The increased mortality rate of smokers is evidence of the dangerous effects. Previous studies showed the development of various diseases, including lung cancer, respiratory tract, heart problems, and problems during pregnancy. Furthermore, about 40% of the total smoking population are known to die prematurely.<sup>6</sup> The increase in complications from diseases have been attributed to the free radicals present in cigarette smoke and are known to directly stimulate the body's inflammatory response.<sup>7</sup> Further exposure potentially instigates a decline in antioxidant levels and propagates immunosuppression.<sup>8</sup> Previous studies also highlighted the tendency for cigarettes to initiate endothelial dysfunction as a result of excessive free radicals and inflammation.<sup>9</sup>

Physiologically, free radicals play an important role in cellular functions related to the enzymatic defence system assumed to prevent further accumulation. The presence of excessive radicals causes cellular changes, and consequently oxidative stress.<sup>10,11</sup> The production of reactive oxygen species, including superoxide ( $O_2 \cdot^-$ ), peroxynitrite ( $ONOO \cdot^-$ ), and hydroxyl ( $OH$ ) is possible during metabolism.<sup>12</sup> The enzymatic antioxidants, in form of superoxide dismutase (SOD), glutathione peroxidase (GSH-Px), and Catalase (CAT) function to neutralize these free radicals.<sup>13</sup> Other variants are possibly derived from food intake, including non-enzymatic antioxidants.<sup>14</sup> In addition, inflammation is another expression of the body's natural defence mechanism, which occurs in response to the entry of various foreign objects, including bacteria, viruses, allergens, toxic chemicals, and cigarettes. Pathogens or other hazardous substances continuously perpetuates chronic oxidative stress and consequently trigger protein oxidation, which releases pro-inflammatory cytokines and stimulates inflammatory responses all over the body.<sup>15</sup>

The intake of non-enzyme antioxidants is necessary to reduce excessive free radicals. These include polyphenols, a natural compound obtained from numerous fruits and vegetables, including broccoli. This constituent serves as a defence system from ultravi-

### Significance for public health

*The increasing number of active smokers is expected to cause serious health problems. This habit has been implicated in the increase in free radicals entering the respiratory tract, which stimulate an inflammatory process in the body. This process triggers cell damage and death, eventually causing health problems. Moreover, Brassica oleracea var. italica is known to contain high antioxidants, hence, there is a postulation of potential capacity to neutralize free radicals and reduce body inflammations.*

oiet radiation as well as various pathogens and also has anti-inflammatory properties with the potential to confer cellular protection against oxidative stress.<sup>16</sup> The broccoli plant (*Brassica oleracea* var. *italica*) is known to contain one of the polyphenol types termed flavonoids and considered to have higher amounts compared to other vegetables.<sup>17,18</sup> Therefore, this research aims to determine the effectiveness of broccoli in reducing free radicals and inflammation resulting from exposure to cigarette smoke.

## Design and Methods

This is an experimental research with a post-test control group design. This involved the use of male Wistar rats (*Rattus norvegicus*) as samples during a 21 days experiment (No: 139/KE/X/2021). The animals were subdivided into 6 groups (one negative control group, one positive control group, and four treatment groups). In addition, basic maintenance and care were provided with reference to the 3R principle (Replacement, Reduction, and Refinement).

Kretek cigarettes, predominantly consumed by Indonesians, was used in this study, and is characterized by 2.4 mg nicotine and 38 mg tar. The smoke exposure was carried out two times daily for 21 days (each for 2 minutes with 50 PPM Carbon Monoxide levels measured using a Carbon Monoxide Meter device).

The broccoli was extracted through a maceration process using 96% ethanol as the solvent. Furthermore, the preparations were obtained by evaporating the extracts with a vacuum rotary evaporator. Based on Laurence and Bacharach's framework, a flavonoid present in Broccoli extract (18 mg/ml) was measured and evaluated by comparing it with the adequacy daily rate of (8 mg/day). Hence, a dose of 0.5 ml; 0.75 ml; 1 ml; 1.25 ml were derived and used for the experiment.

The negative control (Group I) consisted of experimental animals administered a daily intake, while the positive control (II) were only exposed to cigarette smoke. In addition, each of the four treatment groups (Group III, IV, V, VI) comprised of experimental animals exposed to cigarette smoke and the different doses of Broccoli extracts (0.5 ml; 0.75 ml; 1 ml; 1.25 ml). Therefore, the ratio data of malondialdehyde and C-Reactive Protein for each

group were obtained and analysed to ascertain the possible differences and relationships.

## Results and Discussions

The results showed the mean levels of malondialdehyde in each group. Table 1 highlights the highest values ( $11.23 \pm 1.45$ ) in Group II, while the lowest ( $3.36 \pm 0.74$ ) were demonstrated in I. Furthermore, all data obtained were subjected to a normality test ( $p > 0.05$ ) to ensure a well-modelled dataset is used, based on a normal distribution. The homogeneity test ( $p = 0.018$ ) identified the dataset as non-homogeneous ( $p < 0.05$ ). Therefore, differences in both results were calculated using the T-Test. Table 2 showed differences in level within each group, although no significant differences were observed between II and III ( $p = 0.898$ ) as well as between Group I and V ( $p = 0.465$ ).

The free radicals contained in cigarette smoke are assumed to instigate tissue damage through various mechanical processes, and consequently trigger the peroxidation of lipids, proteins, and DNA. This damage reportedly reduces the antioxidant defence system and stimulates the release of pro-inflammatory cytokines.<sup>19</sup> In addition, an increase in the predominance of cellular damage from exposure to cigarette smoke that enters the respiratory tract directly elevates malondialdehyde levels in the blood. This compound is not considered a free radical, but the end product of lipid peroxidation.<sup>20</sup> The intrinsic characteristics include relatively strong stability and potential for use as an indicator of oxidative stress or body radical content.<sup>21,22</sup>

Table 3 shows the highest levels of C-reactive protein ( $6.58 \pm 0.58$ ) in Group II, while I had the lowest values ( $4.44 \pm 0.74$ ). Therefore, a normality test ( $P > 0.05$ ) was performed on all the data obtained to determine if the dataset was well-modelled by a normal distribution. The homogeneity test ( $p = 0.831$ ) showed the samples as homogeneous ( $p > 0.05$ ). The differences in both results for each group were then calculated using the ANOVA test, followed by the Least Significance Different (LSD) assessment.

In addition, free radicals entering the body potentially stimulate the phagocytosis process of the body's immune system. This

**Table 1. Means of malondialdehyde between all groups.**

Groups	Means $\pm$ SD	Normality	Homogeneity
I A group only given daily intake	$3.36 \pm 0.74$	0.754	0.018
II A group only exposed to cigarette smoke	$11.23 \pm 1.45$	0.367	
III A group given daily intake, cigarette smoke, and 0.5 ml of <i>Brassica oleracea</i> var. <i>italica</i> extract	$11.14 \pm 1.62$	0.580	
IV A group given daily intake, cigarette smoke, and 0.75 ml of <i>Brassica oleracea</i> var. <i>italica</i> extract	$5.60 \pm 0.94$	0.237	
V A group given daily intake, cigarette smoke, and 1 ml of <i>Brassica oleracea</i> var. <i>italica</i> extract	$3.87 \pm 0.83$	0.640	
VI A group given daily intake, cigarette smoke, and 1.25 ml of <i>Brassica oleracea</i> var. <i>italica</i> extract	$1.90 \pm 0.70$	0.552	

**Table 2. The results of T-test on malondialdehyde levels between all groups.**

Groups	I	II	III	IV	V	VI
I	-	-	-	-	-	-
II	0.000	-	-	-	-	-
III	0.000	0.898	-	-	-	-
IV	0.003	0.000	0.000	-	-	-
V	0.465	0.000	0.000	0.019	-	-
VI	0.038	0.000	0.000	0.000	0.006	-

process stimulates the secretion of pro-inflammatory cytokines, including interleukin-1, interleukin-6, and tumour necrosis factor-alpha (TNF- $\alpha$ ), known to be responsible for inflammatory responses.<sup>23,24,25</sup> The reaction serves as a defence mechanism against various microorganisms or foreign objects, and there is a need to control this effect to avoid attacks by autoimmune diseases.<sup>26</sup> Furthermore, a gradual change in the cells surrounding the site is observed during chronic attacks, and potentially results in permanent damage.<sup>27</sup> These conditions are also possibly created after chronic cigarette exposure, which initiates a plaque build-up in the artery. This formation is consequently implicated in atherosclerosis, which leads to serious diseases, including cardiovascular and stroke.<sup>28</sup> The increase in free radicals is signalled by an elevation in malondialdehyde levels, which directly triggers inflammatory responses with negative health impacts.

The ANOVA analysis results showed differences in the levels of C-reactive protein in each group. These were further evaluated using the Least Significance Different (LSD) test. Table 4 showed a significant difference ( $p < 0.05$ ) in Group I compared to others, while II demonstrated no substantial variation in contrast with III ( $p = 0.917$ ) and Group IV ( $p = 0.152$ ). However, no significant difference were observed between Group IV and III ( $p = 0.182$ ), as well as V ( $p = 0.139$ ) and VI ( $p = 0.085$ ). The results also indicate no substantial variations between Group V and VI ( $p = 0.794$ ). Table 5 depicts the relationship between increased malondialdehyde and C-reactive protein levels. This was ascertained using the Pearson test, and the results showed a strong correlation between both parameters ( $r = 0.713$ ).

The external antioxidants or exogen antioxidants obtained from food possess different action mechanisms compared to enzymatic antioxidants, despite the intrinsic neutralizing capacity. These compounds are known to reduce free radicals in various ways, including through the protection of cells from lipid peroxidation (Vitamin E), as strong reducing agents to be potentially reduced by enzymes and glutathione (Vitamin C). Previous reports showed proficiency in boosting the immune system ( $\beta$  carotene) and polyphenol groups, known to act as both antioxidant and anti-inflammatory agents.<sup>29,30,31</sup> In addition, non-enzymatic antioxidants reportedly reduce muscle pain and physical strain resulting from oxidative stress reactions.<sup>32</sup> Broccoli (*Brassica oleracea* var. *italica*) contains various types of non-enzymatic antioxidants and high polyphenol, including flavonoids, which is estimated to reduce free radicals and inflammation caused by cigarette smoke exposure.<sup>33,34</sup> This vegetable also contains selenium, and is known to potentially increase various enzymatic antioxidants, including superoxide dismutase (SOD) and glutathione peroxidase (GSH-Px).<sup>35</sup>

## Conclusions

The conclusion of this study is exposure to cigarette smoke increases free radicals and stimulates an inflammatory response in the body. Broccoli (*Brassica oleracea* var. *italica*) extracts plays an intrinsic role in reducing these negative impacts, and the effect is dose-dependent.

**Table 3. Means of C-reactive protein levels between all groups.**

Groups	Means $\pm$ SD	Normality	Homogeneity	P
I A group only given daily intake	4.44 $\pm$ 0.74	0.530	0.831	0.000
II A group only exposed to cigarette smoke	6.58 $\pm$ 0.58	0.642		
III A group given daily intake, cigarette smoke, and 0.5 ml of <i>Brassica oleracea</i> var. <i>italica</i> extract	6.54 $\pm$ 0.50	0.384		
IV A group given daily intake, cigarette smoke, and 0.75 ml of <i>Brassica oleracea</i> var. <i>italica</i> extract	6.02 $\pm$ 0.52	0.758		
V A group given daily intake, cigarette smoke, and 1 ml of <i>Brassica oleracea</i> var. <i>italica</i> extract	5.44 $\pm$ 0.45	0.074		
VI A group given daily intake, cigarette smoke, and 1.25 ml of <i>Brassica oleracea</i> var. <i>italica</i> extract	5.34 $\pm$ 0.74	0.248		

**Table 4. The results of the Least Significance Different (LSD) test between all groups.**

Groups	I	II	III	IV	V	VI
I						
II	0.000					
III	0.000	0.917	-	-	-	-
IV	0.000	0.152	0.182	-	-	-
V	0.014	0.006	0.008	0.139	-	-
VI	0.026	0.003	0.004	0.085	0.794	-

**Table 5. The results of the Pearson test between all groups.**

Groups	P	Pearson Test
Malondialdehyde	0.000	0.713
C-Reactive Protein		

Correspondence: Rivan Virlando Suryadinata, Faculty of Medicine, Universitas Surabaya (UBAYA), Surabaya, Jl. Tenggilis Mejoyo, KaliRungkut, Kec. Rungkut, Kota SBY, Jawa Timur 60293, Indonesia.

Tel.: +62.31.2981000. E-mail: rivan.virlando.s@staff.ubaya.ac.id

Key words: *Brassica oleracea*; C-reactive protein; free radical; inflammation; malondialdehyde.

Acknowledgement: The authors are sincerely grateful to the staff at the Faculty of Medicine, Universitas Surabaya (UBAYA) for the assistance rendered during this research.

Contributions: All authors contributed equally to this research.

Conflict of Interest: The author declare no conflicts of interest.

Funding: This research uses independent funds from all research members.

Clinical trials: This research was approved by the Health Research Ethics Committee of the University of Surabaya.

Availability of data and materials: All data generated or analyzed during this study are included in this published article.

Informed consent: Written informed consent was obtained from a legally authorized representative(s) for anonymized patient information to be published in this article.

Conference presentation: Part of this paper was presented at the 2<sup>nd</sup> International Nursing and Health Sciences Symposium that took place at the Faculty of Medicine, Universitas Brawijaya, Malang, Indonesia.

Received for publication: 3 December 2021.

Accepted for publication: 10 May 2022.

This work is licensed under a Creative Commons Attribution 4.0 License (by-nc 4.0).

©Copyright: the Author(s), 2023

Licensee PAGEPress, Italy

Healthcare in Low-resource Settings 2023; 11(s1):11210

doi:10.4081/hls.2023.11210

*Publisher's note: All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.*

## References

1. Lorensia A, Muntu CM, Suryadinata RV, et al. Effect of lung function disorders and physical activity on smoking and non-smoking students. *J Prev Med Hygiene* 2021;62:E89-E96.
2. West R. Tobacco smoking: Health impact, prevalence, correlates and interventions. *Psychol Health* 2017;32:1018–36.
3. Drope J, Schluger N, Cahn Z, et al. The Tobacco Atlas. Atlanta: American Cancer Society and Vital Strategies. *Am Cancer Soc* 2018;26 p.
4. Jamal A, King BA, Neff LJ, et al. Current Cigarette Smoking Among Adults — United States, 2005–2015. *MMWR Morb Mortal Wkly Rep* 2016;65:1205–1211.
5. Saleheen D, Zhao W, Rasheed A. Epidemiology and Public Health Policy of Tobacco Use and Cardiovascular Disorders in Low- and Middle-Income Countries. *Arterioscler Thromb Vasc Biol* 2014;34:1811–9.
6. National Center for Chronic Disease Prevention and Health Promotion (US) Office on Smoking and Health. The Health Consequences of Smoking—50 Years of Progress: A Report of the Surgeon General. Atlanta (GA): Centers for Disease Control and Prevention (US); 2014. PMID: 24455788.
7. Suryadinata RV, Wirjatmadi B. Selenium Linked to Increased Antioxidant Levels and Decreased Free Radicals in Lung Tissue of Wistar Rats Exposed to E-Cigarette Smoke. *J Global Pharma Technol* 2020;12:32-39.
8. Khanna A, Guo M, Mehra M, et al. Inflammation and Oxidative Stress Induced by Cigarette Smoke in Lewis Rat Brains. *J Neuroimmunol* 2013;254:69–75.
9. Winkelmann BR, von Holt K, Unverdorben M. Smoking and atherosclerotic cardiovascular disease: Part I: Atherosclerotic disease process. *Biomarkers in Medicine* 2009;3:411-428.
10. Suryadinata RV, Wirjatmadi B, Adriani M, et al. Effects of knowledge of vitamin D on attitudes toward sun exposure among middle-aged and elderly Indonesian adults. *Indian J Public Health Res Develop* 2018;9:11-15.
11. Bardaweel SK, Gul M, Alzweiri M, et al. Reactive Oxygen Species: the Dual Role in Physiological and Pathological Conditions of the Human Body. *Eurasian J Med* 2018;50:193–201.
12. Weng M, Xie X, Liu C, et al. The Sources of Reactive Oxygen Species and Its Possible Role in the Pathogenesis of Parkinson's Disease. *Parkinsons Dis* 2018;9163040.
13. Wang Y, Chun O, Song W. Plasma and dietary antioxidant status as cardiovascular disease risk factors: A review of human studies. *Nutrients* 2013;5:2969-3004.
14. Suryadinata RV, Wirjatmadi B, Adriani M. Efektivitas penerunan malondialdehyde dengan kombinasi suplemen antioksidan superoxide dismutase melon dengan gliadin akibat paparan asap rokok. *Global Medical and Health Communication* 2017;5:79-83
15. Biswas S, Das R, Banerjee ER. Role of free radicals in human inflammatory diseases. *AIMS Biophysics* 2017;4:596-614.
16. Hussain T, Tan B, Yin Y, et al. Oxidative Stress and Inflammation: What Polyphenols Can Do for Us?. *Oxidative Medicine and Cellular Longevity* 2016;1-9.
17. Sami FJ, Rahimah S. Uji aktivitas antioksidan ekstrak metanol bunga brokoli (*Brassica oleracea* L. var. *Italica*) dengan metode DPPH (2,2 diphenyl-1-picrylhydrazyl) dan metode ABTS (2,2 azinobis (3-etilbenzotiazolin)-6-asam sulfonat). [Test di attività antiossidante dell'estratto metanolo di fiori di broccoli (*Brassica oleracea* L. var. *Italica*) utilizzando i metodi DPPH (2,2 difenil-1-picrylhydrazyl) e ABTS (2,2 azinobis (3-etilbenzotiazolina)-6-solfonico)]. *Jurnal Fito farmaka Indonesia* 2015;2:107-110.
18. Suryadinata RV, Sukarno DA, Sardjono SC, et al. Antioxidant activity in red mulberries on sperm development exposed by cigarette smoke. *Bali Med J* 2021;10:583-586.
19. Suryadinata RV, Wirjatmadi B. The Molecular Pathways of Lung Damage by E-Cigarette in Experimental Mice. *Sultan Qaboos University Med J* 2021;1(1).
20. Cherian DA, Peter T, Narayanan A, et al. Malondialdehyde as a Marker of Oxidative Stress in Periodontitis Patients. *J Pharm Bioallied Sci* 2019;11: S297-S300.
21. Cui X, Gang J, Han H, et al. Relationship between free and total malondialdehyde, a well-established marker of oxidative stress, in various types of human biospecimens. *J Thorac Dis* 2018;10:3088–3097.
22. Ito F, Sono Y, Ito T. Measurement and Clinical Significance of Lipid Peroxidation as a Biomarker of Oxidative Stress:

- Oxidative Stress in Diabetes, Atherosclerosis, and Chronic Inflammation. *Antioxidants* 2019;8:72.
23. Mantovani A, Dinarello C, Molgora M, et al. Interleukin-1 and Related Cytokines in the Regulation of Inflammation and Immunity. *Immunity* 2019;50:778-795.
  24. Borsini A, Benedetto M, Giacobbe J, et al. Pro- and Anti-Inflammatory Properties of Interleukin in Vitro: Relevance for Major Depression and Human Hippocampal Neurogenesis. *Int J Neuropsychopharmacol* 2020;23:738-750.
  25. Phaniendra A, Jestadi DB. Free Radicals: Properties, Sources, Targets, and Their Implication in Various Diseases. *Ind J Clin Biochem* 2015;30:11-26.
  26. Chen L, Deng H, Cui H, et al. Inflammatory responses and inflammation-associated diseases in organs. *Oncotarget* 2018;9:7204-7218.
  27. Takeuchi O, Akira S. Pattern recognition receptors and inflammation. *Cell* 2010;140:805-20.
  28. Jee Y, Jung KJ, Lee S, et al. Smoking and atherosclerotic cardiovascular disease risk in young men: the Korean Life Course Health Study. *BMJ Open* 2019;9:e024453.
  29. Azlina MFN, Qodriyah MS, Kamisah Y. Tocopherol and Tocotrienol: Therapeutic Potential in Animal Models of Stress. *Curr Drug Targets* 2018;19:1456-1462.
  30. Petermelj TT, Coombes JS. Antioxidant supplementation during exercise training: beneficial or detrimental? *Sports Med* 2011;41:1043-69.
  31. Lohan SB, Vitt K, Scholz P, et al. ROS production and glutathione response in keratinocytes after application of  $\beta$ -carotene and VIS/NIR irradiation. *Chem Biol Interact* 2018;280:1-7.
  32. Simioni C, Zauli G, Martelli AM, et al. Oxidative stress: role of physical exercise and antioxidant nutraceuticals in adulthood and aging. *Oncotarget* 2018;9:17181-17198.
  33. Teixeira J, Chavarria D, Borges F, et al. Dietary Polyphenols and Mitochondrial Function: Role in Health and Disease. *Curr Med Chem* 2019;26:3376-3406.
  34. Suryadinata RV, Lorensia A, Sefania K. Effectiveness of Lime Peel Extract (*Citrus aurantifolia* Swingle) against C-Reactive Protein Levels in Alloxan-Induced Wistar Rats. *Global Medical and Health Communication* 2021;9:23-28.
  35. Suryadinata RV, Wirjatmadi G, Lorensia A. The time pattern of selenomethionine administration in preventing free radicals due to exposure to electric cigarette smoke. *J Public Health Res* 2021;10:2232.

# Brassica oleracea var. italica extract reducing free radicals and inflammation initiated by an exposure to cigarette smoke

## ORIGINALITY REPORT

14%

SIMILARITY INDEX

10%

INTERNET SOURCES

8%

PUBLICATIONS

6%

STUDENT PAPERS

## PRIMARY SOURCES

- 1** **Guillermo Paraje, Daniel Araya, Jeffrey Drope. "Illicit cigarette trade in Metropolitan Santiago de Chile", Tobacco Control, 2020** **1 %**

Publication
- 2** **Sabina Ranjit, Benjamin J. Patters, Kelli A. Gerth, Sanjana Haque, Sanjeev Chaudhary, Santosh Kumar. "Potential neuroprotective role of astroglial exosomes against smoking-induced oxidative stress and HIV-1 replication in the central nervous system", Expert Opinion on Therapeutic Targets, 2018** **1 %**

Publication
- 3** **[bmcvetres.biomedcentral.com](http://bmcvetres.biomedcentral.com)** **1 %**

Internet Source
- 4** **[healtheducationandpublichealth.com](http://healtheducationandpublichealth.com)** **1 %**

Internet Source
- 5** **[www.future-science.com](http://www.future-science.com)** **1 %**

Internet Source
- 6** **[www.scielo.br](http://www.scielo.br)** **1 %**

Internet Source

7	Submitted to Australian National University Student Paper	1 %
8	Giusi Prencipe, Claudia Bracaglia, Fabrizio De Benedetti. "Interleukin-18 in pediatric rheumatic diseases", Current Opinion in Rheumatology, 2019 Publication	1 %
9	<a href="http://ifst.onlinelibrary.wiley.com">ifst.onlinelibrary.wiley.com</a> Internet Source	1 %
10	<a href="http://www.spandidos-publications.com">www.spandidos-publications.com</a> Internet Source	1 %
11	Submitted to University of Iowa Student Paper	1 %
12	Chao Yu Hsu, Yi Sheng Lin, Wei Chun Weng, Lauren Panny et al. "Phloretin Ameliorates Testosterone-Induced Benign Prostatic Hyperplasia in Rats by Regulating the Inflammatory Response, Oxidative Stress and Apoptosis", Life, 2021 Publication	<1 %
13	Submitted to Colorado State University, Global Campus Student Paper	<1 %
14	<a href="http://www.nejm.org">www.nejm.org</a> Internet Source	<1 %

Submitted to Columbia College of Missouri



15

Student Paper

&lt;1 %

16

Submitted to Higher Education Commission  
Pakistan

Student Paper

&lt;1 %

17

Submitted to University of Newcastle

Student Paper

&lt;1 %

18

Submitted to University of Wales Swansea

Student Paper

&lt;1 %

19

encyclopedia.pub

Internet Source

&lt;1 %

20

journals.lww.com

Internet Source

&lt;1 %

21

wiadlek.pl

Internet Source

&lt;1 %

22

"Oxidative Stress in Fibromyalgia Syndrome",  
Reactive Oxygen Species in Biology and  
Human Health, 2016.

Publication

&lt;1 %

23

Sevil Şenkardeş, İhsan Han, Hakan Hançer,  
Mürüvvet Abbak, Özge Çevik, Ş. Güniz  
Küçükgül. "Synthesis, characterization and  
biological evaluation of some novel  
sulfonylthiosemicarbazides", Phosphorus,  
Sulfur, and Silicon and the Related Elements,  
2019

&lt;1 %

---

24	<a href="http://nutritionj.biomedcentral.com">nutritionj.biomedcentral.com</a> Internet Source	<1 %
25	<a href="http://web.citius.technology">web.citius.technology</a> Internet Source	<1 %
26	Submitted to University of York Student Paper	<1 %
27	<a href="http://etheses.whiterose.ac.uk">etheses.whiterose.ac.uk</a> Internet Source	<1 %
28	<a href="http://meltem.ege.edu.tr">meltem.ege.edu.tr</a> Internet Source	<1 %
29	<a href="http://watermark.silverchair.com">watermark.silverchair.com</a> Internet Source	<1 %
30	<a href="http://cmjournal.biomedcentral.com">cmjournal.biomedcentral.com</a> Internet Source	<1 %
31	<a href="http://ppgfsc.posgrad.ufsc.br">ppgfsc.posgrad.ufsc.br</a> Internet Source	<1 %
32	<a href="http://ppi.ubaya.ac.id">ppi.ubaya.ac.id</a> Internet Source	<1 %
33	<a href="http://research.thea.ie">research.thea.ie</a> Internet Source	<1 %
34	<a href="http://www.springermedizin.de">www.springermedizin.de</a> Internet Source	<1 %

---

35

Jorddy Neves Cruz, Mozaniel Santana de Oliveira, Sebastião Gomes Silva, Antonio Pedro da Silva Souza Filho et al. "Insight into the Interaction Mechanism of Nicotine, NNK, and NNN with Cytochrome P450 2A13 Based on Molecular Dynamics Simulation", Journal of Chemical Information and Modeling, 2019

<1 %

Publication

---

Exclude quotes Off

Exclude matches Off

Exclude bibliography Off

# Brassica oleracea var. italica extract reducing free radicals and inflammation initiated by an exposure to cigarette smoke

---

GRADEMARK REPORT

---

FINAL GRADE

**/0**

GENERAL COMMENTS

**Instructor**

---

PAGE 1

---

PAGE 2

---

PAGE 3

---

PAGE 4

---

PAGE 5

---