



Systematic Literature Review: Side Effect of *Moringa oleifera* Lam.

Deborah Felinda Hari Wijaya¹, Alfian Hendra Krisnawan², Oeke Yunita², Krisyanti Budipramana², Karina Citra Rani³, Nikmatul Ikhrom Eka Jayani^{2*}

¹Faculty of Pharmacy, University of Surabaya, Surabaya, Indonesia

²Departement of Pharmaceutical Biology, University of Surabaya, Surabaya, Indonesia

³Departement of Pharmaceutics, Faculty of Pharmacy, University of Surabaya, Surabaya, Indonesia

*Corresponding author. Email: nikmatul.ikhrom@staff.ubaya.ac.id

ABSTRACT

Moringa oleifera Lam. (Drumstick tree) is a herbal plant commonly found in subtropical and tropical areas such as Indonesia. Moringa plants have been widely used because they have many pharmacological effects. In addition, due to the high consumption of Moringa, there still needs to be monitoring regarding the safety of Moringa. This study aimed to identify and analyze the side effects of using Moringa based on the results of a systematic literature review. A systematic literature review using articles with a case report study design (collected from 2011 to 2021). The search databases used are PubMed, Google Scholar, and Science Direct. Journal quality is assessed by JBI (Joanna Briggs Institute) Critical Appraisal. 10 articles met the criteria and were of good quality. Based on the results, the side effects that occur in the use of Moringa are in the respiratory tract, dermal, and rhabdomyolysis. The incidence of side effects in Moringa is associated with chemical compounds contained in Moringa which can trigger hypersensitivity reactions and structural similarities to other synthetic drug compounds so that they can cause the same side effect reactions. Moringa in both leaves and seeds can cause side effects, the manifestations of which are respiratory, dermal, and rhabdomyolysis disorders.

Keywords: *Moringa oleifera* Lam., side effects, atopic dermatitis, rhabdomyolysis.

1. INTRODUCTION

Adverse drug reactions (ADR) are the body's response to an unwanted drug, usually occurring at doses of common drugs used to prevent, diagnose, or treat disease [1]. Of the cases of ADR that occur in the use of pharmaceutical products, about 7.9% are cases of ADR due to the use of herbs. Side effects are any unintended effect of a pharmaceutical product occurring at doses normally used in humans related to the drug's pharmacological properties. Reactions to side effects of using herbs do not always occur in every individual and are often unpredictable, but the factors that cause these side effects will increase a person's chances of experiencing side effects from using herbs. The main factors that cause side effects in the use of herbs are the use of herbs that are not following the indications, the

dosage of herbs is not right, or the consumption of herbs for a long time [2].

Moringa oleifera Lam. (drumstick tree) is a herbal plant that grows in subtropical and tropical climates such as Indonesia and belongs to the Moringaceae family [3]. Moringa is easy to grow and easy to obtain by the community and its use has long been known in the community. Moringa has many benefits and has been widely used by the community both as an herb and as food because Moringa contains many nutritious compounds as well as complex nutritional values [4][5]. Although Moringa herbs are often considered safer than synthetic drugs, in reality, Moringa herbs can cause side effects, either due to side effects from the use of Moringa itself or the interaction of Moringa with other herbs or other synthetic drugs [6]. Monitoring related to

safety in the use of Moringa herbs then becomes important, in addition to the efficacy of Moringa herbs which has long been known in the community [7].

Monitoring is important not only for therapeutic effects but also for side effects. Indications for the use of Moringa that are not appropriate, the period of consumption of Moringa is too long, and the dose of Moringa that is too much can not provide good efficacy and can even trigger unwanted side effects for the user [7][8]. The side effects of using Moringa in patients can be observed from physical examination and laboratory examinations related to the clinical manifestations of the side effects of Moringa experienced by patients.

This study reviews the literature systematically on the side effects that occur due to the use of *Moringa oleifera* Lam. (Moringa). This systematic literature review contains a description of the theory and findings from reference materials to be used as a basis for research activities on the risk of side effects using Moringa (*Moringa oleifera* Lam.). As previously described, the use of Moringa herbs is considered safe by the public, but it is also possible to cause side effects in its use. This study aimed to identify and analyze the side effects of using Moringa based on the results of a literature review. Research related to the evaluation of the side effects of herbal medicine is one part of the pharmacovigilance system of herbal medicines that the WHO has regulated [8]. The most crucial part is that it is hoped that awareness will arise so that people are more aware of the use of herbs, especially moringa, and can detect if signs of side effects appear after consuming moringa so that they can be prevented from causing more severe problems.

2. METHODS

2.1 Literature Search

The literature search aims to find articles with case report study designs that report cases related to side effects after using Moringa. This literature search used the publication year limitation on the search database with the search limitation for the last 10 years. The case reports collected are case reports of side effects of using Moringa from 2011 to 2021.

The literature search database sources used include PubMed, Google Scholar, and Science Direct. Search the database using the keywords "moringa", "moringa oleifera", "adverse drug reaction", "case report", "asthma", "allergy", and "skin irritation". All searches in the database used Boolean Logic, on PubMed, Google Scholar, and Science Direct were used ((moringa OR "moringa oleifera") AND "adverse drug reaction" OR "case report"); "moringa oleifera" AND "asthma" AND "allergy"; "moringa oleifera" AND "skin irritation" AND "allergy". The article search process was carried out using the PubMed, Google Scholar, and Science Direct search databases.

2.2 Article Selection

The articles selected for review have met the inclusion and exclusion criteria. The inclusion criteria in this study included articles from the last 10 years; article using English or Indonesian; articles in the form of case reports or case study; articles that meet the critical assessment criteria using the JBI Critical Appraisal Tools.

Exclusion criteria include duplication of journals; the article does not match the topic of Moringa oleifera natural side effects.; articles in the form of reviews; the plant presented in the article is not *Moringa oleifera* Lam.; drug interactions cause the incidence of side effects related to the use of Moringa; the identity of the library source is incomplete; there are no full-text articles.

2.3 Article Quality Criteria

In determining the quality criteria of the article, an article review is carried out using critical appraisal tools that follow the article research design. This critical assessment aims to assess the quality, credibility, and clinical practice value of a study and to avoid possible bias in a study design.

The article research design used is a case report, for and the appropriate critical appraisal tools are JBI Critical Appraisal tools. The answers in each of the available question checklists will affect the quality of the article. There are four answer columns related to the questions in the checklist, namely "yes", "no", "unclear", and "not applicable". From the results of critical appraisal using JBI Critical Appraisal Tools, articles are considered to have good quality if the article has a "yes" answer >75% of the total answers to questions on JBI Critical Appraisal Tools.

Articles that have been collected are then reviewed or studied by 3 different reviewers. If there are differences in perception, then they are discussed with 3 other reviewers. The article tracing diagram refers to the PRISMA 2020 flow diagram, which is presented in Figure 1.

2.4 Data Analysis

All selected articles found in the search database using keywords that have been designed and have met the inclusion and exclusion criteria will be further examined using the JBI Critical Appraisal Tools. The subjects in this study were all patients who had experienced adverse reactions while using Moringa. Manifestations of side effects have been assessed by physical examinations and laboratory test results of the patient that are relevant to the clinical. Further data analysis was carried out descriptively.

The side effects on the respiratory tract can be seen from the results of physical examination in the form of symptoms of shortness of breath, tachypnea (rapid breathing frequency), and the patient's blood pressure; as well as the results of laboratory tests, namely the value of the Forced Expiratory Volume in one second / Forced Vital Capacity (FEV1/FVC) as well as the value of the FeNO (Fractional Exhaled Nitric Oxide) test,

serum tryptase level, and skin prick test. The side effects on the dermal side can be seen from the results of physical examination in the form of erythema (redness of the skin), vesicles on skin rashes, and itching. Rhabdomyolysis can be seen from the results of laboratory tests in the form of creatine kinase (CK) levels and the results of physical examinations in the form of pain in certain body parts.




Figure 1. PRISMA 2020 flow diagram search

3. RESULTS AND DISCUSSION

The search in the database was performed using keywords combined with Boolean operators. Search the PubMed, Google Scholar, and Science Direct databases from November 2021, to February 2022, using the keywords ((Moringa OR "moringa oleifera") AND "adverse drug reaction" OR "case report"); "moringa oleifera" AND "asthma" AND "allergy"; "moringa oleifera" AND "skin irritation" AND "allergy". Then the articles in the PubMed, Google Scholar, and Science Direct databases were obtained, each with 2 articles; 2,898 articles; and 188 articles. The articles that have

been obtained from the search database are then screened based on their language, title, and abstract. After screening, duplicate articles were eliminated for further analysis. The analysis results based on the inclusion and exclusion criteria obtained 10 articles that are relevant to the topic to be studied and can be used to answer the research topic. Table 1. Shows quality of case report articles with JBI Critical Appraisal Tools.

The ten relevant articles were then checked for quality using the JBI Critical Appraisal Tools as shown

in Table 2. All articles with good quality in the JBI Critical Appraisal Tools assessment will be analyzed further. After the quality assessment was carried out, 10 articles of good quality were obtained which would then be used in research. Of the 10 case reports that have been collected, 2 articles that are side effects of using Moringa seeds where the clinical manifestations of these side effects are disorders of the respiratory tract, while 8 other articles are side effects of using Moringa leaves where 5 articles report disturbances in the parts of the body, dermal and 3 other articles reported muscle disturbances in the form of rhabdomyolysis.

Clinical manifestations of the respiratory tract disorders can be in the form of asthma or anaphylactic reactions that occur in a fast time, caused using Moringa seeds. The side effect reaction of using Moringa leaves on the dermal clinical manifestations can be in the form of skin redness (erythema) with the presence or absence of lesions or prominent skin areas accompanied by

itching, even skin redness in some patients can leave hyperpigmentation. Skin redness can be a clustered maculopapular rash or a morbilliform rash that spreads throughout the body. The side effects of rhabdomyolysis due to the use of Moringa leaves manifest as pain in certain body parts, especially the limbs. Clinical manifestations of side effects of Moringa leaves or seeds, can then be classified based on the dose and duration of exposure. Chronic types classification when Moringa used for six months or up to years [9]. Chronic types side effect classification in case reports of Poussel *et al.* [10] and Olanrewaju *et al.* [11] due to Moringa is consumed for a long time, about 1-2 years. The case reports of Berglund [12]; Adegbidi *et al.* [13]; Witharana *et al.* [14]; Sagrera *et al.* [15]; Deogade *et al.* [16]; Acharya *et al.* [17]; Treidler *et al.* [18] can be classified into the bizarre types of side effects because of the appearance of side effects in the patient unpredictable and the use of Moringa in less than six months.

Table 1. Checklist the quality of case report articles with JBI Critical Appraisal Tools [19]

No.	Author	Subject	Question							
			1	2	3	4	5	6	7	8
1	Poussel <i>et al.</i> (2015)	Human	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
2	Berglund, L. (2018)	Human	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
3	Olanrewaju <i>et al.</i> (2017)	Human	Yes	Yes	Yes	Unclear	Yes	Yes	Yes	Yes
4	Adegbidi <i>et al.</i> (2019)	Human	Yes	Unclear	Yes	No	Yes	Yes	Yes	Yes
5	Witharana <i>et al.</i> (2018)	Human	Yes	Yes	Yes	Yes	Yes	Yes	Yes	Yes
6	Sagrera <i>et al.</i> (2021)	Human	Yes	Yes	Yes	No	Yes	Yes	Yes	Yes
7	Deogade <i>et al.</i> (2019)	Human	Yes	Unclear	Yes	Yes	Yes	No	Yes	Yes
8	Acharya <i>et al.</i> (2020)	Human	Yes	Yes	Yes	Unclear	Yes	Unclear	Yes	Yes
9	Treidler <i>et al.</i> (2017)	Human	Yes	Unclear	Yes	Yes	Yes	Yes	Yes	Yes
10	Treidler <i>et al.</i> (2017)	Human	Yes	Unclear	Yes	Yes	Yes	No	Yes	Yes

The presence of side effect reactions due to the use of Moringa leaves or seeds can be seen from the physical examination results and the results of the patient's laboratory tests presented in Table 2. Disorders of the respiratory tract due to the use of Moringa seeds can be seen from the symptoms of coughing, shortness of breath, and wheezing, patients, as well as the results of the measurement of the FEV1/FVC value where if there is a decrease of more than 20% from the normal value (normal value FEV1/FVC > 75-80%) indicates an obstruction in the airways whose manifestations can be shortness of breath and wheezing such as as well as asthma symptoms complained of by patients in case

reports [20]. In addition to asthma, respiratory disorders due to the use of Moringa seeds can be in the form of an anaphylactic reaction that occurs rapidly, characterized by symptoms of hypotension, high respiratory rate (tachypnea), and angioedema. Investigations for anaphylaxis are a measurement of serum tryptase levels. This serum tryptase level will indicate when a patient has an anaphylactic reaction. Elevated serum tryptase levels were associated with mast cell degranulation due to the anaphylaxis reaction that occurred in the patient. Mast cells contain mature -tryptase so that when mast cells degranulate, β-tryptase will be released into the blood, increasing serum tryptase levels [21]. Disorders

of the dermal part seen from the results of the patient's physical examination in the form of skin redness (erythema) with prominent skin parts (the presence of lesions) or not and accompanied by itching, even redness of the skin which can leave persistent hyperpigmentation [22]. Muscle disorders in the form of rhabdomyolysis can be seen from the symptoms of

muscle pain in certain body parts, as well as the results of laboratory tests in the form of creatine kinase (CK) levels. An increase in CK levels >40 times (normal CK value in men: 55-170 IU/L and women: 30-135 IU/L) indicates the presence of muscle cell necrosis which is a manifestation of ongoing muscle pain and progresses to rhabdomyolysis [23].

Table 2. Manifestation of *Moringa oleifera* Side Effects

No	Part of Moringa	Dosage and duration of exposure	Administration route	Moringa side effects manifestations	Parameters affected	Author
1	Pods of <i>Moringa oleifera</i> (MO)	Dosage: - Duration of exposure: 2 years where the patient was exposed 3 hours/day; 3 days/week	Inhaled	Respiratory tract (shortness of breath and wheezing such as as well as asthma symptoms)	<ul style="list-style-type: none"> • FEV1/FVC= 81%. • FeNO test= 91.21 ppb; • Methacholine test of 1280 g showed a decrease in the value of FEV1/FVC by 22% from the normal value. • Skin prick test, the diameter of the reddish spots on the skin is 10 mm. 	[10]
2	Pods of MO	Dosage: - Exposure time: 20 minutes after eating MO seed cooking	Oral (Food)	Respiratory tract (anaphylactic reaction)	<ul style="list-style-type: none"> • Blood pressure= 85/51 mmHg. • Tachypnea (+) • Serum tryptase levels were 39.6 g/L; • Skin prick test, the diameter of the reddish spots on the skin is 4 mm. 	[12]
3	Leaves of MO	Dosage: - Exposure time: 1 year after consuming MO leaf aqueous extract	Oral (Water extract)	Dermatitis	<ul style="list-style-type: none"> • Physical examination: <ol style="list-style-type: none"> a) red macules with the presence or absence of vesicles; b) Lesions in the form of black spots on the upper and lower arms, and thighs. 	[11]
4	Leaves of MO	Dosage: - Exposure time: 24 hours after eating MO leaf cooking	Oral (Food)	Dermatitis	<ul style="list-style-type: none"> • Physical examination: Grayish patches, not scaly, oval with clear boundaries, and scattered around the neck, back, legs, chest, and abdomen. 	[13]
5	Leaves of MO	Dosage: - Exposure time: 14 hours after eating MO leaf cooking	Oral (Food)	Steven Johnson Syndrome	<ul style="list-style-type: none"> • Fever (+). • Skin redness (macular popular rash) (+); • Canker sores (mouth ulcers) to difficulty swallowing. • Blood pressure= 130/80 mmHg. • Heart rate = 82 x/minute. 	[14]

					<ul style="list-style-type: none"> • Neutrophil content = 90%. • Platelet level= 260,000×1012/L.
6	Leaves of MO	Dosage: - Exposure time: 3 days after consuming MO leaf powder	Oral (Food)	Dermatitis	<ul style="list-style-type: none"> • Mild shortness of breath. • Swollen tongue. • Morbilliform rash (red spots) on the face, trunk, and upper arms accompanied by papules. • Biopsy data revealed the presence of necrotic keratinocytes.
7	Leaves of MO	Dosage: - Exposure time: 3 days after consuming MO leaves	Oral (Food)	Dermatitis	<ul style="list-style-type: none"> • Skin rash that spreads all over the patient's back. • Blood pressure 110/80. • Heart rate 82 beats/minute.
8	Leaves of MO	Dosage: - Exposure time: Immediately after ingestion of MO leaves	Oral (Food)	Rhabdomyolysis	<ul style="list-style-type: none"> • Severe pain in both thighs. • Nausea. • The creatinine kinase level was increased, but the creatinine kinase value was not attached.
9	Leaves of MO	Dosage: - Exposure time: Immediately after ingestion of MO leaves	Oral (Food)	Rhabdomyolysis	<ul style="list-style-type: none"> • Pain in the hands. • Creatine Kinase (CK) 19.210 U/L.
10	Leaves of MO	Dosage: - Exposure time: Immediately after ingestion of MO leaves	Oral (Food)	Rhabdomyolysis	<ul style="list-style-type: none"> • Pain in the hands. • Creatine Kinase (CK) 22.000 U/L.

Asthma that occurs after 2 years of exposure to Moringa leaf powder is associated with the content of niazirin, marumoside A, and sitosterol-3-O- β -D-glucoside contained in the seeds which can increase the overexpression of Th2 T Cells [24]. The increase in T cells will induce the mechanism of a type IV hypersensitivity reaction. Type IV hypersensitivity reaction is a delayed hypersensitivity reaction that takes 24-72 hours or even a long time to show the onset of symptoms [25]. The mechanism of the type IVb hypersensitivity reaction is associated with an asthmatic

reaction (Figure 2) which begins with an epitope on an antigen recognized by T cells memory. The antigens in this case were niazirin, marumoside A, and sitosterol-3-O- β -D-glucoside. Th2 T-cells secrete IL-4, IL-13, and IL-5 cytokines, which promote B cell production of IgE and IgG4, macrophage deactivation and mast cell and eosinophil responses. The high production of IL-5 leads to eosinophilic inflammation, the characteristic inflammatory cell type in many drug hypersensitivity reactions. The inflammatory mediators that can cause asthma reactions [26][27].




Figure 2. Mechanism of type IV hypersensitivity reaction when exposed to Moringa seeds

The anaphylactic reaction that only lasted 20 minutes after ingestion of Moringa seeds was associated with the protein contained in Moringa seeds capable of binding to IgE which then proved the occurrence of a type I hypersensitivity reaction as shown in Figure 3 [28][29]. Type I hypersensitivity reactions occur rapidly within seconds to 2 hours. The mechanism of the type I hypersensitivity reaction starts from Moringa seed protein which is recognized by bronchial epithelial cells and then triggers the formation of IL-25 and IL-33 which will trigger the action of dendritic cells, as well as basophil cells and Natural Killer (NK) cells which

will trigger IL-4. Dendritic cells will carry allergens (proteins) to be processed in the lymph and after those allergens will be recognized by CD4+ T cells together with MHC class II which can then cause T cells to differentiate into Th2 cells [30]. IL-4 will then amplify th2 cells to produce B cells. B cells will then produce IgE. IgE will bind to high-affinity IgE receptors (FcεRI) causing mast cells to degranulate and release inflammatory mediators that can cause vasoconstriction and excessive mucus secretion which can cause shortness of breath reactions in patients [31][32].




Figure 3. Mechanism of type I hypersensitivity reaction when exposed to Moringa seed protein

Dermal disorders due to the use of Moringa leaves are associated with the alkaloid content, namely moringin contained in Moringa leaves shown in Figure 4. Moringin has a similar mechanism of action with ephedrine, a sympathomimetic drug [33][34]. This sympathomimetic drug then triggers the formation of a T cell complex with MHC, triggering a type IVa hypersensitivity reaction mediated by T cells. The

mechanism of the type IVa hypersensitivity reaction begins with the epitope on moringin recognized by T cells memory. T cells memory then together with MHC class II will trigger T cells to differentiate into Th1 cells and Th2 cells. Th1 cells with Th17 cells then stimulate macrophage activation and strengthen the work of CD8+ Tc cells. Activation of these macrophages can release IL-12, which can then stimulate ThCD4+ cells

to produce IL-2 and IFN- γ which are inflammatory mediators whose manifestations are dermatitis [35]. In addition, the epitope on the antigen will be on the surface of the APC (Antigen-presenting cells) which can then be directly recognized by Tc CD8+ cells which

will induce apoptosis or cell death occupied by the antigen from Moringa. Apoptosis then induces activation of macrophage cells which causes the release of IL-12 cells which will then stimulate ThCD4+ cells to produce IL-2 and Interferon gamma (IFN- γ) [32][36].




Figure 4. Type IV hypersensitivity reaction causing atopic dermatitis

Rhabdomyolysis disorders are associated with the content of flavonoids in Moringa leaves, namely quercetin and kaempferol which have a similar structure to the statin class of drugs shown in Figure 5 [37]. The flavonoids in Moringa leaves can inhibit HMG-CoA reductase thereby inhibiting the formation of farnesyl-PP which is the basic ingredient in the formation of ubiquinone (CoQ10) so that CoQ10 levels decrease in the body as depicted in Figure 6. The absence or low levels of ubiquinone (coenzyme Q10) can further cause damage to the mitochondria so that they cannot produce ATP or energy for the muscles so that the manifestation is experiencing muscle pain or myalgia. ATP that is not produced sustainably can result in the death of mitochondrial cells which will lead to muscle tissue damage because the muscles do not get ATP as energy to be used by the muscles, which manifests in the form of rhabdomyolysis which is characterized by a significant increase in creatine kinase levels up to >40 times the normal value limit [23][38].

In addition, the mechanism of action of flavonoids in inhibiting cholesterol synthesis can also result in cholesterol depletion in the plasma membrane of muscle fibers (sarcolemma) which is thought to cause myopathy because the sarcolemma is depleted so that the stability of the sarcolemma can be disturbed and the ion balance in the sarcolemma can change [23].




Figure 5. The structural similarity of kaempferol and quercetin with simvastatin




Figure 6. Mechanism of Action of Statins

CONCLUSION

From the results of case reports collected from 2011 to 2021, it was found that the Moringa plant used for its leaves and seeds can cause side effects with clinical manifestations in the respiratory tract, dermatology, and rhabdomyolysis. The side effects of these herbs can arise depending on the method of consuming the herbs, both the dose consumed and the duration of herbal consumption. Reports related to side effects of using herbs in case reports should be accompanied by complete patient laboratory examination data to identify side effect reactions correctly. If side effects are found in the use of herbs in the community, they should be reported to the Pusat Farmacovigilant / MESO Nasional Badan Pengawasan Obat dan Makanan RI.

AUTHOR'S CONTRIBUTIONS

Deborah Felinda Hari Wijaya contribute for conducting research and preparing the data, contribute for writing original article. Alfian Hendra Krisnawan, Oeke Yunita, Krisyanti Budipramana, contribute for reviewing the article, Karina Citra Rani contribute for writing original article and reviewing the article. Nikmatul Ikrom Eka Jayani contribute for the conceptualization and writing original article, reviewing the article.

ACKNOWLEDGEMENT

The authors would like to thank the Faculty of Pharmacy and LPPM (Research and Community Service

Unit) of the University of Surabaya for supporting this research.

REFERENCES

- [1] Badan Pengawas Obat dan Makanan Republik Indonesia, "Modul Farmakovigilans Bagi Tenaga Profesional Kesehatan," 2020.
- [2] D. H. Phua, A. Zosel, and K. Heard, "Dietary supplements and herbal medicine toxicities-when to anticipate them and how to manage them," *Int. J. Emerg. Med.*, vol. 2, no. 2, pp. 69–76, 2009, doi: 10.1007/s12245-009-0105-z.
- [3] J. A. Parotta, "Moringa oleifera Moringa oleifera," *Agrofor. Database*, vol. 0, no. November 2017, pp. 4–9, 2009.
- [4] K. N. Berawi, R. Wahyudo, and A. A. Pratama, "Potensi Terapi Moringa oleifera (Kelor) pada Penyakit Degeneratif," *J. Kedokt. Univ. Lampung*, vol. 3, no. 1, pp. 210–214, 2019, doi: 10.23960/jkunila31210-214
- [5] S. Kesharwani, P. Prasad, A. Roy, and R. K. Sahu, "An Overview on Phytochemistry and Pharmacological Explorations of Moringa oleifera," *UK J. Pharm. Biosci.*, vol. 2, no. 1, p. 34, 2014, doi: 10.20510/ukjpb/2/i1/91151.
- [6] Badan Pengawas Obat dan Makanan Republik Indonesia, "Peraturan BPOM No 32 Tahun 2019," 2019.
- [7] Katno and S.Pramono, "Tingkat Manfaat dan Keamanan Tanaman Obat dan Obat Tradisional," *Trends in Cognitive Sciences*, vol. 6, no. 12, pp. 538–539, 2002.
- [8] World Health Organization (WHO), "WHO guidelines on safety monitoring of herbal medicines in pharmacovigilance systems," *World Heal. Organ. Geneva*, p. 82, 2004.
- [9] H. Dewoto, "Pengembangan Obat Tradisional Indonesia menjadi Fitofarmaka," *Maj. Kedokt. Indones.*, pp. 205–11, 2007.
- [10] M. Poussel, E. Penven, C. Richard, S. Jacquenet, F. Chabot, and C. Paris, "Occupational asthma to 'the miracle tree' (Moringa oleifera): First description," *J. Allergy Clin. Immunol. Pract.*, vol. 3, no. 5, pp. 813–814, 2015, doi: 10.1016/j.jaip.2015.05.016.
- [11] F. O. Olanrewaju, O. Onayemi, O. A. Olasode, and M. M. Oripelaye, "Fixed drug eruption induced by moringa oleifera leaf extracts – A case report," *Our Dermatology Online*, vol. 8, no. 4, pp. 413–416, 2017, doi: 10.7241/ourd.20174.117.

- [12] L. J. Berglund, "Anaphylaxis to *Moringa oleifera*: First description," *Australas. Med. J.*, vol. 11, no. 3, pp. 176–177, 2018, doi: 10.21767/amj.2018.3344.
- [13] H. Adégbidi et al., "Fixed Pigmented Erythema Induced by," *Clin. Res. Dermatology*, vol. 2, no. 2, pp. 1–3, 2019.
- [14] E. W. R. A. Witharana, W. M. G. A. S. T. B. Wijetunga, and S. K. J. Wijesinghe, "Stevens - Johnson syndrome (SJS) following Moringa leaf (*Moringa oleifera*) consumption," *Ceylon Med. J.*, vol. 63, no. 4, p. 188, 2018, doi: 10.4038/cmj.v63i4.8771.
- [15] A. Sagrera, T. Montenegro, and L. Borrego, "Cutaneous Toxicity Due to *Moringa oleifera*," *Actas Dermo-Sifiliográficas (English Ed.)*, vol. 112, no. 10, pp. 953–954, 2021, doi: 10.1016/j.adengl.2021.09.008.
- [16] M. S. Deogade, V. Kumbhare, and S. Gaidhane, "Moringa oleifera Lam . Enriched Biscuits Induced Adverse Drug Reaction- a Case Report," vol. 16, no. 2, pp. 27–30, 2019.
- [17] A. Acharya, L. Freeman, and M. R. Alvarez, "A Case Of Necrotizing Myopathy After Moringa Use," 2020.
- [18] S. Treidler et al., "Necrotizing myopathy in two patients after the use of Moringa (P2.137)," *Neurology*, vol. 88, no. 16 Supplement, p. P2.137, 2017, [Online]. Available: http://n.neurology.org/content/88/16_Supplement/P2.137.abstract.
- [19] Joanna Briggs Institute, "Checklist for Case Reports," 2016. [Online]. Available: https://joannabriggs.org/ebp/critical_appraisal_tool%0Ahttps://jbi.global/critical-appraisal-tools%0Ahttp://joannabriggs.org/assets/docs/critical-appraisal-tools/JBI_Critical_Appraisal-Checklist_for_Case_Reports2017.pdf.
- [20] K. J. Tietze, *Clinical Skills for Pharmacists*, 3rd ed. Philadelphia: Elsevier, 2012.
- [21] P. Valent et al., "The serum tryptase test: An emerging robust biomarker in clinical hematology," *Expert Rev. Hematol.*, vol. 7, no. 5, pp. 683–690, 2014, doi: 10.1586/17474086.2014.955008.
- [22] S. Shams et al., "Maculopapular skin eruptions associated with Covid-19: A systematic review," *Dermatol. Ther.*, vol. 34, no. 2, pp. 1–14, 2021, doi: 10.1111/dth.14788.
- [23] G. M. Camerino, N. Tarantino, I. Canfora, M. De Bellis, O. Musumeci, and S. Pierno, "Statin-induced myopathy: Translational studies from preclinical to clinical evidence," *Int. J. Mol. Sci.*, vol. 22, no. 4, pp. 1–18, 2021, doi: 10.3390/ijms22042070.
- [24] N. Ma et al., "Three constituents of *moringa oleifera* seeds regulate expression of Th17-relevant cytokines and ameliorate tpa-induced psoriasis-like skin lesions in mice," *Molecules*, vol. 23, no. 12, pp. 7–9, 2018, doi: 10.3390/molecules23123256.
- [25] Y. Zhao, J. Yang, Y. D. Gao, and W. Guo, "Th17 immunity in patients with allergic asthma," *Int. Arch. Allergy Immunol.*, vol. 151, no. 4, pp. 297–307, 2010, doi: 10.1159/000250438.
- [26] A. Zawodniak and W. J. Pichler, *Allergy Frontiers: Clinical Manifestations*, 3rd ed. Springer, 2009.
- [27] J. Adam, W. J. Pichler, and D. Yerly, "Delayed drug hypersensitivity: models of T-cell stimulation," *Br. J. Clin. Pharmacol.*, vol. 71, no. 5, pp. 701–707, 2011, doi: 10.1111/j.1365-2125.2010.03764.x.
- [28] H. Wijaya, Y. Chalid, A. Thaharah, and A. F. Nugroho, "Pengaruh Proses Pengolahan terhadap Karakteristik Protein Alergen Belalang Sawah (*Oxya chinensis*)," *War. IHP/Journal Agro-based Ind.*, vol. 36, no. 1, pp. 11–21, 2019.
- [29] R. Böhm, E. Proksch, T. Schwarz, and I. Cascorbi, "Drug Hypersensitivity," *Dtsch. Arztebl. Int.*, vol. 113, no. 29–30, pp. 501–512, 2018, doi: 10.3238/arztebl.2018.0501.
- [30] A. Martorell et al., "Position document: IgE-mediated allergy to egg protein," *Allergol. Immunopathol. (Madr.)*, vol. 41, no. 5, pp. 320–336, 2013, doi: 10.1016/j.aller.2013.03.005.
- [31] Ó. Palomares et al., "dIvergEnt: How IgE axis contributes to the continuum of allergic asthma and anti-IgE therapies," *Int. J. Mol. Sci.*, vol. 18, no. 6, pp. 1–14, 2017, doi: 10.3390/ijms18061328.
- [32] L. E. Wuryaningsih, *Imunitas: Sarana Organ Beraktivitas*. 2015.
- [33] R.-T. Alphonsine, K. Moumouni, O. Nabere, L. Marius, and P. G. Innocent, "A comparative study of phytochemical profile and antioxidant activity of Sahelian plants used in the treatment of infectious diseases in northern part of Burkina Faso: *Acacia seyal* Delile and *Acacia tortilis* (Forssk.) Hayne subsp. *raddiana* (Savi)," *J. Pharmacogn. Phyther.*, vol. 11, no. 3, pp. 74–79, 2019, doi: 10.5897/jpp2019.0555.
- [34] J. O. Popoola, O. S. Aworunse, O. L. Oyesola, O. O. Akinola, and O. O. Obembe, "A systematic review of pharmacological activities and safety of *Moringa oleifera*," *J. HerbMed Pharmacol.*, vol. 9, no. 3, pp. 174–190, 2020, doi: 10.34172/jhp.2020.24.

- [35] W. J. Pichler, "Immune mechanism of drug hypersensitivity," *Immunol. Allergy Clin. North Am.*, vol. 24, no. 3, pp. 373–397, 2004, doi: 10.1016/j.iaac.2004.03.012.
- [36] M. C. Dispenza, "Classification of hypersensitivity reactions," *Allergy Asthma Proc.*, vol. 40, no. 6, pp. 470–473, 2019, doi: 10.2500/aap.2019.40.4274.
- [37] D. A. O. Dwi Ayu Romadhoni, Sri Murwani, "Efek Pemberian Ekstrak Air Daun Kelor (*Moringa oleifera* lam.) Terhadap Kadar LDL danHDL Serum Tikus Putih (*Rattus norvegicus*) Strain Wistar Yang Diberi Diet Aterogenik," 2012.
- [38] M. E. C. Magalhães, "Mecanismos de rabdomiólise com as estatinas," *Arq. Bras. Cardiol.*, vol. 85, no. SUPPL. 5, pp. 42–44, 2005, doi: 10.1590/s0066-782x2005002400011.

Open Access This chapter is licensed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits any noncommercial use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license and indicate if changes were made.

The images or other third party material in this chapter are included in the chapter's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the chapter's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder.





Series: [Advances in Biological Sciences Research](#)

Proceedings of the Conference on Natural Resources And Life Sciences 2022 (NRLS-BIO 2022)

[HOME](#)

[PREFACE](#)

[ARTICLES](#)

[AUTHORS](#)

[SESSIONS](#)



[ORGANIZERS](#)

[PUBLISHING INFORMATION](#)



Welcome to the Conference on Natural Resources And Life Sciences 2022 (NRLS-BIO 2022). To highlight the rapidly growing research and product development in the field of Biotechnology and Pharmacy, the 4th International Conference on Natural Resources and Life Sciences (NRLS) themed Biotechnology- and pharmacy-driven research and product development is presented. The 4th NRLS seeks to delved into the heart of these advancements, fostering an environment where over 100 researchers from diverse countries and regions can converge, such as Indonesia, Thailand, Poland, Germany, England, and South Korea.

Please click [here](#) for the conference website.

Atlantis Press

Atlantis Press – now part of Springer Nature – is a professional publisher of scientific, technical & medical (STM) proceedings, journals and books. We offer world-class services, fast turnaround times and personalised communication. The proceedings and journals on our platform are Open Access and generate millions of downloads every month.

For more information, please contact us at: contact@atlantis-press.com

- ▶ PROCEEDINGS
- ▶ JOURNALS
- ▶ BOOKS
- ▶ POLICIES
- ▶ MANAGE COOKIES/DO NOT SELL MY INFO
- ▶ ABOUT
- ▶ NEWS
- ▶ CONTACT
- ▶ SEARCH

[Home](#) [Privacy Policy](#) [Terms of use](#)   

Copyright © 2006-2024 Atlantis Press – now part of Springer Nature



Series: [Advances in Biological Sciences Research](#)

Proceedings of the Conference on Natural Resources And Life Sciences 2022 (NRLS-BIO 2022)

[HOME](#)

[PREFACE](#)

[ARTICLES](#)

[AUTHORS](#)

[SESSIONS](#)



[ORGANIZERS](#)

[PUBLISHING INFORMATION](#)

Chair

Tjie Kok

Faculty of Biotechnology–University of Surabaya, Indonesia

Technical Committee

Yulanda Antonius

Faculty of Biotechnology–University of Surabaya, Indonesia

Endang Wahyu Fitriani

Faculty of Pharmacy–University of Surabaya, Indonesia

Steering Committee

Sulistyo Emantoko D. P.

Faculty of Biotechnology—University of Surabaya, Indonesia

Farida Suhud

Faculty of Pharmacy—University of Surabaya, Indonesia

Organizing Committee

Popy Hartatie Hardjo

Faculty of Biotechnology—University of Surabaya, Indonesia

Yayon Pamula Mukti

Faculty of Biotechnology—University of Surabaya, Indonesia

Christina Mumpuni Erawati

Faculty of Biotechnology—University of Surabaya, Indonesia

Ardhia Deasy Rosita Dewi

Faculty of Biotechnology—University of Surabaya, Indonesia

Editors

Johan Sukweenadhi

Faculty of Biotechnology—University of Surabaya, Indonesia

Finna Setiawan

Faculty of Pharmacy—University of Surabaya, Indonesia

Scientific Committee

Mariana Wahjudi

Faculty of Biotechnology—University of Surabaya, Indonesia

Ike Dhiah Rochmawati

Faculty of Pharmacy—University of Surabaya, Indonesia

Miriam Agler-Rosenbaum

Friedrich-Schiller University, Jena, Germany

Jeniffer Silva

Department of Research and Development, the Bridge Biofoundry, Ciudad de Saber, Clayton, Panama

Shadi Rahimi

Chalmers University of Technology, Department of Biology and Biological Engineering Division of Systems and Synthetic Biology Kemivägen 10

Chris Jones

School of Biological Sciences—Institute of Cardiovascular and Metabolic Research, University of Reading, England

Grzegorz Pasternak

Wroclaw University of Science and Technology, Poland

Wouter Hinrichs

Groningen Research Institute of Pharmacy, University of Groningen, Netherlands

Garnpimol C. Ritthidej

Faculty of Pharmaceutical Sciences—Chulalongkorn University, Thailand

Mohamed El-Agamy Farh

Atlantis Press

Atlantis Press – now part of Springer Nature – is a professional publisher of scientific, technical & medical (STM) proceedings, journals and books. We offer world-class services, fast turnaround times and personalised communication. The proceedings and journals on our platform are Open Access and generate millions of downloads every month.

For more information, please contact us at: contact@atlantis-press.com

- ▶ PROCEEDINGS
- ▶ JOURNALS
- ▶ BOOKS
- ▶ POLICIES
- ▶ MANAGE COOKIES/DO NOT SELL MY INFO
- ▶ ABOUT
- ▶ NEWS
- ▶ CONTACT
- ▶ SEARCH



Search



Series: [Advances in Biological Sciences Research](#)

Proceedings of the Conference on Natural Resources And Life Sciences 2022 (NRLS-BIO 2022)

[HOME](#)

[PREFACE](#)

[ARTICLES](#)

[AUTHORS](#)

[SESSIONS](#)

[ORGANIZERS](#)

[PUBLISHING INFORMATION](#)

Search

[+ Advanced search](#)

SEARCH

22 articles

Proceedings Article

Peer-Review Statements

Johan Sukweenadhi, Finna Setiawan

All of the articles in this proceedings volume have been presented at the 4th International Conference on Natural Resources and Life Sciences

(NRLS) 2022 during 24-26th August, 2023 in University of Surabaya (2 days online seminar and 1 day onsite workshop). These articles have been peer reviewed by...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

Systematic Literature Review: Side Effect of *Moringa oleifera* Lam.

Deborah Felinda Hari Wijaya, Alfian Hendra Krisnawan, Oeke Yunita, Krisyanti Budipramana, Karina Citra Rani, Nikmatul Ikhrom Eka Jayani

Moringa oleifera Lam. (Drumstick tree) is a herbal plant commonly found in subtropical and tropical areas such as Indonesia. *Moringa* plants have been widely used because they have many pharmacological effects. In addition, due to the high consumption of *Moringa*, there still needs to be monitoring regarding...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

Meta-Analysis on the effectiveness of the Use of Prophylactic Antibiotics in Preventing Surgical Site Infection in Patients with Hernia Surgery

Heru Wijono, Febryza Kristin, Beryl Bayanaka Agustha Peerera, Fauna Herawati, Rika Yulia

The administration of prophylactic antibiotics in hernia surgery patients is currently controversial, with varying opinions regarding their necessity. The European Hernia Society guideline does not recommend prophylactic antibiotics in patients with hernia surgery, while recent studies found that prophylactic...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

Analysis of Factors Affecting Adherence to Anti-Retroviral Treatment among HIV/AIDS Out-patients at the Dinoyo Public Health Centre

Annisa Lazuardy, Yosi Irawati Wibowo, Adji Prayitno Setiadi

There is an increasing prevalence of HIV/AIDS cases in Indonesia, particularly in Malang City, a tourist and educational destination. This study aimed to identify the characteristics of people living with HIV/AIDS (PLWHA) as well as to analyze their adherence to anti-retroviral (ARV) treatment and the...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

In Silico Evaluation of Soybean Protein as Bioactive Peptide for Anti-Thrombotic Agent by Molecular Docking Study

Fadilla Sherlyna, Muhammad Ilham Fahri, Arief Koeswanto, Dandan Wang

Cardiovascular Diseases (CVDs) are a growing global concern, including in Indonesia. By 2021, strokes were the leading cause of cardiovascular diseases (CVD), affecting 87% of the global population, while in 2018, 33.3% of Indonesians experienced CVD primarily due to strokes. Thrombolysis mediated by...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

Study of the Potential Use of Fermentation Methods to Increase Antioxidant and Antibacterial Activity of Fruit Peels: A Review

Merry Meryam Martgrita, Nehemia Roito Hutajulu, Hanna Gretty Manik, Adelina Manurung

Food industry waste, especially fruit peels, increases every year and their disposal can cause environmental pollution that can affect health. Fruit peels are known to contain antioxidant and antibacterial compounds, and their utilization can provide more value to fruit peel waste. This literature review...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

Estimation of Shelf-life of Porang Glucomannan Analog Rice By Accelerated Shelf-life Testing (ASLT) Method

Michael Surya Dharma, Yayon Pamula Mukti, Gisela Buschle-Diller, Ardhia Deasy Rosita Dewi

Analog rice is rice made from some or all non-rice ingredients and was created as a food diversification product so that it can be an alternative staple food to replace rice produced from rice. The shelf-life of food products can be determined using the ASLT method, which in principle accelerates the...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

Fortification of Dayak Onion Extract (*Eleutherine palmifolia* (L.) Merr.) to Jelly Drink as a Functional Food

Yayon Pamula Mukti, Berliana Yusup, Ardhia Deasy Rosita Dewi, Se Chan Kang

Indonesia is a nation characterized by a diverse array of plant species that possess notable health benefits. Among these botanical resources is the Dayak onion, scientifically known as *Eleutherine palmifolia* (L.) Merr. The Dayak Tribe – an indigenous people of Borneo's island has historically utilized...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

The Effect of *Bajakah Tampala* Stem (*Spatholobus littoralis* Hassk) Extract on Clotting Time in Vitro

Noza N. Moyananda, Rachmad P. Armanto, Mariana Wahjudi

Wounds that are not treated immediately will have a risk for the patient; the risk is infection. The risk of this infection can be reduced by stopping the bleeding as soon as possible. One way to accelerate the bleeding to stop is by giving haemostatic agents. One source of this haemostatic agent can...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

Analysis of Consumer Knowledge and Needs For Herbal Information

Oeke Yunita, Erlin Theterissa

Worldwide, the use of herbal medications is still growing quickly as more customers turn to these treatments for a variety of health issues. It is projected that the global market for Indonesian traditional medicinal products, of which more than half are herbal medicines, will grow. Unfortunately, there...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

Bst polymerase enhancement a bioinformatics approach to improve *Bst* polymerase characteristics

Jonathan, Ernest Suryadjaja, Sulistyo Emantoko D. Putra

DNA polymerase is a remarkably incredible invention in the biotechnology field. Since its discovery, molecular genetic-based research has been growing rapidly. Various methods for molecular-based diagnostics have been developed since. One of which is Loop-Mediated Isothermal Amplification; this method...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

Methylation Specific PCR (MSP): Nested PCR vs Unnested PCR

Farizky Martriano Humardani, Lisa Thalia Mulyanata, Lady Theresa Adeodata Tanaya, Risma Ikawaty, Heru Wijono, Hikmawan Wahyu Sulistomo, Dini Kesuma, Sulistyo Emantoko Dwi Putra

Methylation-specific PCR (MSP) is a valuable technique for studying DNA methylation patterns due to its straightforward design and implementation, high sensitivity in detecting methylated DNA, and ability to analyze large sample sizes cost-effectively rapidly. However, researchers need to be cautious...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

Natural Products Isolated from Various Parts of Mangosteen (*Garcinia mangostana* L.) as Therapeutic Agent: A Review

Arif Nur Muhammad Ansori, Yulanda Antonius, Ahmad Affan Ali Murtadlo, Viol Dhea Kharisma, Bayyinatul Muchtaromah, Muhammad Khaliim Jati Kusala, Dora Dayu Rahma Turista, Imam Rosadi, Vikash Jakhmola, Maksim Rebezov, Tarun Parashar, Rahadian Zainul

This review provides a comprehensive analysis of the therapeutic potential of natural products derived from mangosteen (*Garcinia mangostana* L.). Mangosteen, a tropical fruit native to Southeast Asia, has long been valued for its medicinal properties. The review focuses on the isolation and characterization...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

Identification of ACE1 Inhibitor Derived from Ashitaba's Chalcones: An *in Silico* Approach

Thomas Alessandro, Yulanda Antonius, Ardhia Deasy Rosita Dewi, Sin War Naw, Prita Ayu Kusumawardhany, Lanny Kusuma Widjaja, Hazrul Iswadi, Mariana Wahjudi

The angiotensin-converting enzyme, ACE1, is one of enzymes important to blood pressure modulation. The inhibition of protein responsible for blood pressure regulation, the angiotensin-converting enzyme, ACE1, is considered as a method to alleviate the hypertension condition. Ashitaba plant might be potent...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

Patchouli Alcohol Optimization from *Pogostemon cablin* Benth. cv. Sidikalang Leaves Using Response Surface Methodology

Mochammad Firmansyah, Feri Irwansyah, Krisyanti Budipramana, Mochammad Arbi Hadiyat, Ida Bagus Made Artadana, Popy Hartatie Hardjo

The demand for essential oils in the industrial sector continues to increase, proportional to the number of people using them. *Pogostemon cablin* popularly known as nilam in Indonesia produces patchouli oil with patchouli alcohol as the major compound. Patchouli oil has been used for a long time as perfume...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

Explant surface sterilization protocol for micropropagation of *Amorphophallus muelleri* Blume

Fenny Irawati, Agnes Natalia Wijaya, Anggi Manurung, Michael Anthony Thongiratama, Wina Dian Savitri, Popy Hartatie Hardjo

The success of tissue culture is greatly influenced by the explant surface sterilization technique. The presence of bacterial and fungi contamination, and the occurrence of browning on the explants can interfere with the process of culture propagation. High concentration of sterilant agents will inhibit...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

Utilization of Tempeh Extract as an Organic Supplement

Alternative for Banana Tissue Culture

Alexander Willy Dimaswarabrata, Anastasia Tatik Hartanti, Listya Utami Karmawan

The addition of organic materials to tissue culture media has been known to have a positive impact on plant growth. However, a tissue culture medium utilizing organic supplements originating from Indonesia as its specialty, such as tempeh, has not been discovered. This study aims to determine the effect...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

Anti-Inflammatory and Mucolytic Activity Test of Ethanol Extract Fennel Leaf (*Foeniculum vulgare* Mill.)

Syifatul Lutviani, Ita Nur Anisa, Andreanus A. Soemardji

Chronic Obstructive Pulmonary Disease (COPD) is a progressive lung disease characterized by chronic bronchitis, airway thickening, and emphysema. There are several main mechanisms of COPD, namely chronic inflammatory processes in the airways, oxidative stress, and disturbances in the balance between...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

LFER and 3D-QSAR Analysis of Febrifugine Derivatives against *Plasmodium falciparum* FCR-3 Strain

Nur Aina, Tegar Achsendo Yuniarta, Dini Kesuma

Malaria is a serious disease caused by Plasmodium through the bite of the female Anopheles mosquito. Due to resistance to artemisin, a first-line

antimalarial, new compounds are needed. This study aims to obtain a QSAR model from febrifugine derivatives against the Plasmodium falciparum FCR-3 strain....

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

Natural Dyes as Photosensitizers of *Propionibacterium acnes*

Asmiyenti Djaliasrin Djalil, Aqshal Pramudya Susanto, Rizal Nandha Arisugita, Binar Asrining Dhiani, Muhammad Faris Maulidan, Irfan Zamzani

The patient's quality of life may be negatively impacted by the high prevalence of acne vulgaris among adolescents. Acne vulgaris is a skin condition in which hair follicles become clogged with dead skin cells, bacteria, and natural facial oils. It has been demonstrated that acne antibiotics raise antibacterial...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

Formulation of Chewable Gummy Tablet of *Moringa oleifera* L. Leaf Extract Using Combination Kappa Carrageenan and Iota Carrageenan

Nabilaberty Prisma Gemilang, Nikmatul Ikhrom Eka Jayani, Karina Citra Rani

Moringa leaves were the most commonly used part of the Moringa plant because they were rich in nutrients. Moringa leaves extract was developed into a chewable gummy tablet to improve its acceptability. The main component of the chewable gummy tablet was a gelling agent. This study

aims to determine the...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Proceedings Article

Stability and antioxidant tests of ethanol extract liposome of moringa leaves (*Moringa oleifera*)

Robert Tungadi, Teti Sutriyati Tuloli, Sri Manovita Pateda

Moringa leaf potentially has an antioxidant effect because it contains Quercetin having porsolubility in water. Liposomes as carriers of drug compounds can increase the solubility of quercetin through an entrapment system in the lipid bilayer. This study aimed to determine the stability and antioxidant...

[+ Article details](#)

[+ Download article \(PDF\)](#)

Atlantis Press

Atlantis Press – now part of Springer Nature ¹ – is a professional publisher of scientific, technical & medical (STM) proceedings, journals and books. We offer world-class services, fast turnaround times and personalised communication. The proceedings and journals on our platform are Open Access and generate millions of downloads every month.

For more information, please contact us at: contact@atlantis-press.com

▶ PROCEEDINGS

▶ JOURNALS

▶ BOOKS

▶ POLICIES

▶ ABOUT

▶ NEWS

▶ CONTACT

▶ SEARCH

▶ [MANAGE COOKIES/DO NOT SELL MY](#)

[INFO](#)

[Home](#) [Privacy Policy](#) [Terms of use](#)   

Copyright © 2006-2024 Atlantis Press – now part of Springer Nature