

Cervical cancer screening in former sex workers in ex-localization area

DOI: 10.29063/ajrh2023/v27i2.4

Ardyan Wardhana*, Anita Dahliana and Katharina MA Angkawidjaja

Faculty of Medicine, Universitas Surabaya, Indonesia

*For Correspondence: Email: ardyanwardhana@staff.ubaya.ac.id; Phone: +62 82138433032

Abstract

Despite the lack of commercial sex activities as the localization has been shut down since 2014, former commercial sex workers (CSW) could be at risk for cervical cancer. Therefore, this study aimed to determine the prevalence of cervical cancer using early detection with Papanicolaou (Pap) smears in former CSW in this ex-localization area. This study was conducted with purposive sampling intended to be limited to former CSW and women who still live around this ex-localization area as a control group. This study included 76 women, 52.6% of whom were former CSW. Pap smear results showed no pre-cancerous lesions of the cervix in any of the participants. Comparisons between the CSW and control groups showed no differences in Pap smear results. Infection findings were found in 36.1% of CSW group. Early detection of human papillomavirus (HPV) DNA is necessary for identifying risk factors for cervical cancer in these populations. (*Afr J Reprod Health 2023; 27 [2]: 45-48*).

Keywords: Cervical cancer; early detection; pap smear; public health; women health

Résumé

Malgré le manque d'activités sexuelles commerciales, la localisation étant fermée depuis 2014, les anciennes travailleuses du sexe (TS) pourraient être à risque de cancer du col de l'utérus. Par conséquent, cette étude visait à déterminer la prévalence du cancer du col de l'utérus en utilisant la détection précoce avec des frottis de Papanicolaou (Pap) chez les anciens CSW dans cette ex-zone de localisation. Cette étude a été menée avec un échantillonnage raisonné destiné à être limité aux anciennes travailleuses du sexe et aux femmes qui vivent encore autour de cette ancienne zone de localisation en tant que groupe témoin. Cette étude a inclus 76 femmes, dont 52,6 % étaient d'anciennes travailleuses du sexe. Les résultats des frottis de Pap n'ont montré aucune lésion précancéreuse du col de l'utérus chez aucune des participantes. Les comparaisons entre les groupes de CSW et de contrôle n'ont montré aucune différence dans les résultats des tests de Pap. Des résultats d'infection ont été trouvés dans 36,1% du groupe CSW. La détection précoce de l'ADN du papillomavirus humain (HPV) est nécessaire pour identifier les facteurs de risque du cancer du col de l'utérus dans ces populations. (*Afr J Reprod Health 2023; 27 [2]: 45-48*).

Mots-clés: Cancer du col de l'utérus ; la détection précoce; frottis de pap ; santé publique; santé des femmes

Introduction

Localization, which was once one of the largest in Southeast Asia, has been shut down by the government since 2014. In addition to economic problems faced by former commercial sex workers (CSW) in this area, health problems cannot be ignored. Despite not engaging in commercial sex activities, former CSW could be at risk for cervical cancer. Cervical cancer is the most common gynecological cancer in developing countries¹. Global statistic reported that there were an estimated 565.541 new cases and 280.479 deaths because of cervical cancer in 2019².

Risk factors for cervical cancer include early sexual activity; having multiple partners; not

using condoms; history of chlamydia and human immunodeficiency virus infections; and contraceptive use³. In addition, smoking and high parity are also thought to be associated with the risk for cervical cancer⁴. Therefore, CSW require early screening for cervical cancer, namely using the Papanicolaou (Pap) smear test. This study aimed to determine the prevalence of cervical cancer based on early detection using Pap smears in former CSW in an ex-localization area.

Methods

The research was conducted in an ex-localization area in one of the cities in Indonesia after obtaining ethical approval from the Medical Research Ethics

Committee of Universitas Surabaya. The study was conducted with purposive sampling, which was intended to be limited to former CSW and adult women who still live around this ex-localization area as the control group. The reason for using purposive sampling was because they are hard to reach group of people. Information about this research was disseminated to the neighborhood/hamlet/urban village to collect data on former CSW in the area. Identifying former CSW was carried out together with village cadres to further persuade individuals to participate in research activities. Adult women living nearby were invited to participate in the control group.

This cross-sectional study aimed to identify the prevalence of cervical cancer using Pap smears and identify the risk of cervical cancer in adult women in the ex-localization area. The inclusion criterion was women aged 21-65 years who were still working in ex-localization areas. Women who had a history of pre-cancerous lesion abnormalities based on previous pap smear tests; have undergone hysterectomy, while menstruating; used tampons or local contraceptives 48 hours before the Pap test were excluded from the study.

All participants were recruited at predetermined times and locations. Before study participation, the participants received an oral explanation of the research procedure and provided their consent to participate in the study. Socio-demographic and health data, including age, marital status, number of pregnancies, smoking status, oral contraceptive drug use, and previous working history as CSW were collected.

Cervical samples were collected from the enrolled participants using a cytobrush and spatula. After fixing smears onto slides, conventional cytology was performed using 95% alcohol fixation. Pre-cancerous lesions based on pap smear results were reported using the Bethesda system.

Data are expressed as quantities and percentages; averages; and standard deviations. The primary outcome was the prevalence of cervical cancer lesions. The secondary result was a correlation between sociodemographic status and cervical cancer health. Data analysis was performed using the SPSS 24 software.

Results

In collaboration with village cadres, initially, former CSW (n = 120) who were still in the area were

identified. Since invitations and explanations about activities were carried out in an individual persuasion manner, they voluntarily wanted to participate in research activities. Adult women living nearby were invited to participate in the control group. On the day of activity, only 76 women participated in the research activity held: 52.6% of them were former CSW. One subject was excluded because she was pregnant, resulting in a total of 75 participants.

Their basic characteristics are listed in Table 1. The average age of the participants was 43.3±11.4 years. There was no difference between the basic characteristics in both groups, except for marital status ($p=0.03$) and smoking status ($p=0.02$). The CSW group had more widows and active smokers than did the control group. The age at first sexual intercourse in both groups was not significantly different ($p=0.20$).

Four subjects in each group refused to have a Pap smear procedure. Pap smear results showed no pre-cancerous lesions of the cervix in any of the participants (Table 2). Comparisons between the CSW and control groups showed no differences in Pap smear results. Infection findings based on Pap smears were found in 36.1% of the participants in the CSW group. The majority of the former CSW (41.7%) had non-specific inflammation (Table 3).

Discussion

This study was the first to report early detection of cervical cancer in former CSW in a localization that has been closed for a dozen years. Of note, in this study, pre-cancerous lesions were absent in the former CSW, and it was in contrast with other similar study. In a previous study in Belgium, CSW had a prevalence of pre-cancerous lesions of 38.2% of all ages and 23.6% in those aged over 35 years⁵. Meanwhile, a study in the Dominican Republic reported that 36.1% of CSW had precancerous lesion results⁶.

Risk factors for cervical cancer evaluated in this study included age; age at first sexual intercourse; and history of parity, smoking, and oral contraceptive use. The difference between the two groups was only found in smoking status. However, this study did not detect human papillomavirus (HPV) infection, which is a cause of cervical cancer. The incidence of cervical cancer has consistently been related to the prevalence of HPV infection in Indonesia⁶.

Table: Baseline characteristics

Parameter	Former-sex workers (n=40)	Control (n=35)	p
Age	42.9±8.2	44.3±14.0	0.61
Age first sex	19.6±4.0	21.2±5.0	0.20
Marital status	Single	2	0.03
	Still married	27	
	Divorced	2	
	Widow	4	
	Total	35	
Frequency Visit to health facility	> 8 times a year	3	0.961
	5-8 times a year	6	
	2-4 times a year	5	
	< 2 times a year	13	
	Total	27	
Smoking status	Active smoker	0	0.02
	Former smoker	3	
	Not smoker	30	
	Total	33	
Parity	0 pregnancy	10	0.27
	1-2 pregnancy	14	
	3-5 pregnancy	4	
	>5 pregnancy	4	
	Total	32	
Duration using oral contraception	Never	15	0.74
	< 5 years	8	
	5-10 years	3	
	> 10 years	3	
	Total	29	
Duration become sex worker	6.8±4.7 years		
Duration of non-active as sex worker	13.3±8.9 years		

Tables 2: Pap smear results

Pap Smear result	History becoming sex worker		p
	Yes	No	
No infection	23 (63.9%)	22 (71.0%)	0.608
Infection	12 (36.1%)	9 (29.0%)	
Abnormal	0 (0%)	0 (0%)	

Table 3: Pap smear diagnosis in former-sex worker group

Diagnosis	n (%)
Normal	8 (22.2%)
Non-specific inflammation	15 (41.7%)
Non-specific purulent	9 (25.0%)
Specific infection	4 (11.1%)

The prevalence rate of HPV infection at the age of 35-44 years in three cities in Indonesia was reported to be 18.4%, while the prevalence rate of high-risk HPV of all ages was 8.1%⁷. The incidence rate of high-risk HPV in CSW was reported to be as high as 41.7%, higher than that in the non-CSW population (19.8%)⁵. A meta-analysis reported a prevalence for high-risk HPV of 41.7% in the CSW population⁸. However, the low incidence of pre-

cervical cancerous lesions in the former CSW in this study may have been due to low HPV infections in the cities we studied. Therefore, early detection examination is necessary in subsequent studies. The relatively short duration of exposure as a sex worker and the relatively long duration of stopping engagement as a sex worker also may have played a role in the absence of pre-cancerous lesion findings in this study.

Conclusion

In conclusion, no pre-cancerous lesions were found in the former CSW in the ex-localization area. Further early detection of HPV DNA is necessary for the identification of risk factors for cervical cancer in these populations.

Acknowledgements

This work was performed as part of authors employment at the Faculty of Medicine, Universitas Surabaya, Indonesia. This work received internal funding from Universitas Surabaya. The authors

declare that the research was conducted in the absence of any commercial or financial relationships that could be considered as a potential conflict of interests.

Contribution of authors

AW conceived and designed the study as well as the person who wrote the final manuscript and gave supervision. AD collected and analyzed the data and the person who prepared the manuscript. MA collected the data and prepared the draft of manuscript. All authors mentioned in the article have approved the manuscript.

References

1. Benard VB, Thomas CC, King J, Massetti GM, Doria-Rose VP and Saraiya M; Centers for Disease Control and Prevention (CDC). Vital signs: cervical cancer incidence, mortality, and screening - United States, 2007-2012. *MMWR Morb Mortal Wkly Rep.* 2014; 63(44): 1004-1009.
2. Momenimovahed Z, Mazidimoradi A, Maroofi P, Allahqoli L, Salehiniya H and Alkatout I. Global, regional and national burden, incidence, and mortality of cervical cancer. *Cancer Rep (Hoboken).* 2022 Dec 21:e1756.
3. Tsikouras P, Zervoudis S, Manav B, Tomara E, Iatrakis G, Romanidis C, Bothou A and Galazios G. Cervical cancer: screening, diagnosis and staging. *J BUON Off J Balk Union Oncol.* 2016; 21(2): 320-325.
4. Chelimo C, Wouldes TA, Cameron LD and Elwood JM. Risk factors for and prevention of human papillomaviruses (HPV), genital warts and cervical cancer. *J Infect.* 2013; 66(3): 207-217.
5. Vorsters A, Cornelissen T, Leuridan E, Bogers J, Vanden Broeck D, Benoy I, Goossens H, Hens N and Van Damme P. Prevalence of high-risk human papillomavirus and abnormal pap smears in female sex workers compared to the general population in Antwerp, Belgium. *BMC Public Health.* 2016; 16(1): 477.
6. Richards SD, Stonbraker S, Halpern M and Amesty S. Cervical cancer screening among transactional female sex workers in the Dominican Republic. *Int J STD AIDS.* 2018; 29(12): 1204-1214.
7. Vet JN, de Boer MA, van den Akker BE, Siregar B, Lisnawati, Budiningsih S, Tyasmorowati D, Moestikaningsih, Cornain S, Peters AA and Fleuren GJ. Prevalence of human papillomavirus in Indonesia: a population-based study in three regions. *Br J Cancer.* 2008; 99(1): 214-218.
8. Farahmand M, Moghoofei M, Dorost A, Abbasi S, Monavari SH, Kiani SJ and Tavakoli A. Prevalence and genotype distribution of genital human papillomavirus infection in female sex workers in the world: a systematic review and meta-analysis. *BMC Public Health.* 2020; 20(1): 1455.



African Journal of Reproductive Health

Revue Africaine de Sante de la Reproduction

AJRH



[Home](#)

[About AJRH](#)

[Login](#)

[Register](#)

[Search](#)

[Browse](#)

[WHARC Website](#)

Go

STAFF AND EDITORIAL BOARD

MEMBERS

Editor-in-Chief

Friday Okonofua Nigeria

Editor

Akhere Omonkhua Nigeria

Deputy Editor

Joseph Balogun USA

Editor (French)

Cyril Mokwenye Nigeria

Assistant Editors

Clifford Odimegwu Nigeria

Judith Ann-Walker Nigeria

Loretta Ntoimo Nigeria

Babatunde Ahonsi Nigeria

Albrecht Jahn Germany

Phyllis Kanki USA

Annette Kapaun Germany

Saidi Kapiga Tanzania

Joan Kaufman China

Peju Olukoya Switzerland

Mere Kissekka Ethiopia

Managing Editor

Karl Eromosele Eimuhi Nigeria

O.A Ladipo UK

Ulla Larsen USA

Adetokunbo Lucas Nigeria

Editorial Assistant

Judith Otesiri Obaseki Nigeria

Gernard Msamanga Tanzania

Osato Giwa Osagie Nigeria

Michael Mbizo Switzerland

<i>Computer Typesetter</i>		Ester Mwaikambo	Tanzania
Judith Otesiri Obaseki	Nigeria	Fredrick Naftolin	USA
		Carla Obermeyer	USA
<i>Subscription/Marketing Officer</i>		Grace Wyshak	USA
Joy Odion Nomamiukor	Nigeria	Michael Reich	USA
		Khama Rogo	Kenya
<i>Cover Design</i>		A. Orubuloye	Nigeria
Shereen Siddiqui	New York, USA	Andrzej Kulczycki	USA
		Frank Van Balen	Netherlands
<i>Editorial Advisory Board</i>		Kesley Harrison	Finland
Rachel Snow	USA	Joseph Otubu	Nigeria
Alayne Adams	USA	William Pick	South Africa
Andrew Onokerhorhay	Nigeria	Helen Rees	South Africa
Lawrence Adeokun	Nigeria	John Caldwell	Australia
Simi Afonja	Nigeria	Sarah Castle	USA
Wole Akande	Nigeria	Kole Shettima	Nigeria
Nimi Briggs	Nigeria	Igbal Shah	Switzerland
Pitt Reitmaier	Germany	Richard Turkson	Ghana
Lincoln Chen	USA	Kim Dickson-Tetteh	South Africa
John Cleland	United Kingdom	Staffan Bergstrom	Sweden

Sylvia Deganus	Ghana	Mags Beksinska	South Africa
Ana Langer	USA	Lindsay Edouard	Mauritius
Olufemi Olatunbosun	Canada	Dozie Ikedife	Nigeria
Peter Okebukola	Nigeria	Kunle Odunsi	USA
Allan Hill	USA	Andre Lalonde	Canada
Margaret Hoffman	South Africa		

Table of Contents

Articles


The politics of abortion rights in the 2022 United States midterm election: Lessons for fledgling democracies around the world	PDF
Joseph A. Balogun, Friday E. Okonofua	
A retrospective evaluation of pregnant adolescents presenting to an emergency department	PDF
Raziye Kılıççeker, Mine Yurdakul	
Teachers' reflections on menstrual management among urban and rural schoolgirls in South Africa	PDF
Thelma Fennie, Mokgadi Moletsane, Anita Padmanabhanunni	
Cervical cancer screening in former sex workers in ex-localization area	PDF
Ardyan Wardhana, Anita Dahliana, Katharina MA Angkawidjaja	
In-school adolescents' perception of sexuality education in region 1 of the Gambia	PDF
Phebian I.G. Sagnia, Momodou Jeng	


<p>Bayesian modeling of maternal mortality in Ghana</p> <p>Dioggban Jakperik, Saralees Nadarajah, Michael J. Adjabui</p>	<p>PDF</p>
<p>Factors influencing contraceptive use among women of reproductive age in plantation farming communities in South-South Nigeria</p> <p>Veronica A. Undelikwo, Ntongha E. Ikpi, Glory E. Bassey</p>	<p>PDF</p>
<p>Patriarchal norms, partner pronatalism, and women's fertility intentions in Ghana</p> <p>Nkechi S. Owoo</p>	<p>PDF</p>
<p>The association between the prevalence of HIV infection and ectopic pregnancy</p> <p>Mokoena M. Mohosho</p>	<p>PDF</p>
<p>The educational quality perception, professional interest, and career choices among hospitality postgraduate students</p> <p>Wenhua Xu, Gang Han</p>	<p>PDF</p>
<p>Mental health needs among pregnant and parenting adolescent girls and young women in South Africa: A scoping review</p> <p>Linneth N. Mabila, Livhuwani Muthelo, Masenyani O. Mbombi, Peter M. Mphekgwana, Reneilwe G. Mashaba, Cairo B. Ntimana, Kagiso P. Seakamela, Joseph Tlouyamma, Inos Dhau, Rathani Nemuramba, Katlego Mothapo, Eric Maimela</p>	<p>PDF</p>

African journal of reproductive health

COUNTRY

Nigeria

 Universities and research institutions in Nigeria

 Media Ranking in Nigeria

SUBJECT AREA AND CATEGORY

Medicine

Obstetrics and Gynecology
Public Health, Environmental and Occupational Health
Reproductive Medicine

PUBLISHER

Women's Health and Action Research Centre

H-INDEX

44

PUBLICATION TYPE

Journals

ISSN

11184841

COVERAGE

1997-1998, 2000-2021

INFORMATION


[Homepage](#)

[How to publish in this journal](#)

africanjournalofreproductiveH@gmail.com

SCOPE

African Journal of Reproductive Health is a multidisciplinary and international journal which started in 1997 and published tri-annually (April, August and December) up to December 2008, quarterly from 2009 to 2013, and now quarterly (March, June, September and December) with a Special Edition that comes up any time of the year. The journal focuses on publishing original research, comprehensive review articles, case reports and commentaries that on reproductive health and related in Africa issues in both English and French. It strives to provide a forum for both African and foreign authors working in Africa to share findings in all aspects of reproductive health and also to disseminate innovative, relevant and useful information on reproductive health throughout the continent.

 Join the conversation about this journal

 Quartiles

FIND SIMILAR JOURNALS 

options 

- | | | | | |
|--|--|---|---|--|
| <p>1
Reproductive Health
GBR</p> <p>74%
similarity</p> | <p>2
International Perspectives on Sexual and Reproductive
USA</p> <p>58%
similarity</p> | <p>3
BMC Women's Health
GBR</p> <p>54%
similarity</p> | <p>4
BMC Pregnancy and Childbirth
GBR</p> <p>52%
similarity</p> | <p>5
Health Care for Women International
GBR</p> <p>49%
similarity</p> |
|--|--|---|---|--|



African journal of reproductive health

Obstetrics and Gynecology

Q3 best quartile

SJR 2021 0.36

powered by scimagojr.com

Show this widget in your own website

Just copy the code below and paste within your html code:

```
<a href="https://www.scimagojr.com" >
```

SCImago Graphica

Explore, visually communicate and make sense of data with our new data visualization tool.

Metrics based on Scopus® data as of April 2022

Kitti 1 year ago

Does the Scopus database cover this journal in 2022?

reply

Melanie Ortiz 1 year ago

Dear Kitti,

Thank you very much for your comment.

All the metadata have been provided by Scopus /Elsevier in their last update sent to SCImago, including the Coverage's period data. The SJR for 2020 was released on 17 May 2021. We suggest you consult the Scopus database directly to see the current index status as SJR is a static image of Scopus, which is changing every day.

Best Regards, SCImago Team

Widia shofa ilmiah 3 years ago

How long this journal coverage on the scopus. Thankyou

reply

Melanie Ortiz 3 years ago

Dear Widia,

Thank you very much for your comment.

All the metadata have been provided by Scopus /Elsevier in their last update sent to SCImago, including the Coverage's period data. The SJR for 2019 was updated on June 2020, 11. We suggest you consult the Scopus database directly to see the current index

status as SJR is a static image of Scopus, which is changing every day.
Best Regards, SCImago Team

Y **Yessy** 3 years ago

Dear SCImago team,

Can i get information about wether African journal of reproductive health still indexed by scopus in Juli 2020 until the end of the year 2020?

Thanks

reply

M **Mounir Ouzir** 2 years ago

African journal of reproductive health
Scopus coverage years:from 1997 to 1998, from 2000 to Present



Melanie Ortiz 2 years ago

SCImago Team

Dear Mounir,

Thank you very much for your comment.

All the metadata have been provided by Scopus /Elsevier in their last update sent to SCImago, including the Coverage's period data (check above). The SJR for 2019 was released on 11 June 2020. We suggest you consult the Scopus database directly to see the current index status as SJR is a static image of Scopus, which is changing every day.

Best Regards, SCImago Team



Melanie Ortiz 3 years ago

SCImago Team

Dear Yessy,

Thank you very much for your comment.

All the metadata have been provided by Scopus /Elsevier in their last update sent to SCImago, including the Coverage's period data. The SJR for 2019 was released on June 2020, 11. We suggest you consult the Scopus database directly to see the current index status as SJR is a static image of Scopus, which is changing every day.

For further information, please contact Scopus support: https://service.elsevier.com/app/answers/detail/a_id/14883/kw/scimago/supporthub/scopus/

Best Regards, SCImago Team

Leave a comment

Name

Email

(will not be published)

Submit

The users of Scimago Journal & Country Rank have the possibility to dialogue through comments linked to a specific journal. The purpose is to have a forum in which general doubts about the processes of publication in the journal, experiences and other issues derived from the publication of papers are resolved. For topics on particular articles, maintain the dialogue through the usual channels with your editor.

Developed by:



Powered by:



Follow us on @ScimagoJR

Scimago Lab, Copyright 2007-2022. Data Source: Scopus®

EST MODUS IN REBUS

© Scimago (Scopus) 1.1.100

[Edit Cookie Consent](#)

Source details

African journal of reproductive health

Scopus coverage years: from 1997 to 1998, from 2000 to 2022

Publisher: Women's Health and Action Research Centre

ISSN: 1118-4841

Subject area: [Medicine: Obstetrics and Gynecology](#) [Medicine: Reproductive Medicine](#)

[Medicine: Public Health, Environmental and Occupational Health](#)

Source type: Journal

[View all documents >](#) [Set document alert](#) [Save to source list](#) [Source Homepage](#)

CiteScore 2021
1.3 ⓘ

SJR 2021
0.360 ⓘ

SNIP 2021
0.648 ⓘ

[CiteScore](#) [CiteScore rank & trend](#) [Scopus content coverage](#)

CiteScore **2021** ▼
1.3 = $\frac{354 \text{ Citations 2018 - 2021}}{275 \text{ Documents 2018 - 2021}}$

Calculated on 05 May, 2022

CiteScoreTracker 2022 ⓘ
1.2 = $\frac{454 \text{ Citations to date}}{382 \text{ Documents to date}}$

Last updated on 05 March, 2023 • Updated monthly

CiteScore rank 2021 ⓘ

Category	Rank	Percentile
Medicine		
↳ Obstetrics and Gynecology	#121/188	35th
Medicine		
↳ Reproductive Medicine	#54/76	29th
Medicine		

[View CiteScore methodology >](#) [CiteScore FAQ >](#) [Add CiteScore to your site ↗](#)

About Scopus

[What is Scopus](#)

[Content coverage](#)

[Scopus blog](#)

[Scopus API](#)

[Privacy matters](#)

Language

[日本語版を表示する](#)

[查看简体中文版本](#)

[查看繁體中文版本](#)

[Просмотр версии на русском языке](#)

Customer Service

[Help](#)

[Tutorials](#)

[Contact us](#)

ELSEVIER

[Terms and conditions ↗](#) [Privacy policy ↗](#)

Copyright © Elsevier B.V. ↗. All rights reserved. Scopus® is a registered trademark of Elsevier B.V.

We use cookies to help provide and enhance our service and tailor content. By continuing, you agree to the use of cookies ↗.

