



System optimization and validation to improve thin-layer chromatography of roselle calyces (*Hibiscus sabdariffa* L.) required by the Indonesian Herbal Pharmacopoeia Edition II

[Optimización y validación del sistema para mejorar la cromatografía en capa fina de cálices de rosa mosqueta (*Hibiscus sabdariffa* L.) requerida por la Farmacopea Herbal Indonesia Edición II]

Bhujangga Agung Ayu Sri Kartika Dewi, Kartini Kartini*

Department of Pharmaceutical Biology, Faculty of Pharmacy, University of Surabaya, Surabaya, Indonesia.

*E-mail: kartini@staff.ubaya.ac.id

Abstract

Context: Roselle (*Hibiscus sabdariffa* L.) is one of the traditional crude drugs monographed in the Indonesian Herbal Pharmacopoeia Edition II (IHP II). As per IHP II requirements, a TLC pattern should be used in its authentication. However, the TLC system described for this purpose does not produce a clear reference chromatogram, often leading to inconclusive results.

Aims: To develop and validate TLC systems of *H. sabdariffa* calyces for a better-quality chromatogram than those listed in the IHP II, thereby facilitating crude drugs authentication.

Methods: To optimize TLC for *H. sabdariffa*, sixteen systems, differing in stationary phase, mobile phase composition, and/or visualization reagent were tested. The TLC system was then validated using several parameters such as analyte stability during chromatography, analyte stability in the extract solution and on the TLC plate, stability of the derivatization result, and precision on a plate as well as intermediate precision.

Results: Two systems (I and II) were successfully designed and applied to evaluate *H. sabdariffa* quality. System I used Si gel 60 F₂₅₄ for the stationary phase, ethyl acetate-formic acid-glacial acetic acid-water (100:11:11:2) for the mobile phase, *H. sabdariffa*'s ethanol extract (5%, 20 µL) for the test solution, cyanidin 3-O-glucoside (100 ppm, 4 µL) as a reference, no derivatization, and detection with visible light. System II combined Si gel 60 F₂₅₄ for the stationary phase, ethyl acetate-formic acid-glacial acetic acid-toluene-water (80:11:11:20:19), *H. sabdariffa*'s ethanol extract (5%, 20 µL) for the test solution, chlorogenic acid (1000 ppm, 2 µL) and caffeic acid (50 ppm, 2 µL) as references, the visualizing reagent NP/PEG, and investigation under 366 nm UV light.

Conclusions: Both systems are simple but have good stability and precision, thus facilitating *H. sabdariffa* calyx authentication and paving the way for developing content analysis methods for *H. sabdariffa* markers.

Keywords: authenticity; herbal pharmacopoeia; quality evaluation; roselle; TLC pattern.

Resumen

Contexto: La rosa mosqueta (*Hibiscus sabdariffa* L.) es una de las drogas crudas tradicionales monografiadas en la segunda edición de la Farmacopea Herbal Indonesia (PHI II). Según los requisitos de la PHI II, se debe utilizar un patrón de TLC para su autenticación. Sin embargo, el sistema de TLC descrito para este fin no produce un cromatograma de referencia claro, lo que a menudo conduce a resultados no concluyentes.

Objetivos: Desarrollar y validar sistemas de TLC de cálices de *H. sabdariffa* para obtener un cromatograma de mejor calidad que los listados en el PHI II, facilitando así la autenticación de drogas crudas.

Métodos: Para optimizar la TLC para *H. sabdariffa*, se probaron dieciséis sistemas, que diferían en la fase estacionaria, la composición de la fase móvil y/o el reactivo de visualización. A continuación, se validó el sistema de TLC utilizando varios parámetros, como la estabilidad del analito durante la cromatografía, la estabilidad del analito en la solución del extracto y en la placa de TLC, la estabilidad del resultado de la derivatización y la precisión en una placa, así como la precisión intermedia.

Resultados: Se diseñaron y aplicaron con éxito dos sistemas (I y II) para evaluar la calidad de *H. sabdariffa*. El sistema I utilizó Si gel 60 F₂₅₄ como fase estacionaria, acetato de etilo-ácido fórmico-ácido acético glacial-agua (100:11:11:2) como fase móvil, extracto de etanol de *H. sabdariffa* (5%, 20 µL) como solución de prueba, cianidina 3-O-glucósido (100 ppm, 4 µL) como referencia, sin derivatización, y detección con luz visible. El sistema II combinó Si gel 60 F₂₅₄ para la fase estacionaria, acetato de etilo-ácido fórmico-ácido acético glacial-tolueno-agua (80:11:11:20:19), extracto de etanol de *H. sabdariffa* (5%, 20 µL) para la solución de prueba, ácido clorogénico (1000 ppm, 2 µL) y ácido cafeico (50 ppm, 2 µL) como referencias, el reactivo de visualización NP/PEG, y la investigación bajo luz UV de 366 nm.

Conclusiones: Ambos sistemas son sencillos pero tienen buena estabilidad y precisión, facilitando así la autenticación del cáliz de *H. sabdariffa* y allanando el camino para desarrollar métodos de análisis de contenido para marcadores de *H. sabdariffa*.

Palabras Clave: autenticidad; farmacopea herbal; evaluación de la calidad; rosa mosqueta; patrón TLC.

ARTICLE INFO

Received: December 20, 2022.

Accepted: February 25, 2023.

Available Online: March 8, 2023.

AUTHOR INFO

ORCID: [0000-0002-7685-9933](https://orcid.org/0000-0002-7685-9933) (KK)

INTRODUCTION

Roselle (*Hibiscus sabdariffa* L., family *Malvaceae*) is a flowering plant ubiquitously used as a medicinal and food plant in many countries in Central Africa, West Africa, South Africa, and Southeast Asia, including Indonesia, Thailand, Malaysia, and Vietnam (Ali et al., 2005; Antonia et al., 2019; Chumsri et al., 2008; Pham et al., 2019). The calyx contains various beneficial therapeutic compounds, especially organic acid, phenolic acid, flavonoid, and polysaccharide groups, making it the plant's most important part (Hapsari and Setyaningsih, 2021; Izquierdo-Vega et al., 2020). Anthocyanins are flavonoids responsible for the red sepals, and organic acids (citric acid, malic acid, tartaric acid, and hibiscus acid) for the sour taste. In addition to these typical organoleptic features, the compounds are known to exhibit numerous biological activities; for instance, antioxidative, antipyretic, antinociceptive, anti-inflammatory, antibacterial, antifungal, antiparasitic, anticancer, antidiabetic, antihypertensive, anti-cholesterol, and hepatoprotective (Ali et al., 2005; Izquierdo-Vega et al., 2020).

H. sabdariffa var. *Sabdariffa* and *H. sabdariffa* var. *Altissimac Wester* are the two most traded varieties. Their varied phenotypes and genotypes have also been reported (Antonia et al., 2019; Hapsari and Setyaningsih, 2021; Sukkhaeng et al., 2018; Torres-Morán et al., 2011). In Indonesia alone, there are two known variants: red roselle and purple roselle (Aryanti et al., 2019). As an ingredient in food, beverages, and traditional medicine, dried *H. sabdariffa* is traded in at least three forms: whole calyx, flakes, and powder. The dried calyx has a purplish to blackish-red color.

As part of traditional remedies, *H. sabdariffa* is monographed in several herbal pharmacopoeias, including Indonesian Herbal Pharmacopoeia Edition II (IHP II), Malaysian Herbal Monograph, and European Pharmacopoeia 7.0 (Committee MHM, 2015; EDQM, 2008; Health IMO, 2017). The monographs contain botanical characteristics to be used in the authentication of a suspected *H. sabdariffa* specimen and quality control. Aside from organoleptic and microscopic identifications, IHP II also includes a TLC system and its associated pattern for a more sensitive authentication method. In this case, users can compare their results with the provided pattern to determine similarities (or differences) in the number, color, and Rf or Rx values of spots produced by a particular TLC system. Therefore, the reference chromatogram should have a clear pattern and separation. This is, however, not seen in the one used for *H. sabdariffa* authentication in IHP II (Fig. 1), where the five differ-

ent compound spots show indistinct colors and are separated poorly—hindering its use as a reference in related analyses.

For the above reasons, the research was intended to develop and validate TLC systems for a better-quality chromatogram, where the compound spots are not overlapping and streaking but distinguishable (their amount and color are easily identifiable). This research is essential in fostering the use of *H. sabdariffa* as a traditional crude drug with standardized quality and, thus, maintaining its consistent efficacy and safety.

MATERIAL AND METHODS

Chemicals


Chemicals used were TLC silica gel 60 F₂₅₄ plates, TLC cellulose plates, n-hexane, toluene, ethyl acetate, chloroform, n-butanol, acetone, acetic acid, formic acid, ethanol, methanol, and polyethylene glycol (PEG); all from Merck KGaA (Darmstadt, Germany). In addition, the analyses also used 2-aminoethyl diphenylborinate, quercetin, gallic acid, rutin, chlorogenic acid, caffeic acid, and p-coumaric acid from Sigma Aldrich Co. (St. Louis, MO, USA). Cyanidin 3-O-glucoside was obtained from PhytoLab GmbH & Co. KG (Vestenbergsgreuth, Germany), and demineralized water was from the Pharmaceutical Chemistry Laboratory, Faculty of Pharmacy, University of Surabaya, Indonesia.

Plant material

The plant material used in this research was purple *H. sabdariffa* calyces (Fig. 2) harvested in Magetan Regency, Jawa Timur Province, Indonesia, in April 2021 (7°37'22"S; 111°17'4"E). Botanical identification was conducted by the Center for Traditional Drug Information and Development, the University of Surabaya, with the plant authentication certificate No. 1476/D.T/VI/2022. Whole dried *H. sabdariffa* calyces were ground in a blender (Philips HR 2222) and sifted through a 20 mesh sieve. The powder was then stored in a dark, airtight container until further use.

Extract preparation

The method described in IHP II was adopted to extract the calyces. First, calyx powder weighing 0.5 g was added with 10 mL of 95% ethanol, then shaken in a water bath (Mettler GmbH + Co. KG, Schwabach, Germany) for ten minutes at 60°C to improve the extraction process. Afterward, the derived extract was passed through a sieve into a volumetric flask and



poured with demineralized water until the volume was 10.0 mL.

TLC system optimization

To optimize TLC for *H. sabdariffa*, sixteen systems differing in stationary phase, mobile phase composition, and/or visualization reagent derived from the literature study were tested. The two stationary phases used in this optimization stage were silica gel 60 F₂₅₄ and cellulose. The sample and reference solution were each dotted onto the plate 1.5 cm from the bottom, 1 cm from the right and left side, and 1 cm from the adjacent spots. Dotting or spotting was conducted manually using a 2 µL capillary pipe (CAMAG, Muttenz, Switzerland). Elution was allowed until 8 cm from the lower edge of the plate in a twin-trough chamber (CAMAG, Switzerland). The visualizing agents included FeCl₃ reagent, ammonia vapor, and NP/PEG reagent. After selecting the optimal mobile phase, stationary phase, and visualization reagent, the optimization stage continued with testing the volume of the spotted sample and the reference solution type and volume. The reference solutions optimized at this stage were quercetin, cyanidin 3-O-glucoside, gallic acid, rutin, chlorogenic acid, caffeic acid, p-coumaric acid, and vanillic acid. All tests were conducted in three replicates.

TLC system validation

The TLC system for *H. sabdariffa* was validated using several parameters: analyte stability during chromatography, analyte stability in the extract solution and on the TLC plate, stability of the derivatization result, and precision on a plate as well as intermediate

precision. In the validation process, samples and reference solutions were spotted using a CAMAG 100 µL-sample syringe (Hamilton, Switzerland) automatically with a Linomat 5 applicator (CAMAG, Switzerland) under N₂ gas flow.

Analyte stability during chromatography

A sample solution was applied to the lower-right side of a 10 × 10 cm plate in the form of a spot. The plate was eluted until the mobile phase reached about 8 cm from the start line, then lifted and allowed to dry. Next, it was rotated 90° clockwise, placed in the chamber, and re-eluted with a new mobile phase. After drying, it was derivatized with NP/PEG reagent and documented under 366 nm UV light. An analyte is deemed stable if the compound spots form a diagonal line connecting the spotting position with the intersection of the two mobile-phase fronts (Reich and Schibli, 2007). Analysis was performed in three replicates on three different TLC plates.

Analyte stability in the extract solution and on the plate

On a TLC plate, the *H. sabdariffa* calyx extract was deposited on lanes 1 and 2. The extract and the recently spotted plate were left to stand for three hours. Then, the 3-hour-old extract was applied to lanes 5 and 6. A freshly prepared extract was also spotted onto lanes 3 and 4 right before elution with the optimized system. The chromatogram before and after derivatization using NP/PEG reagent was documented with a TLC visualizer (CAMAG, Muttenz, Switzerland). This process also generates other data, including R_f value, peak height, and peak area of each compound spot appearing on the chromatogram. Of all

Table 1. Optimal thin-layer chromatography systems for *H. sabdariffa*.

System	Stationary phase	Mobile phase	Test solution	Reference solution	Derivatization*	Detection
I	Si gel 60 F ₂₅₄	Ethyl acetate-formic acid-glacial acetic acid-water (100:11:11:2)	Ethanol extract of roselle calyx (5%, 20 µL)	Cyanidin 3-O-glucoside (100 ppm, 4 µL)	N/A	Visible light
II	Si gel 60 F ₂₅₄	Ethyl acetate-formic acid-glacial acetic acid-toluene-water (80:11:11:20:19)	Ethanol extract of roselle calyx (5%, 20 µL)	Chlorogenic acid (1000 ppm, 2 µL) Caffeic acid (50 ppm, 2 µL)	NP/PEG	366 nm UV light

*N/A: not applicable

the spots, one to a few were selected as probable markers, and the average, standard deviation, and relative standard deviation (RSD) of their peak heights were calculated. RSD was calculated from the marker spots on lanes 1, 2, 3, and 4 to determine analyte stability on the plate and from those on lanes 3, 4, 5, and 6 to assess the stability in the extract solution (Reich and Schibli, 2007). Analysis was performed in three replicates on three different TLC plates.

Precision on a plate and intermediate precision

The *H. sabdariffa* extract solution was deposited on a plate in five replicates for the precision on a plate test and on three consecutive days for the intermediate precision test. Each plate was eluted with the optimized mobile phase. Then, the chromatogram before and after derivatization with a visualizing reagent was documented using a TLC visualizer. Rf values were then compared and peak heights of the marker spots were used to calculate their average, standard deviation, and RSD. RSD of the marker spots from one plate was calculated to determine precision on a plate, while the one from the different-day application was for intermediate precision (Spangenberg et al., 2011).

TLC pattern generation for *H. sabdariffa*

The validated TLC systems for *H. sabdariffa* calyx were each used to create a TLC pattern. It was then compared with the pattern depicted in IHP II. Observations were made to see similarities or differences in these parameters: the number, color, Rf or Rx value, and separation of the spots.

Statistical analysis

All data were reported as mean $\pm$ SD ($n = 3-5$), then the relative standard deviations (RSD) were calculated and analyzed using descriptive statistics to determine whether the developed method satisfies the validation parameters.

RESULTS AND DISCUSSION

Optimized TLC system

Two of the 16 TLC systems tested in the research are deemed suitable for creating a clear chromatogram for *H. sabdariffa* calyces, namely systems I and II (Table 1). Figs. 3 and 4 show the chromatograms derived from the two systems.

Under visible light, the *H. sabdariffa* calyx extract tested with the system I was separated into two red spots suspected as anthocyanins (see spots a and b in Fig. 3A). While they shared a similar red chroma with cyanidin 3-O-glucoside, the Rf values differed. Therefore, both spots were proposed as markers for system I and referred to as 'anthocyanin 1' and 'anthocyanin 2'. These compounds could be delphinidin-3-glucoside, delphinidin-3-sambubioside, or cyanidin-3-sambubioside, which are also red to purple in color (Santos et al., 2013; Committee MHM, 2015; Lago et al., 2014). The system I produced a better TLC pattern than IHP II, where only one anthocyanin spot is not separated from other compound spots (Fig. 1). Fig. 3A also shows that 20 µL of extract (lane 2) and 4 µL of reference solution (lane 5) produced a good pattern. For these reasons, it is recommended that the *H. sabdariffa*'s chromatogram be made with system I, combining Si gel 60 F₂₅₄ as the stationary phase, ethyl acetate-formic acid-glacial acetic acid-water (100:11:11:2) as the mobile phase, *H. sabdariffa*'s ethanol extract (5%, 20 µL) as the test solution, cyanidin 3-O-glucoside (100 ppm, 4 µL) as the reference solution, no derivatization, and investigation under visible light.

In addition to anthocyanins, *H. sabdariffa* is a good source of flavonoids from other subclasses like quercetin-3-glucoside, methyl epigallocatechin, myricetin, quercetin, and kaempferol. No less than 95 flavonoids have been discerned from the calyces. Besides, the plant is rich in other phenolic compounds, such as neochlorogenic acid, chlorogenic acid, cryptochloro-

genic acid, methyl chlorogenate, coumaroylquinic acid, dihydroferulic acid-4-O-glucuronide, ethyl chlorogenate, and 5-O-caffeoyl shikimic acid (Hapsari and Setyaningsih, 2021). Therefore, this research also experimented with reference solutions other than the one used in IHP II (cyanidin 3-O-glucoside), namely flavonoids (quercetin and rutin) and phenolic acid (gallic acid, chlorogenic acid, caffeic acid, p-coumaric acid, and vanillic acid). From the varying TLC systems tested for optimization, chlorogenic acid (Fig. 4C, spot a) and caffeic acid (Fig. 4C, spot b) were readily detectable, while other compounds were not. To ensure that spots a and b (Fig. 4C) were chlorogenic acid and caffeic acid, their UV spectrum was scanned and compared with their respective reference compounds. Fig. 5A-B shows an overlap between the former and the latter with a maximum wavelength of 320-330 nm, indicating that spots a and b are indeed chlorogenic acid and caffeic acid (Kim et al., 2015; Orfali et al., 2021). Therefore, both acids can be used as alternative analytical markers for *H. sabdariffa* quality analysis, especially based on TLC patterns.


Chlorogenic and caffeic acids are the two most common phenolic acids found in *H. sabdariffa*, which also possess anti-diabetic and anti-hyperlipidemic properties. Previous research confirmed the presence of chlorogenic acid (67.12 mg/g) and caffeic acid (15.38 mg/g) in *H. sabdariffa* extract and identified their neuroprotective activity by testing it against acetylcholinesterase (AChE), butyrylcholinesterase

(BChE), monoamine oxidase (MAO), and ecto-5'-nucleotidase (E-NTDase) (Hapsari and Setyaningsih, 2021). Furthermore, because of their high contents and biological activities, chlorogenic and caffeic acids are suitable as not only analytical but also active markers for *H. sabdariffa* calyces (Chan et al., 2007).

Fig. 4C also shows that 2 μ L of chlorogenic acid (lane 1), 2 μ L of caffeic acid (lane 4), and 20 μ L of the extract (lane 8) produced interpretable chromatograms. Thus, aside from system I, it is also suggested that the *H. sabdariffa*'s TLC pattern be made using system II, consisting of Si gel 60 F₂₅₄ as the stationary phase, ethyl acetate-formic acid-glacial acetic acid-toluene-water (80:11:11:20:19) as the mobile phase, *H. sabdariffa*'s ethanol extract (5%, 20 μ L) as the test solution, chlorogenic acid (1000 ppm, 2 μ L) and caffeic acid (50 ppm, 2 μ L) as the reference solutions, derivatization with the visualizing reagent NP/PEG, and detection with 366 nm UV light.

Analyte stability during chromatography

To validate the optimized systems I and II, analyte stability in the extract during chromatography was tested with a two-dimensional elution. Fig. 6A-B shows that both systems produced good analyte stability. This is evident in the plate's diagonal line in the elution results (Kartini et al., 2021; Reich and Schibli, 2007).






Figure 4. TLC patterns of *H. sabdariffa* calyx with system II observed under visible light (A), 254 nm UV light (B), and 366 nm UV light after derivatization with NP/PEG (C).

Different lanes from left to right: 2, 4, 6 µL of chlorogenic acid (1000 ppm), 2, 4, 6 µL of caffeic acid (50 ppm), and 10, 20, 30 µL of the purple *H. sabdariffa*.




Figure 5. Spectral comparison between chlorogenic acid standard and sample (A) and caffeic acid standard and sample (B).




Figure 6. Analyte stability during chromatography to validate TLC system I, observed under visible light (A), and system II under 366 nm UV light (B).




Table 2. Peak heights of the marker compounds obtained from testing the analyte stability in the extract solution and on the plate.

Marker	Lanes 1–2	Lanes 3–4	Lanes 5–6	RSD (%) from lanes 1–4	RSD (%) from lanes 3–6
Anthocyanin 1	1632.1	1444.5	1247.6	12.0	9.2
Anthocyanin 2	1191.4	883.8	793.1	5.5	6.4
Chlorogenic acid	1632.1	1566.9	1421.2	11.5	8.1
Caffeic acid	4143.8	5157.4	4416.2	15.4	10.9

Analyte stability in the extract solution and on the plate

To assess stability more objectively (Fig. 7), the peak heights of the four markers (anthocyanin 1, anthocyanin 2, chlorogenic acid, and caffeic acid on lanes 1–2 and 5–6) were compared with those detected on lanes 3–4. There is no difference between the individual lane, indicated by the RSD calculated for each marker was $\leq 20\%$ (Table 2). It means the four analytes have good stability in the extract solution and on the TLC plate (Kartini et al., 2021; Reich and Schibli, 2007).

Stability of the chromatography results

Evaluating the stability of the chromatography results aimed to determine how long after elution or derivatization the chromatogram would still show the same results. Fig. 8B–F shows that anthocyanin 1 (spot a) and anthocyanin 2 (spot b) on lanes 1–6 appeared in the same color and intensity as those in Fig. 8A, indicating that the chromatogram made with the system I has good stability within 30 minutes after elution. Fig. 9B–F also show that chlorogenic acid (spot a) and caffeic acid (spot b) on lanes 1–6 had the same color and intensity as those in Fig. 9A, suggest-

ing that the chromatogram made with system II has good stability until 30 minutes after derivatization.

Then, to achieve more objective results, the peak height average was calculated for each marker (anthocyanin 1, anthocyanin 2, chlorogenic acid, and caffeic acid on lanes 1–6) and was plotted against time of observation to make a curve (Fig. 10). The graph shows no significant decline on all the marker curves during the 30-minute observation, demonstrating good stability of the four analytes within 30 minutes after elution or derivatization. These results indicated that the chromatogram could be either documented immediately or delayed for not more than 30 minutes after elution or derivatization.

Precision on a plate and intermediate precision

Precision represents the closeness between chromatography results from multiple sample replicates and random errors in a method. For qualitative analyses, precision can be linked to the R_f values of the separated compounds (Reich and Schibli, 2007). However, in this current study RSD of the peak height was calculated as well to make the data more comprehensive. Tables 3 and 4 show the markers' R_f and peak heights from five application replicates eluted with systems I and II. These data indicated that


the two systems designed to determine the *H. sabdariffa*'s TLC pattern have good precision on a plate and intermediate precision, with a RSD of smaller than 20%.

Comparison between *H. sabdariffa* chromatograms made with the optimized and the IHP II systems

Fig. 11A-B shows the TLC patterns of *H. sabdariffa* made with the IHP II system. Investigations under visible light (Fig. 11A) revealed one red spot of the extract (spot 1, lane S) with a different R_f from the reference solution, cyanidin 3-O-glucoside (St). The spot was clear but showed tailing, and the rest of the sample did not seem to elute from the application site. When observed under 366 nm UV light (Fig. 11B), the extract (lane S) produced one dark blue fluorescent spot (spot 1) above the tailed bright one. Meanwhile, the reference cyanidin 3-O-glucoside (St) fluoresced a blackish-blue color with a relatively large area. It will

be difficult to use TLC patterns with poor separation as a reference, making misidentification of a crude drug highly likely.

With the optimized TLC systems, the *H. sabdariffa* calyx extract was eluted into more spots with better shape and separation (Fig. 11C-D). The system I produced two well-separated red spots with different R_f values than the reference cyanidin 3-O-glucoside (St). Therefore, both spots are concluded as not cyanidin 3-O-glucoside but suspected as anthocyanins (1 and 2) because they appeared red in visible light. Therefore, anthocyanins 1 and 2 are proposed as analytical markers for *H. sabdariffa*. System II produced several spots, two of which were chlorogenic acid and caffeic acid with clear blue fluorescence, no tails, and were well-separated from other compounds. Because they were easily and clearly detected using TLC, chlorogenic and caffeic acids are also proposed as analytical markers for *H. sabdariffa* calyces.






Figure 9. *H. sabdariffa* chromatograms made with system II under 366 nm UV light at different times of observation: 0 (A), 5 (B), 10 (C), 15 (D), 20 (E), and 30 minutes (F) after derivatization with NP/PEG. Different lanes from left to right: first to sixth replicates, a and b: chlorogenic acid and caffeic acid.




Figure 10. Stability graphs of anthocyanin 1, anthocyanin 2, chlorogenic acid, and caffeic acid after elution or derivatization.

Table 3. Rf values and peak heights of the marker anthocyanin during precision on a plate and intermediate precision.


Lane	Anthocyanin 1						Anthocyanin 2					
	Day 1		Day 2		Day 3		Day 1		Day 2		Day 3	
	Rf	Height	Rf	Height	Rf	Height	Rf	Height	Rf	Height	Rf	Height
1	0.392	2402.7	0.286	2166.3	0.286	2805.2	0.474	1558.3	0.350	1191.6	0.362	1316.1
2	0.384	2392.3	0.287	2192.0	0.294	2965.3	0.473	1509.0	0.356	1199.9	0.367	1461.1
3	0.386	2223.8	0.292	2156.0	0.295	2999.9	0.468	1376.1	0.359	1215.7	0.367	1518.4
4	0.383	2071.3	0.291	2159.7	0.297	2964.3	0.473	1074.8	0.359	1180.5	0.370	1530.1
5	0.381	2146.7	0.297	2140.8	0.298	2969.6	0.461	1306.9	0.362	1208.5	0.369	1544.0
Mean ± SD	2253.3 ± 169.4		2164.7 ± 21.2		2926.1 ± 80.6		1362.3 ± 220.3		1195.1 ± 11.9		1462.8 ± 104.3	
RSD (%)	7.5		1.0		2.8		16.2		1.0		7.1	
Precision on a plate, RSD (%)	1.0–7.5						1.0–16.2					
Intermediate precision, RSD (%)	17.0						10.1					

*Mean, SD, and RSD were calculated from the height of peaks.

Table 4. Rf values and peak heights of the markers chlorogenic acid and caffeic acid during precision on a plate and intermediate precision.

	Chlorogenic acid						Caffeic acid					
Lane	Day 1		Day 2		Day 3		Day 1		Day 2		Day 3	
	Rf	Height	Rf	Height	Rf	Height	Rf	Height	Rf	Height	Rf	Height
1	0.503	614.0	0.455	694.1	0.471	883.3	0.912	4507.9	0.879	4786.7	0.904	4646.8
2	0.503	597.6	0.460	596.8	0.476	741.7	0.905	4530.3	0.879	4848.4	0.907	4843.8
3	0.503	622.8	0.445	795.1	0.479	938.8	0.912	4808.8	0.879	4783.9	0.914	5004.8
4	0.508	613.8	0.447	768.8	0.479	1105.3	0.908	4833.8	0.875	5177.7	0.914	4987.2
5	0.503	661.7	0.449	715.8	0.482	856.3	0.912	4939.7	0.877	4829.6	0.914	5024.6
Mean ± SD	622.0 ± 24.0		714.1 ± 77.0		905.1 ± 133.0		4724.1 ± 193.6		4885.3 ± 165.8		4901.4 ± 159.2	
RSD (%)	3.9		10.8		14.7		4.1		3.4		3.3	
Precision on a plate, RSD (%)	3.9–14.7						3.3–4.1					
Intermediate precision, RSD (%)	19.3						2.0					

*Mean, SD, and RSD were calculated from the height of peaks



CONCLUSION

The research has successfully developed two TLC systems (I and II) for *H. sabdariffa* calyx authentication and quality control better than the one described in IHP II. System I uses Si gel 60 F₂₅₄ for the stationary phase, ethyl acetate-formic acid-glacial acetic acid-water (100:11:11:2) for the mobile phase, *H. sabdariffa*'s ethanol extract (5%, 20 µL) for the test solution, cyanidin 3-O-glucoside (100 ppm, 4 µL) for the reference solution, no derivatization, and visible light detection. System II combines Si gel 60 F₂₅₄ as the stationary phase, ethyl acetate-formic acid-glacial acetic acid-toluene-water (80:11:11:20:19) as the mobile phase, *H. sabdariffa*'s ethanol extract (5%, 20 µL) as the test solution, chlorogenic acid (1000 ppm, 2 µL) and caffeic acid (50 ppm, 2 µL) as the reference solutions, derivatization with the visualization reagent NP/PEG, and detection with 366 nm UV light. These developed systems have good stability and precision, facilitating *H. sabdariffa* calyx authentication and paving the way for developing content analysis methods for *H. sabdariffa* markers.

CONFLICT OF INTEREST

The authors declare no conflicts of interest.

ACKNOWLEDGMENTS

This research was funded by University of Surabaya with the grant number 013/SP-Lit/LPPM-01/FF/Int/FF/V/2022. The authors thank Dr. Nina Dewi Octaviyanti for the donations of cyanidin 3-O-glucoside.

REFERENCES

- Ali BH, Wabel NA, Blunden G (2005) Phytochemical, pharmacological and toxicological aspects of *Hibiscus sabdariffa* L.: A review. *Phytother Res*. 19: 369–375. <https://doi.org/10.1002/ptr.1628>

<https://jppres.com>

- Antonia YT, Nii AA, Nancy C, Achana N (2019) Genetic diversity, variability and characterization of the agro-morphological traits of Northern Ghana Roselle (*Hibiscus sabdariffa* var. *altissima*) accessions. *Afr J Plant Sci* 13: 168–184. <https://doi.org/10.5897/AJPS2019.1783>
- Aryanti N, Nafiunisa A, Wardhani DH (2019) Conventional and ultrasound-assisted extraction of anthocyanin from red and purple roselle (*Hibiscus sabdariffa* L.) calyces and characterisation of its anthocyanin powder. *Int Food Res J* 26: 529–535.
- Chan SS-K, Li S-L, Lin G (2007) Pitfalls of the selection of chemical markers for the quality control of medicinal herbs. *J Food Drug Anal* 15: 365–371. <https://doi.org/10.38212/2224-6614.2398>
- Chumsri P, Sirichote A, Itharat A (2008) Studies on the optimum conditions for the extraction and concentration of roselle (*Hibiscus sabdariffa* Linn.) extract. *Songklanakarin J Sci Technol* 30: 133–139.
- Committee MHM (2015) Malaysian Herbal Monograph 2015: Institute for Medical Research.
- EDQM (2008) European Pharmacopeia 7.0, Strasbourg: The Directorate for the Quality of Medicines of the Council of Europe (EDQM).
- Hapsari BW, Setyaningsih W (2021) Methodologies in the analysis of phenolic compounds in roselle (*Hibiscus sabdariffa* L.): Composition, biological activity, and beneficial effects on human health. *Horticulturae* 7: 35. <https://doi.org/10.3390/horticulturae7020035>
- Health IMO (2017) Farmakope Herbal Indonesia Edisi II, Jakarta: Departemen Kesehatan Republik Indonesia.
- Izquierdo-Vega JA, Arteaga-Badillo DA, Sánchez-Gutiérrez M, Morales-González JA, Vargas-Mendoza N, Gómez-Aldapa CA, Castro-Rosas J, Delgado-Olivares L, Madrigal-Bujaidar E, Madrigal-Santillán E (2020) Organic acids from roselle (*Hibiscus sabdariffa* L.)—A brief review of its pharmacological effects. *Biomedicines* 8: 100. <https://doi.org/10.3390/biomedicines8050100>
- Kartini K, Andriani YA, Priambodo W, Jayani NI, Hadiyat MA (2021) Validating and developing TLC-based fingerprinting for *Curcuma longa* L. *J Pharm Pharmacogn Res*. 9: 704–715. https://doi.org/10.56499/jppres21.1062_9.5.704
- Kim Y-H, Cho ML, Kim D-B, Shin G-H, Lee J-H, Lee JS, Park S-O, Lee S-J, Shin HM, Lee O-H (2015) The antioxidant activity and their major antioxidant compounds from *Acanthopanax*

- senticosus* and *A. koreanum*. *Molecules* 20: 13281–13295. <https://doi.org/10.3390/molecules200713281>
- Lago C, Landoni M, Cassani E, Atanassiu S, Cantaluppi E, Pili R (2014) Development and characterization of a coloured sweet corn line as a new functional food. *Maydica* 59: 191–200.
- Orfali R, Perveen S, Aati HY, Alam P, Noman OM, Palacios J, Al-Kurbi BSS, Al-Taweel AM, Khan A, Mehmood R (2021) High-performance thin-layer chromatography for rutin, chlorogenic acid, caffeic acid, ursolic acid, and stigmasterol analysis in *Periploca aphylla* extracts. *Separations* 8: 44. <https://doi.org/10.3390/separations8040044>
- Pham TN, Nguyen TNP, Lam TD, Tran TH, Nguyen DC, Vo D-VN, Le XT, Do ST, Bach LG (2019) Effects of various solvent concentration, liquid-solid ratio, temperatures and time values on the extraction yield of anthocyanin from Vietnam *Hibiscus sabdariffa* L. (roselle). *IOP Conf Ser: Mater Sci Eng* 542: 012033. <https://doi.org/10.1088/1757-899X/542/1/012033>
- Reich E, Schibli A (2007) High-performance thin-layer chromatography for the analysis of medicinal plants. Stuttgart: Thieme. <https://doi.org/10.1055/b-0034-65188>
- Santos DT, Cavalcanti RN, Rostagno MA, Queiroga CL, Eberlin MN, Meireles MA (2013) Extraction of polyphenols and anthocyanins from the jambul (*Syzygium cumini*) fruit peels. *Food Public Health* 3: 12–20. <https://doi.org/10.5923/j.fph.20130301.02>
- Spangenberg B, Poole CF, Weins C (2011) Quantitative thin-layer chromatography: A practical survey. Springer Berlin, Heidelberg. Pp. XV, 388. <https://doi.org/10.1007/978-3-642-10729-0>
- Sukkhaeng S, Promdang S, Doung-Ngern U (2018) Fruit characters and physico-chemical properties of roselle (*Hibiscus sabdariffa* L.) in Thailand – A screening of 13 new genotypes. *J Appl Res Med Aromat Plants* 11: 47–53. <https://doi.org/10.1016/j.jarmap.2018.10.001>
- Torres-Morán MI, Escoto-Delgadillo M, Ron-Parra J, Parra-Tovar G, Mena-Munguía S, Rodríguez-García A, Rodríguez-Sahagún A (2011) Relationships among twelve genotypes of roselle (*Hibiscus sabdariffa* L.) cultivated in western Mexico. *Ind Crops Prod* 34: 1079–1083. <https://doi.org/10.1016/j.indcrop.2011.03.0205>

AUTHOR CONTRIBUTION:

Contribution	Dewi BAASK	Kartini K
Concepts or ideas		x
Design		x
Definition of intellectual content		x
Literature search	x	x
Experimental studies	x	x
Data acquisition	x	x
Data analysis	x	x
Statistical analysis	x	x
Manuscript preparation	x	x
Manuscript editing		x
Manuscript review	x	x

Citation Format: Dewi BAASK, Kartini K (2023) System optimization and validation to improve thin-layer chromatography of roselle calyces (*Hibiscus sabdariffa* L.) required by the Indonesian Herbal Pharmacopoeia Edition II. *J Pharm Pharmacogn Res* 11(2): 243–254. https://doi.org/10.56499/jppres22.1566_11.2.243

Publisher's Note: All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Open Access: This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits use, duplication, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license and indicate if changes were made.


[Submit Article](#)

About


Description

The Journal of Pharmacy & Pharmacognosy Research (JPPRes) is an international, specialized and peer-reviewed open access journal, under the auspices of GarVal Editorial Ltda., which publishes studies in the pharmaceutical and herbal fields concerned with the physical, botanical, chemical, biological, toxicological properties and clinical applications of molecular entities, active pharmaceutical ingredients, devices and delivery systems for drugs, vaccines and biologicals, including their design, manufacture, evaluation and marketing. This journal publishes research papers, reviews, commentaries and letters to the editor as well as special issues and review of pre-and post-graduate thesis from pharmacists or professionals involved in Pharmaceutical Sciences or Pharmacognosy.

JPPRes has an acceptance rate of 33.6% (2022). The average time between submission and final decision is 100 days and the average time between acceptance and final publication is 15 days.

Manuscripts submitted to JPPRes are only accepted on the understanding that they are subject to editorial review and that they have not been, and will not be, published in whole or in part in any other journal.

The URL of the journal website is jppres.com/jppres. You can write to the editor through the following email:

[Email](#)


Language

JPPRes accepts, in Spanish or English, review articles, articles for educational forum, original research articles (full length and short communications), letter to editor and case reports. Articles concerning all aspects of Pharmacy and Pharmacognosy will be considered.

Charges

In order to provide free access to readers, and to cover the costs of peer review, copyediting, typesetting, long-term archiving, and journal management, an article-processing charge (**APC**) of **600 USD (US Dollars)** applies to papers accepted after peer review.

Articles of general interest (e.g. methods, therapeutics, medical education, interesting websites, new drug information and commentary on a recent topic) are also welcome.

Manuscripts in which language is difficult to understand may be returned to the author for revision before scientific review.

Audience

Biochemists, Biotechnologists, Botanists, Chemical Engineers, Clinical Pharmacologists, Medical Scientists, Medicinal Chemists, Natural Product Chemists, Pharmaceutical Scientists, Pharmacologists, Pathologists, Plant Scientists, and Toxicologists, among others.

Copyright

The copyright of the articles are those of the authors and the rights of editing and publishing to JPPRes. The articles published in the journal may be used freely for educational and scientific purposes as long as proper citation of these is made.

Areas

Alternative and Complementary Medicine, Analytical Toxicology, Biochemical Pharmacology, Biologicals, Chinese Medicine Resources, Clinical Pharmacology, Cosmetic Science, Drug Delivery System, Drug Information, Drug Metabolism, Education in Pharmacy, Pharmacology and Pharmacognosy, Ethnobotany and Ethnopharmacology, Food-Drug Interactions, Herbal-Drug Interactions, Drug-Drug Interactions, Formulations, Functional Foods and Nutraceuticals, Health and Medical Informatics, Herbal Medicine, Immunopharmacology, Medical Anthropology, Medication Management, Medicinal Chemistry, Molecular Medicine, Molecular Modeling, Molecular Pharmacology, Neuropsychopharmacology, Nutrition, Pharmaceutical Care, Pharmaceutical Marketing, Pharmaceutical Microbiology, Pharmaceutical Raw Material Science, Pharmaceutical Research, Pharmaceuticals, Pharmaceutical Raw Materials, Pharmacodynamics, Pharmacoepidemiology, Pharmacogenetics, Pharmacogenomics, Pharmacokinetics, Pharmacology, Pharmacotherapy, Pharmacy Informatic, Pharmacy Practice, Pharmacognosy, Physical Pharmaceutics, Phytochemistry, Phytomedicine, Phytotherapy, Social and Administrative Pharmacy, Toxicology, Traditional Medicine, Vaccines, Veterinary Pharmacology, Zoopharmacognosy

Editorial Policy

JPPRes considers only original contributions submitted exclusively to the journal. Prior and duplicate publications are not allowed. Publication of abstract under conference proceedings will not be considered as prior publication. It is the duty of the authors to inform the JPPRes about all submissions and previous reports that might be regarded as prior or duplicate publication.

Manuscripts for publication will be considered on their individual merits. All manuscripts will be subjected to peer review. Normally manuscripts will be sent to at least two reviewers and their comments along with the editorial board's decision will be forwarded to the contributor for further action.

The JPPRes insists on ethical practices in both human and animal experiments. Evidence for approval by a local

The ethical standards of experiments must be in accordance with the Code of Ethics of the World Medical Association (Declaration of Helsinki) for experiments involving humans and EU Directive 2010/63/EU for animal experiments. The journal will not consider any paper which is ethically unacceptable. A statement on ethics committee permission and ethical practices must be included in all research articles under the 'Materials and Methods' section. Uniform Requirements for manuscripts submitted to Biomedical Journals must be observed.

Authors must be careful when they reproduce text, tables or illustrations from other sources. Plagiarism will be viewed seriously. Please see instructions. All accepted papers are subject to editorial changes.

Ethics Committee must be supplied by the authors on demand. Animal experimental procedures should be as humane as possible and the details of anaesthetics and analgesics used should be clearly stated.

Abstracting and Indexing information:

Databases:

- EMBASE
- Chemical Abstracts Service (CAS), American Chemical Society (ACS)
- MIAR
- SHERPA/RoMEO
- Scopus
- Web of Science (Emerging Sources Citation Index)

Directories:

- DOAJ (Directory of Open Access Journals)
- REDALYC (Red de Revistas Científicas de América Latina y el Caribe, España y Portugal)
- REDIB (Red Iberoamericana de Innovación y Conocimiento Científico)
- ROAD (Directory of Open Access Scholarly Resources)
- Ulrich's International Periodical Directory, New Jersey

Indexes:

- Google Scholar
- Index Copernicus International
- PERIODICA (Índice de Revistas Latinoamericanas en Ciencias - Dirección General de Bibliotecas, UNAM, Mexico)
- Qualis CAPES

Catalogues:

- HINARI (Research in Health Programme from the World Health Organization)
- LATINDEX (Sistema Regional de Información en Línea para Revistas Científicas de América Latina, el Caribe, España y Portugal)
- United States National Library of Medicine (NLM) – National Center for Biotechnology Information

You can also follow the content of the journal and the updates through Mendeley, Academia.org, and ResearchGate.

Note: This site does not host or receive funding from advertising.

Last Update: July 3th, 2023

[Submit Article](#)

Editorial Board

**JPPRes**Journal of Pharmacy
& Pharmacognosy
Research

Gabino Garrido

Editor-in-Chief

[Scholar Google](#)[Scopus](#)[ResearchGate](#)[ORCID](#)

GarVal Editorial Ltda.

[Email](#)

Marisela Valdés González

Executive Editor

[ORCID](#)

GarVal Editorial Ltda.

[Email](#)

Xavier Garrido

Editorial and Design Manager

[LinkedIn](#)[Behance](#)[Web](#)AVAGAX Studio
Antofagasta, Chile[Email](#)

Editorial Board Members



Antonio Vassallo

Dipartimento di Scienze.
Università degli Studi della
Basilicata, Potenza, Italia.[Research Gate](#)[ORCID](#)[Scopus](#)

Bárbara B. Garrido Suárez

Laboratorio de
Farmacología y
Toxicología, Centro de
Investigación y Desarrollo
de Medicamentos, La
Habana, Cuba.[Research Gate](#)[ORCID](#)

Carla Delporte

Facultad de Ciencias
Químicas y Farmacéuticas,
Universidad de Chile,
Santiago, Chile[Research Gate](#)[ORCID](#)

Damaris Silveira

Faculdade de Ciências da
Saúde, Universidade de
Brasília, Brazil[Lattes](#)[ORCID](#)

Douglas S. de Almeida



Edgar Pastene



Etile Spegazzini



Farid Chemat

Chaves

Dep. de Ciências Farmacêuticas (DCFAR), Instituto de Ciências Biológicas e da Saúde, Universidade Federal Rural do Rio de Janeiro, Rodovia, RJ. Brazil

ResearchGate

ORCID

Lattes



Fidel O. Castro

Departamento de Ciencias Pecuarias, Universidad de Concepción, Chillán, Chile

ResearchGate

ORCID



José H. Isaza-Martínez

Universidad del Valle, Cali, Colombia

ResearchGate

ORCID



María Inés Isla

Facultad de Ciencias Naturales e Instituto Miguel Lillo. Universidad Nacional de Tucumán, Tucumán, Argentina.

ResearchGate

ORCID



Universidad de Concepción, Concepción, Chile

ResearchGate

ORCID



Guilherme N. L. do Nascimento

Universidade Federal de Tocantins. Brazil

ORCID



Juan C. Sepúlveda-Arias

Universidad Tecnológica de Pereira, Pereira, Colombia

ResearchGate

ORCID



Marisol Fernández Alfonso

Facultad de Farmacia, Universidad Complutense de Madrid, España

ResearchGate

ORCID



Universidad de Belgrano, Buenos Aires, Argentina

ResearchGate



Jacqueline Sepúlveda

Universidad de Concepción, Concepción, Chile

ResearchGate

ORCID



Madan Mohan Gupta

School of Pharmacy, Faculty of Medical Sciences. The University of the West Indies, Trinidad & Tobago, West Indies

Google Scholar

ORCID



Mayank Gangwar

Department of Pharmacology, Institute of Medical Sciences, Banaras Hindu University, Varanasi (UP) India

Scholar Google

ORCID

Université Avignon et des Pays de Vaucluse, Avignon, France

ResearchGate



Jelena Nadinic

Universidad de Buenos Aires, Argentina.

ResearchGate

ORCID



Mahomoodally M. Fawzi

Department of Health Sciences, Faculty of Science, University of Mauritius. Mauritius.

ORCID

ResearchGate



Silvia Debenedetti

Pharmacy Career. Universidad de Belgrano, Buenos Aires, Argentina

ORCID

ResearchGate

Reviewers

The reviewers will be recruited among researchers and

Vikash Kumar Ruhil

PDM College of Pharmacy,
Bahadurgarh, Haryana,
India

[Google Scholar](#)**Yasser Shahzad**

Department of
Pharmacy. COMSATS
University Islamabad,
Pakistan

[Scopus](#)[ORCID](#)[Google Scholar](#)

clinicians with high international reputation in Pharmacy
and Pharmacognosy.

Last Update: August 20th, 2023

**JPPRes**

Journal of Pharmacy
& Pharmacognosy
Research

[Home](#) [About](#) [Archive](#) [Authors](#)
[Reviewers](#) [Events](#) [Blog](#)

[Twitter](#)[LinkedIn](#)[Facebook](#)

© 2013–2020 by the authors; licensee JPPRes, Antofagasta, Chile. This journal is an open-access journal distributed under the terms and conditions of the Creative Commons Attribution license–Non Commercial 4.0 international. The content on this site is intended for health professionals. If you are not a health professional, please talk to your doctor about any doubts or concerns regarding your health

Made with ♥ by **AVAGAX Studio**

Volume 11, Issue 2

J Pharm Pharmacogn Res 11(2), (Mar-Apr) 2023



IFC (**Journal of Pharmacy & Pharmacognosy Research**).

1.- Original Article

Dini Kesuma, Aguslina Kirtishanti, Citra H.A. Makayasa, I Gede A. Sumartha (2023) **Anticancer activity of N-(4-t-butylbenzoyl)-N'-phenylthiourea: Molecular docking, synthesis, and cytotoxic activity in breast and cervical cancer cells.** | [Actividad anticancerígena de la N-(4-t-butillbenzoil)-N'-feniltiourea: acoplamiento molecular, síntesis y actividad citotóxica en células de cáncer de mama y de cuello uterino]. J Pharm Pharmacogn Res 11(2): 208-215. https://doi.org/10.56499/jppres22.1508_11.2.208 [jppres_pdf_free](#)[640 Kb]

2.- Original Article

Maria R. Ramirez, Ignacio Rintoul, Manuel Leiva, José S. Molli, Juan C. Yori (2023) **Oral supplementation of *Ilex paraguariensis* extract and eggshell membrane exhibit beneficial effects on the skin of mature male rats.** | [La suplementación oral de extracto de *Ilex paraguariensis* y membrana de cáscara de huevo muestran efectos beneficiosos sobre la piel de ratas macho maduras]. J Pharm Pharmacogn Res 11(2): 216-228. https://doi.org/10.56499/jppres22.1509_11.2.216 [jppres_pdf_free](#)[955 Kb]

3.- Original Article

Hassan Sohrabian Kafraj, Maryam Alipour, Abdolhosein Shiravi, Vida Hojati, Mojtaba Khaksarian (2023) **Chitosan-alginate porous scaffold incorporated with hydroalcoholic *Zingiber officinale* Roscoe extract for neural tissue engineering.** | [Estructura porosa de quitosano-alginato incorporada con extracto hidroalcohólico de *Zingiber officinale* Roscoe para la ingeniería del tejido neural]. J Pharm Pharmacogn Res 11(2): 229-242. https://doi.org/10.56499/jppres22.1505_11.2.229 [jppres_pdf_free](#)[1.26 Mb]

4.- Original Article

Bhujangga Agung Ayu Sri Kartika Dewi, Kartini Kartini (2023) **System optimization and validation to improve thin-layer chromatography of roselle calyces (*Hibiscus sabdariffa* L.) required by the Indonesian Herbal Pharmacopoeia Edition II.** | [Optimización y validación del sistema para mejorar la cromatografía en capa fina de cálices de rosa mosqueta (*Hibiscus sabdariffa* L.) requerida por la Farmacopea Herbal Indonesia Edición II]. J Pharm Pharmacogn Res 11(2): 243-254. https://doi.org/10.56499/jppres22.1566_11.2.243 [jppres_pdf_free](#)[699 Kb]

5.- Original Article



STK71 mejora los niveles de HDL-c entre los pacientes con diabetes mellitus tipo 2 recién diagnosticada]. J Pharm Pharmacogn Res 11(2): 255–262. https://doi.org/10.56499/jppres22.1541_11.2.255 jppres_pdf_free [463 Kb]

6.- Original Article

Dewi Fibrini, I Nyoman Ehrich Lister (2023) **Pilot study on the effect of autologous activated platelet-rich plasma on diabetic foot ulcer wound healing and serum HbA1c levels.** | [Estudio piloto sobre el efecto del plasma rico en plaquetas activado autólogo en la cicatrización de la úlcera del pie diabético y los niveles séricos de HbA1c]. J Pharm Pharmacogn Res 11(2): 263–269. https://doi.org/10.56499/jppres22.1555_11.2.263 jppres_pdf_free [619 Kb]

7.- Original Article

Felipe Surco-Laos, Jorge A. Garcia, María R. Bendezú, Angel T. Alvarado, Doris Laos-Anchante, Manuel Valle-Campos, Juan F Panay-Centeno, Juan J. Palomino-Jhong, Paulina Eliades Yarasca-Carlos, Ana María Muñoz, Mario Bolarte-Arteaga, Mario Pineda, Berta Loja (2023) **Characterization of polyunsaturated fatty acids and antioxidant activity of *Vitis vinifera* L. (grape seeds from the Ica Valley, Peru).** | [Caracterización de ácidos grasos poliinsaturados y actividad antioxidante de las semillas de *Vitis vinifera* L. (uva) del valle de Ica, Perú]. J Pharm Pharmacogn Res 11(2): 270–280. https://doi.org/10.56499/jppres23.1575_11.2.270 jppres_pdf_free [910 Kb]

8.- Original Article

Hala F. Kasim (2023) **Prevalence of potential drug interactions among outpatients' prescriptions of community pharmacies in Nineveh Governorate, Iraq.** | [Prevalencia de posibles interacciones farmacológicas entre las recetas de los pacientes ambulatorios de las farmacias comunitarias de la gobernación de Nínive, Iraq]. J Pharm Pharmacogn Res 11(2): 281–290. https://doi.org/10.56499/jppres23.1577_11.2.281 jppres_pdf_free [344 Kb]

9.- Original Article

Monika Kerry Army, Rohimatul Khodijah, Yuli Haryani, Hilwan Yuda Teruna, Rudi Hendra (2023) **Antibacterial *in vitro* screening of *Helminthostachys zeylanica* (L.) Hook. root extracts.** | [Cribado antibacteriano *in vitro* de extractos de raíz de *Helminthostachys zeylanica* (L.) Hook.]. J Pharm Pharmacogn Res 11(2): 291–296. https://doi.org/10.56499/jppres22.1540_11.2.291 jppres_pdf_free [349 Kb]

10.- Original Article

Yurnadi Hanafi Midoen, Syafruddin Ilyas, Putra Santoso, Putri Cahaya Situmorang (2023) **Effect of maximal physical exercise on apoptosis via cytochrome c in hippocampus cells after administration of *Vitis gracilis* Wall.** | [Efecto del ejercicio físico máximo sobre la apoptosis vía citocromo c en células del hipocampo tras la administración de *Vitis gracilis* Wall.]. J Pharm Pharmacogn Res 11(2): 297–307. https://doi.org/10.56499/jppres22.1563_11.2.297 jppres_pdf_free [1.02 Mb]

11.- Original Article

Quranayati Quranayati, Nurdin Saidi, Nurliana Nurliana, Rinaldi Idroes, Naufal Gusti, Rosnani Nasution (2023) **Effect of *Phyllanthus emblica* L. stem bark extract on diabetic nephropathy and hyperlipidemia in rats.** | [Efecto del extracto de corteza de tallo de *Phyllanthus emblica* L. sobre la nefropatía diabética y la hiperlipidemia en ratas]. J Pharm Pharmacogn Res 11(2): 308–314. https://doi.org/10.56499/jppres23.1595_11.2.308 jppres_pdf_free [434 Kb]

12.- Original Article

Fatchiyah Fatchiyah, Dewi Ratih Tirta Sari, Hazna Noor Meidinna, Nia Kurnianingsih, Ema Pristi Yunita, Sinta Murlistyarini (2023) **Stability evaluation and heavy metal quantification of lotions from caprine milk, yogurt, and kefir.** | [Evaluación de la



Syamsu Nur, Muhammad Hanafi, Heri Setiawan, Nursamsiar, Berna Elya (2023) ***In silico* evaluation of the dermal antiaging activity of *Molineria latifolia* (Dryand. ex W.T. Aiton) Herb. Ex Kurz compounds. I** [Evaluación *in silico* de la actividad antienvjecimiento dérmica de compuestos de *Molineria latifolia* (Dryand. ex W.T. Aiton) Herb. ex Kurz]. J Pharm Pharmacogn Res 11(2): 325–345. https://doi.org/10.56499/jppres23.1606_11.2.325 jppres_pdf_free [2.69 Mb]

14.- Original Article

Cristian Valiente, Silvana Alvariza, Diego M. Robaina, Gonzalo Suárez (2023) **Development and validation of UHPLC method the determination of fluazuron in bovine tissues. I** [Desarrollo y validación de método UHPLC para la determinación de fluazuron en tejidos bovinos]. J Pharm Pharmacogn Res 11(2): 346–353. https://doi.org/10.56499/jppres22.1548_11.2.346 jppres_pdf_free [559 Kb]

15.- Original Article

Honesty N. Pinanti, Yuyun I. Christina, Muhaimin Rifa'i, Nashi Widodo, Muhammad S. Djati (2023) **Molecular docking study of ethyl acetate fraction of *Selaginella doederleinii* Hieron extract as anti-Warburg effect in breast cancer. I** [Estudio de acoplamiento molecular de la fracción de acetato de etilo del extracto de *Selaginella doederleinii* Hieron como efecto anti Warburg en cáncer de mama]. J Pharm Pharmacogn Res 11(2): 354–366. https://doi.org/10.56499/jppres22.1549_11.2.354 jppres_pdf_free [1.70 Mb]

© 2023 Journal of Pharmacy & Pharmacognosy Research (JPPRes)

Journal of Pharmacy and Pharmacognosy Research

COUNTRY Chile Universities and research institutions in Chile Media Ranking in Chile	SUBJECT AREA AND CATEGORY Health Professions Pharmacy Medicine Complementary and Alternative Medicine Pharmacology, Toxicology and Pharmaceutics Drug Discovery Pharmaceutical Science Pharmacology	PUBLISHER Asociacion Academica de Ciencias Farmaceuticas de Antofagasta (ASOCIFA)	SJR 2024 0.314Q2 H-INDEX 21
PUBLICATION TYPE Journals	ISSN 07194250	COVERAGE 2013-2025	INFORMATION Homepage How to publish in this journal editor@jppres.com

SCOPE

The Journal of Pharmacy & Pharmacognosy Research (JPPRes) is an international, specialized and peer-reviewed open access journal, under the auspices of AVAGAX – Diseño, Publicidad y Servicios Informáticos, which publishes studies in the pharmaceutical and herbal fields concerned with the physical, botanical, chemical, biological, toxicological properties and clinical applications of molecular entities, active pharmaceutical ingredients, devices and delivery systems for drugs, vaccines and biologicals, including their design, manufacture, evaluation and marketing. This journal publishes research papers, reviews, commentaries and letters to the editor as well as special issues and review of pre-and post-graduate thesis from pharmacists or professionals involved in Pharmaceutical Sciences or Pharmacognosy.

 Join the conversation about this journal




 Quartiles


FIND SIMILAR JOURNALS

options 

1 Advances in Pharmacological and Pharmaceutical Sciences GBR 70% similarity	2 Phytomedicine Plus NLD 70% similarity	3 Journal of Applied Pharmaceutical Science IND 67% similarity	4 Journal of Natural Remedies IND 67% similarity	5 Journal of HerbMed Pharmacology IRN 66% similarity
---	--	---	---	---



Gabino Garrido Garrido 4 years ago

Dear Melanie,
Thank you for your kind and quick reply.
Sincerely

reply



Gabino Garrido Garrido 4 years ago

Dear Scimagojr,
Please, some time ago I contacted Scimago in this way to see the possibility of incorporating my journal (Journal of Pharmacy and Pharmacognosy Research) into the Pharmacy category. I have not received any responses.
Hopefully, you can answer me if this is the appropriate way to make such a request.

Thank you!
Best regards,
Gabino Garrido,
editor-in-chief

← reply



Melanie Ortiz 4 years ago

SCImago Team

Dear Gabino,

Thank you for contacting us.

SJR is a portal with scientometric indicators of journals indexed in Scopus. All the metadata (Title, ISSN, Publisher, Category, etc.) have been provided by Scopus / Elsevier and SCImago doesn't have the authority over these data which are property of Scopus / Elsevier. SCImago is not allowed to do any changes of the metadata unless Scopus authorizes it expressly. SCImago has a signed agreement that limits our performance to the generation of scientometric indicators derived from the metadata sent in the last update. Keep also in mind that the SJR is a static image (the update is made only one time per year) of a database (Scopus) which is changing every day.

Therefore, we suggest you contact Scopus support regarding this matter here:

https://service.elsevier.com/app/answers/detail/a_id/14883/kw/scimago/supporthub/scopus/

Best Regards,
SCImago Team



Leave a comment

Name

Email

(will not be published)

Submit

The users of Scimago Journal & Country Rank have the possibility to dialogue through comments linked to a specific journal. The purpose is to have a forum in which general doubts about the processes of publication in the journal, experiences and other issues derived from the publication of papers are resolved. For topics on particular articles, maintain the dialogue through the usual channels with your editor.

Developed by:



Powered by:



Follow us on @ScimagoJR

Scimago Lab, Copyright 2007-2025. Data Source: Scopus®

EST MODUS IN REBUS
Horacio (Sedre 1.1.1.106)

[Legal Notice](#)

[Privacy Policy](#)



Source details

Journal of Pharmacy and Pharmacognosy Research

Open Access ⓘ

Years currently covered by Scopus: from 2013 to 2025

Publisher: Asociacion Academica de Ciencias Farmaceuticas de Antofagasta (ASOCIFA)

E-ISSN: 0719-4250

Subject area: Health Professions: Pharmacy Medicine: Complementary and Alternative Medicine
Pharmacology, Toxicology and Pharmaceutics: Pharmaceutical Science
Pharmacology, Toxicology and Pharmaceutics: Pharmacology [View all](#) ▾

Source type: Journal

[View all documents](#) > [Set document alert](#)  [Save to source list](#)

CiteScore 2023
3.0 ⓘ

SJR 2023
0.291 ⓘ

SNIP 2023
0.554 ⓘ

[CiteScore](#) [CiteScore rank & trend](#) [Scopus content coverage](#)

CiteScore 2023 ▾

3.0

=

913 Citations 2020 - 2023

308 Documents 2020 - 2023

Calculated on 05 May, 2024

CiteScoreTracker 2024 ⓘ

3.1

=

1,167 Citations to date

371 Documents to date

Last updated on 05 April, 2025 • Updated monthly

CiteScore rank 2023 ⓘ

Category	Rank	Percentile
Health Professions		
Pharmacy	#11/45	76th
Medicine		
Complementary and Alternative Medicine	#47/105	55th

[View CiteScore methodology](#) > [CiteScore FAQ](#) > [Add CiteScore to your site](#) 

About Scopus

[What is Scopus](#)

[Content coverage](#)

[Scopus blog](#)

[Scopus API](#)

[Privacy matters](#)

Language

[日本語版を表示する](#)

[查看简体中文版本](#)

[查看繁體中文版本](#)

[Просмотр версии на русском языке](#)

Customer Service

[Help](#)

[Tutorials](#)

[Contact us](#)

ELSEVIER

[Terms and conditions](#) [Privacy policy](#) [Cookies settings](#)

All content on this site: Copyright © 2025 Elsevier B.V. [↗](#), its licensors, and contributors. All rights are reserved, including those for text and data mining, AI training, and similar technologies. For all open access content, the relevant licensing terms apply.

We use cookies to help provide and enhance our service and tailor content. By continuing, you agree to the use of cookies [↗](#).

