



Induction of Protocorm-Like Bodies (PLBs) *Phalaenopsis* spp. Hybrids Mutation through Ultraviolet Irradiation (UV₂₅₄) and Ethyl Methane Sulfonate (EMS)

Induksi Mutasi Protocorm Like Bodies (PLBs) Phalaenopsis spp. Hybrids melalui Iradiasi Sinar Ultraviolet (UV₂₅₄) dan Ethyl Methane Sulfonate (EMS)

Author(s): Agatha Sullivania Kurniadi ⁽¹⁾; Fenny Irawati ⁽¹⁾;
Sulistyo Emantoko Dwi Putra ⁽¹⁾; Poppy Hartatie Hardjo ^{(1)*}

⁽¹⁾ Universitas Surabaya

* Corresponding author: poppy_hardjo@staff.ubaya.ac.id

Submitted: 31 Dec 2022

Accepted: 5 Feb 2023

Published: 31 Mar 2023

ABSTRACT

Phalaenopsis sp. is the most-produced orchid species in Indonesia. Compared to conventional breeding, mutation induction by using mutagens, such as Ultraviolet Light-C ($\lambda = 254$ nm) (UV₂₅₄) and Ethyl Methane Sulfonate (EMS), could probably result in new superior orchid variants. This research aims to get some mutants with phenotypes that have visual differences in the *Phalaenopsis* spp. hybrids wild type. There were 4 durations of UV₂₅₄ irradiation: 5' on, 85' off; 10' on, 80' off (1 day and 7 days for each treatment); 4 concentrations of EMS used in this research: 0.05%; 0.06%; 0.07%; 0.08% for 6 hours of immersion; selected UV₂₅₄ irradiation (5' on, 85' off (7 days)) combined with these concentrations. UV₂₅₄ irradiation treatment (5' on, 85' off (1 day and 7 days); 10' on, 80' off (7 days)) resulted in some mutants with leaf phenotypes that were visually different from the wild type; 0.05% EMS treatment resulted in PLBs mutant with a visually larger size than the wild type; 0.08% EMS treatment and combination treatments (for EMS 0.05% and 0.08% for each treatment) resulted in non-growing albino PLBs. Hence, mutation induction using UV₂₅₄ and EMS in this research produced several most likely mutants having visual differences that may be more desirable than the wild type.

Keywords:

EMS,

Phenotype,

Mutation,

Combination Treatmen,

UV₂₅₄.

ABSTRAK

Kata Kunci:

EMS,

Fenotipe,

Mutasi,

Perlakuan Kombinasi,

UV254.

Phalaenopsis sp. merupakan spesies anggrek terbanyak yang diproduksi di Indonesia. Dibandingkan dengan pemuliaan secara konvensional, induksi mutasi dengan menggunakan mutagen, seperti sinar ultraviolet-C ($\lambda = 254$ nm) (UV₂₅₄) dan Ethyl Methane Sulfonate (EMS), dapat meningkatkan probabilitas didapatkannya varian anggrek unggul baru. Penelitian ini bertujuan untuk mendapatkan mutan yang secara fenotipe memiliki kenampakan visual yang berbeda dari *Phalaenopsis* spp. hybrids wild type. Terdapat 4 durasi iradiasi UV₂₅₄ yang digunakan: 5' menyala, 85' mati; 10' menyala, 80' mati (masing-masing selama 1 hari dan 7 hari); 4 konsentrasi EMS yang digunakan: 0,05%; 0,06%; 0,07%; 0,08% dengan durasi perendaman selama 6 jam; durasi iradiasi UV₂₅₄ terpilih (5' menyala, 85' mati selama 7 hari) dikombinasikan dengan 4 konsentrasi EMS. Perlakuan iradiasi UV₂₅₄ (5' menyala, 85' mati (1 hari dan 7 hari); 10' menyala, 80' mati (7 hari)) menghasilkan beberapa mutan dengan fenotipe daun yang secara visual berbeda dari wild type; perlakuan 0,05% EMS menghasilkan PLBs mutan yang secara visual berukuran lebih besar dari wild type; perlakuan 0,08% EMS dan perlakuan kombinasi (pada EMS dengan konsentrasi 0,05% dan 0,08%) menghasilkan PLBs albino yang tidak mampu tumbuh. Dalam penelitian ini, induksi mutasi menggunakan UV₂₅₄ dan EMS menghasilkan planlet yang kemungkinan besar merupakan mutan dengan perbedaan visual yang mungkin lebih diinginkan daripada wild type



INTRODUCTION

According to the Indonesian Central Bureau of Statistics, orchids are the most-produced floriculture crop in 2020 when compared to other ornamental plants. *Phalaenopsis* sp., known as the moth orchid, is the most popular species and sales reach 75% of the various types of orchids traded in Indonesia, which have unique characteristics, including appealing blossoms with a butterfly shape, huge flowers that can endure up to two months before dropping off, and flowers with a variety of colors and shapes that allow for future development (Sulistianingsih & Purwantoro, 2012; Zahara, 2017). The cross-breeding method can be used to create a new hybrid of the characteristics of the parents to increase the variety of orchids. However, this approach has a low chance of producing crosses with the desired characteristics and is difficult to transmit consistently (Romiyadi et al., 2018). In vitro plant breeding and propagation is one approach that can be used to modify the nature of orchids, from their appearance to their growing capacity. Using in vitro mutation treatment, both chemical and physical mutagens can be used to develop new, superior varieties of orchids, which can relatively accelerate the formation of mutants while expanding the possibilities for variations in orchids (Qosim et al., 2016). The resulting mutants are expected to increase the variety of orchids as well as their selling price.

UV₂₅₄ light constitutes one of the physical mutagens which can induce mutations because it triggers spontaneous changes and the production of new genetic variations, such as causing color variations in flowers (Li et al., 2021). The physical mutagen commonly used for orchids is gamma rays because they can increase vegetative genetic variation in plants (Damayanti, 2021). However, these mutagens have enormous energy and cannot be used carelessly. Magdalita et al. (2019)

said that germinating embryos of *Phalaenopsis aphrodite* irradiated with 15 Gy gamma rays showed a faster flowering response than the wild type. Irradiating gamma rays on *Phalaenopsis amabilis* plantlets that are ready to be acclimatized and getting a radiation dose of 25 Gy (Widiarsih & Dwimahyani, 2013). Surya et al. (2016) said that gamma rays above 300 Gy cause a reduction in the percentage of germination in the M1 generation of *Rubus lineatus* and *Rubus chrysophytes*. Radiation with UV light of 200–280 nm produces carnation plants that flower quickly and are shorter (Afza & Iriawati, 2016). Research by Castronovo et al. (2017) showed a drastic decrease in chlorophyll content due to exposure to UV-C light after 30; 60; and 120 minutes of exposure to *Echinacea purpurea* and *Taraxacum officinale* flowers.

EMS is of chemical mutagen which can trigger alkylation and is most likely to cause gene mutations (Talebi et al., 2012). According to Li et al. (2021), there have been no studies related to mutations caused by chemical mutagens belonging to alkylating agents that cause high heterozygosity in orchids. Mutation induction, which has been proven successful in several studies, utilizes the polyploidization breeding method, using colchicine and nitric oxide (NO) mutagens in *Cymbidium*, *Dendrobium*, *Oncidium*, and *Phalaenopsis* orchids (Li et al., 2021). Romiyadi et al. (2018) induced mutations by using EMS concentrations shown indicated interactions with three *Phalaenopsis* spp. hybrid species on root variables. Qosim et al. (2016) showed that soaking EMS for 12 hours with a concentration of 0.025% and 0.05% affected the formation of shoots from *Phalaenopsis* spp. hybrids. In other crops, such as sugarcane, EMS is used to improve genetics so as to produce high yields (Avivi et al., 2022).

This research aims to study and determine the effect of mutation induction treatment with UV₂₅₄ irradiation and



immersion in EMS with several concentrations on measurable and visible morphological differences. The orchids whose PLBs were used as explants were hybrid orchids from a cross between *Doritaenopsis* sp. × *Phalaenopsis elegant* 'Karin Aloha'. Through mutation treatment with UV irradiation and immersion in EMS in vitro on PLBs from *Phalaenopsis* spp. hybrids, it is expected to obtain new superior orchid variants.

METHODOLOGY

This research was carried out at the University of Surabaya. Some tools were used in this research, such as Laminar Air Flow (LAF) Cabinet, petri dish, UV box incubator, tweezers, analytical scale, culture bottles, and Bunsen burner, whereas the material used in this research were PLBs from *Phalaenopsis* spp. hybrids from a cross between *Doritaenopsis* sp. × *Phalaenopsis elegant* 'Karin Aloha', UV-C light ($\lambda = 254$ nm), EMS (Sigma), DMSO, MS medium, 6-Benzyl Amino Purine (BAP), Naphthalene Acetic Acid (NAA), sucrose, gel agent, and 96% alcohol.

PLBs were grown in vitro condition using an MS medium with the addition of BAP and NAA hormones. Then the PLBs were exposed to four different UV₂₅₄ exposure durations: 5' on, 85' off; 10' on, and 80' off (1 day and 7 days for each treatment). Each treatment was repeated five times with twenty clumps per petri dish. Parameters measured included: the average increase in the number of PLBs, PLBs color change, the percentage of regeneration, the survival rate of PLBs, the average percentage of albino PLBs, and the percentage of mutants. Based on these parameters, explants from one duration with the most optimum results were selected to be immersed in EMS as a combination treatment.

By calculating the LC₅₀ value using the probit analysis method and getting an

LC₅₀ value of 0.05% after 6 hours of soaking, the treatment was started. Then, with a 6-hour immersion period, four concentrations were chosen: 0.05%, 0.06%, 0.07%, and 0.08%. According to the following procedure, the EMS reagent was made by dissolving EMS in distilled water and 2% DMSO (Hadebe et al., 2017).

Required EMS rate (mL)

$$\frac{\text{EMS concentration} \times (\text{aquadex} + \text{DMSO} 2\%)}{100}$$

Example: 0.02% EMS

$$= \frac{0.02 \times (39.2 + 0.8)}{100} \\ = 0.016 \text{ mL EMS}$$

PLBs were immersed for 6 hours in EMS reagent with a concentration of 0.05%; 0.06%; 0.07%; and 0.08%. After that, the PLBs were rinsed and drained, then planted on *in vitro* medium. Each treatment was repeated five times with twenty clumps per petri dish. The parameters measured were the same as the UV₂₅₄ irradiation treatment.



Combination treatment was carried out by immersing the PLBs that had been irradiated with the optimum duration of UV₂₅₄ (5' on, 85' off (7 days)) in EMS with concentrations of: 0.05%; 0.06%; 0.07%; and 0.08%. Then the PLBs were rinsed and drained, then planted on *in vitro* medium. Each treatment was repeated five times with twenty clumps per petri dish. The parameters measured were the same as the UV₂₅₄ irradiation treatment.

Kruskal-Wallis and Mann-Whitney test ($\alpha=5\%$) for identifying the differences between treatments were used in data analysis using the SPSS program for the parameter of the average increase in the number of PLBs and PLBs color change. The PLBs color reference scale uses the standards that have been made (Table 1).



Table 1. Color scale reference for PLBs

Tabel 1. Referensi skala warna untuk PLBs

Score Skor	Color Warna	Color Reference Referensi Warna	Score Skor	Color Warna	Color Reference Referensi Warna
1	Black		6	Chartreuse Green	
2	Grey		7	Lime Green	
3	Brown		8	Pear Green	
4	White		9	Shamrock Green	
5	Yellow		10	Emerald Green	

Other parameters (percentage of regeneration, average percentage of live PLBs, and average percentage of albino PLBs) were obtained by averaging the results per treatment. The percentage of mutants was obtained by calculating the number of mutants with the same character compared to all surviving mutants per treatment. Explants that have visually different phenotypes to the wild type are referred to as mutants and are analyzed descriptively.

RESULT AND DISCUSSION

UV₂₅₄ Irradiation Treatment

At various durations, there was a decrease in measured phenotype parameters (mean number of PLBs, PLBs color, PLBs regenerated, survival rate, and percentage of albino) compared to the wild

type (Table 2). UV₂₅₄ irradiation in 5' on, 85' off for 7 days treatment had a decrease that was not significantly different from the wild type and had the highest average survival rate compared to other durations. This can be due to PLB mutations which occur randomly and form pyrimidine dimers which have an impact on increasing their multiplicative ability although not as well as the wild type. Radiation with UV-C light ($\lambda=100-280$ nm) causes chromosome damage due to the wavelength energy absorbed by purines and pyrimidines which results in transitions between guanine-cytosine and adenine-thymine nitrogen base pairs (Yulianto et al., 2019).

Damage that occurs due to UV₂₅₄ radiation causes 10 minutes to be more severe as seen from the presence of browning. The mutations that occur cause



Table 2. Effect of duration of exposure to UV₂₅₄ on measured parameters in 7 weeks after planting (WAP) explantsTabel 2. Pengaruh durasi paparan UV₂₅₄ pada parameter terukur pada eksplan berusia 7 minggu setelah tanam (MST)

Treatment Perlakuan	Average Increase in Number of PLBs ¹ <i>Rerata Peningkatan Jumlah PLB¹</i>	Color ^{1,2} Warna ^{1,2}	Regeneration (%) <i>Regenerasi (%)</i>	Survival Rate (%) <i>Tingkat Survival (%)</i>	Albino (%) <i>Albino (%)</i>
Wild type	14 ± 6.10 ^a	6 ^a	3	70 ± 14.22	2 ± 1.5
5' on, 85' off for 1 day	4 ± 2.10 ^b	6 ^a	30	44 ± 4.64	0
10' on, 80' off for 1 day	0 ^c	4 ^b	49	16 ± 10.74	0
5' on, 85' off for 7 days	13 ± 6.86 ^a	6 ^a	9	66 ± 4.89	1 ± 0.83
10' on, 80' off for 7 days	0 ^d	4 ^b	26	15 ± 4.38	0

Note:

1= numbers followed by the same letter in the same column indicate results that are not significantly different, with the Mann-Whitney test ($\alpha = 5\%$); 2 = refers to Table 1.1= Angka-angka yang diikuti huruf yang sama pada kolom yang sama mengindikasi hasil yang tidak berbeda nyata menurut uji Mann-Whitney ($\alpha = 5\%$); 2 = merujuk pada Tabel 1.

the cells to produce secondary metabolites in the epidermal tissue in the form of phenolic compounds (Müller-Xing et al., 2014). Explant mortality can be carried on by browning from cell damage that results in the production of phenolic chemicals as well as by disruptions in the genome brought on by the induction of mutations.

A decrease in the capacity to produce chlorophyll may be the cause of albino PLBs. This result is similar to the research from Afza & Iriawati (2016), UV₂₅₄ radiation has an inhibitory effect on carnation plants, resulting in reduced chlorophyll levels, shorter plant segments, and suppression of axillary shoots.

Many absorbing hairs were grown on the surface of PLBs that had been given the UV₂₅₄ treatment (5' on, 85' off for 7 days) (Figure 1). PLBs contain absorbing hairs that have a fine hair-like appearance at the base and do not cover the bipolar poles (Mose et al., 2017). Absorbing hair can optimize the absorption of nutrients and water (Ningrum et al., 2017; Novak et al., 2014). This complimentary tool also plays a role in the ability of PLBs to recover as

well as correlated with the speed of growth and multiplication of PLBs.

Figure. 1. Absorbing hair on PLBs surface induced by UV₂₅₄ (5' on, 85' off (7 days))Gambar 1. Absorbing hair pada permukaan PLBs yang diinduksi dengan UV₂₅₄ (5' menyala, 85' mati (7 hari))

PLBs that generate new leaves and/or roots are referred to as regenerating PLBs. As can be observed, the amount of PLBs that regenerate depends on how long they are exposed to UV₂₅₄, with a one-day



treatment resulting in a higher proportion of PLBs than a seven-day treatment (Table 2). This may be related to the reasons that PLBs take longer to recover from long exposures, which can cause more severe damage, and that cells are more concerned with maintaining their viability than with regenerating.

Immersion of EMS

In this research, the EMS single and combination treatments were compared to the wild type using a mutagen solvent control (distilled water + DMSO 2%) in both cases. DMSO has an antibacterial effect, can enhance cell division and cell biomass, and is employed as a solvent for water-insoluble chemicals and mutagens like EMS (Xia Chen & Thibeault, 2013).

Table 3. Comparison of wild type with mutagen solvent (distilled water + DMSO 2%) to the measured parameters in 7 weeks after planting (WAP) explants

Tabel 3. Perbandingan dari wild type dengan pelarut mutagen (akuades + DMSO 2%) pada parameter terukur pada eksplan berusia 7 minggu setelah tanam (MST)

Treatment Perlakuan	Average Increase in Number of PLBs ¹ Rerata Peningkatan Jumlah PLB ¹	Color ^{1,2} Warna ^{1,2}	Regeneration (%) Regenerasi (%)	Survival Rate (%) Tingkat Survival (%)	Albino (%) Albino (%)
Wild type	14 ± 6.10 ^a	6 ^a	3	70 ± 14.22	2 ± 1.5
0% EMS	16 ± 8.91 ^a	6 ^a	4	96 ± 1.25	1 ± 0.80
0% EMS for the combination treatment	10 ± 5.38 ^b	6 ^a	37	82 ± 9.71	7 ± 4.18

Note:

1= numbers followed by the same letter in the same column indicate results that are not significantly different, with the Mann-Whitney test ($\alpha = 5\%$); 2 = refers to Table 1.

1= Angka-angka yang diikuti huruf yang sama pada kolom yang sama mengindikasi hasil yang tidak berbeda nyata menurut uji Mann-Whitney ($\alpha = 5\%$); 2 = merujuk pada Tabel 1.

Table 4. Effect of EMS concentrations on measured parameters in 7 weeks after planting (WAP) explants

Tabel 4. Pengaruh konsentrasi EMS pada parameter terukur pada eksplan berusia 7 minggu setelah tanam (MST)

Treatment Perlakuan	Average Increase in Number of PLBs ¹ Rerata Peningkatan Jumlah PLB ¹	Color ^{1,2} Warna ^{1,2}	Regeneration (%) Regenerasi (%)	Survival Rate (%) Tingkat Survival (%)	Albino (%) Albino (%)
Wild type	14 ± 6.10 ^a	6 ^a	3	70 ± 14.22	2 ± 1.5
0% (solvent control)	16 ± 8.91 ^a	6 ^a	4	96 ± 1.25	1 ± 0.80
0.05%	4 ± 2.48 ^{bcd}	4 ^c	0	69 ± 4.32	53 ± 20.58
0.06%	4 ± 2.14 ^{cd}	4 ^c	13	42 ± 15.32	37 ± 19.92
0.07%	5 ± 2.44 ^{bc}	5 ^{bc}	6	54 ± 12.80	63 ± 22.54
0.08%	3 ± 1.56 ^d	4 ^c	8	45 ± 18.80	96 ± 3.70

Note:

1= numbers followed by the same letter in the same column indicate results that are not significantly different, with the Mann-Whitney test ($\alpha = 5\%$); 2 = refers to Table 1.

1= Angka-angka yang diikuti huruf yang sama pada kolom yang sama mengindikasi hasil yang tidak berbeda nyata menurut uji Mann-Whitney ($\alpha = 5\%$); 2 = merujuk pada Tabel 1.



The function of solvent control is to guarantee that the solvent has no mutation effect so that the mutagen effect can be accurately assessed. The results indicate that none of the two solvent controls has a mutation effect, allowing the solvent to be used (Table 3).

Giving EMS at a specific concentration affects the observed parameters since the concentration used exceeds the LC₅₀ EMS value and results in increased PLB mortality (Table 4). High levels of EMS exposure can interfere with the function of enzymes involved in the germination process and inhibit auxin production from taking (Arisha et al., 2014). The optimal EMS concentration is 0.05% EMS, with a higher survival rate than other concentrations. The percentage of albinos at various EMS concentrations is high, indicating that this response is a response to cell damage. PLBs that survive will exhibit growth. Albinism can be caused by chlorophyll mutations, which result in discoloration due to a lack of chlorophyll. PLBs are unable to form chlorophyll or lose the ability to do so in all parts of the cell exposed to the mutagen (Romeida et al., 2012). 0.05% EMS can cause changes in the color of the leaves of *Aerides crispa* Lindl. from green to white due to a lack of chlorophyll (Srivastava et al., 2018).

Secondary PLB grew and regenerated from the leaf tips of PLBs exposed to 0.05% EMS as a result of the leaves coming into contact with *in vitro* medium (Figure 2). PLBs can be induced to proliferate by using leaves, and these PLBs grow from the epidermal cells of the leaves, which can then be regenerated to form plantlets (Chookoh et al., 2019; Huang et al., 2014).




Figure. 2. Mutant secondary PLB formed from leaf tips induced by 0.05% EMS. Remarks: marked with an orange circle.

Gambar 2. PLB sekunder mutan yang terbentuk pada ujung daun yang diinduksi dengan 0,05% EMS. Keterangan: ditandai dengan lingkaran oranye.

Combination Treatment of UV₂₅₄ Light Irradiation and EMS

The mutagens used in this research can cause random mutagenesis. Although the dose and duration can be planned, the results are difficult to predict (Mullins et al., 2021). The survival rate increased as the concentration increased, but the average albino percentage increased as immersion concentration increased (Table 5). These albino PLBs did not exhibit any signs of growth or death. The percentage of albinos in the combination treatment was generally lower than in the single EMS treatment, which could be due to PLBs already receiving UV₂₅₄ irradiation treatment, which may affect their increasing tolerance to EMS toxicity, allowing for less depigmentation.

This combination treatment was carried out continuously by exposing PLBs to UV₂₅₄ light, then observing them for 7 weeks before immersing them in EMS, causing PLBs to require time to recover and adapt from cell damage caused by mutagens. Furthermore, it may cause more PLBs to develop chlorosis as a result of ongoing cell damage.



Table 5. Effect of combination treatment of UV₂₅₄ exposure and various EMS concentrations on measured parameters in 7 weeks after planting (WAP) explantsTabel 5. Pengaruh perlakuan kombinasi paparan UV₂₅₄ dan variasi konsentrasi EMS pada parameter terukur pada eksplan berusia 7 minggu setelah tanam (MST)

Treatment Perlakuan	Average Increase in Number of PLBs ¹ Rerata Peningkatan Jumlah PLB ¹	Color ^{1,2} Warna ^{1,2}	Regeneration (%) Regenerasi (%)	Survival Rate (%) Tingkat Survival (%)	Albino (%) Albino (%)
Wild type	14 ± 6.10 ^a	6 ^a	3	70 ± 14.22	2 ± 1.5
UV ₂₅₄ (5' on, 85' off for 7 days) + 0% EMS (solvent control)	10 ± 5.38 ^b	6 ^{ab}	37	82 ± 9.71	7 ± 4.18
UV ₂₅₄ (5' on, 85' off for 7 days) + 0.05% EMS	6 ± 2.99 ^b	4 ^b	10	50 ± 10.57	17 ± 10.36
UV ₂₅₄ (5' on, 85' off for 7 days) + 0.06% EMS	4 ± 1.98 ^{bc}	4 ^b	10	56 ± 10.33	57 ± 17.69
UV ₂₅₄ (5' on, 85' off for 7 days) + 0.07% EMS	5 ± 2.69 ^{bc}	4 ^b	20	62 ± 9.07	43 ± 13.63
UV ₂₅₄ (5' on, 85' off for 7 days) + 0.08% EMS	5 ± 1.95 ^{bc}	4 ^b	0	69 ± 6.32	71 ± 10.94

Note:

1= numbers followed by the same letter in the same column indicate results that are not significantly different, with the Mann-Whitney test ($\alpha = 5\%$); 2 = refers to Table 1.I= Angka-angka yang diikuti huruf yang sama pada kolom yang sama mengindikasi hasil yang tidak berbeda nyata menurut uji Mann-Whitney ($\alpha = 5\%$); 2 = merujuk pada Tabel 1.

The results of this research revealed that EMS 0.06% performed better as compared to EMS 0.07% (in both single treatment and treatment). This can be caused by EMS, which is unstable at 0.07% concentrations, reducing its mutagenic effect. EMS is a water-soluble reagent, but it will be decomposed by water during storage, rendering it unstable (Amini, 2014).

Most Likely Mutant Phenotype Character

Most likely mutant explants at 7 WAP were classified based on their proliferative ability, and after forming plantlets, they were classified based on the visual difference in plantlet phenotype to

the wild type (Table 6; Table 7). In this research, plantlets with characters that are visually different from the wild type are referred to as mutants because they have the potential to become mutants, but the further characterization is needed. At 7 WAP, the UV₂₅₄ treatment (5' on, 85' off for 1 day) produced the highest percentage of mutants with various multiplication abilities. In general, the UV₂₅₄ radiation treatment produced mutants with different leaf characters from the wild type, some EMS treatments produced mutants with visual differences from the wild type, and some combination treatments and EMS produced albino mutants that did not grow (Table 7).

Table 6. Percentage of mutant PLBs with various characteristics of multiplication capability resulting from various mutation induction treatments in 7 weeks after planting (WAP) explants

Tabel 6. Persentase mutan PLBs dengan karakteristik kemampuan multiplikasi dari berbagai perlakuan induksi mutasi pada eksplan berusia 7 minggu setelah tanam (MST)

Treatment Perlakuan	Mutant Multiplication Characteristics			Number of Mutant Jumlah Mutant	Mutant Percentage (%) (Ratio)		
	Karakteristik Multiplikasi Mutan						
	Slow Lambat	Stagnant Stagnan	Fast Cepat				
UV ₂₅₄ (5' on, 85' off for 1 day)	5	10	2	17	45 ($\frac{17}{38}$)		
UV ₂₅₄ (10' on, 80' off for 1 day)	2	4	1	7	23 ($\frac{7}{30}$)		
UV ₂₅₄ (5' on, 85' off for 7 days)	14	2	4	20	27 ($\frac{20}{75}$)		
UV ₂₅₄ (10' on, 80' off for 7 days)	4	0	2	6	33 ($\frac{6}{18}$)		
0.05% EMS	0	2	5	7	32 ($\frac{7}{22}$)		
0.06% EMS	0	2	2	4	17 ($\frac{4}{23}$)		
0.07% EMS	0	0	3	3	14 ($\frac{3}{22}$)		
0.08% EMS	2	1	1	4	19 ($\frac{4}{21}$)		
UV ₂₅₄ (5' on, 85' off for 7 days) + 0.05% EMS	2	1	2	5	22 ($\frac{5}{23}$)		
UV ₂₅₄ (5' on, 85' off for 7 days) + 0.06% EMS	0	3	2	5	18 ($\frac{5}{28}$)		
UV ₂₅₄ (5' on, 85' off for 7 days) + 0.07% EMS	0	3	1	4	16 ($\frac{4}{25}$)		
UV ₂₅₄ (5' on, 85' off for 7 days) + 0.08% EMS	6	5	1	12	35 ($\frac{12}{34}$)		

Table 7. Percentage of mutant plantlets from various mutation induction treatments

Tabel 7. Persentase mutan planlet dari berbagai perlakuan induksi mutasi

Treatment Perlakuan	Mutant Age (WAP ¹)	Mutant Percentage (%) (Ratio ²)	Mutant Characteristic Karakteristik Mutan
	Umur Mutan (MST ¹)		
UV ₂₅₄ (5' on, 85' off for 1 day)		15 ($\frac{2}{13}$)	Wavy leaf edges (Figure 3B)
UV ₂₅₄ (10' on, 80' off for 1 day)		0 ($\frac{0}{2}$)	-
	44	3 ($\frac{1}{37}$)	Leaves bases with purplish color and have rounded roots (Figure 3C)
UV ₂₅₄ (5' on, 85' off for 7 days)		3 ($\frac{1}{37}$)	Elongated and twisted leaves (Figure 3D)
		3 ($\frac{1}{37}$)	Inward-curved leaf shape (Figure 3E)



Treatment <i>Perlakuan</i>	Mutant Age (WAP ¹) <i>Umur Mutan</i> (MST ¹)	Mutant Percentage (%) (Ratio ²)	Mutant Characteristic <i>Karakteristik Mutan</i>
UV ₂₅₄ (10' on, 80' off for 7 days)		3 ($\frac{1}{37}$) 3 ($\frac{1}{37}$) 3 ($\frac{1}{37}$) 25 ($\frac{1}{4}$) 25 ($\frac{1}{4}$)	The leaves are shaped like a heart. (Figure 3F) Explant growth is slower and has a flat leaf tip (Figure 3G) Rounded leaf (Figure 3H) White spots on the leaf (Figure 3I) The yellow color on half of the leaf side and root (Figure 3J) Roots with 3 different colors (Figure 3K)
0.05% EMS		10 ($\frac{2}{21}$)	PLBs that are larger than the wild type (Figure 4B)
0.06% EMS	14	0 ($\frac{0}{2}$)	-
0.07% EMS		0 ($\frac{0}{19}$)	-
0.08% EMS		37 ($\frac{11}{30}$)	Non-growing albino PLBs (Figure 4C)
UV ₂₅₄ (5' on, 85' off for 7 days) + 0.05% EMS		27 ($\frac{3}{11}$)	Non-growing albino PLBs
UV ₂₅₄ (5' on, 85' off for 7 days) + 0.06% EMS	17	0 ($\frac{0}{3}$)	-
UV ₂₅₄ (5' on, 85' off for 7 days) + 0.07% EMS		0 ($\frac{0}{11}$)	-
UV ₂₅₄ (5' on, 85' off for 7 days) + 0.08% EMS		8 ($\frac{3}{37}$)	Non-growing albino PLBs

Note:

1 = Week After Planting; 2 = presented in the percentage of mutants with different phenotypic characters from the wild type from the total number of surviving mutants.

1 = Minggu Setelah Tanam; 2 = disajikan dalam persentase mutan dengan karakter fenotip berbeda dari tipe liar dari jumlah total mutan yang bertahan

The mutants produced by UV₂₅₄ irradiation showed differences in leaf phenotypes when compared to the wild type, with the resulting various forms (Figure 3). The UV₂₅₄ treatment (10' on, 80' off for 1 day) prevented mutants from forming and surviving. Treatment with various EMS concentrations revealed the presence of PLBs that were larger than the wild type as well as the presence of non-growing albino PLBs (Figure 4). The

combination treatment also produced non-growing albino PLBs. Aside from mutagens, factors influencing PLB growth response include differences in PLB characteristics because one orchid contains hundreds to thousands of seeds that are sown and grown together *in vitro* and are not uniform in nature. Seed propagation results in heterozygous offspring (Utami et al., 2022).



The mutants obtained must be selected and further characterized in terms of genetic stability and mutant tolerance to extreme conditions that are not appropriate for their agro climate. The mutants that

have been obtained must first be raised for the flowers that form to be identified, and it is hoped that the mutants that form will also produce unique flowers. The mutations induced in PLBs in this study resulted in




Figure 3. Mutant plantlets obtained from UV₂₅₄ irradiation treatment: wild type (a), mutants from 5' on, 85' off (1 day) treatment (b), mutants from 5' on, 85' off (7 days) treatment (c to i), mutants from 10' on, 80' off (7 days) treatment (j and k).

Gambar 3. Planlet mutan yang didapatkan dari perlakuan iradiasi UV₂₅₄: wild type (a), mutan dari perlakuan 5' menyala, 85' mati (1 hari) (b), mutan dari perlakuan 5' menyala, 85' mati (7 hari) (c sampai i), mutan dari perlakuan 10' menyala, 80' mati (j dan k).




Figure 4. Mutant PLBs obtained from EMS treatment: wild type (a), mutants from 0.05% EMS (b), mutants from 0.08% EMS (c).

Gambar 4. PLBs mutan dari perlakuan EMS: wild type (a), mutan dari 0,05% EMS (b), mutan dari 0,08% EMS (c).

undirected mutations; the results obtained should be subjected to molecular analysis to ensure that mutations occur.

The mutants obtained must be selected and further characterized in terms of genetic stability and mutant tolerance to extreme conditions that are not appropriate for their agro climate. The mutants that have been obtained must first be raised for the flowers that form to be identified, and it is hoped that the mutants that form will also produce unique flowers. The mutations induced in PLBs in this study resulted in undirected mutations; the results obtained should be subjected to molecular analysis to ensure that mutations occur.

CONCLUSION

Mutation induction through UV₂₅₄ irradiation (5' on, 85' off (1 day and 7 days) and 10' on, 80' off (7 days)) produced mutants with phenotype leaf shape and color which were visually different from the wild type (wavy leaf edges, leaves bases with purplish color, elongated and twisted leaves, inward-curved leaf, flat leaf tip, rounded leaf, leaf with white spots, and yellow color on half of the leaf side); 0.05% EMS resulted in PLBs mutant with a visually larger size than the wild type; 0.08% EMS and combination treatments (0.05% and 0.08% EMS for each treatment) resulted in non-growing albino PLBs.

ACKNOWLEDGEMENT

This research was funded by Campus Intellectual Product Business Development Program Grant [*Hibah Program Pengembangan Usaha Produk Intelektual Kampus (PPUPIK)*] Kemendikbud-Ristek 2021 with contract no. 004/SPP-PPM/LPPM-02/DRPM/FTB/IV/ 2021 (on behalf of Dr.rer.nat. Sulistyo Emantoko Dwi Putra). All experiments were conducted in the Faculty of Biotechnology, University of Surabaya.

DAFTAR PUSTAKA

- Afza, H., & Iriawati, N. (2016). Pengaruh Iradiasi Ultraviolet terhadap Multiplikasi Tunas Aksiler dan Kadar Klorofil Anyelir (*Dianthus caryophyllus* L.). *Buletin Plasma Nutfah*, 21(1), 39. <https://doi.org/10.21082/blpn.v21n1.2015.p39-46>
- Amini, M. (2014). Ethyl Methanesulfonate. In *Encyclopedia of Toxicology* (Third, pp. 522–524). Elsevier. <https://doi.org/10.1016/B978-0-12-386454-3.01124-6>
- Arisha, M. H., Liang, B.-K., Muhammad Shah, S. N., Gong, Z.-H., & Li, D.-W. (2014). Kill curve analysis and response of first generation *Capsicum annuum* L. B12 cultivar to ethyl methane sulfonate. *Genetics and Molecular Research*, 13(4), 10049–10061. <https://doi.org/10.4238/2014.November.28.9>
- Avivi, S., Hariyono, K., Hartatik, S., Pertanian, F., & Jember, U. (2022). Analisis Pendugaan Parameter Genetik pada Genotipe Tebu Mutan. *Agriprima : Journal of Applied Agricultural Sciences*, 6(2), 124–134. <https://agriprima.polije.ac.id/index.php/journal/article/view/v6i2-c>
- Castronuovo, D., Sofo, A., Lovelli, S., Candido, V., & Scopa, A. (2017). Effects of UV-C Radiation on Common Dandelion and Purple Coneflower: First Results. *International Journal of Plant Biology*, 8(1), 7255. <https://doi.org/10.4081/pb.2017.7255>
- Chookoh, N., Chiu, Y.-T., Chang, C., Hu, W.-H., & Dai, T.-E. (2019). Micropropagation of Tolumnia Orchids through Induction of Protocorm-like Bodies from Leaf Segments. *HortScience*, 54(7), 1230–1236.



- <https://doi.org/10.21273/HORTSCI13747-18>
- Damayanti, F. (2021). Potensi Pemuliaan Mutasi Radiasi sebagai upaya Peningkatan Variasi Genetik pada Tanaman Hias. *EduBiologia: Biological Science and Education Journal*, 1(2), 78. <https://doi.org/10.30998/edubiologia.v1i2.9300>
- Hadebe, S. T., Modi, A. T., & Shimelis, H. A. (2017). Determination of optimum ethylmethanesulfonate conditions for chemical mutagenesis of selected vernonia (Centrapalus pauciflorus) accessions. *South African Journal of Plant and Soil*, 34(4), 311–317. <https://doi.org/10.1080/02571862.2017.1317851>
- Huang, Y.-W., Tsai, Y.-J., Cheng, T.-C., Chen, J.-J., & Chen, F.-C. (2014). Physical wounding and ethylene stimulated embryogenic stem cell proliferation and plantlet regeneration in protocorm-like bodies of Phalaenopsis orchids. *Genetics and Molecular Research*, 13(4), 9543–9557. <https://doi.org/10.4238/2014.November.1.2.3>
- Li, C., Dong, N., Zhao, Y., Wu, S., Liu, Z., & Zhai, J. (2021). A review for the breeding of orchids: Current achievements and prospects. *Horticultural Plant Journal*, 7(5), 380–392. <https://doi.org/10.1016/j.hpj.2021.02.006>
- Magdalita, P., San Pascual, A., & Villareal, R. (2019). Characterization and Flowering Behavior of Eleven Philippine Native Phalaenopsis Species and Gamma Irradiation Effects on *Phalaenopsis aphrodite*. *Philippine Journal of Science*, 149(S1), 1–10. <https://doi.org/10.56899/149.S1.01>
- Mose, W., Indrianto, A., Purwantoro, A., & Semiarti, E. (2017). The Influence of Thidiazuron on Direct Somatic Embryo Formation from Various Types of Explant in *Phalaenopsis amabilis* (L.) Blume Orchid. *HAYATI Journal of Biosciences*, 24(4), 201–205. <https://doi.org/10.1016/j.hjb.2017.11.005>
- Müller-Xing, R., Xing, Q., & Goodrich, J. (2014). Footprints of the sun: memory of UV and light stress in plants. *Frontiers in Plant Science*, 5(SEP), 474. <https://doi.org/10.3389/fpls.2014.00474>
- Mullins, E., Bresson, J.-L., Dalmay, T., Dewhurst, I. C., Epstein, M. M., Firbank, L. G., Guerche, P., Hejatko, J., Moreno, F. J., Naegeli, H., Nogu , F., S nchez Serrano, J. J., Savoini, G., Veromann, E., Veronesi, F., Casacuberta, J., Lenzi, P., Munoz Guajardo, I., Raffaello, T., & Rostoks, N. (2021). In vivo and in vitro random mutagenesis techniques in plants. *EFSA Journal*, 19(11). <https://doi.org/10.2903/j.efsa.2021.6611>
- Ningrum, E. F. C., Rosyidi, I. N., Puspasari, R. R., & Semiarti, E. (2017). Perkembangan Awal Protocorm Anggrek *Phalaenopsis amabilis* secara In Vitro setelah Penambahan Zat Pengatur Tumbuh α -Naphthaleneacetic Acid dan Thidiazuron. *Biosfera*, 34(1), 9. <https://doi.org/10.20884/1.mib.2017.34.1.393>
- Novak, S. D., Luna, L. J., & Gamage, R. N. (2014). Role of Auxin in Orchid Development. *Plant Signaling & Behavior*, 9(10), e972277. <https://doi.org/10.4161/psb.32169>
- Qosim, W. A., Istifadah, N., Djatnika, I., & -, Y. (2016). Pengaruh Mutagen Etil Metan Sulfonat terhadap Kapasitas Regenerasi Tunas Hibrida *Phalaenopsis*



- In Vitro. *Jurnal Hortikultura*, 22(4), 360. <https://doi.org/10.21082/jhort.v22n4.2012.p360-365>
- Romeida, A., Stujahjo, S. H., Purwito, A., Sukma, D., & Rurstikawati. (2012).  Induksi Mutasi Protocorm Like Bodies (PLB) Anggrek *Spathoglottis plicata* Blume. Akses Bengkulu pada Sebelas Taraf Dosis Iradiasi Sinar Gamma. *Prosiding Simposium Dan Seminar Bersama PERAGI-PERHORTI-PERIPI-HIGI Mendukung Kedaulatan Pangan Dan Energi Yang Berkelanjutan*, 381–387. <https://repository.ipb.ac.id/handle/123456789/59898>
- Romiyadi, R., Komariah, A., & Amien, S.  (2018). Keragaan tiga jenis planlet anggrek *Phalaenopsis* asal Protocorm yang diinduksi Ethyl Methyl Sulfonate (EMS) secara in vitro. *Kultivasi*, 17(1), 596–607. <https://doi.org/10.24198/kultivasi.v17i1.16077>
- Srivastava, D., Gayatri, M. C., & Sarangi, S. K. (2018). In vitro mutagenesis and characterization of mutants through morphological and genetic analysis in orchid *Aerides crispa* Lindl. *Indian Journal of Experimental Biology*, 56(6), 385–394. http://nopr.niscpr.res.in/bitstream/123456789/44540/1/IJEB_56%286%29_385-394.pdf
- Sulistianingsih, R., & Purwantoro, A.  (2012). Variasi genetik anggrek alam *Phalaenopsis amabilis* (L .) Blume hasil iradiasi sinar gamma (Genetic variation of natural orchid *Phalaenopsis amabilis* (L .) Blume produce gamma ray irradiation). *Jurnal Ilmiah Aplikasi Isotop Dan Radiasi*, 8(1), 1–10. <https://jurnal.batan.go.id/index.php/jir/article/download/488/418>
-  Surya, M. I., Ismaini, L., Destri, D., & Normasiwi, S. (2016). An Effort of Mutation Breeding by Oryzalin and Gamma Rays on Wild Raspberry (*Rubus* sp.) in Cibodas Botanical Garden. *Biosaintifika: Journal of Biology & Biology Education*, 8(3), 331. <https://doi.org/10.15294/biosaintifika.v8i3.6559>
-  Talebi, A. B., Talebi, A. B., & Shahrokhifar, B. (2012). Ethyl Methane Sulphonate (EMS) Induced Mutagenesis in Malaysian Rice (cv. MR219) for Lethal Dose Determination. *American Journal of Plant Sciences*, 03(12), 1661–1665. <https://doi.org/10.4236/ajps.2012.312202>
-  Utami, N. R., Rahayuningsih, M., Suwarsi, E., Alighiri, D., & Yuwono, S. (2022). Aklimatisasi Anggrek Species Hasil Kultur Jaringan Melalui Pemberdayaan Masyarakat Dusun Gempol. *Sarwahita*, 19(01), 171–181. <https://doi.org/10.21009/sarwahita.19.1.15>
-  Widiarsih, S., & Dwimahyani, I. (2013). Aplikasi iradiasi gamma untuk pemuliaan mutasi anggrek bulan (*Phalaenopsis amabilis* B1.) umur genjah. *Jurnal Ilmiah Aplikasi Isotop Dan Radiasi*, 9(1), 59–66. <https://jurnal.batan.go.id/index.php/jir/article/view/1202/1147>
- Xia Chen, & Thibeault, S. (2013). Effect of  DMSO concentration, cell density and needle gauge on the viability of cryopreserved cells in three dimensional hyaluronan hydrogel. *2013 35th Annual International Conference of the IEEE Engineering in Medicine and Biology Society (EMBC)*, 6228–6231. <https://doi.org/10.1109/EMBC.2013.6610976>



- Yulianto, T. B., Taufiq, A. J., & Suyadi, A. (2019). Rancang Bangun Pengaturan Intensitas Sinar Uv (Ultraviolet) Dengan Mikrokontroler PIC Untuk Tanaman. *Jurnal Riset Rekayasa Elektro*, 1(1), 54–70. <https://doi.org/10.30595/jrre.v1i1.4929>
- Zahara, M. (2017). A Review: Micropropagation Of Phalaenopsis sp. From Leaf And Flower Stalk Explants. *Jurnal Natural*, 17(2), 91. <https://doi.org/10.24815/jn.v0i0.8130>



AGRIPRIMA

Journal of Applied Agricultural Sciences

VOL.7 NO.1
MARCH 2023



Editorial Team

Editor in Chief

1. Netty Ermawati, SP., Ph.D

Politeknik Negeri Jember, Indonesia



Editors

1. Dwi Rahmawati, SP., MP

Politeknik Negeri Jember, Indonesia



2. Sepdian Luri Asmono, S.ST, MP

Politeknik Negeri Jember, Indonesia



3. Abdurrahman Salim, S.Si, M.Si

Politeknik Negeri Jember, Indonesia



4. Dr. Ir. Didik Pudji Restanto, MP

Universitas Jember, Indonesia



Assistant Editor

1. Putri Santika, S.ST, M.Sc

Politeknik Negeri Jember, Indonesia



2. Anni Nuraisyah, SP, M.Si

Politeknik Negeri Jember, Indonesia



3. Irma Harlianingtyas, S.Si, M.Si

Politeknik Negeri Jember, Indonesia



4. Dian Nur Aini, S.ST

Politeknik Negeri Jember, Indonesia



IT Support

1. Afif Sugi Hendrianto

Politeknik Negeri Jember, Indonesia



AIMS AND SCOPE JOURNAL

AUTHOR GUIDELINES

MAKE A SUBMISSION

EDITORIAL BOARD

OPEN ACCESS POLICY


AUTHOR FEE

PUBLICATION ETHICS

PLAGIARISM CHECK



Vol. 7 No. 1 (2023): MARCH



The Thirteenth Edition of
Agriprima: Journal of Applied Agricultural Sciences.
Published: March 31, 2023
Consists of 9 Articles and 99 Pages.

All content license under:



Creative Commons Attribution ShareAlike 4.0 International License

DOI: [10.25047/agriprima.v7i1](https://doi.org/10.25047/agriprima.v7i1)

[Articles](#)

[Full Issue](#)

 **Published:** Mar 31, 2023

Induction of Protocorm-Like Bodies (PLBs) Phalaenopsis spp. Hybrids Mutation through Ultraviolet Irradiation (UV254) and Ethyl Methane Sulfonate (EMS)

Agatha Sullivania Kurniadi  , Fenny Irawati  , Sulistyo Emantoko Dwi Putra  ,
Poppy Hartatie Hardjo 

1-15

DOI: [10.25047/agriprima.v7i1.512](https://doi.org/10.25047/agriprima.v7i1.512)

Statistics:  248 |  216

 PDF (ENGLISH)**Investigation of In Vitro Apocarotenoid Expression in Perianth of Saffron (*Crocus sativus L.*) Under Different Soil EC**Mandana Mirbakhsh  , Zahra Zahed , Sepideh Mashayekhi , Monire Jafari 

16-24

DOI: [10.25047/agriprima.v7i1.508](https://doi.org/10.25047/agriprima.v7i1.508)Statistics:  215 |  101 PDF (ENGLISH)**Keragaman Genetik Tanaman dan Produktivitas Tebu Klon Unggul Harapan SB 01, SB 03, SB 12 (*Saccharum Officinarum L.*) Di Lahan Kering Sambiroto Mojokerto**Setyobudi , Wiharyanti Nur Lailiyah , Descha Giatri Cahyaningrum  , Rizqa Yuhda Rohmah

25-39

DOI: [10.25047/agriprima.v7i1.494](https://doi.org/10.25047/agriprima.v7i1.494)Statistics:  351 |  195 PDF (INDONESIA)

Pemetaan Kesehatan Kebun Kelapa Sawit Berdasarkan Nilai Normalized Difference Vegetation Index (NDVI) Menggunakan Citra Landsat-8 Di Kebun PT. Wanapotensi Guna

Nur Hadi Ageng Pangestu , Galuh Banowati
40-49

DOI: [10.25047/agriprima.v7i1.513](https://doi.org/10.25047/agriprima.v7i1.513)

Statistics:  590 |  1054

 Citations

1

 [PDF \(INDONESIA\)](#)

Pengaruh Biopestisida Fobio dan Agens Hayati Trichoderma sp., terhadap Penyakit Layu Fusarium pada Bawang Merah

Farisa  , Dita Megasari , Sri Wiyatiningsih 
50-57

DOI: [10.25047/agriprima.v7i1.522](https://doi.org/10.25047/agriprima.v7i1.522)

Statistics:  553 |  353

 Citations

0

 [PDF \(INDONESIA\)](#)

Evaluasi Penambahan Kalium pada AB-Mix Terhadap Pertumbuhan Tiga Varietas Selada (*Lactuca sativa L.*) Hidroponik

Olandino Tome Francisco Dorosario de Sousa , Kacung Hariyono  , Parawita Dewanti 
58-71

DOI: [10.25047/agriprima.v7i1.478](https://doi.org/10.25047/agriprima.v7i1.478)

Statistics:  324 |  197

 Citations

0

 [PDF \(INDONESIA\)](#)

Pengaruh Kehadiran Gulma pada Tanaman Sawi (*Brassica juncea L.*) Terhadap sebelum dan setelah Pemberian Pupuk Limbah Udang

Aditya Murtilaksono  , Ratna Presanthy , Sri Andini Lestari , Muh. Adiwena 
72-78

DOI: [10.25047/agriprima.v7i1.487](https://doi.org/10.25047/agriprima.v7i1.487)

Statistics:  394 |  352



0

 [PDF \(INDONESIA\)](#)

Karakteristik Pupuk Cair Eco-Enzyme Berbahan Dasar Limbah Sayur Dan Buah Terhadap Kandungan Nutrisi Dan Bahan Organik

Ari Istanti , Aldy Bahaduri Indraloka , Sari Wiji Utami

79-85

DOI: [10.25047/agripriama.v7i1.503](https://doi.org/10.25047/agripriama.v7i1.503)

Statistics: 435 | 645



1

 [PDF \(INDONESIA\)](#)

Keterkaitan Umur Panen dan Lama Waktu Curing dengan Produksi dan Mutu Benih Mentimun (*Cucumis sativus L.*) Galur MTH 15

Rahmat Ali Syaban , Suwardi , Sri Rahayu , Indrianingsih

86-99

DOI: [10.25047/agripriama.v7i1.500](https://doi.org/10.25047/agripriama.v7i1.500)

Statistics: 398 | 241



0

 [PDF \(INDONESIA\)](#)

AIMS AND SCOPE JOURNAL

AUTHOR GUIDELINES

MAKE A SUBMISSION


EDITORIAL BOARD

OPEN ACCESS POLICY

AUTHOR FEE

PUBLICATION ETHICS

PLAGIARISM CHECK



3,732 Pageviews
Jun 25th - Jul 25th



	ID	56.062		RU	232
	US	4.393		CA	182
	SG	1.636		NL	124
	CN	457		JP	103
	IN	322		DE	87

Pageviews: 242.794



JOURNAL PUBLISHER

Agriprima is published by Politeknik Negeri Jember and Managed by Department of Agricultural Production. Politeknik Negeri Jember is a higher education institution under the Ministry of Education and Culture of Indonesia.

CONTACT US

- 📞 Phone. 0331 333532 ex.260
- ✉️ Mail. agriprima@polije.ac.id
- 🌐 Web. <https://agriprima.polije.ac.id>

INFORMATION

For Authors

For Readers

For Librarians



AGRIPROMA : JOURNAL OF APPLIED AGRICULTURAL SCIENCES

📍 [POLITEKNIK NEGERI JEMBER](#)

✳️ P-ISSN : 25492934 < E-ISSN : 25492942



0.875
Impact



680
Google Citations




Sinta 3
Current
Accreditation

<> [Google Scholar](#) <> [Garuda](#) [Website](#) [Editor URL](#)

History Accreditation

2017 2019 2020 2021 2022 2023 2024 2025

Citation Per Year By Google Scholar



Journal By Google Scholar

	All	Since 2019
Citation	680	663
h-index	12	12
i10-index	20	19

Garuda

[Google Scholar](#)

[Evaluasi Penambahan Kalium pada AB-Mix Terhadap Pertumbuhan Tiga Varietas Selada \(*Lactuca sativa L.*\) Hidroponik](#)

Politeknik Negeri Jember [Agriproma : Journal of Applied Agricultural Sciences Vol 7 No 1 \(2023\): MARCH 58-71](#)

[2023](#) [DOI: 10.25047/agriproma.v7i1.478](#) [Accred : Sinta 3](#)

[Pengaruh Kehadiran Gulma pada Tanaman Sawi \(*Brassica juncea L.*\) Terhadap sebelum dan setelah Pemberian Pupuk Limbah Udang](#)

Politeknik Negeri Jember [Agriproma : Journal of Applied Agricultural Sciences Vol 7 No 1 \(2023\): MARCH 72-78](#)

[2023](#) [DOI: 10.25047/agriproma.v7i1.487](#) [Accred : Sinta 3](#)

[Keragaman Genetik Tanaman dan Produktivitas Tebu Klon Unggul Harapan SB 01, SB 03, SB 12 \(*Saccharum Officinarum L.*\) Di Lahan Kering Sambiroto Mojokerto](#)

Politeknik Negeri Jember [Agriproma : Journal of Applied Agricultural Sciences Vol 7 No 1 \(2023\): MARCH 25-39](#)

[2023](#) [DOI: 10.25047/agriproma.v7i1.494](#) [Accred : Sinta 3](#)

[Keterkaitan Umur Panen dan Lama Waktu Curing dengan Produksi dan Mutu Benih Mentimun \(*Cucumis sativus L.*\) Galur MTH 15](#)

Politeknik Negeri Jember [Agriproma : Journal of Applied Agricultural Sciences Vol 7 No 1 \(2023\): MARCH 86-99](#)

[2023](#) [DOI: 10.25047/agriproma.v7i1.500](#) [Accred : Sinta 3](#)

[Karakteristik Pupuk Cair Eco-Enzyme Berbahan Dasar Limbah Sayur Dan Buah Terhadap Kandungan Nutrisi Dan Bahan Organik](#)

[Investigation of In Vitro Apocarotenoid Expression in Perianth of Saffron \(*Crocus sativus L.*\) Under Different Soil EC](#)

Politeknik Negeri Jember Agriproma : Journal of Applied Agricultural Sciences Vol 7 No 1
(2023): MARCH 16-24


2023 DOI: 10.25047/agriproma.v7i1.508 Accred : Sinta 3

[Induction of Protocorm-Like Bodies \(PLBs\) Phalaenopsis spp. Hybrids Mutation through Ultraviolet Irradiation \(UV254\) and Ethyl Methane Sulfonate \(EMS\)](#)

Politeknik Negeri Jember Agriproma : Journal of Applied Agricultural Sciences Vol 7 No 1
(2023): MARCH 1-15

2023 DOI: 10.25047/agriproma.v7i1.512 Accred : Sinta 3

Citation Per Year By Google Scholar



[Pemetaan Kesehatan Kebun Kelapa Sawit Berdasarkan Nilai Normalized Difference Vegetation Index \(NDVI\) Menggunakan Citra Landsat-8 Di Kebun PT. Wanapotensi Guna](#)

Politeknik Negeri Jember Agriproma : Journal of Applied Agricultural Sciences Vol 7 No 1
(2023): MARCH 40-49

2023 DOI: 10.25047/agriproma.v7i1.513 Accred : Sinta 3

[Pengaruh Biopestisida Fobio dan Agens Hayati Trichoderma sp., terhadap Penyakit Layu Fusarium pada Bawang Merah](#)

Politeknik Negeri Jember Agriproma : Journal of Applied Agricultural Sciences Vol 7 No 1
(2023): MARCH 50-57

2023 DOI: 10.25047/agriproma.v7i1.522 Accred : Sinta 3

Journal By Google Scholar

	All	Since 2019
Citation	680	663
h-index	12	12
i10-index	20	19

[Pengaruh Dosis dan Waktu Aplikasi Pupuk KCI Terhadap Pertumbuhan dan Hasil Tanaman Buncis \(*Phaseolus vulgaris L.*\)](#)

Politeknik Negeri Jember Agriproma : Journal of Applied Agricultural Sciences Vol 6 No 1
(2022): MARCH 85-97

2022 DOI: 10.25047/agriproma.v6i1.431 Accred : Sinta 3

[View more ...](#)