

Anti-inflammatory Potential of Mangosteen Pericarp Extract (*Garcinia mangostana* L.)

Auw Natasha Felicia Karnadi, Tjie Kok*

Faculty of Biotechnology, University of Surabaya, Jalan Raya Kalirungkut, Surabaya-60293

*Corresponding author: tjie_kok@staff.ubaya.ac.id

Submit: March 27th, 2022

Revised: May 19th, 2023

Accept: June 16th, 2023

Copyright © 2023 Universitas Pakuan



FITOFARMAKA: Jurnal Ilmiah Farmasi is licensed under a Creative Commons Attribution-ShareAlike 4.0 International License

ABSTRACT

Mangosteen pericarp (*Garcinia mangostana* L.) has been used as an active ingredient in numerous traditional medicine formulations. However, their function as an anti-inflammatory agent has yet to be fully understood. Macrophage migration inhibitory factor (MIF) is a mediator that is essential for the development of pro-inflammatory reactions. Hence, MIF activity is often associated with inflammatory diseases and its inhibition could therefore be used as an approach to assess the potential of a material as anti-inflammatory agent. In this study, MIF was exposed with mangosteen pericarp extracts in order to assess the possible anti-inflammatory effect of the extract. The type of mangosteen pericarp extract that possesses the greatest potential as MIF inhibitor and its IC₅₀ value were determined. Based on the results, n-hexane extract of mangosteen pericarp possesses the activity as MIF inhibitor with an IC₅₀ of 4.24 mg/L, and therefore could potentially be used as an anti-inflammatory agent.

Keywords: *Garcinia mangostana* (L); Pericarp extract; Anti-inflammation; MIF

INTRODUCTION

Living mammals respond to inflammation as a local body defense mechanism to prevent the spread of injurious agents (Malleshappa et al., 2018). Several soluble molecular messengers and immune cells are involved in the inflammation process, including cytokines, chemokines, and innate and adaptive immune cells (Nahand et al., 2020). Inflammation should be disabled after the triggering stimulus has been eliminated and the injury has been resolved. However, if the body fails to resolve the inflammation, it may lead to several health problems that involves chronic inflammation.

MIF is a pro-inflammatory cytokine that circulates in human plasma as a homotrimeric protein (Sumaiya et al., 2022). MIF is involved in various biological functions, such as leukocyte recruitment, inflammation, immune responses, cell proliferation, tumorigenesis, and the counter-regulation of glucocorticoids. Due to its

multifaceted nature, MIF plays a central role in the progression of many chronic diseases that involve chronic inflammation, such as cancer, diabetes mellitus, and autoimmune diseases (Purvis et al., 2019; Sumaiya et al., 2022). Several studies have highlighted MIF's role in enhancing the production of inflammatory molecules (IL-1, IL-6, IL-8, nitric oxide, and TNF- α) as a critical factor in the progression of chronic diseases (Cao et al., 2023; Cardoso et al., 2018; Sumaiya et al., 2022), underlining the importance of finding materials that could be used as anti-inflammatory agents by inhibiting MIF activity.

Several studies have suggested that anti-inflammatory potential of a material could be investigated using MIF tautomerase activity inhibition approach (Kok et al., 2018; Nyotohadi & Kok, 2023). MIF enhances the production of inflammatory molecules by interacting with cluster of differentiation 74 (CD74) (Cheng et al., 2022; Cirillo et al., 2020). It is already known

that the interaction site of MIF-CD74 is situated near the p-hydroxyphenylpyruvate tautomerase active site of MIF and the enzyme active site is located at the interfaces of MIF monomer subunits (Wen et al., 2021). It has been identified that the binding of compounds near the active site could disturb the MIF-CD74 interaction (Cirillo et al., 2020; Zapatero et al., 2016).

According to several epidemiological studies, mangosteen pericarp is often used as an active ingredient in traditional medicines in South East Asia for chronic diseases that involve chronic inflammation (Ansori et al., 2020; Li et al., 2020). Mangosteen (*Garcinia mangostana* L.) is one of commonly grown fruit trees in South East Asia countries, such as Thailand, Indonesia, and Malaysia (Ansori et al., 2020; Failla & Gutiérrez-Orozco, 2017; Syahputra et al., 2021). It is reported that the pericarp of mangosteen fruit contains numerous phytochemicals with various health benefits. Some of the phytochemicals, such as garcinone E, gartanin, γ -mangostin, and smeaxanthone A, have been shown to possess anti-inflammatory properties (Gondokesumo et al., 2019; Xu et al., 2017). However, it is not fully understood how mangosteen pericarp exerts its anti-inflammatory effect. Thus, the mangosteen pericarp anti-inflammatory potential was evaluated in this study by its extract inhibition on MIF tautomerase activity. The extracts were evaluated as a whole without further component separation. From the screening process, n-hexane extract of mangosteen pericarp is identified as the one that possess the greatest potential as MIF inhibitor. The extract gave an IC_{50} of 4.24 mg/L, and could probably disrupt MIF pro-inflammatory activity and be used as an anti-inflammatory agent.

MATERIALS AND METHODS

Instruments

A rotary evaporator (Heidolph, Germany) was used to prepare the mangosteen pericarp extracts. The equipments used to produce and purify the recombinant MIF were incubator shaker (Yihder, Taiwan), refrigerated centrifuge (Thermo Scientific, Germany), and ultra

sonicator (Tefic, China). A plate reader (BMG LABTECH, Germany) was used to measure the MIF tautomerase activity.

Chemicals and Reagents

The solvents from Merck (n-hexane, ethyl acetate, and ethanol 99.9%) were used to prepare the mangosteen pericarp extracts. Recombinant MIF was purified using Profinity™ IMAC Ni-charged resin (Biorad). SDS PAGE analysis was conducted with Precision Plus Protein™ dual color standards (Biorad). 4-hydroxyphenylpyruvate (4-HPP) from Difco Laboratories was used as the substrate of MIF. Ampicillin from Sigma Aldrich was used as the antibiotic in the 2YT medium.

Collection of Plant Material

The mangosteen fruits (*Garcinia mangostana* L.) originated in Blitar, East Java, Indonesia. This study used perfectly mature mangosteen fruits with purple and black outer skin. The fruits were authenticated by taxonomist from The Center for Information and Development of Traditional Medicine of the Faculty of Pharmacy, University of Surabaya, Surabaya, East Java, Indonesia.

Mangosteen Pericarp Extracts Preparation

Mangosteen fruits were washed with distilled water and peeled. The pericarp part of the mangosteen fruit was chopped, dried, milled, and sifted. Then the mangosteen pericarp powder was subjected to an extraction procedure using various organic solvents, namely ethanol (99.9%), ethyl acetate, and n-hexane. The extraction was carried out with maceration technique, using 1:10 (w/v) powder:solvent ratio, and incubated for 3 days at 25 °C. The extracts were then evaporated using a rotary evaporator to further eliminate the solvents. Until further use, the concentrated extracts of mangosteen pericarp were stored at 4 °C.

Recombinant MIF Production and Purification

Cultures of *Escherichia coli* BL21 (DE3) pET20b(+)-MIF were prepared by inoculating glycerol stocks into 500 mL of 2YT-ampicillin media (11 g/L yeast extract, 11 g/L NaCl, 22 g/L bacto™ tryptone, 0.1 g/L ampicillin). Then the cultures were incubated at 37 °C until their OD₆₀₀ reached ~ 0.5. The intracellular MIF production by *E. coli* BL21 (DE3) pET20b(+)-MIF was induced by adding isopropyl β-thiogalactopyranoside (IPTG) at a final concentration of 0.05 mM and incubating the cultures overnight (37 °C, 175 rpm).

MIF purification process was started by separating the bacterial cells from the media with centrifugation method (4 °C, 4500 rpm, 15 min). The intracellular MIF was obtained by disrupting the bacterial cells with ultrasonication method (10 cycles at 30% amplitude with 15 s cycle duration and 45 s rest duration) while being cooled in the ice bath. Cell debris was removed by centrifugation and the cell-free extract was incubated overnight with 2 mL of Profinity™ IMAC Ni-charged resin in a gravity flow chromatography column (4 °C, 10 rpm). The chromatography column was then washed with 30 mL of washing buffer (10% glycerol, 50 mM Tris, pH 7.4) to remove the non-bound proteins. MIF was eluted from the chromatography resin with elution buffer (10% glycerol, 50 mM Tris, 500 mM imidazole, pH 7.4). The total protein concentration of each elution fraction was determined using Bradford method. Fractions with high protein concentration were pooled and stored at -80 °C for further use.

SDS PAGE Analysis of Purified Recombinant MIF

SDS PAGE analysis was used to separate and estimate the molecular weight of proteins contained in a solution. It was specifically performed in this study to observe the purity of the obtained recombinant MIF solution. The recombinant MIF solution was first diluted with 1x SDS sample buffer and incubated for 8 min at 60 °C. Together with the protein ladder, sample solution was then loaded onto 18% SDS

gel and electrophoresis was performed in SDS-Tris-glycine buffer system (100 V for 60 min). The gel was then stained with Coomassie brilliant blue R-250 to estimate the molecular weight of the proteins.

Screening of Extracts That Have The Potential as MIF Inhibitor

An assay was carried out in triplicate to screen the extracts with potential as MIF inhibitor. The assay began with the preparation of test and control solutions. Test solutions were made by mixing 45 µL of MIF solution (1.1 µM in boric acid buffer 0.4 M, pH 6.2) with 5 µL of the extract solutions (1346 mg/L for the ethanol extract, 948 mg/L for the ethyl acetate extract, and 257 mg/L for the n-hexane extract; in DMSO). 45 µL of boric acid buffer (0.4 M, pH 6.2) was mixed with 5 µL of DMSO to make the negative control. The inhibitor control was made by mixing 45 µL of MIF solution (1.1 µM in boric acid buffer 0.4 M, pH 6.2) with CuSO₄ solution (50 µM). Meanwhile, the positive control was made as the negative control, except that the boric acid buffer was replaced with MIF solution (1.1 µM in boric acid buffer 0.4 M, pH 6.2). To begin the tautomerase reaction, 50 µL of 4-HPP solution (0.5 mM) was added to a UV-Star® half area 96-well microplate containing test and control solutions. The MIF residual tautomerase activity was observed at 306 nm (Nyotohadi & Kok, 2023). The extracts were then screened on the basis of their MIF residual tautomerase activity percentage. Extract with MIF residual tautomerase activity below 20% was chosen to be subjected to IC₅₀ assay (protocol modified after Nyotohadi & Kok, 2023).

The IC₅₀ Assay

The IC₅₀ assay was conducted in triplicate, using the same volume for test and control solutions as the screening assay, but the test solutions were conditioned to have final extract concentrations of 2.15–8.21 mg/L in 5% DMSO (1.25 fold serial dilution). As in the screening assay, 50 µL of test and control solutions were transferred into a UV-Star® half

area 96-well microplate and added with 50 μ L of 4-HPP solution (0.5 mM) to start the tautomerase reaction by MIF. The tautomerase activity of MIF was measured at 306 nm.

Data Analysis

The analysis of data was conducted using GraphPad Prism 8.0. Absorbance vs. time graph was first constructed. Calculating the ascending slope of absorbance vs time graph gave the reaction rate of MIF tautomerase activity. The residual tautomerase activity was then obtained by normalizing the reaction rate of MIF activity to the controls. Then, the percentage of residual tautomerase activity vs log [extract concentration] was constructed using the curve fitting feature to get the sigmoidal graph and IC₅₀ value of the extract.

RESULTS AND DISCUSSION

Authentication of Plant Material

Center for Information and Development of Traditional Medicine of the University of Surabaya, Surabaya, East Java, Indonesia verified that the collected plant materials were mangosteen (*Garcinia mangostana* L.) fruits.

Mangosteen Pericarp Extracts Preparation

Ethanol (99.9%), ethyl acetate, and n-hexane extracts were produced after the extraction process of mangosteen pericarp. Each extract gave a different yield: 10% is obtained from extraction with ethanol (99.9%), 9% from extraction with ethyl acetate, and 0.5% from extraction with n-hexane.

SDS PAGE Analysis of Purified Recombinant MIF

Figure 1A indicated that recombinant MIF was successfully purified. From the visualization of SDS gel electrophoresis results, it could be seen that the sample solution (SL) only gave one protein band with ~13 kDa molecular weight. The recombinant MIF has a molecular weight of ~13 kDa, hence this process was considered to yield a relatively pure recombinant MIF solution. MIF was reported to take part in the development of several chronic inflammatory diseases by enhancing the production of inflammatory molecules (Cao et al., 2023). Thus the purified MIF solution was then used in further assays as an approach to evaluate the anti-inflammatory potential of the produced mangosteen extracts.

Screening of Extracts That Have The Potential As MIF Inhibitor

The assay was conducted to measure the MIF residual tautomerase activity by detecting the presence of borate and enzymatic enol product of 4-HPP complexes in the solution (Nyotohadi & Kok, 2023). The extract with MIF residual tautomerase activity below 20% was chosen for further assays. The screening assay showed that all types of mangosteen pericarp extracts can inhibit MIF tautomerase activity (Figure 1B), but only n-hexane extract of mangosteen pericarp that gave residual tautomerase activity below 20% (3.25% \pm 0.16%). Thus, the n-hexane extract of mangosteen pericarp (HMP) was chosen for IC₅₀ assay. N-hexane extract of mangosteen pericarp was reported to contain various anti-inflammatory phytochemicals, such as gartanin and β -mangostin (Kameng et al., 2020). Thus, it may have the potential as an anti-inflammatory agent (Gondokesumo et al., 2019).




Figure 1. A is the visualization of SDS-PAGE gel electrophoresis (18% acrylamide), containing protein ladder (L) and sample solution (SL). One protein band (~13 kDa) was formed at SL lane. B is screening assay results of the positive control (MIF), inhibitor control (MIF + inhibitor Cu²⁺), and the test solutions (MIF + EMP (ethanol extract of mangosteen pericarp), MIF + EAMP (ethyl acetate extract of mangosteen pericarp), and MIF + HMP (n-hexane extract of mangosteen pericarp)).

IC₅₀ Assay

In this assay, the MIF residual tautomerase activity was measured for the construction of the sigmoidal graph and determination of IC₅₀ value of HMP extract. The result of the assay could be seen in Figure 2, in which the HMP inhibits MIF tautomerase activity with an IC₅₀ of 4.24 mg/L. The IC₅₀ is an important parameter that is often measured in many pharmaceutical studies (Bag, 2020; Chtita et al., 2021). The IC₅₀ is described as the required concentration of an inhibitor to achieve 50% inhibition of an enzyme (Lei et al., 2020; Liu et al., 2020; Wang et al., 2019). In previous work with multi-strain probiotics extract, an IC₅₀ of 7.80 mg/L was obtained for MIF tautomerase activity inhibition. Meanwhile, in other study, acetaminophen was reported to inhibit MIF tautomerase activity, with an IC₅₀ of 10 mM (Senter et al., 2002). Hence, it was indicated that HMP has the potential as MIF tautomerase activity inhibitor and could possibly be developed as an anti-inflammatory agent.




Figure 2. Sigmoidal graph of the percentage of MIF residual activity vs log [extract concentration] for MIF + HMP. It was shown that HMP gives an IC₅₀ of 4.24 mg/L.

CONCLUSIONS

Mangosteen pericarp has been used as active ingredient in numerous traditional medicines in South East Asia, to treat chronic diseases that involve chronic inflammation. The results of this study indicated the possible anti-inflammatory effect of the mangosteen pericarp extract. It was shown that n-hexane extract of mangosteen pericarp has the potential as MIF tautomerase activity inhibitor, with an IC₅₀ of 4.24 mg/L. Hence, mangosteen pericarp could possibly be developed as an anti-inflammatory agent.

List of Abbreviations: MIF: macrophage migration inhibitory factor, CD74: cluster of differentiation 74, SDS-PAGE: sodium dodecyl sulfate-polyacrylamide gel electrophoresis, 4-HPP: 4-hydroxyphenyl pyruvate, DMSO: dimethyl sulfoxide, EMP: ethanol extract of mangosteen pericarp, EAMP: ethyl acetate extract of mangosteen pericarp, HMP: n-hexane extract of mangosteen pericarp.

Acknowledgements: TK conceived the project, designed and supervised the experiments, and critically reviewed the manuscript; ANFK performed the experiments, analyzed the data

and wrote the manuscript. All authors have read and agreed to the published version of the manuscript.

REFERENCES

- Ansori, A., Fadholly, A., Hayaza, S., Susilo, R., Inayatillah, B., Winami, D., & Husen, S. (2020). A Review on Medicinal Properties of Mangosteen (*Garcinia mangostana* L.). *Research Journal of Pharmacy and Technology*, *13*, 974–982. <https://doi.org/10.5958/0974-360X.2020.00182.1>
- Ansori, A. N. M., Susilo, R. J. K., Fadholly, A., Hayaza, S., Nugraha, A. P., & Husen, S. A. (2020). Antidiabetes type 2 phytomedicine: Mangosteen (*Garcinia Mangostana* L.)-A review. *Biochemical and Cellular Archives*, *20*. <https://doi.org/10.35124/bca.2020.20.S1.3173>
- Bag, A. (2020). DFT based Computational Methodology of IC50 Prediction. *Current Computer-Aided Drug Design*, *17*(2). <https://doi.org/10.2174/1573409916666200219115112>
- Cao, L., Wang, X., Liu, X., Meng, W., Guo, W., Duan, C., Liang, X., Kang, L., Lv, P., Lin, Q., Zhang, R., Zhang, X., & Shen, H. (2023). Tumor Necrosis Factor α -Dependent Lung Inflammation Promotes the Progression of Lung Adenocarcinoma Originating From Alveolar Type II Cells by Upregulating MIF-CD74. *Laboratory Investigation*, *103*(3), 100034. <https://doi.org/10.1016/J.LABINV.2022.100034>
- Cardoso, E. M., Reis, C., & Manzanares-Céspedes, M. C. (2018). Chronic periodontitis, inflammatory cytokines, and interrelationship with other chronic diseases. In *Postgraduate Medicine* (Vol. 130, Issue 1, pp. 98–104). Taylor and Francis Inc. <https://doi.org/10.1080/00325481.2018.1396876>
- Cheng, L., Chang, T. K. H., & Wong, H. (2022). Drug-Drug Interactions With a Pharmacokinetic Basis. *Comprehensive Pharmacology*, 698–715. <https://doi.org/10.1016/B978-0-12-820472-6.00179-1>
- Chtita, S., Belhassan, A., Bakhouch, M., Taourati, A. I., Aouidate, A., Belaidi, S., Moutaabbid, M., Belaaouad, S., Bouachrine, M., & Lakhlifi, T. (2021). QSAR study of unsymmetrical aromatic disulfides as potent avian SARS-CoV main protease inhibitors using quantum chemical descriptors and statistical methods. *Chemometrics and Intelligent Laboratory Systems*, *210*, 104266. <https://doi.org/10.1016/J.CHEMOLAB.2021.104266>
- Cirillo, P. F., Asojo, O. A., Khire, U., Lee, Y., Mootien, S., Hegan, P., Sutherland, A. G., Peterson-Roth, E., Ledizet, M., Koski, R. A., & Anthony, K. G. (2020). Inhibition of Macrophage Migration Inhibitory Factor by a Chimera of Two Allosteric Binders. *ACS Medicinal Chemistry Letters*, *11*(10), 1843–1847. <https://doi.org/10.1021/acsmmedchemlett.9b00351>
- Failla, M. L., & Gutiérrez-Orozco, F. (2017). Mangosteen Xanthones: Bioavailability and bioactivities. In *Fruit and Vegetable Phytochemicals: Chemistry and Human Health: Second Edition* (Vol. 1). <https://doi.org/10.1002/9781119158042.ch9>
- Gondokesumo, M. E., Pardjianto, B., Sumitro, S. B., Widowati, W., & Handono, K. (2019). Xanthones analysis and antioxidant activity analysis (Applying ESR) of six different maturity levels of mangosteen rind extract (*Garcinia mangostana* Linn.). *Pharmacognosy Journal*, *11*(2), 369–373. <https://doi.org/10.5530/pj.2019.11.56>
- Karneng, S., Fadlan, A. & Ersam, T. (2021). Isolation and antidiabetic Activity of Prenylated Xanthones from Pericarp of Mangosteen (*Garcinia Mangostana* Linn.). *IPTEK Journal of Proceedings Series*, (6), 95-101.

- <http://dx.doi.org/10.12962/j23546026.y2020i6.8940>
- Kok, T., Wapenaar, H., Wang, K., Neochoritis, C. G., Zarganes-Tzitzikas, T., Proietti, G., Eleftheriadis, N., Kurpiewska, K., Kalinowska-Thüscik, J., Cool, R. H., Poelarends, G. J., Dömling, A., & Dekker, F. J. (2018). Discovery of chromenes as inhibitors of macrophage migration inhibitory factor. *Bioorganic and Medicinal Chemistry*, 26(5), 999–1005.
<https://doi.org/10.1016/j.bmc.2017.12.032>
- Lei, C., Fu, W., Qian, K., Li, T., Zhang, S., Ding, M., & Hu, S. (2020). Potent neutralization of 2019 novel coronavirus by recombinant ACE2-Ig. *BioRxiv*.
- Li, P., Yang, Z., Tang, B., Zhang, Q., Chen, Z., Zhang, J., Wei, J., Sun, L., & Yan, J. (2020). Identification of Xanthenes from the Mangosteen Pericarp that Inhibit the Growth of *Ralstonia solanacearum*. *ACS Omega*, 5(1), 334–343.
<https://doi.org/10.1021/acsomega.9b02746>
- Liu, L., To, K. K. W., Chan, K. H., Wong, Y. C., Zhou, R., Kwan, K. Y., Fong, C. H. Y., Chen, L. L., Choi, C. Y. K., Lu, L., Tsang, O. T. Y., Leung, W. S., To, W. K., Hung, I. F. N., Yuen, K. Y., & Chen, Z. (2020). High neutralizing antibody titer in intensive care unit patients with COVID-19. *Emerging Microbes and Infections*.
<https://doi.org/10.1080/22221751.2020.1791738>
- Mallesappa, P., Kumaran, R. C., Venkatarangaiah, K., & Parveen, S. (2018). Peels of Citrus fruits: A potential source of anti-inflammatory and anti-nociceptive agents. *Pharmacognosy Journal*, 10(6).
<https://doi.org/10.5530/pj.2018.6s.30>
- Nahand, S. J., Moghoofei, M., Salmaninejad, A., Bahmanpour, Z., Karimzadeh, M., Nasiri, M., Mirzaei, H. R., Pourhanifeh, M. H., Bokharaei-Salim, F., Mirzaei, H., & Hamblin, M. R. (2020). Pathogenic role of exosomes and microRNAs in HPV-mediated inflammation and cervical cancer: A review. In *International Journal of Cancer* (Vol. 146, Issue 2).
<https://doi.org/10.1002/ijc.32688>
- Nyotohadi, D., & Kok T. (2023). Potential of multi-strain probiotics extract as an anti-inflammatory agent through inhibition of macrophage migration inhibitory factor activity. *Functional Foods in Health and Disease*, 13(1), 1.
<https://doi.org/10.31989/ffhd.v13i1.1033>
- Purvis, I. J., Avilala, J., Guda, M. R., Venkataraman, S., Vibhakar, R., Tsung, A. J., Velpula, K. K., & Asuthkar, S. (2019). Role of MYC-miR-29-B7-H3 in medulloblastoma growth and angiogenesis. *Journal of Clinical Medicine*, 8(8).
<https://doi.org/10.3390/jcm8081158>
- Senter, P. D., Al-Abed, Y., Metz, C. N., Benigni, F., Mitchell, R. A., Chesney, J., Han, J., Gartner, C. G., Nelson, S. D., Todaro, G. J., & Bucala, R. (2002). Inhibition of macrophage migration inhibitory factor (MIF) tautomerase and biological activities by acetaminophen metabolites. *Proceedings of the National Academy of Sciences*, 99(1), 144–149.
<https://doi.org/10.1073/pnas.011569399>
- Sumaiya, K., Langford, D., Natarajaseenivasan, K., & Shanmughapriya, S. (2022). Macrophage migration inhibitory factor (MIF): A multifaceted cytokine regulated by genetic and physiological strategies. *Pharmacology & Therapeutics*, 233, 108024.
<https://doi.org/10.1016/J.PHARMTHER.2021.108024>
- Syahputra, M. R., Setiada, H., Siregar, L. A. M., & Damanik, R. I. (2021). Morphological characteristics of Mangosteen plants (*Garcinia mangostana* L.) in Langkat District, North Sumatera, Indonesia. *IOP Conference Series: Earth and Environmental Science*, 782(4).
<https://doi.org/10.1088/1755-1315/782/4/042056>

- Wang, C. Y., Chen, Y. W., & Hou, C. Y. (2019). Antioxidant and antibacterial activity of seven predominant terpenoids. *International Journal of Food Properties*, 22(1).
<https://doi.org/10.1080/10942912.2019.1582541>
- Wen, Y., Cai, W., Yang, J., Fu, X., Putha, L., Xia, Q., Windsor, J. A., Phillips, A. R., Tyndall, J. D. A., Du, D., Liu, T., & Huang, W. (2021). Targeting Macrophage Migration Inhibitory Factor in Acute Pancreatitis and Pancreatic Cancer. In *Frontiers in Pharmacology* (Vol. 12). Frontiers Media S.A.
<https://doi.org/10.3389/fphar.2021.638950>
- Xu, X. H., Liu, Q. Y., Li, T., Liu, J. L., Chen, X., Huang, L., Qiang, W. A., Chen, X., Wang, Y., Lin, L. G., & Lu, J. J. (2017). Garcinone e induces apoptosis and inhibits migration and invasion in ovarian cancer cells. *Scientific Reports*, 7(1).
<https://doi.org/10.1038/s41598-017-11417-4>
- Zapatero, M. C., Pérez, P., Vázquez, M. J., Colmenarejo, G., De Los Frailes, M., & Ramón, F. (2016). Discovery of novel inhibitors of the tautomerase activity of macrophage Migration Inhibitory Factor (MIF). *Journal of Biomolecular Screening*, 21(5), 446-458.
<https://doi.org/10.1177/1087057116633997>

FITOFARMAKA: **Jurnal Ilmiah Farmasi**

Artikel Penelitian

**FORMULASI DAN EVALUASI KOSMETIK DEKORATIF PERONA PIPI DARI
EKSTRAK ANGKAK (*Monascus purpureus*) SEBAGAI PEWARNA
DENGAN MENGGUNAKAN LESITIN SEBAGAI PELEMBAB KULIT**

Anna Yuliana, Lusi Nurdianti, Fitriani Fitriani, Saeful Amin

**UJI AKTIVITAS ANTIBAKTERI SABUN WAJAH EKSTRAK DAUN BELUNTAS (*Pluchea indica* L.)
TERHADAP *Propionibacterium acnes***

Oom Komala, Septia Andini, Fatimah Zahra

**PENGETAHUAN PENGGUNAAN OBAT ANTIBIOTIK PADA MASYARAKAT YANG TINGGAL
DI KELURAHAN BABAKAN MADANG**

Sara Nurmala, Dewi Oktavia Gunawan

**PENGARUH PEMBERIAN EKSTRAK BIJI KOPI ARABIKA (*Coffea arabica* L.)
TERHADAP HISTOPATOLOGI LAMBUNG TIKUS PUTIH GALUR WISTAR**

Tovani Sri, Rani Rubiyanti

**SKRINING TOKSISITAS AKUT BEBERAPA FRAKSI BUAH KARONDA (*Carissa carandas* L.)
PADA EMBRIO ZEBRAFISH (*Danio rerio*)**

Zaldy Rusli, Bina Lohita Sari, Sri Wardatun, Wildan Aristyo

**EVALUASI PEMBERIAN ANTIBIOTIK PADA PASIEN DEMAM TIFOID
DI INSTALASI RAWAT INAP RUMAH SAKIT AZRA KOTA BOGOR**

Dewi Oktavia, Lusi Indriani, Mira Dewi

**EFEK EDUKASI MELALUI BROSUR TERHADAP
KONTROL TEKANAN DARAH DAN KEPATUHAN PASIEN HIPERTENSI**

Emy Oktaviani, Oktaviana Zunnita, Marlia Handayani

**PENGARUH BERBAGAI METODE EKSTRAKSI PADA PENENTUAN KADAR FLAVONOID
EKSTRAK ETANOL DAUN ILER (*Plectranthus scutellarioides*)**

Novi Fajar Utami, Sutanto, Sely Meidi Nurdayanty, Usep Suhendar

**FORMULASI DAN UJI ANTI BAKTERI SEDIAAN GEL
EKSTRAK DAUN MANGGA ARUMANIS (*Mangifera indica* L.)
SEBAGAI ANTIBAKTERI *Staphylococcus aureus* DAN *Propionibacterium acnes***

Prasetyorini, Isna Diana, Dwi Indriati



Editorial Team

EDITOR IN CHIEF

1. **Bina Lohita Sari**, Pharmacy Study Program, Faculty of Math and Science, Universitas Pakuan, Indonesia

EDITORIAL BOARD

1. **Nor Amlizan binti Ramli**, Department of Pharmaceutics, Faculty of Pharmacy, Universiti Teknologi MARA, Malaysia
2. **Enade Perdana Istyastono**, Faculty of Pharmacy, Universitas Sanata Dharma, Indonesia
3. **Sapto Yuliani**, Fakultas Farmasi, Universitas Ahmad Dahlan, Indonesia
4. **Eli Halimah**, Faculty of Pharmacy, Universitas Padjadjaran, Indonesia
5. **Vienna Saraswaty**, Pusat Penelitian Bioteknologi LIPI, Indonesia
6. **Ahmad Najib**, Faculty of Pharmacy, Universitas Muslim Indonesia Makassar, Indonesia
7. **Dadan Rohdiana**, Fakultas Teknologi Pertanian, Universitas Al-Ghifari, Indonesia
8. **Anna Yuliana**, Faculty of Pharmacy, Sekolah Tinggi Ilmu Kesehatan Bakti Tunas Husada, Indonesia
9. **Bustanussalam .**, Badan Riset dan Inovasi Nasional (BRIN), Indonesia

MANAGING EDITOR

1. **Zaldy Rusli**, Universitas Pakuan, Indonesia
2. **Siti Mahyuni**, Pharmacy Study Program, Faculty of Math and Science, Universitas Pakuan, Indonesia

IT SUPPORT

1. **Sulaeman SKom**, Universitas Pakuan

Sertifikat Akreditasi



Vol 13, No 1 (2023)

FITOFARMAKA: Jurnal Ilmiah Farmasi




DOI: <https://doi.org/10.33751/jf.v13i1> (<https://doi.org/10.33751/jf.v13i1>)

Articles

Antibacterial Activity of *Ocimum citriodorum* Leave Extracts Against *Shigella dysenteriae*
(<https://journal.unpak.ac.id/index.php/fitofarmaka/article/view/FJIF.v13i1.5643>)

Oom Komala, Novi Fajar Utami, Saraswati Darmawan




 PDF (<https://journal.unpak.ac.id/index.php/fitofarmaka/article/view/FJIF.v13i1.5643/pdf>) |
1-8

 [10.33751/jf.v13i1.5643](https://doi.org/10.33751/jf.v13i1.5643) (<http://dx.doi.org/10.33751/jf.v13i1.5643>)  Abstract views : 9  views : 10

Inventory of Medicinal Plants The Treatment and Prevention of Malaria in The East Sentani District, Jayapura Regency (<https://journal.unpak.ac.id/index.php/fitofarmaka/article/view/FJIF.v13i1.6396>)

Brechkers Lieske Angruni Tukayo, Rahayu Samalo




 PDF (<https://journal.unpak.ac.id/index.php/fitofarmaka/article/view/FJIF.v13i1.6396/pdf>) |
9-19

 [10.33751/jf.v13i1.6396](https://doi.org/10.33751/jf.v13i1.6396) (<http://dx.doi.org/10.33751/jf.v13i1.6396>)  Abstract views : 17  views : 10

Standardization of Semanggi (*Marsilea crenata* C. Presl.) Leaves from Benowo District, Surabaya for Standardized Herbal Raw Material (<https://journal.unpak.ac.id/index.php/fitofarmaka/article/view/FJIF.v13i1.7459>)


Burhan Ma'arif, Almay Abidzar Elpasha, Arief Suryadinata, Tanaya Jati Dharma Dewi, Novia Maulina, Mangestuti Agil




 PDF (<https://journal.unpak.ac.id/index.php/fitofarmaka/article/view/FJIF.v13i1.7459/pdf>) |
20-30

 [10.33751/jf.v13i1.7459](https://doi.org/10.33751/jf.v13i1.7459) (<http://dx.doi.org/10.33751/jf.v13i1.7459>)  Abstract views : 8  views : 7

Application of Alkyletanolamide Surfactants Based on Nyamplung (*Callophylum Inophyllum* L.) Seed Oil in Cream Base (<https://journal.unpak.ac.id/index.php/fitofarmaka/article/view/FJIF.v13i1.6196>)

Dita Auliya Nurfalah, Erin Ryantin Gunawan, Dedy Suhendra, Wahidah Fajrin

 PDF (<https://journal.unpak.ac.id/index.php/fitofarmaka/article/view/FJIF.v13i1.6196/pdf>) |
31-41




 [10.33751/jf.v13i1.6196](https://doi.org/10.33751/jf.v13i1.6196) (<http://dx.doi.org/10.33751/jf.v13i1.6196>)  Abstract views : 6  views : 6

Anti-inflammatory Potential of Mangosteen Pericarp Extract (*Garcinia mangostana* L.)

(<https://journal.unpak.ac.id/index.php/fitofarmaka/article/view/FJIF.v13i1.7360>)

Auw Natasha Felicia Karnadi, Tjie Kok




 PDF (<https://journal.unpak.ac.id/index.php/fitofarmaka/article/view/FJIF.v13i1.7360/pdf>) | 42-49

 [10.33751/jf.v13i1.7360](http://dx.doi.org/10.33751/jf.v13i1.7360) (<http://dx.doi.org/10.33751/jf.v13i1.7360>)  Abstract views : 15 
views : 10

Formulation of Mouthwash Honje Laka (*Etlingera Hemisphaerica* (Blume) R.M.Sm.) Flower Extract Against *Streptococcus Mutans* (<https://journal.unpak.ac.id/index.php/fitofarmaka/article/view/FJIF.v13i1.6830>)

Ira Rahmiyani, Aneu Nia Rolita Pasa, Saeful Amin, Anna Yuliana




 PDF (<https://journal.unpak.ac.id/index.php/fitofarmaka/article/view/FJIF.v13i1.6830/pdf>) | 50-60

 [10.33751/jf.v13i1.6830](http://dx.doi.org/10.33751/jf.v13i1.6830) (<http://dx.doi.org/10.33751/jf.v13i1.6830>)  Abstract views : 12 
views : 7

Analysis of Urinary Fibronectin Levels in Patients with Type 2 Diabetes Mellitus and Non-Diabetes Mellitus (<https://journal.unpak.ac.id/index.php/fitofarmaka/article/view/FJIF.v13i1.7752>)

Muh Anshar J, Liong Boy Kurniawan, Nurahmi Nurahmi, Andi Alfian Zainuddin, Husaini Umar, Mansyur Arif




 PDF (<https://journal.unpak.ac.id/index.php/fitofarmaka/article/view/FJIF.v13i1.7752/pdf>) | 61-69

 [10.33751/jf.v13i1.7752](http://dx.doi.org/10.33751/jf.v13i1.7752) (<http://dx.doi.org/10.33751/jf.v13i1.7752>)  Abstract views : 13 
views : 7

Pancreatic Protection Effects of Butterfly Pea (*Clitoria Ternatea*) Flower Extract Against White *Rattus Novergicus* Induced By Alloxan (<https://journal.unpak.ac.id/index.php/fitofarmaka/article/view/FJIF.v13i1.6346>)

Farid Fani Temarwut, Suprpto Prayitno, Saenal Edi Kamal, Skolastika Biana Sari, Evi Ayu Hartanti




 PDF (<https://journal.unpak.ac.id/index.php/fitofarmaka/article/view/FJIF.v13i1.6346/pdf>) | 70-76

 [10.33751/jf.v13i1.6346](http://dx.doi.org/10.33751/jf.v13i1.6346) (<http://dx.doi.org/10.33751/jf.v13i1.6346>)  Abstract views : 11 
views : 5

Antioxidant Activity of *Passiflora edulis* (Passion fruit) Seed Extracts Obtained from Maceration and Ultrasonic Assisted Extraction Method (<https://journal.unpak.ac.id/index.php/fitofarmaka/article/view/FJIF.v13i1.7303>)

Aktsar Roskiana Ahmad, Abd. Malik

 PDF (<https://journal.unpak.ac.id/index.php/fitofarmaka/article/view/FJIF.v13i1.7303/pdf>) | 77-81

 [10.33751/jf.v13i1.7303](http://dx.doi.org/10.33751/jf.v13i1.7303) (<http://dx.doi.org/10.33751/jf.v13i1.7303>)  Abstract views : 11 
views : 8

Sertifikat Akreditasi



FITOFARMAKA: JURNAL ILMIAH FARMASI

[UNIVERSITAS PAKUAN](#)

★ [P-ISSN : 20879164](#) < > [E-ISSN : 2622755X](#)



0.914286

Impact Factor



785

Google Citations



Sinta 3

Current Accreditation

[Google Scholar](#) [Garuda](#) [Website](#) [Editor URL](#)

History Accreditation

2018

2019

2020

2021

2022

Garuda

[Google Scholar](#)

[Formulation and Characterization of Transfersomal Containing Breadfruit Leaves \(*Artocarpus altilis* \(Park.\) Fsb.\) Ethanolic Extract](#)

Universitas Pakuan

[FITOFARMAKA: Jurnal Ilmiah Farmasi Vol 12, No 2 \(2022\)](#); [FITOFARMAKA: Jurnal Ilmiah Farmasi 112-121](#)

2022

[DOI: 10.33751/jf.v12i2.4736](#)

[Accred : Sinta 3](#)

[EXTRACTION OF ROBUSTA COFFEE BEANS \(*Coffea canephora*\) FROM WONOSOBO BY ULTRASONIC WAVES AND ANTICANCER TESTS](#)

Universitas Pakuan

[FITOFARMAKA: Jurnal Ilmiah Farmasi Vol 12, No 1 \(2022\)](#); [FITOFARMAKA: Jurnal Ilmiah Farmasi 89-99](#)

2022

[DOI: 10.33751/jf.v12i1.4933](#)

[Accred : Sinta 3](#)

[PENGARUH GASTROPROTEKTOR KOMBINASI TEPUNG GARUT \(*Marantha arundinaceae*\) DAN KUNYIT \(*Curcuma domestica* Val.\) PADA TIKUS MODEL TUKAK LAMBUNG](#)

Universitas Pakuan

[FITOFARMAKA: Jurnal Ilmiah Farmasi Vol 12, No 1 \(2022\)](#); [FITOFARMAKA: Jurnal Ilmiah Farmasi 24-34](#)

2022

[DOI: 10.33751/jf.v12i1.3194](#)

[Accred : Sinta 3](#)

[\(Cover; from the editor; Editorial Team; Table of Contents\)](#)

Universitas Pakuan

[FITOFARMAKA: Jurnal Ilmiah Farmasi Vol 12, No 1 \(2022\)](#); [FITOFARMAKA: Jurnal Ilmiah Farmasi i-iv](#)

2022

[DOI: 10.33751/jf.v12i1.5558](#)

[Accred : Sinta 3](#)

[ANTIOXIDANT AND ANTI-INFLAMMATORY EFFECT OF *Morinda citrifolia*: A META-ANALYSIS AND SYSTEMATIC REVIEW](#)

Universitas Pakuan [FITOFARMAKA: Jurnal Ilmiah Farmasi Vol 12, No 1 \(2022\): FITOFARMAKA: Jurnal Ilmiah Farmasi 35-45](#)

📅 2022 [DOI: 10.33751/jf.v12i1.3359](#) [Accred : Sinta 3](#)

[EFEKTIVITAS ANTIFUNGI MINYAK ATSIRI KENANGA \(*Cananga odorata*\) TERHADAP *Aspergillus flavus*](#)

Universitas Pakuan [FITOFARMAKA: Jurnal Ilmiah Farmasi Vol 12, No 1 \(2022\): FITOFARMAKA: Jurnal Ilmiah Farmasi 14-23](#)

📅 2022 [DOI: 10.33751/jf.v12i1.3476](#) [Accred : Sinta 3](#)

[TOKSISITAS AKUT SEDIAAN CAIR BERBASIS BAWANG PUTIH DENGAN METODE ZEBRAFISH EMBRYO TOXICITY \(ZFET\)](#)

Universitas Pakuan [FITOFARMAKA: Jurnal Ilmiah Farmasi Vol 12, No 1 \(2022\): FITOFARMAKA: Jurnal Ilmiah Farmasi 78-88](#)

📅 2022 [DOI: 10.33751/jf.v12i1.3883](#) [Accred : Sinta 3](#)

[STUDI PENAMBATAN MOLEKUL DAN PREDIKSI ADMET SENYAWA METABOLIT SEKUNDER TANAMAN KELOR \(*Moringa oleifera* L.\) SEBAGAI INHIBITOR BACE1 PADA PENYAKIT ALZHEIMER](#)

Universitas Pakuan [FITOFARMAKA: Jurnal Ilmiah Farmasi Vol 12, No 1 \(2022\): FITOFARMAKA: Jurnal Ilmiah Farmasi 58-67](#)

📅 2022 [DOI: 10.33751/jf.v12i1.4351](#) [Accred : Sinta 3](#)

[VALIDASI DAN PENETAPAN KADAR SENYAWA RUTIN PADA EKSTRAK ETANOL DAUN BINAHONG \(*Anredera cordifolia* \(Ten.\) Steenis\) DENGAN METODE KCKT](#)

Universitas Pakuan [FITOFARMAKA: Jurnal Ilmiah Farmasi Vol 12, No 1 \(2022\): FITOFARMAKA: Jurnal Ilmiah Farmasi 1-13](#)

📅 2022 [DOI: 10.33751/jf.v12i1.3596](#) [Accred : Sinta 3](#)

[THE ACTIVITY OF GEL EXTRACT MAHKOTA DEWA FRUITS \[\(*Phaleria macrocarpa* \(Scheff.\) Boerl\] THROUGH TYROSINASE ENZYME INHIBITOR](#)

Universitas Pakuan [FITOFARMAKA: Jurnal Ilmiah Farmasi Vol 12, No 1 \(2022\): FITOFARMAKA: Jurnal Ilmiah Farmasi 68-77](#)

📅 2022 [DOI: 10.33751/jf.v12i1.4874](#) [Accred : Sinta 3](#)

[View more ...](#)