

CASE REPORT

Identifying the cause of intoxication case without a clear history using flowcharts

Ardyan Wardhana^{1*} 

¹Faculty of Medicine, Universitas Surabaya, Mitra Keluarga Hospital, Surabaya, Indonesia

Article Info

Article history:

Received: 05-01-2023

Revised: 17-02-2023

Accepted: 28-02-2023

Published: 28-05-2023

Keywords:

medical care

intoxication

toxidrome

vital sign

ORCID ID

Ardyan Wardhana

<https://orcid.org/0000-0003-4048-2414>

ABSTRACT

Background: Suspicion of intoxication and its causative agents are required in the clinical practice of intensive care units. Patients with loss of consciousness and no history of leading causation present obstacles in emergency room management. An understanding of toxidromes and epidemiological evidence can help clinicians. However, classic toxidromes are not always presented. **Objectives:** This case was presented to determine the flow of approaches to identify the cause of intoxication and to design early strategies for supportive care in intoxication case management. **Case:** A woman with loss of consciousness accompanied by seizures, deep and rapid breathing, and hypotension. Since the search for collateral information did not produce any meaningful findings, the toxin diagnosis was given based on vital signs, pupil size, and mental status. Methanol intoxication was suspected based on hypothermia, decreased consciousness, normal pupil size, and Kussmaul breathing. Forensic analysis showed her serum was positive for ethanol and methanol. **Conclusion:** When facing a case without the classic presentation of the toxidrome, the toxin diagnosis can be directed based on vital signs, pupil size, mental status, and muscle tone. Introduction to epidemiology is also important in tracking intoxication causation.



Citation:

Wardhana, A. (2023) 'Identifying the cause of intoxication case without a clear history using flowchart', Surabaya Medical Journal, 1(1), p. 35-39. doi: 1059747/smjidisurabaya.v1i1.7

*Corresponding Author:

Ardyan Wardhana Universitas Surabaya/Mitra Keluarga Hospital. Address: Jl. Raya Kalirungkut, Surabaya, Indonesia. Email: ardyanwardhana@staff.ubaya.ac.id

Highlights

1. Intoxication sign such as weakness and blurred vision and unclear causative agent are necessary in clinical practice in the emergency department (ED) for medical treatment and management of the patients.
2. The priority in patients with suspicion of intoxication of initially unknown cause remains the act of resuscitation.



BACKGROUND

Almost 92% of poisoning events are minor, with 79.4% occurring accidentally, and a mortality rate of less than 0.1%. Exposure agents including analgesic agents (11.3%), cleaning agents (7.7%), cosmetics (7.7%), antidepressants (4.4%), and remaining agents are antipsychotic and sedative-hypnotic drugs (5.9%) (Mowry et al., 2015). Death is usually caused by carbon monoxide poisoning, analgesics, sedative-hypnotic agents, antipsychotics, antidepressants, alcohol, and illegal drugs.

Self-inflicted poisoning incidence (suicide attempts, overdoses, intoxications) in the emergency department totaled 3,533 cases during the 2005-2012 period with less than 2% of all cases in the ED. Alcohol was the majority cause (67.3%), with casualties in four cases due to intoxication (Sorge et al., 2015), followed by benzodiazepines (38.3%), analgesics (24.5%), antidepressants (27.2%), antipsychotics (15.7%), opioids (5.5%), and non-opioid banned substances (24.3%) (Ambrosius et al., 2012). However, data on poisoning incidence in Indonesia are still inadequate, especially on intoxicated individuals.

Intoxicated patients sometimes present in loss of consciousness with no causative leading history. Clinicians could become 'blind' to what they are faced with when dealing with intoxication cases or suspected overdoses, especially in cases of infrequent exposure to the substances. As a result, suspicion of intoxication and its causative agents is necessary in clinical practice in the emergency department (ED).

Identification of specific toxic syndromes (toxidromes) is a rational approach to poisoning cases. The introduction of toxidromes can help, but sometimes the symptoms are not specific or masked in other conditions (e.g., myocardial ischemia in cases of carbon monoxide poisoning). The course of the disease can also change during evaluation. For example, antidepressant poisoning initially undergoes several phases of multiorgan failure leading to cardiovascular collapse. An understanding of epidemiology, the flow of approaches to finding intoxication causes, and early strategies for supportive care are very important in managing intoxication cases.

OBJECTIVE

This case report aimed to understand the flow of approaches to identify intoxication causes and early strategies for supportive care in intoxication case management.

CASE

A 26-year-old woman was brought to hospital with complaints of weakness and blurred vision, reported approximately six hours earlier. The patient was still conscious in the ED, but she appeared agitated. When examined by an ophthalmologist, the patient suddenly fainted and was taken to the resuscitation room. Then, the patient had recurrent seizures.

Preliminary studies showed rapid breathing with over 30 breaths per minute, odorless breathing, 97% saturation with 15 liters of oxygen per minute via non-rebreathing mask (NRM), a blood pressure of 76/35mmHg, a pulse of 139 beats/min, and a temperature of 35°C. The physical examination showed an E2V2M4 level of consciousness, 4/4 mm pupils, +/+ light reflexes, clean pulmonary sounds, a non-distended abdomen, and normal bowel sounds.

An electrocardiography examination obtained a tachycardia sine wave pattern. No QRS complex dilation or elevated T waves were found. The ED physicians maintained the airway by tracheal intubation and the provision of mechanical ventilation assistance. Circulation was improved with an infusion of Ringer's lactate solution and gelatin, measuring 500 ml each, supplemented with dobutamine 10-15 mcg/kg/min. The patient was given midazolam 5 mg bolus iv followed by continued doses of 3-5 mg/hour to terminate the seizure.

After initial treatment, the patient's level of consciousness decreased to GCS E1VtM1 and vital signs showed a TD of 98/56 mmHg, a heart rate of 121 beats/min, a breath rate of 28 breaths/min, and spO2 99% on ventilator mode SIMV TV400mL RR22 PS12 PEEP5 FiO2 50% with a minute volume of 10-

13L/min. Post-intubation blood gas analysis still showed severe metabolic acidosis. The minute volume was increased to >13 L/min and given the addition of intravenous sodium bicarbonate measuring 100 mEq within 30 minutes, followed by 100 mEq within 3-4 hours. Hemodialysis was carried out for 3-4 hours and treatment was continued in the high care unit.




Figure 1. The flowchart to identify the cause of intoxication in cases without a clear history

DISCUSSION

Resuscitation remains the priority for patients with suspicion of intoxication with initially unknown causes. These patients come with a loss of consciousness and seizures, thus, intubation is performed. Mechanical ventilation assistance is provided while maintaining hyperventilation as compensation for acidosis. Gas analysis of arterial and/or venous blood is carried out as often as possible to evaluate the pH and ensure the presence of adequate ventilation and oxygenation. Hypotension is initially treated with the administration of a crystalloid bolus of 10-20 ml/kg. However, hypotension in intoxicated patients is rarely due to lack of fluid. Hence, dobutamine is administered according to the suspicion of the toxin type and its effect on hemodynamics (failure of the heart pump vs vasodilation) (Greene, 2019).

Furthermore, the initial assessment is typically in the form of a blood glucose level, vital sign, and mental status assessment. A cocktail consisting of oxygen supplementation, thiamine 100 mg, glucose 25 g, and naloxone 0.4-2.0 mg iv is often taught in literature for cases with loss of consciousness. However, this approach is not cost-effective in developing countries. Patients do not develop hypoglycemia based on rapid measurements of bed-side glucometers, whereas opioid poisoning is quite rare given the limitations of access. There is no hard evidence for thiamine administration before dextrose to prevent Wernicke's encephalopathy (Schabelman and Kuo, 2012). Due to this, cocktails were not administered in this case.

This patient did not show the classic toxidrome presentation; therefore, the authors used a diagnostic approach based on their vital signs, pupil size, mental status, and muscle tone (Pohjola-Sintonen et al., 2000). Mental status assessment can be divided into two sections: excitation/agitative and depressive (Figure 1). Excitation can be in the form of CNS stimulation and increased heart rate, blood pressure, breath rate, and temperature. It is usually caused by anticholinergic agents, sympathomimetics, and hallucinogenic agents. Meanwhile, depressive mental status presents in the form of CNS depression, decreasing blood pressure, heart rate, breath rate and depth, and temperature. It is usually caused by cholinergic agents, sympatholytic agents, opioids, sedative-hypnotic agents, or alcohol (Greene, 2019).

The availability of agents/substances and the incidence rate in an area can narrow the search for toxin types. For instance, salicylates are very rarely used in Indonesia. Organophosphates and alcohols are the most reported agents. Psychiatric drugs are sometimes also abused. Figure 2 shows the possibility of certain drugs being detected in poisoning cases. However, Pohjola-Sintonen et al. (2000) reported a

discrepancy between the history of directing the agent type (collateral information, empty containers, or known prescriptions) in drug poisoning cases and 73% of toxicologically detected drugs, of which 18% were clinically meaningful (Pohjola-Sintonen et al., 2000). Neither the victim's discoverers nor those close to these patients were able to provide collateral information for the identification of the toxin's substance, dosage, route of exposure, and time of exposure. That important information could be covered up because of the fear of being linked to criminal acts. Empty containers/bottles were not reported at the victim's premises. Medications prescribed to patients or medications accessible from family or friends were also disclaimed. A history of psychiatric disorders as well as previous suicide attempts were also refuted. Although it was denied, the use of illicit substances could not be ruled out due to the increase in abuse cases in entertainment venues.




Figure 2. Detected drugs in poisoning cases (Schabelman and Kuo, 2012).

This patient was suspected of drinking alcohol six hours before going to the hospital. It is important to emphasize that the onset of methanol toxicity is delayed when ethanol is also consumed alongside it (Kerns et al., 2002). Ethanol competitively inhibits alcohol dehydrogenase, thereby reducing the formation of toxic metabolites. The possibility of concomitant ethanol and methanol consumption should always be considered, especially in alcoholics who may ingest alcohol in any form.

Based on the findings of hypothermia, non-constricted pupils, and early symptoms of depressive mental status coupled with epidemiological basis, the patient was suspected of alcohol intoxication as well as sedative-hypnotic drugs. The urine drug screening test did not detect amphetamines, benzodiazepines, or cannabis. The blood gas analysis suggested metabolic acidosis with high anion gaps, leading to methanol/ethylene glycol intoxication. Blood samples sent to forensic laboratories suggested the presence of methanol and ethanol.

Strengths and Limitations

Further studies are needed to explore other possibility.

CONCLUSION

When facing a case without the classic presentation of the toxidrome, the toxin diagnosis can be directed based on vital signs, pupil size, mental status, and muscle tone. Introduction to epidemiology is also important in tracking intoxication causation.

Acknowledgment

The authors would like to thank the patient's family for permitting this publication.

Conflict of Interest

All authors have no conflict of interest.

Funding

None.

Author Contribution

The author contributed to all processes in this study, including preparation, data gathering and analysis, drafting, and approval for the manuscript's publication.

Patient Consent for Publication

This case report has been approved by the patient and his/her guardian.

REFERENCES

- Ambrosius, R.G.A., Vroegop, M.P., Jansman, F.G.A., Hoedemaekers, C.W., Aarnoutse, R.E., Van Der Wilt, G.J., Kramers, C., 2012. Acute intoxication patients presenting to an emergency department in the Netherlands: Admit or not? Prospective testing of two algorithms. *Emerg. Med. J.* 29(6):467–472. doi: 10.1136/emj.2010.106500.
- Greene, S., 2019. General Management of Poisoned Patients. In: Tintinalli, J.E. (Ed.), *Tintinalli's Emergency Medicine: A Comprehensive Study Guide*. pp. 1187–1194. Mc Graw Hill, New York, US,
- Kerns, W., Tomaszewski, C., McMartin, K., Ford, M., Brent, J., Bogdan, G., Dart, R., Heard, K., Kokan, L., Kulig, K., Phillips, S., Wells, M., Akhtar, J., Ahsan, N., Burkhart, K., Donovan, J., Zimmerman, H., Curry, S., Wallace, K., Aaron, C., Burns, M., Graudins, C., Hartigan, S., Hantsch, C., Seger, D., Berlin, R., Douglas, D., White, S., Kirk, M., Hollander, J., McKay, C., Wax, P., 2002. Formate kinetics in methanol poisoning. *J. Toxicol. - Clin. Toxicol.* 40(2):137–143. doi: 10.1081/ct-120004401.
- Mowry, J.B., Spyker, D.A., Brooks, D.E., Mcmillan, N., Schauben, J.L., 2015. 2014 annual report of the American association of poison control centers National Poison Data System (NPDS): 32nd Annual Report. *Clin. Toxicol.* 53(10):962–1147. doi: 10.3109/15563650.2015.1102927.
- Pohjola-Sintonen, S., Kivistö, K.T., Vuori, E., Lapatto-Reiniluoto, O., Tiula, E., Neuvonen, P.J., 2000. Identification of drugs ingested in acute poisoning: Correlation of patient history with drug analyses. *Ther. Drug Monit.* 22(6):749–752. doi: 10.1097/00007691-200012000-00016.
- Schabelman, E., Kuo, D., 2012. Glucose before thiamine for Wernicke encephalopathy: A literature review. *J. Emerg. Med.* 42, 488–494.
- Sorge, M., Weidhase, L., Bernhard, M., Gries, A., Petros, S., 2015. Selbstvergiftung in der Akutmedizin – 2005–2012. *Anaesthesist* 64, 456–462.



SURABAYA MEDICAL JOURNAL

Editor-in-Chief

Brahmana Askandar Tjokropawiro

Editorial Boards Coordinator:

Viskasari Pintoko Kalanjati

Secretary Editor:

Arif Tunjung Seto

Editorial Boards:

Ridzky Firmansyah Hardian

Chandrashekhar E. Deopujari

Diah Purwaningsari

Gadis Meinar Sari

Irwanto

M. Ilham Aldika Akbar

Soedarsono Hadipranata

Assistant Editor

Eva Ardianah

Eki Anjangsari

Reviewer :

1. Pradana Zaky Romadhon
2. Nurrizzah Farahiyah Sofia
3. Satriyo Dwi Suryantoro
4. Choirina Windradi
5. Bagus Aulia Mahdi
6. Ida Bagus Subanada
7. I Gusti Ayu Putu Eka Pratiwi
8. I Gusti Lanang Sidiartha
9. Kurniawan Taufiq Kadafi
10. Saptadi Yulianto
11. Ronald Pratama

Surabaya Medical Journal (SMJ) is a scientific journal cover all medical area including basic research in medical, medical management, emergency and critical medicine, medical surgical, mental health, maternity, child health, community and sanitation related to health, health care services, healthcare policy, medical management, nutrition and development, tropical disease, complementary and alternative medicine (CAM), and other fields related published by Ikatan Dokter Indonesia Cabang Surabaya, Indonesia

EDITORIAL ADDRESS:

Ikatan Dokter Indonesia

Mayjen Prof. Dr. Moestopo 117 Street, Surabaya - 60285

East Java, Indonesia Phone: ||62-31| 5938878

E-mail: smj@idisurabaya.or.id

Website: <https://surabayamedicaljournal.or.id> |indonesia



IKATAN DOKTER INDONESIA – SURABAYA SURABAYA MEDICAL JOURNAL

Jl. Mayjen Prof. Dr. Moestopo No. 117, Mojo, Kec. Gubeng, Surabaya, Jawa Timur 60285
Telp. (031) 5938878/ 08123454695 e-mail : smj@idisurabaya.or.id



Editorial Team

Editor in-Chief

Brahmana Askandar Tjokroprawiro, Department of Obstetrics and Gynecology, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia
0000-0003-1658-3477 57217087804

Associate Editor in-Chief

Viskasari Pintoko Kalanjati, Department of Anatomy, Histology and Pharmacology, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia
0000-0002-7005-0025 54388384000

Managing Editor

Arif Tunjungseto, Department of Obstetrics and Gynecology, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia
0000-0003-0079-1394 57217087804

Editorial Boards

Ridzky Firmansyah Hardian, Department of Neurosurgery, Shinshu University, Matsumoto, Japan

0000-0002-2735-8040 57213151036

Chandrashekar Eknath Deopujari, Department of Neurosurgery, Bombay Hospital Institute of Medical Sciences, Mumbai, India

0000-0003-3112-0825 35546316900

Irwanto, Department of Child Health, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia

0000-0002-7573-8793 56705347700

M. Ilham Aldika Akbar, Department of Obstetrics and Gynecology, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia

0000-0002-2003-9282 55676207100

Gadis Meinar Sari, Department of Physiology, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia

0000-0002-9178-8926 57205472893

Soedarsono Hadipranata, Department of Obstetrics and Gynecology, Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia

0000-0000-0000-0000

000000

Diah Purwaningsari, Department of Anatomy and Histology, Faculty of Medicine , Hang Tuah University, Surabaya, Indonesia

0000-0002-9953-6272

57214824615

Assistant Editors

Eva Ardianah
Eki Anjarsari

Sidebar



**MAKE A ✓
SUBMISSION**

Announcements

Focus and Scope

Publication Ethics

Peer Review Process

Article Processing Charge

Open Access Statement

Plagiarism

Meet Our Editorial team



Brahmana Askandar Tjokroprawiro

Editor in-Chief

Universitas Airlangga, Indonesia

57217087804



Viskasari Pintoko Kalanjati

Associate Editor in-Chief

Universitas Airlangga, Indonesia

54388384000



Arif Tunjungseto
Managing Editor
Universitas Airlangga, Indonesia
57217846871

[Read More](#)

Language

Bahasa Indonesia

English

Indexing

Indexed In



Information

For Readers

For Authors

For Librarians



Reference Tools



**Address**

Secretariat of Surabaya
Medical Journal
Jl. Mayjen Prof. Dr. Moestopo No.
117, Surabaya, Jawa Timur

Contact

Phone : 031-5938878
Email : smj@idisurabaya.or.id

Social Media

This journal (p-ISSN:2986-2469; e-ISSN:2986-7584) is licensed under a Creative Commons Attribution 4.0 International License.



IKATAN DOKTER INDONESIA – SURABAYA SURABAYA MEDICAL JOURNAL

Jl. Mayjeri Prof. Dr. Moestopo No. 117, Mojo, Kec. Gubeng, Surabaya, Jawa Timur 60285
Telp. (031) 5938878 / 08123454695 e-mail : smj@idisurabaya.or.id



Vol. 1 No. 1 (2023): Surabaya Medical Journal



Published: 2023-05-27

Front Matter

FRONT MATTER

FRONT MATTER

i-iv



Abstract Views: 19 , File Views: 20

Editorial

Challenge for medical doctor in the future

Challenge for medical doctor

Brahmana Askandar Tjokropawiro

1-4



Abstract Views: 139 , File Views: 94

Original Research Article

The effect of high calorie formula on head circumference and body length on young children with stunted due to feeding problems

High calorie formula and growth parameters

Ni Made Indah Dwijayanti Atmaja, Roedi Irawan

5-13



Abstract Views: 366 , File Views: 85

Clinical profile of patients with COVID-19 at Raden Mattaher General Regional Hospital Jambi

Clinical profile in patients with COVID-19

Olina Novritasari, Sotianingsih, Tia Wida Ekaputri

14-19



Abstract Views: 53 , File Views: 24

Metabolic syndrome, HOMA-IR and adiponectin in obese adolescents

Metabolic syndrome, HOMA-IR and adiponectin in obese adolescents

Nur Aisyah Widjaja, Rendi , Meta Hanindita, Retno, Ugrasena

20-31



Abstract Views: 180 , File Views: 67

Case report/Case series/Image article

Pica in a child with iron deficiency anemia

Pica in a child with iron deficiency anemia

Irwanto Irwanto, Zakky

32-34



Abstract Views: 247 , File Views: 106

Identifying the cause of intoxication case without a clear history using flowcharts

Identifying the cause of intoxication case without a clear history

Ardyan Wardhana

35-39



Abstract Views: 110 , File Views: 33

The safety and efficacy of intravenous thrombolysis for ischemic stroke in an elderly patient with thrombocytopenia

The safety and efficacy of intravenous thrombolysis for ischemic stroke

Ricky Candra Wijaya

40-45



pdf

Abstract Views: 175 , File Views: 65

Scoping review/ systematic review/ meta-analysis

The outcome of percutaneous coronary intervention compared to coronary artery bypass grafting in left main coronary artery stenosis: a systematic review

The outcomes of PCI compared CABG

Faiq Murteza, Chabib Fachry Albab, Almas Talida Habibah, Arya Satya Rajanagara, Zaha El-Ma'i

46-61



pdf

Abstract Views: 298 , File Views: 95

Community Service Report

The implementation of community services of glaucoma counseling in Citra Medika Hospital of Sidoarjo

The implementation of community services of glaucoma counselling

Rini Kusumawardhany, Rani, Raynaldo

62-65



pdf

Abstract Views: 61 , File Views: 42

Viewpoint

Side effect of mRNA-based COVID-19 vaccination in renal: pros- and cons- in pathophysiological view on renal function

Side effect of mRNA-based COVID-19 vaccination in renal

Ari Baskoro, Pranawa Pranawa

66-75



pdf

Abstract Views: 192 , File Views: 39

Back Matter

BACK MATTER

BACK MATTER

77-78



Abstract Views: 20 , File Views: 7

Sidebar



**MAKE A
SUBMISSION**

Announcements

Focus and Scope

Publication Ethics

Peer Review Process

Article Processing Charge

Open Access Statement

Plagiarism

Meet Our Editorial team



Brahmana Askandar Tjokroprawiro

Editor in-Chief

Universitas Airlangga, Indonesia

57217087804



Viskasari Pintoko Kalanjati

Associate Editor in-Chief

Universitas Airlangga, Indonesia

54388384000