



Risk of Vitamin D Deficiency in Smokers: Mix-Method

Amelia Lorensia^{1*}, Marthy Meliana Ariyanti Jalmav², Zahwa Dhiba³, Rivan Virlando Suryadinata⁴

¹Department of Clinical-Community Pharmacy, Faculty of Pharmacy, University of Surabaya, Surabaya, Indonesia

²Petra Christian University, Surabaya, Indonesia

³Faculty of Pharmacy, University of Surabaya, Surabaya, Indonesia

⁴Department of Public Health, Faculty of Medicine, University of Surabaya, Surabaya, Indonesia

*Authors Correspondence: amelia.lorensia@gmail.com; amelia.lorensia@staff.ubaya.ac.id/08155020707

ARTICLE INFO

Article History:

Received Mar, 20th, 2024

Accepted May, 12th, 2014

Published online Jun, 30th, 2024

Keywords:

Smokers;
vitamin D;
deficiency risk;

ABSTRACT

Cigarettes are one of the leading causes of death in the world and Indonesia. Cigarettes emit cigarette smoke, which contains toxins and has pro-inflammatory properties that interfere the metabolism of vitamin D in the lungs. The study aimed to determine the risk of vitamin D deficiency in active smokers in Rungkut Village, Surabaya City. This research method is a mix-method with a Convergent Parallel Design type strategy, which is a way of collecting quantitative (cross-sectional) and qualitative data (interpretative phenomenological analysis) simultaneously, with purposive sampling technique. The variable was the risk of vitamin D deficiency and vitamin D level. Subjects were active smokers aged ≥ 19 years in the Rungkut Subdistrict. The number of samples involved in this research was 125 samples in the quantitative phase, and 15 people in the qualitative phase. The highest level of smoking severity was light smokers (45.60%). Most of respondents had normal vitamin D levels (≥ 20 $\mu\text{g/mL}$) of 73.33%, averaging of 37.66 ± 5.89 $\mu\text{g/mL}$. Most active smokers are at risk of vitamin D deficiency. Risk factors for deficiency in heavy smokers are not consuming foods containing vitamin D (such as fish, eggs, and milk), not consuming fish oil supplements and vitamins, and not sunbathing. However, the level of physical activity tended to be moderate-heavy. The conclusion is that smokers must pay attention to their intake of foods containing vitamin D and exposure to sunlight to achieve sufficient vitamin D levels.

INTRODUCTION

Indonesia is in third place with the most significant number of smokers in the world after the People's Republic of China (PRC) and India. It is estimated that by 2030 the number of smokers in the world and the death rate will reach up to 10 million, of which 70% will come from developing countries. Currently, 50% of the death rate among smokers in the world has come from developing countries.¹ The number of deaths due to smoking is predicted to continue to increase every year because the main ingredient in cigarettes is tobacco.² The main substance in tobacco, namely nicotine, has a relaxing effect, making it addictive for users. The chemical poisons contained in cigarette smoke can be easily transferred from the lungs to the bloodstream and even spread to almost every organ of the human body.³

Systemically, exposure to cigarette smoke can disrupt vitamin D metabolism and damage tubules in the kidneys and liver. Cigarette exposure reduces 25(OH)D and 1,25(OH)2D levels by 10%.⁴ Cigarettes can also cause dryness and shrinkage of the skin, where skin synthesis is the primary source of vitamin D in humans.⁴ Cigarette smoke contains toxic aldehydes that can influence the inflammatory response by increasing the production of pro-inflammatory cytokines.⁵ Additionally, it has pro-inflammatory properties that can weaken host cells in the human body and disrupt the metabolism of vitamin D in the lungs,^{6,7} leading to an accelerated decline in lung function.⁸

In recent years, research has shown an increase in the prevalence of vitamin D deficiency in countries near the equator, where there is abundant exposure to sunlight.⁹

Sun exposure is the most sustainable solution to the widespread issue of vitamin D deficiency, particularly in tropical regions. Despite the abundant sunlight in countries considered a paradox,⁹ like Indonesia, there is a high prevalence of vitamin D deficiency.^{10,11} A Hartung-Knapp-Sidik-Jonkman method meta-analysis revealed that vitamin D deficiency is a significant public health concern in Indonesia.¹² Research by Suryadinata and Lorensia in Indonesia demonstrated a link between vitamin D deficiency and smoking habits.¹³ Respiratory diseases are commonly associated with vitamin D deficiency. This study aimed to assess the risk

of vitamin D deficiency among active smokers in the Rungkut District of Surabaya City, located in the eastern part of Surabaya, with a large population of 833,426 people. Rungkut District is densely populated and includes worker residential areas, such as cigarette factories.¹⁴

MATERIAL AND METHOD

This research is a mixed-methods study that combines quantitative and qualitative data. The design strategy employed is convergent parallel design. The quantitative aspect involves descriptive analysis using a Cross-Sectional Study approach, while the qualitative aspect utilizes Interpretative Phenomenological Analysis (IPA) techniques for data analysis. The study was conducted in Rungkut Village, Surabaya City, from April to July 2023. The research adhered to the code of ethics with No. 160/KE/VI/2023 by Universitas Surabaya, Surabaya.

Smokers are adults who have smoked at least 100 cigarettes in their lifetime and currently smoke every day.¹⁵ Smoking classification based on the Brinkman Index is divided into three categories: light smokers (0-199), moderate smokers (200-599), and heavy smokers (≥ 600). The Brinkman index formula is calculated by multiplying the number of cigarettes smoked daily (sticks) by the number of years of smoking (years).^{16,17} Vitamin D deficiency occurs when the body lacks sufficient vitamin D intake. Most experts consider a 25(OH)D concentration below 25 or 30 nmol/L indicating an increased risk of vitamin D deficiency. A concentration of 50 nmol/L or higher is considered within the normal health range.^{18,19} The study population was adult male active smokers in Rungkut Subdistrict, Surabaya City. The sample included male active smokers aged 19-60 years, with no chronic lung disease, who had been smoking for the last 5 years, and were willing to participate in the research. The total number of active adult smokers in the Rungkut Subdistrict was 125 respondents. Data was collected from residents who attended a health education session at the RW hall facilitated by the local RT head in April 2023. For this quantitative study, the minimum sample size was calculated using the Slovin formula, $n = N / [1 + Ne^2]$ with a sampling error (e) of 5%. The minimum sample size (n) for this research was 95 respondents. The sampling method used was purposive sampling. In the

qualitative phase, data collection continued until data saturation was achieved.

The method used in this research is Mixed-Method, which can simultaneously collect quantitative and qualitative data. The qualitative phase of the mix-method aims to produce a more complete and in-depth picture of the phenomenon. In the initial stage, quantitative methods were used to measure active smokers' vitamin D deficiency risk. These risk factors were described from domains, including smoking history, diet, vitamin/supplement intake, sun exposure, frequency of gadget screens, and physical activity. Data collection began based on adapting the Vitamin D Deficiency Risk Questionnaire.^{4,10,20,21} This research used a questionnaire consisting of 12 closed questions and several open questions. Questions in the vitamin D deficiency risk questionnaire included cigarette use, consumption of foods containing vitamin D (consumption of food from fish, eggs, and milk, which was assessed from the question points in the questionnaire, including the frequency and method of serving food), vitamin D supplements, exposure to sunlight, exposure to gadget radiation, and physical activity. The validity test in this questionnaire only extends to the development of sentences and language because each question item only contains short (closed) questions. In filling out the questionnaire in this study, respondents can fill in each question that has been provided.

Measurement of Vitamin D (25(OH)D) levels in human serum uses VIDAS® 25 OH Vitamin D TOTAL (VIT D), an automated quantitative test employing the ELFA (Enzyme-Linked Fluorescent Assay) technique (Figure 1). VIDAS® 25 OH Vitamin D TOTAL is employed to evaluate Vitamin D sufficiency. A certified laboratory collected 3 ml blood samples from respondents, followed by a blood sample taken at a standardized laboratory in Surabaya.

Quantitative methods are used to obtain measurable data that can be descriptive. The quantitative data in this research is for descriptive analysis. Measuring the percentage value of respondents regarding the risk of deficiency involves using simple statistical data, with the resulting percentage value calculated using the Statistical Package for Social Science (SPSS) program. After the quantitative data collection is complete, qualitative data is gathered in the form of an in-depth examination of the risk factors for vitamin D deficiency in various domains, including cigarette smoking, intake of foods containing vitamin D (such as fish, eggs, milk, and fish oil supplements), vitamin intake, skin exposure to sunlight, screen time on gadgets, and physical activity. Qualitative data is obtained through interviews conducted with the help of an interview guide instrument.

BIOMÉRIEUX

REF 30463 054788 - 04 - 2022-07 

VIDAS® 25 OH Vitamin D TOTAL (VITD)



Intended Use

VIDAS® 25 OH Vitamin D TOTAL (VITD) is an automated quantitative test for use on the instruments of the VIDAS® family for the determination of 25-hydroxyvitamin D Total in human serum or plasma using the ELFA technique (Enzyme Linked Fluorescent Assay).

The VIDAS® 25 OH Vitamin D TOTAL assay is to be used as an aid in the assessment of Vitamin D sufficiency.

Content of the Kit (60 TESTS)

60 VITD Strips ^(A)	STR	Ready-to-use. Stabilizer of human origin*.
60 VITD Solid Phase Receptacles 2 x 30	SPR	Ready-to-use. Interior of SPR device coated with vitamin D.
VITD Calibrator 1 x 1.2 mL (liquid)	S1	Ready-to-use. 25-(OH) Vitamin D diluted in human serum* + preservative. MLE data indicate the calibrator concentration in ng/mL ("Calibrator (S1) Dose Value") and the confidence interval in "Relative Fluorescence Value" (Calibrator (S1) RFV Range).
VITD Control 1 x 1.1 mL (liquid)	C1	Ready-to-use. 25-(OH) Vitamin D diluted in human serum* + preservative. MLE data indicate the confidence interval in ng/mL ("Control C1 Dose Value Range").

Specifications for the factory master data required to calibrate the assay: MLE (Master Lot Entry) barcode printed on the box label.

1 package insert downloadable from www.biomerieux.com

Source: Primary Data, 2023

Figure 1. VIDAS® 25 OH Vitamin D TOTAL (VIT-D) Procedure

RESULTS

This research and data collection was conducted from April 2023 to July 2023 in Rungkut Village. Data were collected by filling in questionnaire questions and conducting interviews with respondents included in the research sample. The number of samples involved in this research was 125. All active smokers were willing to fill out a questionnaire and have their vitamin D levels measured (n=125). However, in the qualitative phase, only 15 respondents were able to be followed up and were willing to continue with in-depth interviews. Based on Table 1, 125 respondents were active smokers, with 65 respondents (52%) starting smoking in adulthood, 22 (40%) having an average normal body weight, and the highest percentage of respondents (92%) having no history of the disease.

Based on Table 1 it showed that 85 respondents (68%) were active smokers with an average of ≥ 10 years of smoking.²³ The maximum number of cigarettes smoked per day was 11-24, with 69 respondents (55.20%). Additionally, 99 respondents (79.20%) smoked filter cigarettes. The Brinkman Index categorized 57 respondents (45.50%) as light smokers.

Based on Table 2, there were 21 respondents (16.80%) who smoked tobacco through direct smoking. Additionally, 108 respondents (86.40%) consumed fish, with 100 respondents (80%) frying it. Moreover, 117 respondents (93.60%) consumed eggs, out of which 115 respondents (92%) used chicken eggs and fried them, as reported by 94 respondents (75.20%). Furthermore, 79 respondents consumed milk (63.20%), and 25 respondents (20.00%) used packaged liquid milk. The highest frequency of respondents who did not consume fish oil supplements was 118 respondents (94.40%). Similarly, the highest frequency of respondents who did not consume vitamins was 73 respondents (58.40%). The main reason respondents consumed vitamins was due to the health benefits, with 45 respondents (36%) citing this reason. The highest frequency of respondents who often looked at gadget screens was 118 respondents (94.40%), with 49 respondents (39.20%) spending more than 7

hours on this activity. Additionally, 98 respondents (78%) were unaware of the ideal time to look at monitor screens (40%). The highest frequency of respondents who rarely engaged in physical activity was 96 respondents (76.80%), with 49 respondents (39.20%) spending more than 7 hours inactive and 96 respondents (76.80%) falling into the moderate activity category (Table 2).

Table 1. Frequency Distribution of Respondents Based on Age, BMI, and Smoking Profiles

Characteristics	n = 125	%
Age (Years)		
19-44 (Adult)	65	52
45-59 (Pre-elderly)	48	38.40
60 (Seniors)	12	9.60
Body Mass Index (BMI)²⁴		
Underweight (<18.5)	12	9.60
Normal (18.5-22.9)	50	40
Overweight with risks (23- 24.9)	25	20
Obesity 1 (25-29.9)	31	24.80
Obesity 2 (≥ 30)	7	5.60
Disease History		
Hypertension	3	2.40
Dyslipidemia	3	2.40
Diabetes Mellitus Type 2	3	2.40
Rheumatoid Arthritis	1	0.80
No History of Illness	115	92
Long Time Smoking Cigarettes (Years)		
< 10	40	32
≥ 10	85	68
Number of Cigarettes Smoked		
1-10 cigarettes per day	47	37.60
11-24 cigarettes per day	69	55.20
≥ 24 cigarettes per day	9	7.20
Types of Cigarettes		
Filter	99	79.20
Kretek	20	16.00
Mixture	6	4.80
Brinkman Index		
Light smoker (0-199)	57	45.60
Medium smoker (200-600)	52	41.60
Heavy smoker (>600)	16	12.80

Source: Primary Data, 2023

Table 2. Distribution Profile of Respondents' Answers

Domain	Question	Respondent's Answer	n = 125	%
Cigarette	Have you ever consumed tobacco directly?	No	104	83.20
		Yes	21	16.80
	How do you use this tobacco?	No	104	83.20
		Inhale directly	21	16.80
		Chewed	0	0
	How do you know this?	Don't know	104	83.20
		Friends/family	13	10.40
		Environment	8	6.40
Fish	Have you ever consumed fish?	Consume	108	86.40
		Do not consume	17	13.60
	What fish do you usually eat?	Mackarel tuna	35	28
		Pindang fish	32	25.60
		Salted fish	25	20
		Milkfish	7	5.60
		Catfish	5	4
		Mujaer fish	4	3.20
	How do you serve the fish you eat?	Fried	100	80
		Burned	5	4
		Steamed	3	2.40
	What is your reason for consuming fish?	Kitchen dishes	40	32
		Love the taste	32	25.60
		Get health benefits	21	16.80
		Do not like	16	12.80
		Affordable prices	10	8
		Diet	5	4
		Allergy	1	0.80
Egg	Have you ever consumed eggs?	Consume	117	93.60
		Do not consume	8	6.40
	What eggs do you usually eat?	Chicken	115	92
		Duck	2	1.60
	How do you serve the eggs you consume?	Fried	94	75.20
		Boiled	23	18.40
	What is the reason you consume eggs?	Love the taste	82	65.60
		Get health benefits	23	18.40
		Kitchen dishes	9	7.20
		Don't consume	7	5.60
		Cheap/affordable prices	3	2.40
		Diet	1	0.80
Milk	Do you consume milk?	Consume	79	63.20
		Do not consume	46	36.80
	What milk do you usually consume?	Packaging (liquid)	25	20
		Condensed milk (sachet)	22	17.60
		Real milk	20	16
		Powder	12	9.60
	What is the reason you consume milk?	Health benefits	50	40
		Love the taste	29	23.20
Fish Oil Supplements	Do you take fish oil supplements?	Do not consume	118	94.40
		Consume	7	5.60
	What type and brand of fish oil supplement do you consume?	Omega-3	4	3.20
		Forgot the brand	3	2.40
	How many fish oil supplements do you consume?	Capsule	7	5.60
		Teaspoon	0	0
		Tablespoon	0	0

Table 2. Distribution Profile of Respondents' Answers

Domain	Question	Respondent's Answer	n = 125	%
	Why do you take fish oil supplements?	Not accustomed to Stench	63	50.40
		Expensive price	40	32
		Health benefits	15	12
			7	5.60
Vitamin	Do you take vitamins?	Do not consume	73	58.40
		Consume	52	41.60
	What type and brand of vitamins do you consume?	Vitacimin®	27	21.60
		Enervon-C®	7	5.60
		Hemaviton®	5	4
		Vitamin D (1000 IU)	5	4
		Imbost®	4	3.20
	The reason you take vitamins?	Don't know	73	58.40
		Health benefits	45	36
		Recommended	4	3.20
		friend/doctor	3	2.40
Changes in Skin Color	Does your skin color change when exposed to direct sunlight?	No	68	54.40
		Yes	57	45.60
	What changes does your skin color look like when exposed to sunlight?	No	40	32
		Blackish red	20	16
Sunbathe	Do you take free time to sunbathe directly?	No	67	53.60
		Yes (eg, tanned)	58	46.40
	What time are you usually exposed to sunlight (sunbathing)?	07.00-09.00	60	48
		10.00-11.00	32	25.60
		12.00-14.00	26	20.80
		15.00-17.00	7	5.60
	Do you sunbathe using personal protective equipment?	Hat	58	46.40
		Jacket	40	32
		Without personal protection	19	15.20
		Sunscreen/sunscreen	6	4.80
		Umbrella	2	1.60
	Do you wear closed clothes?	Shirt and trousers	82	65.60
		Short t-shirt	43	34.40
Gadget Screen	Are you staring at a gadget screen such as a cellphone, laptop or tablet?	Often	118	94.40
		Rarely/not at all	7	5.60
	How long (per day) do you stare at a monitor screen such as a cellphone, laptop or tablet?	≥7 hours	49	39.20
		0-1.5 hours	32	25.60
		4-6 hours	20	16
		2-3 hours	17	13.60
	In your opinion, how long is the ideal time to stare at a monitor screen?	Don't know	98	78.40
		1-2 hours	18	14.40
Physical Activity	Do you often do physical activity?	20-30 minutes	9	7.20
		Seldom	96	76.80
	How often do you do physical activity?	Often	29	23.20
		Light	0	0
		Moderate	96	76.80
		Heavy	29	23.20

Source: Primary Data, 2023

Most of respondents had normal vitamin D levels (sufficiency) of 73.33%, with an average vitamin D level of 37.66 ± 5.89 mg/mL. The results of the cross-tabulation between the description of vitamin D risk factors and vitamin D levels can be seen in Table 3.

When conducting interviews regarding the risk of vitamin D deficiency, 15 respondents out

of 125 respondents were willing to conduct further interviews. The researcher then made interview transcripts and summarized them in table form as well as interview excerpts to further understand the interview themes so that they were in accordance with the research title (Table 4).

Table 3. Cross Tabulation Between the Risk of Vitamin D Deficiency and Vitamin D Levels

Risk of Vitamin D Deficiency		Vitamin D Serum Category ($\mu\text{g/mL}$) ³¹			Total
Domain		Deficiency ($<12\mu\text{g/mL}$) (n = 7)	Insufficiency ($12\text{-}20\mu\text{g/mL}$) (n = 26)	Sufficiency ($>20\mu\text{g/mL}$) (n = 92)	
Cigarette (Brinkman Index)²⁴	Light smoker (0-199)	0	4	53	57
	Medium smoker (200-600)	2	12	38	52
	Heavy smoker (>600)	5	10	1	16
Fish	Consume fish that contain high levels of vitamin D	3	10	52	65
	Consume fish that do not contain high levels of vitamin D	2	7	34	43
	Do not consume fish	3	9	5	17
Egg	Consume egg	1	25	91	117
	Do not consume egg	6	1	1	8
Milk	Consume milk	2	9	68	79
	Do not consume milk	5	17	24	46
Fish Oil Supplements	Consume fish oil	4	24	90	118
	Do not consume fish oil	3	2	2	7
Vitamin	Vitamin C	0	1	36	37
	Vitamin D	0	0	5	5
	Immunomodulator	0	2	2	4
	Not taking any vitamins	7	23	49	79
Changes in Skin Color	Yes	3	13	41	57
	No	4	13	51	68
Sunbathe	Yes	4	11	43	58
	No	3	15	49	67
Gadget Screen	Often	4	24	90	118
	Rarely/not at all	3	2	2	7
Physical Activity	Light	0	0	0	0
	Moderate	4	14	78	96
	Heavy	3	12	14	29

Source: Primary Data, 2023

Table 4. In-depth Interview Results

Domain	Findings	Answer Findings	Quotes From The Interview	Source
Smoke	Respondents knew the severity of smoking	Frequency of cigarette use by respondents	"I've been smoking for more than 15 years, so what's strange is that I can't really stop. Just stop when you cough, then try again with one cigarette, if you cough, throw it away and change to another brand of cigarette, and so on, sis."	Respondent 7
			"Usually I use official cigarettes from factories such as salt warehouses and every day I use up more than 20 cigarettes. I've been smoking for more than 15 years."	Respondent 1
			"I use cigarettes and the brand varies depending on the price, but I usually use 1 pack of solar a day, and I smoked when I was single, around 25 years after graduating from high school, sis."	Respondent 9
			"You don't know how to smoke, sis, sometimes Surya sometimes uses DJI Samsoe. "I've been smoking since I was 18 years old, sis, and now I'm 52 years old."	Respondent 11
	Respondents' views on cigarette use	Views of cigarettes on health by respondents	"For example, if I don't smoke, sis, I'm stressed because my job is uncertain. So, if I smoke, it just feels calmer in my mind, sis."	Respondent 2
			"I don't smoke, or if I just stop for a day, I'll get a headache, but if I smoke just one cigarette, my body will immediately feel better, sis, so I think that smoking isn't completely dangerous because according to my father, cigarettes are medicine."	Respondent 4
Tobacco	Respondents know their knowledge and behavior regarding tobacco use	Tobacco use by respondents	"Previously, I used to smoke tobacco straight away, sis, because in the past tobacco was cheaper than cigarettes, and I knew how to do it when my friends told me to use tobacco."	Respondent 13
			"In the past, I used tobacco directly, sis, by smoking it. "I knew about tobacco because I was told by a friend and forced by a friend to try it. After a while, I became addicted to it and then I tried to replace it with cigarettes."	Respondent 8
			"When I'm in a crisis, I usually use tobacco, sis, because it's cheaper and the way to do it is to smoke it directly. I know that method, I was given tips from my father for using tobacco too."	Respondent 5
Fish	Respondents know the behavior and amount of fish consumed	Respondents explained their fish consumption habits	"I like fish but I don't know how to eat it, sis, sometimes once a week, often twice at most. "I like it when the fish is fried, but more often I like to eat fish steamed in banana leaves."	Respondent 6

Table 4. In-depth Interview Results

Domain	Findings	Answer Findings	Quotes From The Interview	Source
	Respondents know their views on fish consumption habits	Respondents explained the benefits of consuming fish	"I like to eat fish, but I can't guarantee that it will be every day depending on my wife's business in the kitchen, sis, usually once a week."	Respondent 8
			"Actually, there is no benefit from fish, it's just a complement to food and just increases your appetite, sis."	Respondent 15
			"I like fish, sis, because the benefits of fish are that it is delicious as a complement to food and makes you full, sis."	Respondent 13
			"Because you really like eating fish because the benefits are actually good for your health, sis."	Respondent 11
Egg	Respondents' views on behavior and habits of consuming eggs	Respondents explained their egg consumption habits and views on eggs	"I like eggs, sis, but they contain diabetes, so at my age I cut back on them, sis. So I just eat boiled egg whites and discard the yolks and I eat them often every day, usually mixed with vegetables, because my wife makes that menu, sis."	Respondent 7
			"I like eggs, but I usually eat salted eggs when they are served because I like them. Eggs are good for the strength of your voice and are good for your health."	Respondent 3
			"I also often eat fried eggs, usually twice a week, and now I've switched to salted eggs, sis. "The reason is that I like eggs and they have good health benefits, but if the portion is too big, it's not good, sis."	Respondent 11
			"Eggs are just a food supplement, so they have no health benefits."	Respondent 2
Milk	Respondents understand the behavior and benefits of milk	Respondents explained their milk drinking habits and views on the benefits of milk	"Sometimes I drink Frisian flag milk twice a week and if I drink too much I'm afraid because it contains sugar, sis. I actually prefer whole milk but I don't buy it because it's more expensive, sis. And I think the benefits of milk are good for health and increase energy."	Respondent 5
			"I also drink Anlene milk 3x a week because of its benefits for bones and usually relieves fatigue."	Respondent 11
	Respondents' views on milk consumption behavior	Respondents explained the	"I consume milk almost every day, sis, and usually I consume canned milk. "Drinking milk is just for warming the body and the benefits are only for warming the body, sis."	Respondent 12
			"I also consume the milk from Indomaret, usually it's the Frisian flag	Respondent 9

Table 4. In-depth Interview Results

Domain	Findings	Answer Findings	Quotes From The Interview	Source
	behavior and benefits of milk	behavior and benefits of consuming milk	brand which is low fat, but that's rare, sis, especially if you want it, sis. "It has good health benefits because it contains lots of vitamins."	
Supplement oil fish	Respondents knew the behavior and benefits of fish oil supplements	Views on fish oil supplements by respondents	"I don't take supplements because I don't know the benefits and I don't have the time, sis, because I think I already get enough vitamins from vegetables and fruit."	Respondent 8
		Views on the taste and benefits of oil supplements by respondents	"I don't like drinking fish oil because it tastes fishy even though it has benefits for increasing appetite."	Respondent 1
			"I know that it is useful for stamina, but I don't like taking fish oil supplements because I don't like the fishy taste."	Respondent 5
			"I don't take fish oil supplements because I have a stomach ailment, sis, so I don't like anything strange, especially fish oil supplements because it's fishy and makes me nauseous, even though the benefits are actually good for the body, the effect is only short-lived, sis."	Respondent 7

Source: Primary Data, 2023

DISCUSSION

The results obtained from filling out a questionnaire by 125 respondents were followed by interviews with 15 respondents regarding the risk of vitamin D deficiency in active smokers. This research focuses on exploring and exploring the risk of vitamin D deficiency, starting with the domain regarding smoking, the domain regarding sources of vitamin D, the sunlight exposure domain, the radiation domain, and the physical activity domain.

Based on the data obtained, 125 respondents were active smokers, most of whom smoked for more than 10 years, with the highest number of cigarettes smoked in a day ranging from 11-24 cigarettes, so it can be seen that active smokers in Rungkut Village can be categorized as light smokers, with a respondent frequency of 57 respondents. This relatively high number of smokers is in line with previous research by Salsabila et al.²⁵ Central Java, Yogyakarta, East Java, Bali, NTB, South Kalimantan and South Sulawesi), there were 12,591 research subjects. Based on age, smokers in Indonesia have a median of 38 years. Based on gender, the

majority of smokers in Indonesia are men, namely 11,908 people (95%). The type of cigarette most frequently used is filtered kretek cigarettes with an average of 12 cigarettes/day. Indonesia is the country with the highest number of smokers in ASEAN. In South Asia, smoking tobacco (25.2%) and smokeless tobacco (24.7%) is a widespread habit among men. Data from a nationally representative cross-sectional study (Demographic and Health Survey) conducted in Afghanistan, India, Maldives, Nepal and Pakistan from 2015–2018. The subjects were men between 15 and 49 years old. Results found that higher age, lower education, lower wealth status, and involvement in any occupation were strongly associated with smoking (p-value <0.001).²⁶

Cigarette smoke has pro-inflammatory properties which can weaken host cells in the body and can interfere with the metabolism of vitamin D in the lungs. This is supported by previous research which shows that cigarette smoke can disrupt the potential pathways of the endocrine system so that it can trigger vitamin D deficiency.⁶ Previous research by Yang et al., shows smokers have lower vitamin D

concentrations than non-smokers. So, smokers can trigger a more significant vitamin D deficiency than non-smokers.⁴ Based on interviews conducted, respondents were asked for their reasons for smoking cigarettes. As for respondents who said that cigarettes are something that can relieve and reduce pain or have a calming effect on users (Table 4). From the results of the quote above, several respondents experience stress and even dizziness if they do not smoke and think that cigarettes can relieve stress or dizziness. This is supported by previous research that cigarettes can have a positive impact by eliminating stress and dizziness and providing a sense of calm against disturbing thoughts. The nicotine content in cigarettes gives users peace of mind.²⁷

In general, conventional tobacco can be used as smoked tobacco or smokeless tobacco. Apart from cigarettes, tobacco is consumed in other forms such as loosely chewed tobacco leaves, snus, naswar, gutka tobacco, and tobacco paste. Scientific analysis of these smokeless tobacco products has revealed more than 20 chemical compounds known to be cancer-causing agents, which include tobacco-specific nitrosamines, N-nitrosamine acids, volatile N-nitrosamines, polycyclic triggering hydrocarbons, and aldehydes.²⁸ There are several interview quotes above that the price of using tobacco is cheaper than the price of cigarettes from the factory. Based on previous research, tobacco prices are more affordable because the weather makes tobacco production in Indonesia uncertain. If the quality of the tobacco is not good, the price of the tobacco will be sold cheaper.²⁹ Besides the price of tobacco being cheaper than factory cigarettes, rising cigarette prices will influence the high prevalence in Indonesia. This is because cigarette consumption, especially among adults, is greatly influenced by cigarette prices. The increase in cigarette prices is a preventative effort carried out by the government to reduce cigarette consumption, which currently has the highest number of smokers in the Asian region.²²

The results showed most respondents consumed fish such as tuna, pindang fish, salted fish, milkfish, catfish, and tilapia fish. The most common way of serving fish is by frying rather than grilling or steaming. It can be seen from the interview excerpt above that respondents 6, 8, 15, 13, and 11 mostly consume fish even though

some respondents have inconsistent views regarding the benefits of consuming fish (Table 4). Fish is one of the biggest contributors to vitamin D intake, such as catfish, milkfish, and snapper, which contain around 400 IU per 85 grams of fish. The frequency of fish consumption can affect the concentration of vitamin D, so frequent fish consumption can be recommended to prevent vitamin D deficiency. The process of cooking fish does not affect the vitamin D content in food products because vitamin D is resistant to heat, such as by grilling. However, the amount of vitamin D will decrease by 50% if frying uses cooking oil due to the fat-soluble nature of vitamin D.³⁰ Different cooking processes (fried, boiled, grilled) will affect the contents of the fish. Changes in the amount of vitamins and protein are significantly higher when fried. The change in the amount of fat will be higher in frying fish than in boiling or grilling. It was stated in this research that the cooking method by boiling is the best for healthy food.³¹ A previous study by Sridonpai et al., determined the vitamin D content of commonly consumed fish in Thailand and the effect of different cooking methods on vitamin D retention.³² The results showed that vitamin D3 was the only form of vitamin D detected in fish. The vitamin D content of raw freshwater fish ranges from 2.42 to 48.5 µg per 100 g Edible Portion (EP), higher than that of raw sea fish (2.94 to 4.69 µg per 100 g EP).³² Previous research by Andrade et al., a cross-sectional online study, was conducted among 1,637 adults.³³ Participants were considered vitamin D deficient and consumed an average of 347.05 ± 307.8 IUs of vitamin D through foods/beverages. Overall, though, participants, on average, did not meet the vitamin D RDAs (Recommended Daily Allowances) for healthy people. This is mainly related to participants rarely or never consuming foods high in vitamin D such as fish and dairy products.

The results show that most respondents consume eggs such as chicken and duck. The most common way of serving fish is frying compared to boiling (Table 4). On average, respondents consume eggs and are aware of their habits regarding egg consumption. Additionally, respondents understand the benefits of consuming eggs. Eggs are a rich source of protein containing ample amounts of

vitamin D and antioxidants, which help prevent aging by aiding in cell degradation. Vitamin D-rich eggs are found in egg yolks at 20 IU per yolk. The cooking process significantly affects the vitamin D content in eggs. According to USDA data from 2015, raw and boiled eggs contain 41 IU/50 grams of vitamin D3, while fried eggs contain 40 IU/46 grams of vitamin D3.³⁴ Eggs are among the few rich dietary sources of vitamin D. The results of the study by Vičič et al., revealed that fortifying eggs and milk (including yogurt) could effectively increase vitamin D intake in Slovenian women aged 44 to 65 by almost fivefold, thereby reducing the prevalence of vitamin D deficiency.³⁴ However, a previous study by Daly et al., evaluated the dose-response effect of consuming 2, 7, or 12 commercially available eggs per week on serum 25(OH)D concentrations. The results indicated that consuming 7 eggs per week for 12 weeks effectively reduced the decline in circulating vitamin D concentrations during winter in young Australian adults, with 12 eggs per week not providing any additional benefit.³⁵

Based on the results of the questionnaire answered by respondents, it was found that 79 respondents consumed milk most frequently, using liquid packaged milk (ready to eat) and condensed milk (sachets) (Table 4). Respondents consume milk such as ready-to-eat packaged milk. Milk is a fortified product containing vitamin D. Fortified milk must contain 400 IU of vitamin D per litre, whereas pure cow's milk only has 40 IU per litre. Therefore, the best milk is milk with vitamin D fortification in the form of ready-to-eat liquid packaged milk. This research is also similar to other research by Lorensia et al., who analyzed the intake of foods containing vitamin D in 100 adults, showing that eggs and milk contain vitamin D that are most consumed almost every day (98.50%). The incidence of vitamin D deficiency is widespread worldwide, affecting individuals of all ages, including young and healthy subjects, as well as pregnant and elderly women, due to various factors such as inadequate sun exposure, skin pigmentation, adiposity, lifestyle, and low food intake.³⁰ To address this issue, fortification of daily consumed foods, such as milk, is highly recommended.³⁶ A theoretical model applied vitamin D3 fortifications of 1 µg, 1.5 µg and 2

µg/100g to simulate improvements in vitamin D intakes. Mean $\pm$ SD vitamin D3 in whole milk was 0.06 ± 0.02 µg/100g.³⁷ Previous research by Zhang et al., explored interaction between passive smoking and dietary undernutrition on the risk of vitamin D deficiency. The interaction between passive smoking and inadequate nutrition influences vitamin D risk D deficiency.³⁸ Passive smoking may amplify the effect of poor nutrition on the risk of vitamin D deficiency. The more often cigarettes one consumes, nicotine content enters CNS and CNS, putting an individual's food intake.³⁹ Research has been conducted on consuming foods containing vitamin D on vitamin to prevent deficiency,³⁰ but there has been no research linking the impact of active on the population and population, and no research no assessed that influences influencing levels through the intake of foods containing vitamin D.

Based on the results obtained, the highest frequency of respondents was found among those who did not consume fish oil supplements. Some reasons for not consuming fish oil supplements included being unaccustomed to them and disliking the fishy smell or taste associated with them (Table 4). Respondents elaborated on their reasons for not taking fish oil supplements, leading to most of respondents not taking supplements containing vitamin D. According to the results, the highest frequency of respondents was among those who did not consume vitamins (Table 4). The majority of respondents, as per the interview excerpt, do not take vitamins, despite some being aware of the benefits they offer to the body. Previous research by Lorensia and Suryadinata,⁴⁰ indicated that many individuals are hesitant to use fish oil supplements independently due to the large size of the soft capsules, which can be uncomfortable to swallow, and the fishy odor. However, this research also demonstrates that fish oil supplements are effective in enhancing lung function in smokers.⁴⁰

The questionnaire results showed that as many as 68 respondents experienced color changes when exposed to sunlight, with the most frequent color changes occurring in blackish red. From the interview excerpts of the two respondents above, the respondents experienced changes in skin color when exposed to sunlight. This is supported by research

conducted by Carlberg,⁴¹ skin color will change to pink or even black if the skin has received enough vitamin D. As for several other respondents who experienced something different from the previous respondents, respondents 3 and 9 did not experience changes in skin color when exposed to sunlight (Table 4). The respondent did not experience changes in skin color. The darker a person's skin color, the longer it takes to form vitamin D in the skin. According to Raymond-Lezman and Riskin,⁴² in their research, they explain that someone with dark skin has large amounts of melanin in the epidermis, which will make melanin oppose vitamin D receptors in the skin. Melanin will absorb UVB light exposure, reducing the absorption of UVB light exposure for vitamin D receptors in the skin. A person with dark skin takes 10-50 times more exposure to sunlight than a person with light skin to produce the same amount of vitamin D.⁴²

The results of the questionnaire showed that 67 respondents who took the time to sunbathe directly, which was mostly found at 07.00-09.00 WIB and by using personal protective equipment such as hats/jackets and long clothes (Table 4). Many respondents spend time sunbathing, but when the intensity of sun exposure is low, many respondents also use personal protective equipment and long clothing, which is at risk of experiencing a deficiency. Respondents took time to sunbathe between 07.00-09.00 WIB and wore personal protective equipment and closed clothing. Previous research explained that the UVB intensity of sunlight is low at 07.00-08.00 WIB, while at 09.00 WIB, the intensity achieved increases, but it takes a long time of around 25 minutes to sunbathe directly. A good time to be exposed to sunlight is from 10 AM to 2 PM, because at that time UVB rays are more than UVA with a relatively stable and high intensity. However, to be able to see the adequacy of sun exposure not only at the time of exposure, the use of long clothing and personal protection also has an influence. So, it can be concluded that someone who has insufficient sun exposure and has body protection such as sunscreen, hats, jackets and long clothes will be at risk of vitamin D deficiency or vitamin D deficiency.⁴³

From the results obtained, almost all respondents use gadget screens frequently

(Table 4). Respondents use gadget screens for a long time, up to 7 hours or more. In the research of Shan et al.⁴⁴ the use of gadget screens is associated with physical activity on vitamin D concentrations. The time of gadget screen use has a significant relationship with a person's level of physical activity. So, screen use and physical activity are both markers of exposure to UVB rays. This is also supported by research by Rocka et al.⁴⁵ that there is a significant relationship between gadget screen use and physical activity on a person's vitamin status. A good duration of time to stare at a gadget screen is ≤ 2 hours/per day.

Based on the results obtained, the highest frequency of respondents was 96 respondents who carried out moderate levels of physical activity (Table 4). Respondents carried out moderate levels of physical activity such as doing housework as explained by respondents 10, 3, 13, and 14, while respondents 4 and 8 carried out heavy levels of physical activity. Vitamin D levels in the blood are related to muscles and bones and play a role in energy metabolism, oxidative stress, maintenance and increasing physical fitness.⁴⁶ The measurement of physical activity in this study only assessed physical activity from respondents' confessions but did not measure in quantitative detail the level of physical activity. Measuring physical activity quantitatively generally uses the IPAQ (International Physical Activity Questionnaire) questionnaire which was used in Indonesia by Lorensia et al.^{47,48} Previous research by Lorensia et al., determine differences in lung function and levels of physical activity between smokers and non-smokers. The results show that a significant difference in lung function values was observed between smokers and non-smokers ($p=0.00$).⁴⁸

CONCLUSION AND RECOMMENDATION

Most of the respondents were light smokers with normal blood levels of vitamin D. Most respondents often consume mackerel and pindang fish, even though they are served fried. Most respondents also consumed a lot of fried chicken eggs and drank bottled cow's milk. Smokers with vitamin D deficiency and insufficiency tend to have a low intake of foods containing vitamin D, such as fish, eggs and milk. Intake of fish oil and vitamin supplements is still rare. Most respondents rarely sunbathe and are

only exposed to sunlight in the morning. Respondents also use gadgets a lot and the frequency of physical activity tends to be rare and moderate.

Therefore, to reduce the risk of vitamin D deficiency, a smoker needs to improve his lifestyle by reducing the frequency of smoking, increasing his intake of foods high in vitamin D (fish, eggs, milk), increasing the frequency of exposure to sunlight, reducing the frequency of using gadgets and increasing activity physique.

ACKNOWLEDGMENTS

This research was funded by Institute of Research and Community Service by Universitas Surabaya.

AUTHOR CONTRIBUTIONS

From the research process until the writing of this article, all authors played a role in this research. The correspondent author plays a role in compiling and designing research, the 2nd acts as a data analyzer, the 3rd author acts as data collection in the field, and 4th author acts as an examiner of the clinical condition of each respondent in the field.

CONFLICTS OF INTEREST

The authors declare no conflict of interest.

REFERENCES

1. Tobacco Control Support Center-Ikatan Ahli Kesehatan Masyarakat Indonesia (TCSC-IAKMI). *Atlas tembakau Indonesia* 2020; 2020.
2. West R. Tobacco Smoking: Health Impact, Prevalence, Correlates and Interventions. *Psychology & Health*. 2017;32(8):1018–1036.
<https://doi.org/10.1080/08870446.2017.1325890>
3. Varghese J, Gharde PM. A Comprehensive Review on the Impacts of Smoking on the Health of an Individual. *Cureus*. 2023;15(10):e46532.
<https://doi.org/10.7759/cureus.46532>
4. Yang L, Zhao H, Liu K, et al. Smoking Behavior and Circulating Vitamin D Levels in Adults: A Meta-Analysis. *Food Science & Nutrition*. 2021;9(10):5820–5832.
<https://doi.org/10.1002/fsn3.2488>
5. Dahdah A, Jagers RM, Sreejit G, et al. Immunological Insights into Cigarette Smoking-Induced Cardiovascular Disease Risk. *Cells*. 2022;11(20):3190.
<https://doi.org/10.3390/cells11203190>
6. Nikniaz L, Ghojatzadeh M, Nateghian H, et al. The Interaction Effect of Aerobic Exercise and Vitamin D Supplementation on Inflammatory Factors, Anti-Inflammatory Proteins, and Lung Function in Male Smokers: A Randomized Controlled Trial. *BMC Sports Science, Medicine Rehabilitation*. 2021;13(1):102.
<https://doi.org/10.1186/s13102-021-00333-w>
7. Chen YC, Sung HC, Chuang TY, et al. Vitamin D3 Decreases TNF- α -Induced Inflammation in Lung Epithelial Cells Through a Reduction in Mitochondrial Fission and Mitophagy. *Cell Biology and Toxicology*. 2022;38(3):427–450.
<https://doi.org/10.1007/s10565-021-09629-6>
8. Tian T, Jiang X, Qin R, et al. Effect of Smoking on Lung Function Decline in a Retrospective Study of a Health Examination Population in Chinese Males. *Front Med (Lausanne)*. 2023;9:843162.
<https://doi.org/10.3389/fmed.2022.843162>
9. Augustine LF, Nair KM, Kulkarni B. Sun Exposure as a Strategy for Acquiring Vitamin D in Developing Countries of Tropical Region: Challenges & Way Forward. *Indian J Med Res*. 2021;154(3):423–432.
https://doi.org/10.4103/ijmr.IJMR_1244_18
10. Lorensia A, Suryadinata RV, Inu IA. Comparison of Vitamin D Status And Physical Activity Related With Obesity in Student. *Journal of Applied Pharmaceutical Science*. 2022;12(4):108–118.
<https://doi.org/10.7324/JAPS.2022.120412>
11. Lorensia A, Suryadinata RV. Smoking Cessa-

- tion Experience and Socioeconomic Status of Online Motorcycle Taxi Drivers in Surabaya. *Media Kesehatan Masyarakat Indonesia*. 2021;17(3):80–87.
<https://doi.org/10.30597/mkmi.v17i3.14059>
12. Octavius GS, Shakila A, Meliani M, et al. Vitamin D Deficiency is a Public Health Emergency among Indonesian Children and Adolescents: A Systematic Review and Meta-Analysis of Prevalence. *Annals of Pediatric Endocrinology & Metabolism: apem*. 2023;28(1):10-19.
<https://doi.org/10.6065/apem.2244170.085>
 13. Suryadinata RV, Lorensia A. Frekuensi Asupan Makanan, Pengetahuan Vitamin D dan Obesitas Pada Kelompok Usia Lanjut Food Frequency, Knowledge about Vitamin D and Obesity among Elderly. *Amerta Nutrition*. 2020;4(1):43–48.
<https://doi.org/10.2473/amnt.v4i1.2020>
 14. Widyastuty AASA, Jihan JC. Tingkat Kekumuhan dan Analisis Spasial Permukiman Kumuh Perkotaan (Studi Kasus: Surabaya Timur). *Jurnal Teknik Waktu*. 2018;16(2):47–55.
<https://doi.org/10.36456/waktu.v16i02.1667>
 15. Holipah H, Sulistomo HW, Maharani A. Tobacco Smoking and Risk of All-Cause Mortality in Indonesia. *PLoS One*. 2020;15(12):e0242558.
<https://doi.org/10.1371/journal.pone.0242558>
 16. Herath P, Wimalasekera S, Amarasekara T, et al. Effect of Cigarette Smoking on Smoking Biomarkers, Blood Pressure and Blood Lipid Levels Among Sri Lankan Male Smokers. *Postgraduate Medical Journal*. 2022;98(1165):848–854.
<https://doi.org/10.1136/postgradmedj-2021-141016>
 17. Sugiura T, Dohi Y, Takagi Y, et al. Close Association between Subclinical Atherosclerosis and Pulmonary Function in Middle-Aged Male Smokers. *Journal of Atherosclerosis and Thrombosis*. 2020;27(11):1230–1242.
<https://doi.org/10.5551/jat.55996>
 18. Sutherland JP, Zhou A, Hyppönen E. Vitamin D Deficiency Increases Mortality Risk in the UK Biobank: A Nonlinear Mendelian Randomization Study. *Annals of Internal Medicine*. 2022;175(11):1552–1559.
<https://doi.org/10.7326/M21-3324>
 19. AlSubai A, Baqai MH, Agha H, et al. Vitamin D and Preeclampsia: A Systematic Review And Meta-Analysis. *SAGE Open Medicine*. 2023;11:20503121231212093.
<https://doi.org/10.1177/20503121231212093>
 20. Sherief LM, Ali A, Gaballa A, et al. Vitamin D Status and Healthy Egyptian Adolescents: Where Do We Stand?. *Medicine (Baltimore)*. 2021;100(29):e26661.
<https://doi.org/10.1097/MD.00000000000026661>
 21. Moghadasi N, Alimohammadi I, Safari Variani A, et al. The Effect of Mobile Radiation on the Oxidative Stress Biomarkers in Pregnant Mice. *Journal of Family and Reproductive Health*. 2021 Sep;15(3):172-178.
<https://doi.org/10.18502/jfrh.v15i3.7134>
 22. Nguyen CV, Le TT, Nguyen NH. The Impact of Cigarette Prices on Smoking Participation and Tobacco Expenditure in Vietnam. *PLoS One*. 2021;16(12):e0260415.
<https://doi.org/10.1371/journal.pone.0260415>
 23. Septiani R. Hubungan Lama Merokok dan Frekuensi Merokok dengan Kadar Hemoglobin (HB) Pada Perokok Aktif. *Babul Ilmi: Jurnal Ilmiah Multi Science Kesehatan*. 2022;14(1):30–40. <https://jurnal.stikes-aisyiyah-palembang.ac.id/index.php/Kep/article/view/809>
 24. Weir CB, Jan A. *BMI Classification Percentile And Cut Off Points*. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2023.
 25. Salsabila NN, Indraswari N, Sujatmiko B. Gambaran Kebiasaan Merokok Di Indonesia Berdasarkan *Indonesia Family Life Survey 5 (IFLS 5)*. *Jurnal Ekonomi Kesehatan Indonesia*. 2022;7(1):13-22.
<https://journal.fkm.ui.ac.id/jurnal-eki/article/view/5394>

26. Islam MS, Rashid M, Sizear MI, et al. Cigarette Smoking and Associated Factors Among Men in Five South Asian Countries: A Pooled Analysis of Nationally Representative Surveys. *PLoS One*. 2022;17(11):e0277758.
<https://doi.org/10.1371/journal.pone.0277758>
27. Foulds J, Veldheer S, Pachas G, et al. The Effects of Reduced Nicotine Content Cigarettes on Biomarkers of Nicotine and Toxicant Exposure, Smoking Behavior and Psychiatric Symptoms in Smokers with Mood or Anxiety Disorders: A Double-Blind Randomized Trial. *PLoS One*. 2022;17(11):e0275522.
<https://doi.org/10.1371/journal.pone.0275522>
28. Omare MO, Kibet JK, Cherutoi JK, et al. A Review of Tobacco Abuse And Its Epidemiological Consequences. *Zeitschrift Fur Gesundheitswissenschaften = Journal of Public Health*. 2022;30(6):1485-500.
<https://doi.org/10.1007/s10389-020-01443-4>
29. Golestan YP, Kalan ME, Taleb ZB, et al. The Effect of Price on Cigarette Consumption, Distribution, and Sale In Tehran: A Qualitative Study. *BMC Public Health*. 2021;21(1):1720.
<https://doi.org/10.1186/s12889-021-11733-5>
30. Lorensia A, Suryadinata RV, Arganitya GN. Relationship of Food Vitamin D Intake with Obesity in Adolescent in Surabaya. *Global Medical & Health Communication*. 2022;10(2):104–110.
<https://ejournal.unisba.ac.id/index.php/gmhc/article/view/9024>
31. AlFaris NA, Alshammari GM, AlTamimi JZ, et al. Evaluating The Effects of Different Processing Methods on The Nutritional Composition of Shrimp and The Antioxidant Activity of Shrimp Powder. *Saudi Journal of Biological Sciences* 2022;29(1):640-9.
<https://doi.org/10.1016/j.sjbs.2021.09.029>
32. Sridonpai P, Judprasong K, Tirakomonpong N, et al. Effects of Different Cooking Methods on the Vitamin D Content of Commonly Consumed Fish in Thailand. *Foods*. 2022;11(6):819.
<https://doi.org/10.3390/foods11060819>
33. Andrade JM, Grandoff PG, Schneider ST. Vitamin D Intake and Factors Associated With Self-Reported Vitamin D Deficiency Among US Adults: A 2021 Cross-Sectional Study. *Frontiers in Nutrition*. 2022;9:899300.
<https://doi.org/10.3389/fnut.2022.899300>
34. Vičič V, Mikuš RP, Kugler S, et al. Vitamin D Fortification of Eggs Alone and in Combination with Milk in Women Aged 44-65 Years: Fortification Model and Economic Evaluation. *Slovenian Journal of Public Health*. 2022;62(1):30-33.
<https://doi.org/10.2478/sjph-2023-0005>
35. Daly RM, De Ross B, Gianoudis J, et al. Dose-Response Effect of Consuming Commercially Available Eggs on Wintertime Serum 25-Hydroxyvitamin D Concentrations in Young Australian Adults: a 12-Week Randomized Controlled Trial. *The Journal of Nutrition*. 2022;152(7):1702-1710.
<https://doi.org/10.1093/jn/nxac044>
36. Pellegrino L, Marangoni F, Muscogiuri G, et al. Vitamin D Fortification of Consumption Cow's Milk: Health, Nutritional and Technological Aspects. A Multidisciplinary Lecture of the Recent Scientific Evidence. *Molecules*. 2021;26(17):5289.
<https://doi.org/10.3390/molecules26175289>
37. Weir RR, Johnston M, Lowis C, et al. Vitamin D₃ Content of Cows' Milk Produced in Northern Ireland and Its Efficacy As A Vehicle For Vitamin D Fortification: A UK Model. *International Journal of Food Sciences and Nutrition*. 2021;72(4):447-455.
<https://doi.org/10.1080/09637486.2020.1837743>
38. Zhang Y, Lin S, Li J, et al. Interaction of Passive Smoking and Diet Habits on Vitamin D Deficiency among Women of Reproductive Age in Rural Central China. *Nutrients*. 2022;15(1):126.
<https://doi.org/10.3390/nu15010126>

39. Tiwari RK, Sharma V, Pandey RK, et al. Nicotine Addiction: Neurobiology and Mechanism. *Journal of Pharmacopuncture*. 2020;23(1):1–7.
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7163392/>
40. Lorensia A, Suryadinata RV. Profil of Omega-3 Food Intake and Its Association with Socioeconomic Status on Online Motorcycle Driver Smokers. *Healthcare in Low-resource Settings*. 2023;11(s1):11164.
<https://www.pagepressjournals.org/index.php/hls/article/view/11164>
41. Carlberg C. Vitamin D and Pigmented Skin. *Nutrients*. 2022;14(2):325.
<https://doi.org/10.3390/nu14020325>
42. Raymond-Lezman JR, Riskin SI. Benefits and Risks of Sun Exposure to Maintain Adequate Vitamin D Levels. *Cureus*. 2023;15(5):e38578.
<https://doi.org/10.7759/cureus.38578>
43. Sultana N. Sun Awareness and Sun Protection Practices. *Clinical, Cosmetic and Investigational Dermatology*. 2020;13:717–730.
<https://doi.org/10.2147/CCID.S265477>
44. Shan L, Dong H, Wang T, et al. Screen Time, Age and Sunshine Duration Rather Than Outdoor Activity Time Are Related to Nutritional Vitamin D Status in Children With ASD. *Frontiers in Pediatrics*. 2022;9:806981.
<https://doi.org/10.3389/fped.2021.806981>
45. Rocka A, Jasielska F, Madras D, et al. The Impact of Digital Screen Time on Dietary Habits and Physical Activity in Children and Adolescents. *Nutrients*. 2022;14(14):2985.
<https://doi.org/10.3390/nu14142985>
46. Zhang J, Cao ZB. Exercise: A Possibly Effective Way to Improve Vitamin D Nutritional Status. *Nutrients*. 2022;14(13):2652.
<https://doi.org/10.3390/nu14132652>
47. Lorensia A, Suryadinata RV, Chandra NLMR. Profile of Vitamin D Status, Physical Activity, and Lung Health in Construction Workers. *Community Medicine and Public Health of Indonesia Journal*. 2020;1(2):117–24.
<http://comphi.sinergis.org/index.php/comphi/article/view/19>
48. Lorensia A, Muntu CM, Suryadinata RV, et al. Effect of lung function disorders and physical activity on smoking and non-smoking students. *Journal of Preventive Medicine and Hygiene*. 2021; 62(1):89–96.
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8283647>

Home (<https://journal.unhas.ac.id/index.php/mkmi/index>)

/ Archives (<https://journal.unhas.ac.id/index.php/mkmi/issue/archive>)

/ Vol. 20 No. 2: JUNE 2024 (<https://journal.unhas.ac.id/index.php/mkmi/issue/view/1518>) /

Risk of Vitamin D Deficiency in Smokers: Mix-Method

✉ (<https://orcid.org/0000-0002-9746-0606>) Amelia Lorensia, (<https://orcid.org/0000-0002-1213-1484>) Marthy Meliana Ariyanti Jalmav, Zahwa Dhiba, (<https://orcid.org/0000-0001-8679-7871>) Rivan Virlando Suryadinata

<https://doi.org/10.30597/mkmi.v20i2.32775> (<https://doi.org/10.30597/mkmi.v20i2.32775>)

  (<https://badge.dimensions.ai/details/doi/10.30597/mkmi.v20i2.32775?domain=https://journal.unhas.ac.id>)

Issue

Vol. 20 No. 2: JUNE 2024 (<https://journal.unhas.ac.id/index.php/mkmi/issue/view/1518>)

Keywords:

Smokers (<https://journal.unhas.ac.id/index.php/mkmi/search?query=smokers>)

vitamin D (<https://journal.unhas.ac.id/index.php/mkmi/search?query=vitamin%20d>)

deficiency risk (<https://journal.unhas.ac.id/index.php/mkmi/search?query=deficiency%20risk>)

PDF (<https://journal.unhas.ac.id/index.php/mkmi/article/view/32775/11816>)



Abstract



Authors



Citation



Licensing

Abstract

Cigarettes are one of the leading causes of death in the world and Indonesia. Cigarettes emit cigarette smoke, which contains toxins and has pro-inflammatory properties that interfere the metabolism of vitamin D in the lungs. The study aimed to determine the risk of vitamin D deficiency in active smokers in Rungkut Village, Surabaya City. This research method is a mix-method with a Convergent Parallel Design type strategy, which is a way of collecting quantitative (cross-sectional) and qualitative data (interpretative phenomenological analysis) simultaneously, with purposive sampling technique. The variable was the risk of vitamin D deficiency and vitamin D level. Subjects were active smokers aged ≥ 19 years in the Rungkut Subdistrict. The number of samples involved in this research was 125 samples in the quantitative phase, and 15 people in the qualitative phase. The highest level of smoking severity was light smokers (45.60%). Most of respondents had normal vitamin D levels (≥ 20 mg/mL) of 73.33%, averaging of 37.66 ± 5.89 mg/mL. Most active smokers are at risk of vitamin D deficiency. Risk factors for deficiency in heavy smokers are not consuming foods containing vitamin D (such as fish, eggs, and milk), not consuming fish oil supplements and vitamins, and not sunbathing. However, the level of physical activity tended to be moderate-heavy. The conclusion is that smokers must pay attention to their intake of foods containing vitamin D and exposure to sunlight to achieve sufficient vitamin D levels.

References

1. Tobacco Control Support Center-Ikatan Ahli Kesehatan Masyarakat Indonesia (TCSC-IAKMI). Atlas tembakau Indonesia 2020; 2020.
2. West R. Tobacco Smoking: Health Impact, Prevalence, Correlates and Interventions. *Psychology & Health*. 2017;32(8):1018–1036. <https://doi.org/10.1080/08870446.2017.1325890>

3. Varghese J, Gharde PM. A Comprehensive Review on the Impacts of Smoking on the Health of an Individual. *Cureus*. 2023;15(10):e46532. <https://doi.org/10.7759/cureus.46532>
4. Yang L, Zhao H, Liu K, et al. Smoking Behavior and Circulating Vitamin D Levels in Adults: A Meta-Analysis. *Food Science & Nutrition*. 2021;9(10):5820–5832. <https://doi.org/10.1002/fsn3.2488>
5. Dahdah A, Jaggars RM, Sreejit G, et al. Immunological Insights into Cigarette Smoking-Induced Cardiovascular Disease Risk. *Cells*. 2022;11(20):3190. <https://doi.org/10.3390/cells11203190>
6. Niksaian J, Chaharadab M, Motakabian H, et al. The Interaction Effect of Aerobic Exercise and Vitamin D Supplementation on

[Read More](#)

Downloads

Download data is not yet available.

[Share Now](#)

Address

Jl. Perintis Kemerdekaan Km.10,
Faculty of Public Health, Hasanuddin University,
Tamalanrea, Makassar, South Sulawesi, 90245, Indonesia
[View on Google Map \(https://goo.gl/maps/8A3mBnQTonD2\)](https://goo.gl/maps/8A3mBnQTonD2)

Contact Info

Principal Contact : Ida Leida M.Thaha
HP : +628114440454
Email : journal.mkmi@unhas.ac.id (<mailto:journal.mkmi@unhas.ac.id>)
: jurnal.mkmi@gmail.com (<mailto:jurnal.mkmi@gmail.com>)

JOURNAL ISSUE



Published by Public Health Faculty, Hasanuddin University
Quarterly Publication
View Current Issue (<https://journal.unhas.ac.id/index.php/mkmi/issue/current>)

SOCIAL SHARE



(<https://api.whatsapp.com/send?phone=+628114440454&text=Assalamualaikum%20admin%20MKMI,%20saya%20ingin%20bertanya%20mengenai>)



(<https://www.instagram.com/jurnalmkmi/>)

Published Volume By Year

2024	▼
2023	▼
2022	▼
2021	▼
2020	▼
2019	▼
2018	▼
2017	▼
2016	▼
2015	▼
2014	▼
2013	▼

HIGHLY ACCESSED

Kualitas Hidup Penderita Insomnia Pada Mahasiswa (https://journal.unhas.ac.id/index.php/mkmi/article/view/3464) 1081 16 June 2018 Muhammad Akbar Nurdin, Andi...
MODEL TIKUS DIABETES YANG DIINDUKSI STREPTOZOTOCIN-SUKROSA UNTUK PENDEKATAN PENELITIAN DIABETES MELITUS GESTASIONAL (https://journal.unhas.ac.id/index.php/mkmi/article/view/550) 819 22 June 2016 Firdaus Firdaus, Rimbawan R...
Pengaruh Program Makan Siang terhadap Asupan Makanan, Status Anemia dan Perilaku Gizi Santri Perempuan (https://journal.unhas.ac.id/index.php/mkmi/article/view/5769) 758 17 March 2019 Dewi Kusumawati, Rimbawan R...
FAKTOR PENYEBAB PENYAKIT AKIBAT KERJA PADA PEKERJA LAS (https://journal.unhas.ac.id/index.php/mkmi/article/view/1583) 641 29 March 2017 Husaini Husaini, Ratna Sety...
PENGARUH POLA MAKAN TERHADAP KADAR KOLESTEROL TOTAL (https://journal.unhas.ac.id/index.php/mkmi/article/view/2132) 573 9 December 2017

 (https://journal.unhas.ac.id/index.php/mkmi/CitednessScopus)

Visitors

Country	Visitors
United States	246,536
Germany	14,535
United Kingdom	9,178
India	771
France	615
Japan	381
Italy	316
Spain	282
Canada	271
United Kingdom	249
United States	193
Canada	177
France	159
United States	148
Taiwan	134
United States	130
Turkey	123
United States	106
South Korea	105
United States	93
United States	89
France	77
United States	70
United States	64

<http://info.flagcounter.com/OqN9>

0382162 (<http://statcounter.com/>)

[View My Stats \(https://statcounter.com/p10091631/?guest=1\)](https://statcounter.com/p10091631/?guest=1)

turnitin®
(https://www.turnitin.com/login_page.asp?lang=en_us)



state=2778aeef1a79ae21844b109617f824f5&prompt=login&scope=openid%20email%20profile%20els_auth_info%20els_analytics_info%20urn%3Acom%3Aelsevier%3Aaidp%3Aapolicy%3Aproduct%


Home (<https://journal.unhas.ac.id/index.php/mkmi/index>) / Archives (<https://journal.unhas.ac.id/index.php/mkmi/issue/archive>) / Vol. 20 No. 2: JUNE 2024

Media Kesehatan Masyarakat Indonesia accepts scientific papers in the form of research reports (original research papers) with a focus on the development of public health issues problems in Indonesia, including the developments and main problems in the field of epidemiology; Health Promotion; Environmental Health, Occupational Health, and Safety, Health Administration and Policy, Biostatistics, Reproductive Health, Hospital Management, Nutrition Science, Health Information Systems in Regional of Indonesia. Media Kesehatan Masyarakat Indonesia collaborates with the Professional Organization of the Ikatan Ahli Kesehatan Masyarakat Indonesia (IAKMI) in terms of assisting the advancement of public health science and the dissemination of research results. Although focused on the Indonesia region, MKMI does not preclude manuscript beyond the region which has correlative and/or comparable issues within that geographical scope. Articles published in Media Kesehatan Masyarakat Indonesia go through a double-blind peer-review process. Therefore, the decision to accept scientific articles is in the right of the Editorial Board based on peer reviewers' recommendations.

DOI: <https://doi.org/10.30597/mkmi.v20i2> (<https://doi.org/10.30597/mkmi.v20i2>)

Supports and Barriers Regarding The Iron-Folic Acid Supplementation Adherence Level in Anemic Pregnant Women: Indonesian's Perspective (<https://journal.unhas.ac.id/index.php/mkmi/article/view/32569>)

Rizka Novia Atmadani, Akrom Akrom, Dhea Ananda, Laila Yoga Saputri, Alvina Arum Puspitasari,...




 46-56

10.30597/mkmi.v20i2.32569 (<https://journal.unhas.ac.id/index.php/mkmi/article/view/32569>)

PDF (<https://journal.unhas.ac.id/index.php/mkmi/article/view/32569/11812>)Problems Scanning Related to Stunting Using Iceberg Theory Model in Salakbrojo Village Central Java (<https://journal.unhas.ac.id/index.php/mkmi/article/view/32672>)

Yuniarti Yuniarti, Ristiawati Ristiawati, Dewi Nugraheni Restu Mastuti, Muhammad Fatih




 57-66

10.30597/mkmi.v20i2.32672 (<https://journal.unhas.ac.id/index.php/mkmi/article/view/32672>)

PDF (<https://journal.unhas.ac.id/index.php/mkmi/article/view/32672/11813>)

Navigating the Storm: Unraveling the Factors Shaping Adolescent Anxiety Amid the COVID-19 Pandemic (<https://journal.unhas.ac.id/index.php/mkmi/article/view/28275>)

Rini Nur Diana, Mamnuah Mamnuah, Dwi Ernawati



67-75

10.30597/mkmi.v20i2.28275 (<https://journal.unhas.ac.id/index.php/mkmi/article/view/28275>)

PDF (<https://journal.unhas.ac.id/index.php/mkmi/article/view/28275/11814>)

Factors Affecting The Success of Exclusive Breastfeeding Among Mothers Recovered From Covid-19 (<https://journal.unhas.ac.id/index.php/mkmi/article/view/31400>)

Ulfatul Latifah, Nora Rahmanindar, Ardhi Henda Karmandika

Citations ?

(<https://badge.dimensions.ai/details/doi/10.30597/mkmi.v20i2.31400?domain=https://journal.unhas.ac.id>)

76-84

10.30597/mkmi.v20i2.31400 (<https://journal.unhas.ac.id/index.php/mkmi/article/view/31400>)

PDF (<https://journal.unhas.ac.id/index.php/mkmi/article/view/31400/11815>)

Risk of Vitamin D Deficiency in Smokers: Mix-Method (<https://journal.unhas.ac.id/index.php/mkmi/article/view/32775>)

Amelia Lorensia, Marthy Meliana Ariyanti Jalmav, Zahwa Dhiba, Rivan Virlando Suryadinata

Citations ?

(<https://badge.dimensions.ai/details/doi/10.30597/mkmi.v20i2.32775?domain=https://journal.unhas.ac.id>)

85-101

10.30597/mkmi.v20i2.32775 (<https://journal.unhas.ac.id/index.php/mkmi/article/view/32775>)

PDF (<https://journal.unhas.ac.id/index.php/mkmi/article/view/32775/11816>)

INTERNATIONAL INDEXING



ELSEVIER
Scopus

(<https://www.scopus.com/sourceid/21101176724>)



(<https://www.scimagojr.com/journalsearch.php?q=21101176724&tip=sid&exact=no>)

NATIONAL ACCREDITATION



(<https://sinta.kemdikbud.go.id/journals/profile/782>)

View Sertificate (https://drive.google.com/file/d/1aqVnulhDKILscC4n69LseuzB0H_M45aH/view)

MENU

Editorial Team (<https://journal.unhas.ac.id/index.php/mkmi/EditorialTeam>)

Peer Reviewers (<https://journal.unhas.ac.id/index.php/mkmi/Reviewer>)

Peer Review Process (<https://journal.unhas.ac.id/index.php/mkmi/PeerReviewProcess>)

Focus and Scope (<https://journal.unhas.ac.id/index.php/mkmi/focusandscope>)

Publication Ethics (<https://journal.unhas.ac.id/index.php/mkmi/PublicationEthic>)

Online Submission Guidelines (<https://journal.unhas.ac.id/index.php/mkmi/about/submissions#authorGuidelines>)

Plagiarism Check (<https://journal.unhas.ac.id/index.php/mkmi/Plagiarismcheck>)

Article Processing Charge (<https://journal.unhas.ac.id/index.php/mkmi/ArticleProcessingCharges>)

Policies (<https://journal.unhas.ac.id/index.php/mkmi/JournalPolicies>)

License Term (<https://journal.unhas.ac.id/index.php/mkmi/Licenseterm>)

Statistics (https://statcounter.com/p10091631/summary/?account_id=5107254&login_id=3&code=ce60dcf244d946de14f6c4adc0032acd&guest_login=1)

Template (https://docs.google.com/document/d/1HJExAdP9Z_JtAY_UzPkZ_FA184jTf0tR/edit?usp=share_link&oid=112843235217211379447&rtopf=true&sd=true)

Journal History (<https://journal.unhas.ac.id/index.php/mkmi/JournalHistory>)

Indexing (<https://journal.unhas.ac.id/index.php/mkmi/indexing2>)

EDITORIAL BOARD

Editor in Chief

Ida Leida Maria

Department of Epidemiology, Faculty of Public Health, Universitas Hasanuddin, Indonesia (SCOPUS ID : 57062982100 (<https://www.scopus.com/authid/detail.uri?authorId=57062982100>))

Managing Editor

Indra Dwinata

Department of Epidemiology, Faculty of Public Health, Universitas Hasanuddin, Indonesia (SCOPUS ID : 57194555111 (<https://www.scopus.com/authid/detail.uri?authorId=57194555111>))

Editorial Board

Cordia Ming Yeuk Chu, (SCOPUS ID : 56435044600 (<https://www.scopus.com/authid/detail.uri?authorId=56435044600>)) Centre for Environment and Population Health, Griffith University, Australia

Mahdi Hasan Suhail, (SCOPUS ID : 55399070800 (<https://www.scopus.com/authid/detail.uri?authorId=55399070800>)) Department of Physics, University of Baghdad, Iraq

Isam Mohammed Ibrahim, (SCOPUS ID : 56200248100 (<https://www.scopus.com/authid/detail.uri?authorId=56200248100>)) Department of Physics, University of Baghdad, Iraq

Kristen Marie Hurley, (SCOPUS ID : 8502630200 (<https://www.scopus.com/authid/detail.uri?authorId=8502630200>)) Center for Human Nutrition, International Health, The Johns Hopkins Bloomberg School of Public Health, Baltimore, Maryland, USA

Peter Davey, (SCOPUS ID : 57192956383 (<https://www.scopus.com/authid/detail.uri?authorId=57192956383>)) School of Environment and Science, Griffith University, Australia

Wesam Al Madhoun, (SCOPUS ID : 26430526800 (<https://www.scopus.com/authid/detail.uri?authorId=26430526800>)) Global Ambassadors of Sustainability, Engineering, Gaza University, Palestine

Aini Ahmad, (SCOPUS ID : 57203645325 (<https://www.scopus.com/authid/detail.uri?authorId=57203645325>)) School of Nursing, KPJ Healthcare University College, Malaysia

Veni Hadju, (SCOPUS ID : 8684558700 (<https://www.scopus.com/authid/detail.uri?authorId=8684558700>)), Department of Nutrition, Faculty of Public Health, Universitas Hasanuddin, Indonesia

Oktia Woro Kasmini Handayani, (SCOPUS ID : 57196057104 (<https://www.scopus.com/authid/detail.uri?authorId=57196057104>)), Nutrition, Department of Public Health, Faculty of Sport Science, Universitas Negeri Semarang, Indonesia

Ede Surya Darmawan, (SCOPUS ID : 57215313462 (<https://www.scopus.com/authid/detail.uri?authorId=57215313462>)), Department of Health Policy and Administration, Faculty of Public Health, Universitas Indonesia, Indonesia

IN COLLABORATION WITH



VIEW MoU (<https://drive.google.com/open?id=1RaMPA4mm2-S-bgYJQv7-pylPZ47I9e1E>)



Address

Jl. Perintis Kemerdekaan Km.10,
Faculty of Public Health, Hasanuddin University,
Tamalanrea, Makassar, South Sulawesi, 90245, Indonesia
View on Google Map (<https://goo.gl/maps/8A3mBnQTonD2>)

Contact Info

Principal Contact : Ida Leida M.Thaha
HP : +628114440454
Email : journal.mkmi@unhas.ac.id (<mailto:journal.mkmi@unhas.ac.id>)
: jurnal.mkmi@gmail.com (<mailto:jurnal.mkmi@gmail.com>)

JOURNAL ISSUE



Published by Public Health Faculty, Hasanuddin University
Quarterly Publication
View Current Issue (<https://journal.unhas.ac.id/index.php/mkmi/issue/current>)

SOCIAL SHARE



(<https://api.whatsapp.com/send?phone=+628114440454&text=Assalamualaikum%20admin%20MKMI,%20saya%20ingin%20bertanya%20mengenai>)



(<https://www.instagram.com/jurnalmkmi/>)

Published Volume By Year

2024	▼
2023	▼
2022	▼
2021	▼
2020	▼
2019	▼
2018	▼
2017	▼
2016	▼
2015	▼
2014	▼
2013	▼

HIGHLY ACCESSED

Kualitas Hidup Penderita Insomnia Pada Mahasiswa (<https://journal.unhas.ac.id/index.php/mkmi/article/view/3464>)

1081

16 June 2018

Muhammad Akbar Nurdin, Andi...

MODEL TIKUS DIABETES YANG DIINDUKSI STREPTOZOTOCIN-SUKROSA UNTUK PENDEKATAN PENELITIAN DIABETES MELITUS GESTASIONAL
(<https://journal.unhas.ac.id/index.php/mkmi/article/view/550>)

819

22 June 2016

Firdaus Firdaus, Rimbawan R...

Pengaruh Program Makan Siang terhadap Asupan Makanan, Status Anemia dan Perilaku Gizi Santri Perempuan (<https://journal.unhas.ac.id/index.php/mkmi/article/view/5769>)

758

17 March 2019

Dewi Kusumawati, Rimbawan R...

FAKTOR PENYEBAB PENYAKIT AKIBAT KERJA PADA PEKERJA LAS (<https://journal.unhas.ac.id/index.php/mkmi/article/view/1583>)

641

29 March 2017


Husaini Husaini, Ratna Sety...

PENGARUH POLA MAKAN TERHADAP KADAR KOLESTEROL TOTAL (<https://journal.unhas.ac.id/index.php/mkmi/article/view/2132>)

573

9 December 2017

CITATION



STAT COUNTER

Visitors



0382162 (http://statcounter.com/)

View My Stats (https://statcounter.com/p10091631/?guest=1)

TOOLS



(https://www.turnitin.com/login_page.asp?lang=en_us)



grammarly (https://www.grammarly.com/signin)



(https://id.elsevier.com/as/authorization.oauth2?

state=2778aee1a79ae21844b109617f824f5&prompt=login&scope=openid%20email%20profile%20els_auth_info%20els_analytics_info%20urn%3Acom%3Aelsevier%3Aidp%3Apolycy%3Aproduct%

Home (https://journal.unhas.ac.id/index.php/mkmi/index) / Editorial Team

Editorial Team

Editor in Chief

Ida Leida Maria, (SCOPUS ID : 57062982100 (https://www.scopus.com/authid/detail.uri?authorId=57062982100)), Department of Epidemiology, Faculty of Public Health, Hasanuddin University, Indonesia

Managing Editor

Indra Dwinata, (SCOPUS ID : 57194555111 (https://www.scopus.com/authid/detail.uri?authorId=57194555111)), Department of Epidemiology, Faculty of Public Health, Hasanuddin University, Indonesia

Editorial Board

Cordia Ming Yeuk Chu, (SCOPUS ID : 56435044600 (https://www.scopus.com/authid/detail.uri?authorId=56435044600)) Centre for Environment and Population Health, Griffith University, Australia

Mahdi Hasan Suhail, (SCOPUS ID : 55399070800 (https://www.scopus.com/authid/detail.uri?authorId=55399070800)) Department of Physics, University of Baghdad, Iraq

Isam Mohammed Ibrahim, (SCOPUS ID : 56200248100 (https://www.scopus.com/authid/detail.uri?authorId=56200248100)) Department of Physics, University of Baghdad, Iraq

Kristen Marie Hurley, (SCOPUS ID : 8502630200 (https://www.scopus.com/authid/detail.uri?authorId=8502630200)) Center for Human Nutrition, International Health, The Johns Hopkins Bloomberg School of Public Health, Baltimore, Maryland, USA

Peter Davey, (SCOPUS ID : 57192956383 (https://www.scopus.com/authid/detail.uri?authorId=57192956383)) School of Environment and Science, Griffith University, Australia

Wesam Al Madhoun, (SCOPUS ID : 26430526800 (https://www.scopus.com/authid/detail.uri?authorId=26430526800)) Global Ambassadors of Sustainability, Engineering, Gaza University, Palestine

Aini Ahmad, (SCOPUS ID : 57203645325 (https://www.scopus.com/authid/detail.uri?authorId=57203645325)) School of Nursing, KPJ Healthcare University College, Malaysia

Veni Hadju, (SCOPUS ID : 8684558700 (https://www.scopus.com/authid/detail.uri?authorId=8684558700)), Department of Nutrition, Faculty of Public Health, Hasanuddin University, Indonesia

Oktia Woro Kasmini Handayani, (SCOPUS ID : 57196057104 (https://www.scopus.com/authid/detail.uri?authorId=57196057104)), Nutrition, Department of Public Health, Faculty of Sport Science, Universitas Negeri Semarang, Indonesia

Ede Surya Darmawan, (SCOPUS ID : 57215313462 (https://www.scopus.com/authid/detail.uri?authorId=57215313462)), Department of Health Policy and Administration, Faculty of Public Health, Universitas Indonesia, Indonesia

INTERNATIONAL INDEXING

ELSEVIER
Scopus

(https://www.scopus.com/sourceid/21101176724)



(<https://www.scimagojr.com/journalsearch.php?q=21101176724&tip=sid&exact=no>)

NATIONAL ACCREDITATION



(<https://sinta.kemdikbud.go.id/journals/profile/782>)

View Sertificate (https://drive.google.com/file/d/1aqVnulhDKILscC4n69LseuzB0H_M45aH/view)

MENU

- Editorial Team (<https://journal.unhas.ac.id/index.php/mkmi/EditorialTeam>)
- Peer Reviewers (<https://journal.unhas.ac.id/index.php/mkmi/Reviewer>)
- Peer Review Process (<https://journal.unhas.ac.id/index.php/mkmi/PeerReviewProcess>)
- Focus and Scope (<https://journal.unhas.ac.id/index.php/mkmi/focusandscope>)
- Publication Ethics (<https://journal.unhas.ac.id/index.php/mkmi/PublicationEthic>)
- Online Submission Guidelines (<https://journal.unhas.ac.id/index.php/mkmi/about/submissions#authorGuidelines>)
- Plagiarism Check (<https://journal.unhas.ac.id/index.php/mkmi/Plagiarismcheck>)
- Article Processing Charge (<https://journal.unhas.ac.id/index.php/mkmi/ArticleProcessingCharges>)
- Policies (<https://journal.unhas.ac.id/index.php/mkmi/JournalPolicies>)
- License Term (<https://journal.unhas.ac.id/index.php/mkmi/Licenseterm>)
- Statistics (https://statcounter.com/p10091631/summary/?account_id=5107254&login_id=3&code=ce60dcf244d946de14f6c4adc0032acd&guest_login=1)
- Template (https://docs.google.com/document/d/1HJExAdP9Z_JtAY_UzPkZ_FA184jTf0tR/edit?usp=share_link&ouid=112843235217211379447&rtopf=true&sd=true)
- Journal History (<https://journal.unhas.ac.id/index.php/mkmi/JournalHistory>)
- Indexing (<https://journal.unhas.ac.id/index.php/mkmi/indexing2>)

EDITORIAL BOARD

- Editor in Chief**
 - Ida Leida Maria
Department of Epidemiology, Faculty of Public Health, Universitas Hasanuddin, Indonesia (SCOPUS ID : 57062982100 (<https://www.scopus.com/authid/detail.uri?authorId=57062982100>))
- Managing Editor**
 - Indra Dwinata
Department of Epidemiology, Faculty of Public Health, Universitas Hasanuddin, Indonesia (SCOPUS ID : 57194555111 (<https://www.scopus.com/authid/detail.uri?authorId=57194555111>))
- Editorial Board**
 - Cordia Ming Yeuk Chu, (SCOPUS ID : 56435044600 (<https://www.scopus.com/authid/detail.uri?authorId=56435044600>)) Centre for Environment and Population Health, Griffith University, Australia
 - Mahdi Hasan Suhail, (SCOPUS ID : 55399070800 (<https://www.scopus.com/authid/detail.uri?authorId=55399070800>)) Department of Physics, University of Baghdad, Iraq
 - Isam Mohammed Ibrahim, (SCOPUS ID : 56200248100 (<https://www.scopus.com/authid/detail.uri?authorId=56200248100>)) Department of Physics, University of Baghdad, Iraq
 - Kristen Marie Hurley, (SCOPUS ID : 8502630200 (<https://www.scopus.com/authid/detail.uri?authorId=8502630200>)) Center for Human Nutrition, International Health, The Johns Hopkins Bloomberg School of Public Health, Baltimore, Maryland, USA
 - Peter Davey, (SCOPUS ID : 57192956383 (<https://www.scopus.com/authid/detail.uri?authorId=57192956383>)) School of Environment and Science, Griffith University, Australia
 - Wesam Al Madhoun, (SCOPUS ID : 26430526800 (<https://www.scopus.com/authid/detail.uri?authorId=26430526800>)) Global Ambassadors of Sustainability, Engineering, Gaza University, Palestine
 - Aini Ahmad, (SCOPUS ID : 57203645325 (<https://www.scopus.com/authid/detail.uri?authorId=57203645325>)) School of Nursing, KPJ Healthcare University College, Malaysia

Veni Hadju, (SCOPUS ID : 8684558700 (<https://www.scopus.com/authid/detail.uri?authorId=8684558700>)), Department of Nutrition, Faculty of Public Health, Universitas Hasanuddin, Indonesia

Oktia Woro Kasmini Handayani, (SCOPUS ID : 57196057104 (<https://www.scopus.com/authid/detail.uri?authorId=57196057104>)), Nutrition, Department of Public Health, Faculty of Sport Science, Universitas Negeri Semarang, Indonesia

Ede Surya Darmawan, (SCOPUS ID : 57215313462 (<https://www.scopus.com/authid/detail.uri?authorId=57215313462>)), Department of Health Policy and Administration, Faculty of Public Health, Universitas Indonesia, Indonesia

IN COLLABORATION WITH



VIEW MoU (<https://drive.google.com/open?id=1RaMPA4mm2-S-bgYJQv7-pylPZ47l9e1E>)



Address

Jl. Perintis Kemerdekaan Km.10,
Faculty of Public Health, Hasanuddin University,
Tamalanrea, Makassar, South Sulawesi, 90245, Indonesia
[View on Google Map \(https://goo.gl/maps/8A3mBnQTonD2\)](https://goo.gl/maps/8A3mBnQTonD2)

Contact Info

Principal Contact : Ida Leida M.Thaha
HP : +628114440454
Email : journal.mkmi@unhas.ac.id (<mailto:journal.mkmi@unhas.ac.id>)
: jurnal.mkmi@gmail.com (<mailto:jurnal.mkmi@gmail.com>)

JOURNAL ISSUE



Published by Public Health Faculty, Hasanuddin University
Quarterly Publication
View Current Issue (<https://journal.unhas.ac.id/index.php/mkmi/issue/current>)

SOCIAL SHARE



(<https://api.whatsapp.com/send?phone=+628114440454&text=Assalamualaikum%20admin%20MKMI,%20saya%20ingin%20bertanya%20mengenai>)



(<https://www.instagram.com/jurnalmkmi/>)

Published Volume By Year

2024	▼
2023	▼
2022	▼
2021	▼
2020	▼
2019	▼
2018	▼
2017	▼
2016	▼
2015	▼
2014	▼
2013	▼

HIGHLY ACCESSED

Kualitas Hidup Penderita Insomnia Pada Mahasiswa (https://journal.unhas.ac.id/index.php/mkmi/article/view/3464) 1081 16 June 2018 Muhammad Akbar Nurdin, Andi...
MODEL TIKUS DIABETES YANG DIINDUKSI STREPTOZOTOCIN-SUKROSA UNTUK PENDEKATAN PENELITIAN DIABETES MELITUS GESTASIONAL (https://journal.unhas.ac.id/index.php/mkmi/article/view/550) 819 22 June 2016 Firdaus Firdaus, Rimbawan R...
Pengaruh Program Makan Siang terhadap Asupan Makanan, Status Anemia dan Perilaku Gizi Santri Perempuan (https://journal.unhas.ac.id/index.php/mkmi/article/view/5769) 758 17 March 2019 Dewi Kusumawati, Rimbawan R...
FAKTOR PENYEBAB PENYAKIT AKIBAT KERJA PADA PEKERJA LAS (https://journal.unhas.ac.id/index.php/mkmi/article/view/1583) 641 29 March 2017 Husaini Husaini, Ratna Sety...
PENGARUH POLA MAKAN TERHADAP KADAR KOLESTEROL TOTAL (https://journal.unhas.ac.id/index.php/mkmi/article/view/2132) 573 9 December 2017

Carol Hutchinson, (SCOPUS ID : 55166522600 (<https://www.scopus.com/authid/detail.uri?authorId=55166522600>)), Department of Nutrition, Faculty of Public Health, Mahidol University, Thailand

Evawany Yunita Aritonang, (SCOPUS ID : 56688571200 (<https://www.scopus.com/authid/detail.uri?authorId=56688571200>)), Department of Public Health Nutrition, Faculty of Public Health, Universitas Sumatera Utara, Indonesia

Muhammad Nadjib Bustan, (SCOPUS ID : 6602741608 (<https://www.scopus.com/authid/detail.uri?authorId=6602741608>)), Department of Statistics, Faculty of Mathematics and Natural Sciences, Universitas Negeri Makassar, Indonesia

Haerawati Idris, (SCOPUS ID : 57194584807 (<https://www.scopus.com/authid/detail.uri?authorId=57194584807>)), Department of Health Policy Administration, Faculty of Public Health, Universitas Sriwijaya, Indonesia

Suriah Suriah, (SINTA ID : 5999021 (<https://sinta.ristekbrin.go.id/authors/detail?id=5999021&view=overview>)), Department of Health Promotion, Faculty of Public Health, Universitas Hasanuddin, Indonesia

Agus Bintara Birawida, (SCOPUS ID : 57193116626 (<https://www.scopus.com/authid/detail.uri?authorId=57193116626>)), Department of Environmental Health, Faculty of Public Health, Universitas Hasanuddin, Indonesia

Lalu Muhammad Saleh, (SCOPUS ID : 57200575006 (<https://www.scopus.com/authid/detail.uri?authorId=57200575006>)), Department of Occupational Health and Safety, Faculty of Public Health, Universitas Hasanuddin, Indonesia

Abdul Salam, (SCOPUS ID : 57208671878 (<https://www.scopus.com/authid/detail.uri?authorId=57208671878>)), Departement of Nutrition, Faculty of Public Health, Universitas Hasanuddin, Indonesia

Syamsuar Manyullei, (SCOPUS ID : 56429686000 (<https://www.scopus.com/authid/detail.uri?authorId=56429686000>)), Department of Environmental Health, Faculty of Public Health, Universitas Hasanuddin, Indonesia

Yahya Thamrin, (SCOPUS ID : 35410790900 (<https://www.scopus.com/authid/detail.uri?authorId=35410790900>)), Department of Occupational Health and Safety, Public Health Faculty, Universitas Hasanuddin, Indonesia

Ede Surya Darmawan, (SCOPUS ID: (<https://www.scopus.com/authid/detail.uri?authorId=57215313462>)57215313462 (<https://www.scopus.com/authid/detail.uri?authorId=57215313462>)), Departemen Administrasi dan Kebijakan Kesehatan, Fakultas Kesehatan Masyarakat, Universitas Indonesia, Indonesia

Dian Ayubi, (SCOPUS ID: 57201491749 (<https://www.scopus.com/authid/detail.uri?authorId=57201491749>)), Departemen Pendidikan Kesehatan dan Ilmu Perilaku, Fakultas Kesehatan Masyarakat Universitas Indonesia, Indonesia

Tri Martiana, (SCOPUS ID: 57188930379) (<https://www.scopus.com/authid/detail.uri?authorId=57188930379>), Departemen Keselamatan dan Kesehatan Kerja, Fakultas Kesehatan Masyarakat, Universitas Airlangga, Indonesia

Alimin Maidin, (SCOPUS ID: (<https://www.scopus.com/authid/detail.uri?authorId=6505544947>)6505544947 (<https://www.scopus.com/authid/detail.uri?authorId=6505544947>)), Departemen Manajemen Rumah Sakit, Fakultas Kesehatan Masyarakat, Indonesia

INTERNATIONAL INDEXING



ELSEVIER
Scopus

(<https://www.scopus.com/sourceid/21101176724>)



(<https://www.scimagojr.com/journalsearch.php?q=21101176724&tip=sid&exact=no>)

NATIONAL ACCREDITATION



(<https://sinta.kemdikbud.go.id/journals/profile/782>)

View Certificate (https://drive.google.com/file/d/1aqVnulhDKLscC4n69LseuzB0H_M45aH/view)

MENU

Editorial Team (<https://journal.unhas.ac.id/index.php/mkmi/EditorialTeam>)

Peer Reviewers (<https://journal.unhas.ac.id/index.php/mkmi/Reviewer>)

Peer Review Process (<https://journal.unhas.ac.id/index.php/mkmi/PeerReviewProcess>)

Focus and Scope (<https://journal.unhas.ac.id/index.php/mkmi/focusandscope>)

Publication Ethics (<https://journal.unhas.ac.id/index.php/mkmi/PublicationEthic>)

Online Submission Guidelines (<https://journal.unhas.ac.id/index.php/mkmi/about/submissions#authorGuidelines>)

Plagiarism Check (<https://journal.unhas.ac.id/index.php/mkmi/Plagiarismcheck>)

Article Processing Charge (<https://journal.unhas.ac.id/index.php/mkmi/ArticleProcessingCharges>)

Policies (<https://journal.unhas.ac.id/index.php/mkmi/JournalPolicies>)

License Term (<https://journal.unhas.ac.id/index.php/mkmi/Licenseterm>)

Statistics (https://statcounter.com/p10091631/summary/?account_id=5107254&login_id=3&code=ce60dcf244d946de14f6c4adc0032acd&guest_login=1)

Template (https://docs.google.com/document/d/1HJExAdP9Z_JtAY_UzPkZ_FA184jTf0tR/edit?usp=share_link&oid=112843235217211379447&rtpof=true&sd=true)

Journal History (<https://journal.unhas.ac.id/index.php/mkmi/JournalHistory>)

Indexing (<https://journal.unhas.ac.id/index.php/mkmi/indexing2>)

EDITORIAL BOARD

Editor in Chief

Ida Leida Maria

Department of Epidemiology, Faculty of Public Health, Universitas Hasanuddin, Indonesia (SCOPUS ID : 57062982100 (<https://www.scopus.com/authid/detail.uri?authorId=57062982100>))

Managing Editor

Indra Dwinata

Department of Epidemiology, Faculty of Public Health, Universitas Hasanuddin, Indonesia (SCOPUS ID : 57194555111 (<https://www.scopus.com/authid/detail.uri?authorId=57194555111>))

Editorial Board

Cordia Ming Yeuk Chu, (SCOPUS ID : 56435044600 (<https://www.scopus.com/authid/detail.uri?authorId=56435044600>)) Centre for Environment and Population Health, Griffith University, Australia

Mahdi Hasan Suhail, (SCOPUS ID : 55399070800 (<https://www.scopus.com/authid/detail.uri?authorId=55399070800>)) Department of Physics, University of Baghdad, Iraq

Isam Mohammed Ibrahim, (SCOPUS ID : 56200248100 (<https://www.scopus.com/authid/detail.uri?authorId=56200248100>)) Department of Physics, University of Baghdad, Iraq

Kristen Marie Hurley, (SCOPUS ID : 8502630200 (<https://www.scopus.com/authid/detail.uri?authorId=8502630200>)) Center for Human Nutrition, International Health, The Johns Hopkins Bloomberg School of Public Health, Baltimore, Maryland, USA

Peter Davey, (SCOPUS ID : 57192956383 (<https://www.scopus.com/authid/detail.uri?authorId=57192956383>)) School of Environment and Science, Griffith University, Australia

Wesam Al Madhoun, (SCOPUS ID : 26430526800 (<https://www.scopus.com/authid/detail.uri?authorId=26430526800>)) Global Ambassadors of Sustainability, Engineering, Gaza University, Palestine

Aini Ahmad, (SCOPUS ID : 57203645325 (<https://www.scopus.com/authid/detail.uri?authorId=57203645325>)) School of Nursing, KPJ Healthcare University College, Malaysia

Veni Hadju, (SCOPUS ID : 8684558700 (<https://www.scopus.com/authid/detail.uri?authorId=8684558700>)), Department of Nutrition, Faculty of Public Health, Universitas Hasanuddin, Indonesia

Oktia Woro Kasmini Handayani, (SCOPUS ID : 57196057104 (<https://www.scopus.com/authid/detail.uri?authorId=57196057104>)), Nutrition, Department of Public Health, Faculty of Sport Science, Universitas Negeri Semarang, Indonesia

Ede Surya Darmawan, (SCOPUS ID : 57215313462 (<https://www.scopus.com/authid/detail.uri?authorId=57215313462>)), Department of Health Policy and Administration, Faculty of Public Health, Universitas Indonesia, Indonesia

IN COLLABORATION WITH



VIEW MoU (<https://drive.google.com/open?id=1RaMPA4mm2-S-bgYJQv7-pyIPZ47I9e1E>)



Address

Jl. Perintis Kemerdekaan Km.10,
Faculty of Public Health, Hasanuddin University,
Tamalanrea, Makassar, South Sulawesi, 90245, Indonesia
View on Google Map (<https://goo.gl/maps/8A3mBnQTonD2>)

Contact Info

Principal Contact : Ida Leida M.Thaha

HP : +628114440454

Email : journal.mkmi@unhas.ac.id (mailto:journal.mkmi@unhas.ac.id)

: jurnal.mkmi@gmail.com (mailto:jurnal.mkmi@gmail.com)



Source details

Media Kesehatan Masyarakat Indonesia

Years currently covered by Scopus: from 2019 to 2024
Publisher: Fakultas Kesehatan Masyarakat Universitas Hasanuddin
ISSN: 0216-2482 E-ISSN: 2356-4067

Subject area: Medicine: Health Policy Medicine: Epidemiology
Medicine: Public Health, Environmental and Occupational Health Medicine: Health Informatics

Source type: Journal

[View all documents >](#) [Set document alert](#) [Save to source list](#)

CiteScore 2023 0.3 ⓘ
SJR 2023 0.111 ⓘ
SNIP 2023 0.117 ⓘ

CiteScore CiteScore rank & trend Scopus content coverage

CiteScore 2023

0.3 = $\frac{32 \text{ Citations } 2020 - 2023}{108 \text{ Documents } 2020 - 2023}$

Calculated on 05 May, 2024

CiteScoreTracker 2024 ⓘ

0.2 = $\frac{14 \text{ Citations to date}}{65 \text{ Documents to date}}$

Last updated on 08 July, 2024 • Updated monthly

CiteScore rank 2023 ⓘ

Category	Rank	Percentile
Medicine		
Health Policy	#289/310	6th
Medicine		
Epidemiology	#139/148	6th
Medicine		
Public Health, Environmental and Occupational Health	#628/665	5th

[View CiteScore methodology >](#) [CiteScore FAQ >](#) [Add CiteScore to your site](#)

About Scopus

- What is Scopus
- Content coverage
- Scopus blog
- Scopus API
- Privacy matters

Language

- 日本語版を表示する
- 查看简体中文版本
- 查看繁體中文版本
- Просмотр версии на русском языке

Customer Service

- Help
- Tutorials
- Contact us

ELSEVIER

[Terms and conditions ↗](#) [Privacy policy ↗](#)

All content on this site: Copyright © 2024 Elsevier B.V. ↗, its licensors, and contributors. All rights are reserved, including those for text and data mining, AI training, and similar technologies. For all open access content, the Creative Commons licensing terms apply. We use cookies to help provide and enhance our service and tailor content.By continuing, you agree to the use of cookies ↗.



SERTIFIKAT

Direktorat Jenderal Penguatan Riset dan Pengembangan,
Kementerian Riset, Teknologi dan Pendidikan Tinggi



Kutipan dari Keputusan Direktur Jenderal Penguatan Riset dan Pengembangan
Kementerian Riset, Teknologi dan Pendidikan Tinggi Republik Indonesia

Nomor: 36/E/KPT/2019

Peringkat Akreditasi Jurnal Ilmiah Periode VII Tahun 2019

Nama Jurnal Ilmiah

Media Kesehatan Masyarakat Indonesia

E-ISSN: 23564067

Penerbit: Fakultas Kesehatan Masyarakat Universitas Hasanuddin

Ditetapkan Sebagai Jurnal Ilmiah

TERAKREDITASI PERINGKAT 2

Akreditasi Berlaku Selama 5 (lima) Tahun, Yaitu
Volume 15 Nomor 3 Tahun 2019 sampai Volume 20 Nomor 2 Tahun 2024

Jakarta, 13 Desember 2019

Direktur Jenderal Penguatan Riset dan Pengembangan



[Signature]
Dr. Muhammad Dimyati
NIP. 195912171984021001

Risk of Vitamin D Deficiency in Smokers: Mix-Method

by Amelia Lorensia

Submission date: 26-Aug-2025 09:23AM (UTC+0700)

Submission ID: 2735354164

File name: Risk_of_Vitamin_D_Deficiency_in_Smokers.pdf (639.66K)

Word count: 9340

Character count: 46055

Risk of Vitamin D Deficiency in Smokers: Mix-Method

ORIGINALITY REPORT

14%

SIMILARITY INDEX

12%

INTERNET SOURCES

8%

PUBLICATIONS

3%

STUDENT PAPERS

PRIMARY SOURCES

1	www.mdpi.com Internet Source	2%
2	japsonline.com Internet Source	1%
3	www.ncbi.nlm.nih.gov Internet Source	1%
4	link.springer.com Internet Source	1%
5	www.researchgate.net Internet Source	1%
6	www.tandfonline.com Internet Source	1%
7	www.frontiersin.org Internet Source	<1%
8	Submitted to Badan PPSDM Kesehatan Kementerian Kesehatan Student Paper	<1%
9	journals.plos.org Internet Source	<1%
10	www.omicsdi.org Internet Source	<1%
11	Widya Ramadhaniati, Erna Kusumawardhani, Haryati Haryati, Dwi Laksono Adi Putro, Moh. Isa, Ali Assagaf. "The Effect of Platinum-Based Chemotherapy in Advanced Non-Small Cell Carcinoma Lung Cancer (NSCLC) on	<1%

Cardiovascular Events in Ulin Hospital Banjarmasin", Jurnal Ners, 2025

Publication

12	Haniyyah Isman, Noto Dwimartutie, Siti Setiati, Pringgodigdo Nugroho. "Prevalence and Factors Associated with Vitamin D Inadequacy in the Elderly Population of South Sulawesi", Springer Science and Business Media LLC, 2025	<1 %
----	--	------

Publication

13	repository.unikal.ac.id	<1 %
----	---	------

Internet Source

14	Geissler, Catherine, Powers, Hilary. "Human Nutrition", Human Nutrition, 2023	<1 %
----	---	------

Publication

15	doaj.org	<1 %
----	---	------

Internet Source

16	Markus Herrmann, Christopher-John L. Farrell, Irene Pusceddu, Neus Fabregat-Cabello, Etienne Cavalier. "Assessment of vitamin D status – a changing landscape", Clinical Chemistry and Laboratory Medicine (CCLM), 2017	<1 %
----	---	------

Publication

17	Submitted to Heriot-Watt University	<1 %
----	-------------------------------------	------

Student Paper

18	academic.oup.com	<1 %
----	---	------

Internet Source

19	jurnalfkip.unram.ac.id	<1 %
----	---	------

Internet Source

20	pubmed.ncbi.nlm.nih.gov	<1 %
----	---	------

Internet Source

21	Submitted to Athlone Institute of Technology	<1 %
----	--	------

Student Paper

22	www2.mdpi.com Internet Source	<1 %
23	1library.net Internet Source	<1 %
24	Submitted to University of Wales Swansea Student Paper	<1 %
25	journal.unnes.ac.id Internet Source	<1 %
26	ueaeprints.uea.ac.uk Internet Source	<1 %
27	essay.utwente.nl Internet Source	<1 %
28	www.emro.who.int Internet Source	<1 %
29	Efsane Yavuz, Mustafa Örkmez, Mehmet Akif Bildirici, İsmet Gamze Bozdayı, Murat Usta. "Geographical differences in 25(OH)D reference intervals", The Journal of Steroid Biochemistry and Molecular Biology, 2025 Publication	<1 %
30	Houda Zoubiri, Amina Tahar, Samir AitAbderrhmane, Messaoud Saidani, Elhadj-Ahmed Koceir. "Oral Cholecalciferol Supplementation in Sahara Black People with Chronic Kidney Disease Modulates Cytokine Storm, Oxidative Stress Damage and Athero-Thromboembolic Risk", Nutrients, 2022 Publication	<1 %
31	adihusada.ac.id Internet Source	<1 %
32	journalrbgo.org Internet Source	<1 %
33	nutrition.org Internet Source	

<1 %

34

whatsfordinnerthisweek.blogspot.com

Internet Source

<1 %

35

Vania Dwi Ramadhani, Laili Rahayuwati, Mamat Lukman. "Smoking Behavioral Dependence during the COVID-19 Pandemic Among University Student", Jurnal Berita Ilmu Keperawatan, 2023

Publication

<1 %

36

comphi.sinergis.org

Internet Source

<1 %

37

www.davaodoctors.edu.ph

Internet Source

<1 %

38

www.e-apem.org

Internet Source

<1 %

39

www.meno-me.co.nz

Internet Source

<1 %

40

www.quickanswer.blog

Internet Source

<1 %

41

Geir Bjørklund, Olha Storchylo, Monica Butnariu, Maryam Dadar, Salvatore Chirumbolo. "Milk Thistle (Silybum marianum): Potential Role in Cancer Prevention", Current Medicinal Chemistry, 2025

Publication

<1 %

42

Wen Lei. "INFLUENCES OF GYMNASTICS ON BODY COMPOSITION AND CARDIOPULMONARY FUNCTION OF ATHLETES", Revista Brasileira de Medicina do Esporte, 2023

Publication

<1 %

43

f1000research.com

Internet Source

<1 %

44 oamjms.eu
Internet Source

<1 %

45 onlinelibrary.wiley.com
Internet Source

<1 %

46 pmc.ncbi.nlm.nih.gov
Internet Source

<1 %

47 vitamindwiki.com
Internet Source

<1 %

48 www.vitamind-journal.it
Internet Source

<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography On



Risk of Vitamin D Deficiency in Smokers: Mix-Method

Amelia Lorensia^{1*}, Marthy Meliana Ariyanti Jalmav², Zahwa Dhiba³, Rivan Virlando Suryadinata⁴

¹Department of Clinical-Community Pharmacy, Faculty of Pharmacy, University of Surabaya, Surabaya, Indonesia

²Petra Christian University, Surabaya, Indonesia

³Faculty of Pharmacy, University of Surabaya, Surabaya, Indonesia

⁴Department of Public Health, Faculty of Medicine, University of Surabaya, Surabaya, Indonesia

*Authors Correspondence: amelia.lorensia@gmail.com; amelia.lorensia@staff.ubaya.ac.id/08155020707

ARTICLE INFO

Article History:

Received Mar, 20th, 2024

Accepted May, 12th, 2024

Published online Jun, 30th, 2024

Keywords:

Smokers;
vitamin D;
deficiency risk;

ABSTRACT

Cigarettes are one of the leading causes of death in the world and Indonesia. Cigarettes emit cigarette smoke, which contains toxins and has pro-inflammatory properties that interfere the metabolism of vitamin D in the lungs. The study aimed to determine the risk of vitamin D deficiency in active smokers in Rungkut Village, Surabaya City. This research method is a mix-method with a Convergent Parallel Design type strategy, which is a way of collecting quantitative (cross-sectional) and qualitative data (interpretative phenomenological analysis) simultaneously, with purposive sampling technique. The variable was the risk of vitamin D deficiency and vitamin D level. Subjects were active smokers aged ≥ 19 years in the Rungkut Subdistrict. The number of samples involved in this research was 125 samples in the quantitative phase, and 15 people in the qualitative phase. The highest level of smoking severity was light smokers (45.60%). Most of respondents had normal vitamin D levels (≥ 20 $\mu\text{g/mL}$) of 73.33%, averaging of 37.66 ± 5.89 $\mu\text{g/mL}$. Most active smokers are at risk of vitamin D deficiency. Risk factors for deficiency in heavy smokers are not consuming foods containing vitamin D (such as fish, eggs, and milk), not consuming fish oil supplements and vitamins, and not sunbathing. However, the level of physical activity tended to be moderate-heavy. The conclusion is that smokers must pay attention to their intake of foods containing vitamin D and exposure to sunlight to achieve sufficient vitamin D levels.

INTRODUCTION

Indonesia is in third place with the most significant number of smokers in the world after the People's Republic of China (PRC) and India. It is estimated that by 2030 the number of smokers in the world and the death rate will reach up to 10 million, of which 70% will come from developing countries. Currently, 50% of the death rate among smokers in the world has come from developing countries.¹ The number of deaths due to smoking is predicted to continue to increase every year because the main ingredient in cigarettes is tobacco.² The main substance in tobacco, namely nicotine, has a relaxing effect, making it addictive for users. The chemical poisons contained in cigarette smoke can be easily transferred from the lungs to the bloodstream and even spread to almost every organ of the human body.³

Systemically, exposure to cigarette smoke can disrupt vitamin D metabolism and damage tubules in the kidneys and liver. Cigarette exposure reduces 25(OH)D and 1,25(OH)2D levels by 10%.⁴ Cigarettes can also cause dryness and shrinkage of the skin, where skin synthesis is the primary source of vitamin D in humans.⁴ Cigarette smoke contains toxic aldehydes that can influence the inflammatory response by increasing the production of pro-inflammatory cytokines.⁵ Additionally, it has pro-inflammatory properties that can weaken host cells in the human body and disrupt the metabolism of vitamin D in the lungs,^{6,7} leading to an accelerated decline in lung function.⁸

In recent years, research has shown an increase in the prevalence of vitamin D deficiency in countries near the equator, where there is abundant exposure to sunlight.⁹

Sun exposure is the most sustainable solution to the widespread issue of vitamin D deficiency, particularly in tropical regions. Despite the abundant sunlight in countries considered a paradox,⁹ like Indonesia, there is a high prevalence of vitamin D deficiency.^{10,11} A Hartung-Knapp-Sidik-Jonkman method meta-analysis revealed that vitamin D deficiency is a significant public health concern in Indonesia.¹² Research by Suryadinata and Lorensia in Indonesia demonstrated a link between vitamin D deficiency and smoking habits.¹³ Respiratory diseases are commonly associated with vitamin D deficiency. This study aimed to assess the risk

of vitamin D deficiency among active smokers in the Rungkut District of Surabaya City, located in the eastern part of Surabaya, with a large population of 833,426 people. Rungkut District is densely populated and includes worker residential areas, such as cigarette factories.¹⁴

MATERIAL AND METHOD

This research is a mixed-methods study that combines quantitative and qualitative data. The design strategy employed is convergent parallel design. The quantitative aspect involves descriptive analysis using a Cross-Sectional Study approach, while the qualitative aspect utilizes Interpretative Phenomenological Analysis (IPA) techniques for data analysis. The study was conducted in Rungkut Village, Surabaya City, from April to July 2023. The research adhered to the code of ethics with No. 160/KE/VI/2023 by Universitas Surabaya, Surabaya.

Smokers are adults who have smoked at least 100 cigarettes in their lifetime and currently smoke every day.¹⁵ Smoking classification based on the Brinkman Index is divided into three categories: light smokers (0-199), moderate smokers (200-599), and heavy smokers (≥ 600). The Brinkman index formula is calculated by multiplying the number of cigarettes smoked daily (sticks) by the number of years of smoking (years).^{16,17} Vitamin D deficiency occurs when the body lacks sufficient vitamin D intake. Most experts consider a 25(OH)D concentration below 25 or 30 nmol/L indicating an increased risk of vitamin D deficiency. A concentration of 50 nmol/L or higher is considered within the normal health range.^{18,19} The study population was adult male active smokers in Rungkut Subdistrict, Surabaya City. The sample included male active smokers aged 19-60 years, with no chronic lung disease, who had been smoking for the last 5 years, and were willing to participate in the research. The total number of active adult smokers in the Rungkut Subdistrict was 125 respondents. Data was collected from residents who attended a health education session at the RW hall facilitated by the local RT head in April 2023. For this quantitative study, the minimum sample size was calculated using the Slovin formula, $n = N / [1 + Ne^2]$ with a sampling error (e) of 5%. The minimum sample size (n) for this research was 95 respondents. The sampling method used was purposive sampling. In the

qualitative phase, data collection continued until data saturation was achieved.

The method used in this research is Mixed-Method, which can simultaneously collect quantitative and qualitative data. The qualitative phase of the mix-method aims to produce a more complete and in-depth picture of the phenomenon. In the initial stage, quantitative methods were used to measure active smokers' vitamin D deficiency risk. These risk factors were described from domains, including smoking history, diet, vitamin/supplement intake, sun exposure, frequency of gadget screens, and physical activity. Data collection began based on adapting the Vitamin D Deficiency Risk Questionnaire.^{4,10,20,21} This research used a questionnaire consisting of 12 closed questions and several open questions. Questions in the vitamin D deficiency risk questionnaire included cigarette use, consumption of foods containing vitamin D (consumption of food from fish, eggs, and milk, which was assessed from the question points in the questionnaire, including the frequency and method of serving food), vitamin D supplements, exposure to sunlight, exposure to gadget radiation, and physical activity. The validity test in this questionnaire only extends to the development of sentences and language because each question item only contains short (closed) questions. In filling out the questionnaire in this study, respondents can fill in each question that has been provided.

Measurement of Vitamin D (25(OH)D) levels in human serum uses VIDAS® 25 OH Vitamin D TOTAL (VIT D), an automated quantitative test employing the ELFA (Enzyme-Linked Fluorescent Assay) technique (Figure 1). VIDAS® 25 OH Vitamin D TOTAL is employed to evaluate Vitamin D sufficiency. A certified laboratory collected 3 ml blood samples from respondents, followed by a blood sample taken at a standardized laboratory in Surabaya.

Quantitative methods are used to obtain measurable data that can be descriptive. The quantitative data in this research is for descriptive analysis. Measuring the percentage value of respondents regarding the risk of deficiency involves using simple statistical data, with the resulting percentage value calculated using the Statistical Package for Social Science (SPSS) program. After the quantitative data collection is complete, qualitative data is gathered in the form of an in-depth examination of the risk factors for vitamin D deficiency in various domains, including cigarette smoking, intake of foods containing vitamin D (such as fish, eggs, milk, and fish oil supplements), vitamin intake, skin exposure to sunlight, screen time on gadgets, and physical activity. Qualitative data is obtained through interviews conducted with the help of an interview guide instrument.

BIOMÉRIEUX

REF 30463

054788 - 04 - 2022-07

VIDAS® 25 OH Vitamin D TOTAL (VITD)

IVD

Intended Use

VIDAS® 25 OH Vitamin D TOTAL (VITD) is an automated quantitative test for use on the instruments of the VIDAS® family for the determination of 25-hydroxyvitamin D Total in human serum or plasma using the ELFA technique (Enzyme Linked Fluorescent Assay).

The VIDAS® 25 OH Vitamin D TOTAL assay is to be used as an aid in the assessment of Vitamin D sufficiency.

Content of the Kit (60 TESTS)

60 VITD Strips TM	STR	Ready-to-use. Stabilizer of human origin ¹ .
60 VITD Solid Phase Receptacles 2 x 30	SPR	Ready-to-use. Interior of SPR device coated with vitamin D.
VITD Calibrator 1 x 1.2 mL (liquid)	S1	Ready-to-use. 25-(OH) Vitamin D diluted in human serum ¹ + preservative. M.E. data indicate the calibrator concentration in ng/mL, ("Calibrator (S1) Dose Value") and the confidence interval in "Relative Fluorescence Value" (Calibrator (S1) RFV Range).
VITD Control 1 x 1.1 mL (liquid)	C1	Ready-to-use. 25-(OH) Vitamin D diluted in human serum ¹ + preservative. M.E. data indicate the confidence interval in ng/mL, ("Control C1 Dose Value Range").

Specifications for the factory master data required to calibrate the assay: M.E. (Master Lot Entry) barcode printed on the box label.

¹ package insert downloadable from www.biomerieux.com

Source: Primary Data, 2023

Figure 1. VIDAS® 25 OH Vitamin D TOTAL (VIT-D) Procedure

RESULTS

This research and data collection was conducted from April 2023 to July 2023 in Rungkut Village. Data were collected by filling in questionnaire questions and conducting interviews with respondents included in the research sample. The number of samples involved in this research was 125. All active smokers were willing to fill out a questionnaire and have their vitamin D levels measured (n=125). However, in the qualitative phase, only 15 respondents were able to be followed up and were willing to continue with in-depth interviews. Based on Table 1, 125 respondents were active smokers, with 65 respondents (52%) starting smoking in adulthood, 22 (40%) having an average normal body weight, and the highest percentage of respondents (92%) having no history of the disease.

Based on Table 1 it showed that 85 respondents (68%) were active smokers with an average of ≥10 years of smoking.23 The maximum number of cigarettes smoked per day was 11-24, with 69 respondents (55.20%). Additionally, 99 respondents (79.20%) smoked filter cigarettes. The Brinkman Index categorized 57 respondents (45.50%) as light smokers.

Based on Table 2, there were 21 respondents (16.80%) who smoked tobacco through direct smoking. Additionally, 108 respondents (86.40%) consumed fish, with 100 respondents (80%) frying it. Moreover, 117 respondents (93.60%) consumed eggs, out of which 115 respondents (92%) used chicken eggs and fried them, as reported by 94 respondents (75.20%). Furthermore, 79 respondents consumed milk (63.20%), and 25 respondents (20.00%) used packaged liquid milk. The highest frequency of respondents who did not consume fish oil supplements was 118 respondents (94.40%). Similarly, the highest frequency of respondents who did not consume vitamins was 73 respondents (58.40%). The main reason respondents consumed vitamins was due to the health benefits, with 45 respondents (36%) citing this reason. The highest frequency of respondents who often looked at gadget screens was 118 respondents (94.40%), with 49 respondents (39.20%) spending more than 7

hours on this activity. Additionally, 98 respondents (78%) were unaware of the ideal time to look at monitor screens (40%). The highest frequency of respondents who rarely engaged in physical activity was 96 respondents (76.80%), with 49 respondents (39.20%) spending more than 7 hours inactive and 96 respondents (76.80%) falling into the moderate activity category (Table 2).

Table 1. Frequency Distribution of Respondents Based on Age, BMI, and Smoking Profiles

Characteristics	n = 125	%
Age (Years)		
19-44 (Adult)	65	52
45-59 (Pre-elderly)	48	38.40
45-60 (Seniors)	12	9.60
Body Mass Index (BMI)²⁴		
Underweight (<18.5)	12	9.60
Normal (18.5-22.9)	50	40
Overweight with risks (23- 24.9)	25	20
Obesity 1 (25-29.9)	31	24.80
Obesity 2 (≥30)	7	5.60
Disease History		
Hypertension	3	2.40
Dyslipidemia	3	2.40
Diabetes Mellitus Type 2	3	2.40
Rheumatoid Arthritis	1	0.80
No History of Illness	115	92
Long Time Smoking Cigarettes (Years)		
< 10	40	32
≥ 10	85	68
Number of Cigarettes Smoked		
1-10 cigarettes per day	47	37.60
11-24 cigarettes per day	69	55.20
≥24 cigarettes per day	9	7.20
Types of Cigarettes		
Filter	99	79.20
Kretek	20	16.00
Mixture	6	4.80
Brinkman Index		
Light smoker (0-199)	57	45.60
Medium smoker (200-600)	52	41.60
Heavy smoker (>600)	16	12.80

Source: Primary Data, 2023

Table 2. Distribution Profile of Respondents' Answers

Domain	Question	Respondent's Answer	n = 125	%
Cigarette	Have you ever consumed tobacco directly?	No	104	83.20
		Yes	21	16.80
	How do you use this tobacco?	No	104	83.20
		Inhale directly	21	16.80
		Chewed	0	0
	How do you know this?	Don't know	104	83.20
		Friends/family	13	10.40
		Environment	8	6.40
Fish	Have you ever consumed fish?	Consume	108	86.40
		Do not consume	17	13.60
	What fish do you usually eat?	Mackarel tuna	35	28
		Pindang fish	32	25.60
		Salted fish	25	20
		Milkfish	7	5.60
		Catfish	5	4
	How do you serve the fish you eat?	Mujaer fish	4	3.20
		Fried	100	80
		Burned	5	4
		Steamed	3	2.40
	What is your reason for consuming fish?	Kitchen dishes	40	32
		Love the taste	32	25.60
		Get health benefits	21	16.80
		Do not like	16	12.80
		Affordable prices	10	8
		Diet	5	4
		Allergy	1	0.80
Egg	Have you ever consumed eggs?	Consume	117	93.60
		Do not consume	8	6.40
	What eggs do you usually eat?	Chicken	115	92
		Duck	2	1.60
	How do you serve the eggs you consume?	Fried	94	75.20
		Boiled	23	18.40
	What is the reason you consume eggs?	Love the taste	82	65.60
		Get health benefits	23	18.40
		Kitchen dishes	9	7.20
		Don't consume	7	5.60
		Cheap/affordable prices	3	2.40
		Diet	1	0.80
Milk	Do you consume milk?	Consume	79	63.20
		Do not consume	46	36.80
	What milk do you usually consume?	Packaging (liquid)	25	20
		Condensed milk (sachet)	22	17.60
		Real milk	20	16
	What is the reason you consume milk?	Powder	12	9.60
		Health benefits	50	40
		Love the taste	29	23.20
Fish Oil Supplements	Do you take fish oil supplements?	Do not consume	118	94.40
		Consume	7	5.60
	What type and brand of fish oil supplement do you consume?	Omega-3	4	3.20
		Forgot the brand	3	2.40
	How many fish oil supplements do you consume?	Capsule	7	5.60
		Teaspoon	0	0
		Tablespoon	0	0

Table 2. Distribution Profile of Respondents' Answers

Domain	Question	Respondent's Answer	n = 125	%
	Why do you take fish oil supplements?	Not accustomed to	63	50.40
		Stench	40	32
		Expensive price	15	12
		Health benefits	7	5.60
Vitamin	Do you take vitamins?	Do not consume	73	58.40
		Consume	52	41.60
	What type and brand of vitamins do you consume?	Vitacimin®	27	21.60
		Enervon-C®	7	5.60
		Hemaviton®	5	4
		Vitamin D (1000 IU)	5	4
		Imbost®	4	3.20
	The reason you take vitamins?	Don't know	73	58.40
		Health benefits	45	36
		Recommended friend/doctor	4	3.20
			3	2.40
Changes in Skin Color	Does your skin color change when exposed to direct sunlight?	No	68	54.40
		Yes	57	45.60
	What changes does your skin color look like when exposed to sunlight?	No	40	32
		Blackish red	20	16
Sunbathe	Do you take free time to sunbathe directly?	Red	8	6.40
		No	67	53.60
		Yes (eg, tanned)	58	46.40
		07.00-09.00	60	48
		10.00-11.00	32	25.60
	What time are you usually exposed to sunlight (sunbathing)?	12.00-14.00	26	20.80
		15.00-17.00	7	5.60
	Do you sunbathe using personal protective equipment?	Hat	58	46.40
		Jacket	40	32
		Without personal protection	19	15.20
		Sunscreen/sunscreen	6	4.80
		Umbrella	2	1.60
	Do you wear closed clothes?	Shirt and trousers	82	65.60
		Short t-shirt	43	34.40
Gadget Screen	Are you staring at a gadget screen such as a cellphone, laptop or tablet?	Often	118	94.40
		Rarely/not at all	7	5.60
	How long (per day) do you stare at a monitor screen such as a cellphone, laptop or tablet?	≥7 hours	49	39.20
		0-1.5 hours	32	25.60
		4-6 hours	20	16
		2-3 hours	17	13.60
	In your opinion, how long is the ideal time to stare at a monitor screen?	Don't know	98	78.40
		1-2 hours	18	14.40
Physical Activity	Do you often do physical activity?	20-30 minutes	9	7.20
		Seldom	96	76.80
	How often do you do physical activity?	Often	29	23.20
		Light	0	0
		Moderate	96	76.80
		Heavy	29	23.20

Source: Primary Data, 2023

Most of respondents had normal vitamin D levels (sufficiency) of 73.33%, with an average vitamin D level of 37.66±5.89 mg/mL. The results of the cross-tabulation between the description of vitamin D risk factors and vitamin D levels can be seen in Table 3.

When conducting interviews regarding the risk of vitamin D deficiency, 15 respondents out

of 125 respondents were willing to conduct further interviews. The researcher then made interview transcripts and summarized them in table form as well as interview excerpts to further understand the interview themes so that they were in accordance with the research title (Table 4).

32

Table 3. Cross Tabulation Between the Risk of Vitamin D Deficiency and Vitamin D Levels					
Risk of Vitamin D Deficiency		Vitamin D Serum Category (µg/mL) ³¹			Total
Domain		Deficiency (<12µg/mL) (n = 7)	Insufficiency (12-20 µg/mL) (n = 26)	Sufficiency (>20 µg/mL) (n = 92)	
Cigarette (Brinkman Index) ²⁴	Light smoker (0-199)	0	4	53	57
	Medium smoker (200-600)	2	12	38	52
	Heavy smoker (>600)	5	10	1	16
Fish	Consume fish that contain high levels of vitamin D	3	10	52	65
	Consume fish that do not contain high levels of vitamin D	2	7	34	43
	Do not consume fish	3	9	5	17
		1	25	91	117
Egg	Consume egg	6	1	1	8
	Do not consume egg				
Milk	Consume milk	2	9	68	79
	Do not consume milk	5	17	24	46
Fish Oil Supplements	Consume fish oil	4	24	90	118
	Do not consume fish oil	3	2	2	7
Vitamin	Vitamin C	0	1	36	37
	Vitamin D	0	0	5	5
	Immunomodulator	0	2	2	4
	Not taking any vitamins	7	23	49	79
Changes in Skin Color	Yes	3	13	41	57
	No	4	13	51	68
Sunbathe	Yes	4	11	43	58
	No	3	15	49	67
Gadget Screen	Often	4	24	90	118
	Rarely/not at all	3	2	2	7
Physical Activity	Light	0	0	0	0
	Moderate	4	14	78	96
	Heavy	3	12	14	29

Source: Primary Data, 2023

Table 4. In-depth Interview Results

Domain	Findings	Answer Findings	Quotes From The Interview	Source
Smoke	Respondents knew the severity of smoking	Frequency of cigarette use by respondents	"I've been smoking for more than 15 years, so what's strange is that I can't really stop. Just stop when you cough, then try again with one cigarette, if you cough, throw it away and change to another brand of cigarette, and so on, sis."	Respondent 7
			"Usually I use official cigarettes from factories such as salt warehouses and every day I use up more than 20 cigarettes. I've been smoking for more than 15 years."	Respondent 1
			"I use cigarettes and the brand varies depending on the price, but I usually use 1 pack of solar a day, and I smoked when I was single, around 25 years after graduating from high school, sis."	Respondent 9
			"You don't know how to smoke, sis, sometimes Surya sometimes uses DJI Samsoe. I've been smoking since I was 18 years old, sis, and now I'm 52 years old."	Respondent 11
	Respondents' views on cigarette use	Views of cigarettes on health by respondents	"For example, if I don't smoke, sis, I'm stressed because my job is uncertain. So, if I smoke, it just feels calmer in my mind, sis."	Respondent 2
			"I don't smoke, or if I just stop for a day, I'll get a headache, but if I smoke just one cigarette, my body will immediately feel better, sis, so I think that smoking isn't completely dangerous because according to my father, cigarettes are medicine."	Respondent 4
Tobacco	Respondents know their knowledge and behavior regarding tobacco use	Tobacco use by respondents	"Previously, I used to smoke tobacco straight away, sis, because in the past tobacco was cheaper than cigarettes, and I knew how to do it when my friends told me to use tobacco."	Respondent 13
			"In the past, I used tobacco directly, sis, by smoking it. I knew about tobacco because I was told by a friend and forced by a friend to try it. After a while, I became addicted to it and then I tried to replace it with cigarettes."	Respondent 8
			"When I'm in a crisis, I usually use tobacco, sis, because it's cheaper and the way to do it is to smoke it directly. I know that method, I was given tips from my father for using tobacco too."	Respondent 5
Fish	Respondents know the behavior and amount of fish consumed	Respondents explained their fish consumption habits	"I like fish but I don't know how to eat it, sis, sometimes once a week, often twice at most. I like it when the fish is fried, but more often I like to eat fish steamed in banana leaves."	Respondent 6

Table 4. In-depth Interview Results

Domain	Findings	Answer Findings	Quotes From The Interview	Source
	Respondents know their views on fish consumption habits	Respondents explained the benefits of consuming fish	"I like to eat fish, but I can't guarantee that it will be every day depending on my wife's business in the kitchen, sis, usually once a week."	Respondent 8
			"Actually, there is no benefit from fish, it's just a complement to food and just increases your appetite, sis."	Respondent 15
			"I like fish, sis, because the benefits of fish are that it is delicious as a complement to food and makes you full, sis."	Respondent 13
			"Because you really like eating fish because the benefits are actually good for your health, sis."	Respondent 11
Egg	Respondents' views on behavior and habits of consuming eggs	Respondents explained their egg consumption habits and views on eggs	"I like eggs, sis, but they contain diabetes, so at my age I cut back on them, sis. So I just eat boiled egg whites and discard the yolks and I eat them often every day, usually mixed with vegetables, because my wife makes that menu, sis."	Respondent 7
			"I like eggs, but I usually eat salted eggs when they are served because I like them. Eggs are good for the strength of your voice and are good for your health."	Respondent 3
			"I also often eat fried eggs, usually twice a week, and now I've switched to salted eggs, sis. "The reason is that I like eggs and they have good health benefits, but if the portion is too big, it's not good, sis."	Respondent 11
			"Eggs are just a food supplement, so they have no health benefits."	Respondent 2
Milk	Respondents understand the behavior and benefits of milk	Respondents explained their milk drinking habits and views on the benefits of milk	"Sometimes I drink Frisian flag milk twice a week and if I drink too much I'm afraid because it contains sugar, sis. I actually prefer whole milk but I don't buy it because it's more expensive, sis. And I think the benefits of milk are good for health and increase energy."	Respondent 5
			"I also drink Anlene milk 3x a week because of its benefits for bones and usually relieves fatigue."	Respondent 11
			"I consume milk almost every day, sis, and usually I consume canned milk. "Drinking milk is just for warming the body and the benefits are only for warming the body, sis."	Respondent 12
			"I also consume the milk from Indomaret, usually it's the Frisian flag	Respondent 9

Table 4. In-depth Interview Results

Domain	Findings	Answer Findings	Quotes From The Interview	Source
	behavior and benefits of milk	behavior and benefits of consuming milk	brand which is low fat, but that's rare, sis, especially if you want it, sis. "It has good health benefits because it contains lots of vitamins."	
Supplement oil fish	Respondents knew the behavior and benefits of fish oil supplements	Views on fish oil supplements by respondents	"I don't take supplements because I don't know the benefits and I don't have the time, sis, because I think I already get enough vitamins from vegetables and fruit."	Respondent 8
		Views on the taste and benefits of oil supplements by respondents	"I don't like drinking fish oil because it tastes fishy even though it has benefits for increasing appetite."	Respondent 1
			"I know that it is useful for stamina, but I don't like taking fish oil supplements because I don't like the fishy taste."	Respondent 5
			"I don't take fish oil supplements because I have a stomach ailment, sis, so I don't like anything strange, especially fish oil supplements because it's fishy and makes me nauseous, even though the benefits are actually good for the body, the effect is only short-lived, sis."	Respondent 7

Source: Primary Data, 2023

DISCUSSION

The results obtained from filling out a questionnaire by 125 respondents were followed by interviews with 15 respondents regarding the risk of vitamin D deficiency in active smokers. This research focuses on exploring and exploring the risk of vitamin D deficiency, starting with the domain regarding smoking, the domain regarding sources of vitamin D, the sunlight exposure domain, the radiation domain, and the physical activity domain.

Based on the data obtained, 125 respondents were active smokers, most of whom smoked for more than 10 years, with the highest number of cigarettes smoked in a day ranging from 11-24 cigarettes, so it can be seen that active smokers in Rungkut Village can be categorized as light smokers, with a respondent frequency of 57 respondents. This relatively high number of smokers is in line with previous research by Salsabila et al.²⁵ Central Java, Yogyakarta, East Java, Bali, NTB, South Kalimantan and South Sulawesi), there were 12,591 research subjects. Based on age, smokers in Indonesia have a median of 38 years. Based on gender, the

majority of smokers in Indonesia¹⁹ are men, namely 11,908 people (95%). The type of cigarette most frequently used is filtered kretek cigarettes with an average of 12 cigarettes/day. Indonesia is the country with the highest number of smokers in ASEAN. In South Asia, smoking tobacco (25.2%) and smokeless tobacco (24.7%) is a widespread habit among men. Data from a nationally representative cross-sectional study (Demographic and Health Survey) conducted in Afghanistan, India, Maldives, Nepal and Pakistan from 2015–2018. The subjects were men between 15 and 49 years old. Results found that higher age, lower education, lower wealth status, and involvement in any occupation were strongly associated with smoking (p-value <0.001).²⁶

Cigarette smoke has pro-inflammatory properties which can weaken host cells in the body and can interfere with the metabolism of vitamin D in the lungs. This is supported by previous research which shows that cigarette smoke can disrupt the potential pathways of the endocrine system so that it can trigger vitamin D deficiency.⁶ Previous research by Yang et al, shows smokers have lower vitamin D

concentrations than non-smokers. So, smokers can trigger a more significant vitamin D deficiency than non-smokers.⁴ Based on interviews conducted, respondents were asked for their reasons for smoking cigarettes. As for respondents who said that cigarettes are something that can relieve and reduce pain or have a calming effect on users (Table 4). From the results of the quote above, several respondents experience stress and even dizziness if they do not smoke and think that cigarettes can relieve stress or dizziness. This is supported by previous research that cigarettes can have a positive impact by eliminating stress and dizziness and providing a sense of calm against disturbing thoughts. The nicotine content in cigarettes gives users peace of mind.²⁷

In general, conventional tobacco can be used as smoked tobacco or smokeless tobacco. Apart from cigarettes, tobacco is consumed in other forms such as loosely chewed tobacco leaves, *gus*, *naswar*, *gutka* tobacco, and tobacco paste. Scientific analysis of these smokeless tobacco products has revealed more than 20 chemical compounds known to be cancer-causing agents, which include tobacco-specific nitrosamines, N-nitrosamine acids, volatile N-nitrosamines, polycyclic triggering hydrocarbons, and aldehydes.²⁸ There are several interview quotes above that the price of using tobacco is cheaper than the price of cigarettes from the factory. Based on previous research, tobacco prices are more affordable because the weather makes tobacco production in Indonesia uncertain. If the quality of the tobacco is not good, the price of the tobacco will be sold cheaper.²⁹ Besides the price of tobacco being cheaper than factory cigarettes, rising cigarette prices will influence the high prevalence in Indonesia. This is because cigarette consumption, especially among adults, is greatly influenced by cigarette prices. The increase in cigarette prices is a preventative effort carried out by the government to reduce cigarette consumption, which currently has the highest number of smokers in the Asian region.²²

The results showed most respondents consumed fish such as tuna, *pindang* fish, salted fish, milkfish, catfish, and tilapia fish. The most common way of serving fish is by frying rather than grilling or steaming. It can be seen from the interview excerpt above that respondents 6, 8, 15, 13, and 11 mostly consume fish even though

some respondents have inconsistent views regarding the benefits of consuming fish (Table 4). Fish is one of the biggest contributors to vitamin D intake, such as catfish, milkfish, and snapper, which contain around 400 IU per 85 grams of fish. The frequency of fish consumption can affect the concentration of vitamin D, so frequent fish consumption can be recommended to prevent vitamin D deficiency. The process of cooking fish does not affect the vitamin D content in food products because vitamin D is resistant to heat, such as by grilling. However, the amount of vitamin D will decrease by 50% if frying uses cooking oil due to the fat-soluble nature of vitamin D.³⁰ Different cooking processes (fried, boiled, grilled) will affect the contents of the fish. Changes in the amount of vitamins and protein are significantly higher when fried. The change in the amount of fat will be higher in frying fish than in boiling or grilling. It was stated in this research that the cooking method by boiling is the best for healthy food.³¹ A previous study by Sridonpai et al., determined the vitamin D content of commonly consumed fish in Thailand and the effect of different cooking methods on vitamin D retention.³² The results showed that vitamin D3 was the only form of vitamin D detected in fish. The vitamin D content of raw freshwater fish ranges from 2.42 to 48.5 µg per 100 g Edible Portion (EP), higher than that of raw sea fish (2.94 to 4.69 µg per 100 g EP).³² Previous research by Andrade et al., a cross-sectional online study, was conducted among 1,637 adults.³³ Participants were considered vitamin D deficient and consumed an average of 347.05 ± 307.8 µg of vitamin D through foods/beverages. Overall, though, participants, on average, did not meet the vitamin D RDAs (Recommended Daily Allowances) for healthy people. This is mainly related to participants rarely or never consuming foods high in vitamin D such as fish and dairy products.

The results show that most respondents consume eggs such as chicken and duck. The most common way of serving fish is frying compared to boiling (Table 4). On average, respondents consume eggs and are aware of their habits regarding egg consumption. Additionally, respondents understand the benefits of consuming eggs. Eggs are a rich source of protein containing ample amounts of

vitamin D and antioxidants, which help prevent aging by aiding in cell degradation. Vitamin D-rich eggs are found in egg yolks at 20 IU per yolk. The cooking process significantly affects the vitamin D content in eggs. According to USDA data from 2015, raw and boiled eggs contain 41 IU/50 grams of vitamin D3, while fried eggs contain 10 IU/46 grams of vitamin D3.³⁴ Eggs are among the few rich dietary sources of vitamin D. The results of the study by Vičič et al., revealed that fortifying eggs and milk (including yogurt) could effectively increase vitamin D intake in Slovenian women aged 44–65 by almost fivefold, thereby reducing the prevalence of vitamin D deficiency.³⁴ However, a previous study by Daly et al., evaluated the dose-response effect of consuming 2, 7, or 12 commercially available eggs per week on serum 25(OH)D concentrations. The results indicated that consuming 7 eggs per week for 12 weeks effectively reduced the decline in circulating vitamin D concentrations during winter in young Australian adults, with 12 eggs per week not providing any additional benefit.³⁵

Based on the results of the questionnaire answered by respondents, it was found that 79 respondents consumed milk most frequently, using liquid packaged milk (ready to eat) and condensed milk (sachets) (Table 4). Respondents consume milk such as ready-to-eat packaged milk. Milk is a fortified product containing vitamin D. Fortified milk must contain 400 IU of vitamin D per litre, whereas pure cow's milk only has 40 IU per litre. Therefore, the best milk is milk with vitamin D fortification in the form of ready-to-eat liquid packaged milk. This research is also similar to other research by Lorensia et al., who analyzed the intake of foods containing vitamin D in 100 adults, showing that eggs and milk contain vitamin D that are most consumed almost every day (98.50%). The incidence of vitamin D deficiency is widespread worldwide, affecting individuals of all ages, including young and healthy subjects, as well as pregnant and elderly women, due to various factors such as inadequate sun exposure, skin pigmentation, adiposity, lifestyle, and low food intake.³⁰ To address this issue, fortification of daily consumed foods such as milk, is highly recommended.³⁶ A theoretical model applied vitamin D3 fortifications of 1 µg, 1.5 µg and 2

µg/100g to simulate improvements in vitamin D intakes. Mean ± SD vitamin D3 in whole milk was 0.06 ± 0.02 µg/100g.³⁷ Previous research by Zhang et al., explored interaction between passive smoking and dietary undernutrition on the risk of vitamin D deficiency. The interaction between passive smoking and inadequate nutrition influences vitamin D risk D deficiency.³⁸ Passive smoking may amplify the effect of poor nutrition on the risk of vitamin D deficiency. The more often cigarettes one consumes, nicotine content enters CNS and CNS, putting an individual's food intake.³⁹ Research has been conducted on consuming foods containing vitamin D on vitamin to prevent deficiency,³⁰ but there has been no research linking the impact of active on the population and population, and no research no assessed that influences influencing levels through the intake of foods containing vitamin D.

Based on the results obtained, the highest frequency of respondents was found among those who did not consume fish oil supplements. Some reasons for not consuming fish oil supplements included being unaccustomed to them and disliking the fishy smell or taste associated with them (Table 4). Respondents elaborated on their reasons for not taking fish oil supplements, leading to most of respondents not taking supplements containing vitamin D. According to the results, the highest frequency of respondents was among those who did not consume vitamins (Table 4). The majority of respondents, as per the interview excerpt, do not take vitamins, despite some being aware of the benefits they offer to the body. Previous research by Lorensia and Suryadinata,⁴⁰ indicated that many individuals are hesitant to use fish oil supplements independently due to the large size of the soft capsules, which can be uncomfortable to swallow, and the fishy odor. However, this research also demonstrates that fish oil supplements are effective in enhancing lung function in smokers.⁴⁰

The questionnaire results showed that as many as 68 respondents experienced color changes when exposed to sunlight, with the most frequent color changes occurring in blackish red. From the interview excerpts of the two respondents above, the respondents experienced changes in skin color when exposed to sunlight. This is supported by research

conducted by Carlberg,⁴¹ skin color will change to pink or even black if the skin has received enough vitamin D. As for several other respondents who experienced something different from the previous respondents, respondents 3 and 9 did not experience changes in skin color when exposed to sunlight (Table 4). The respondent did not experience changes in skin color. The darker a person's skin color, the longer it takes to form vitamin D in the skin. According to Raymond-Lezman and Riskin,⁴² in their research, they explain that someone with dark skin has large amounts of melanin in the epidermis, which will make melanin oppose vitamin D receptors in the skin. Melanin will absorb UVB light exposure, reducing the absorption of UVB light exposure for vitamin D receptors in the skin. A person with dark skin takes 10-50 times more exposure to sunlight than a person with light skin to produce the same amount of vitamin D.⁴²

The results of the questionnaire showed that 67 respondents who took the time to sunbathe directly, which was mostly found at 07.00-09.00 WIB and by using personal protective equipment such as hats/jackets and long clothes (Table 4). Many respondents spend time sunbathing, but when the intensity of sun exposure is low, many respondents also use personal protective equipment and long clothing, which is at risk of experiencing a deficiency. Respondents took time to sunbathe between 07.00-09.00 WIB and wore personal protective equipment and closed clothing. Previous research explained that the UVB intensity of sunlight is low at 07.00-08.00 WIB, while at 09.00 WIB, the intensity achieved increases, but it takes a long time of around 25 minutes to sunbathe directly. A good time to be exposed to sunlight is from 10 AM to 2 PM, because at that time UVB rays are more than UVA with a relatively stable and high intensity. However, to be able to see the adequacy of sun exposure not only at the time of exposure, the use of long clothing and personal protection also has an influence. So, it can be concluded that someone who has insufficient sun exposure and has body protection such as [39] sunscreen, hats, jackets and long clothes will be at risk of vitamin D deficiency or vitamin D deficiency.⁴³

From the results obtained, almost all respondents use gadget screens frequently

(Table 4). Respondents use gadget screens for a long time, up to 7 hours or more. In the research of Shan et al.⁴⁴ the use of gadget screens is associated with physical activity on vitamin D concentrations. The time of gadget screen use has a significant relationship with a person's level of physical activity. So, screen use and physical activity are both markers of exposure to UVB rays. This is [45] supported by research by Rocka et al.⁴⁵ that there is a significant relationship between gadget screen use and physical activity on a person's vitamin status. A good duration of time to stare at a gadget screen is ≤ 2 hours/per day.

Based on the results obtained, the highest frequency of respondents was 96 respondents who carried out moderate levels of physical activity (Table 4). Respondents carried out moderate levels of physical activity such as doing housework as explained by respondents 10, 3, 13, and 14, while respondents 4 and 8 carried out heavy levels of physical activity. Vitamin D levels in the blood are related to muscles and bones and play a role in energy metabolism, oxidative stress, maintenance and increasing physical fitness.⁴⁶ The measurement of physical activity in this study only assessed physical activity from respondents' confessions but did not measure in quantitative detail the level of physical activity. Measuring physical activity quantitatively generally uses the IPAQ (International Physical Activity Questionnaire) questionnaire which was used in Indonesia by Lorensia et al.^{47,48} Previous research by Lorensia et al., determine differences in lung function and levels of physical activity between smokers and non-smokers. The results show that a significant difference in lung function values was observed between smokers and non-smokers ($p=0.00$).⁴⁸

CONCLUSION AND RECOMMENDATION

Most of the respondents were light smokers with normal blood levels of vitamin D. Most respondents often consume mackerel and pindang fish, even though they are served fried. Most respondents also consumed a lot of fried chicken eggs and drank bottled cow's milk. Smokers with vitamin [42] deficiency and insufficiency tend to have a low intake of foods containing vitamin D, such as fish, eggs and milk. Intake of fish oil and vitamin supplements is still rare. Most respondents rarely sunbathe and are

only exposed to sunlight in the morning. Respondents also use gadgets a lot and the frequency of physical activity tends to be rare and moderate³

Therefore, to reduce the risk of vitamin D deficiency, a smoker needs to improve his lifestyle by reducing the frequency of smoking, increasing his intake of foods high in vitamin D (fish, eggs, milk), increasing the frequency of exposure to sunlight, reducing the frequency of using gadgets and increasing activity physique.

ACKNOWLEDGMENTS

This research was funded by Institute of Research and Community Service by Universitas Surabaya.

AUTHOR CONTRIBUTIONS

From the research process until the writing of this article, all authors played a role in this research. The correspondent author plays a role in compiling and designing research, the 2nd acts as a data analyzer, the 3rd author acts as data collection in the field, and 4th author acts as an examiner of the clinical condition of each respondent in the field.

CONFLICTS OF INTEREST

The authors declare no conflict of interest.

REFERENCES

1. Tobacco Control Support Center-Ikatan Ahli Kesehatan Masyarakat Indonesia (TCSC-IAKMI). *Atlas tembakau Indonesia* 2020; 2020.
2. West R. Tobacco Smoking: Health Impact, Prevalence, Correlates and Interventions. *Psychology & Health*. 2017;32(8):1018–1036. <https://doi.org/10.1080/08870446.2017.1325890>
3. Varghese J, Gharde PM. A Comprehensive Review on the Impacts of Smoking on the Health of an Individual. *Cureus*. 2023;15(10):e46532. <https://doi.org/10.7759/cureus.46532>
4. Yang L, Zhao H, Liu K, et al. Smoking Behavior and Circulating Vitamin D Levels in Adults: A Meta-Analysis. *Food Science & Nutrition*. 2021;9(10):5820–5832. <https://doi.org/10.1002/fsn.3.2488>
5. Dahdah A, Jagers RM, Sreejit G, et al. Immunological Insights into Cigarette Smoking-Induced Cardiovascular Disease Risk. *Cells*. 2022;11(20):3190. <https://doi.org/10.3390/cells11203190>
6. Nikniaz L, Ghajazadeh M, Nateghian H, et al. The Interaction Effect of Aerobic Exercise and Vitamin D Supplementation on Inflammatory Factors, Anti-Inflammatory Proteins, and Lung Function in Male Smokers: A Randomized Controlled Trial. *BMC Sports Science, Medicine Rehabilitation*. 2021;13(1):102. <https://doi.org/10.1186/s13102-021-00333-w>
7. Chen YC, Sung HC, Chuang TY, et al. Vitamin D3 Decreases TNF- α -Induced Inflammation in Lung Epithelial Cells Through a Reduction in Mitochondrial Fission and Mitophagy. *Cell Biology and Toxicology*. 2022;38(3):427–450. <https://doi.org/10.1007/s10565-021-09629-6>
8. Tian T, Jiang X, Qin R, et al. Effect of Smoking on Lung Function Decline in a Retrospective Study of a Health Examination Population in Chinese Males. *Front Med (Lausanne)*. 2023;9:843162. <https://doi.org/10.3389/fmed.2022.843162>
9. Augustine LF, Nair KM, Kulkarni B. Sun Exposure as a Strategy for Acquiring Vitamin D in Developing Countries of Tropical Region: Challenges & Way Forward. *Indian J Med Res*. 2021;154(3):423–432. https://doi.org/10.4103/ijmr.IJMR_1244_18
10. Lorensia A, Suryadinata RV, Inu IA. Comparison of Vitamin D Status And Physical Activity Related With Obesity in Student. *Journal of Applied Pharmaceutical Science*. 2022;12(4):108–118. <https://doi.org/10.7324/JAPS.2022.120412>
11. Lorensia A, Suryadinata RV. Smoking Cessa-

- tion Experience and Socioeconomic Status of Online Motorcycle Taxi Drivers in Surabaya. *Media Kesehatan Masyarakat Indonesia*. 2021;17(3):80-87. <https://doi.org/10.30597/mkmi.v17i3.14059>
12. Octavius GS, Shakila A, Meliani M, et al. Vitamin D Deficiency is a Public Health Emergency among Indonesian Children and Adolescents: A Systematic Review and Meta-Analysis of Prevalence. *Annals of Pediatric Endocrinology & Metabolism: apem*. 2023;28(1):10-19. <https://doi.org/10.6065/apem.2244170.085>
 13. Suryadinata RV, Lorensia A. Frekuensi Asupan Makanan, Pengetahuan Vitamin D dan Obesitas Pada Kelompok Usia Lanjut Food Frequency, Knowledge about Vitamin D and Obesity among Elderly. *Amerta Nutrition*. 2020;4(1):43-48. <https://doi.org/10.2473/amnt.v4i1.2020>
 14. Widyastuty AASA, Jihan JC. Tingkat Kekumuhan dan Analisis Spasial Permukiman Kumuh Perkotaan (Studi Kasus: Surabaya Timur). *Jurnal Teknik Waktu*. 2018;16(2):47-55. <https://doi.org/10.36456/waktu.v16i02.1667>
 15. Holipah H, Sulistomo HW, Maharani A. Tobacco Smoking and Risk of All-Cause Mortality in Indonesia. *PLoS One*. 2020;15(12):e0242558. <https://doi.org/10.1371/journal.pone.0242558>
 16. Herath P, Wimalasekera S, Amarasekara T, et al. Effect of Cigarette Smoking on Smoking Biomarkers, Blood Pressure and Blood Lipid Levels Among Sri Lankan Male Smokers. *Postgraduate Medical Journal*. 2022;98(1165):848-854. <https://doi.org/10.1136/postgradmedj-2021-141016>
 17. Sugiura T, Dohi Y, Takagi Y, et al. Close Association between Subclinical Atherosclerosis and Pulmonary Function in Middle-Aged Male Smokers. *Journal of Atherosclerosis and Thrombosis*. 2020;27(11):1230-1242. <https://doi.org/10.5551/jat.55996>
 18. Sutherland JP, Zhou A, Hyppönen E. Vitamin D Deficiency Increases Mortality Risk in the UK Biobank: A Nonlinear Mendelian Randomization Study. *Annals of Internal Medicine*. 2022;175(11):1552-1559. <https://doi.org/10.7326/M21-3324>
 19. AlSubai A, Baqai MH, Agha H, et al. Vitamin D and Preeclampsia: A Systematic Review And Meta-Analysis. *SAGE Open Medicine*. 2023;11:20503121231212093. <https://doi.org/10.1177/20503121231212093>
 20. Sherief LM, Ali A, Gaballa A, et al. Vitamin D Status and Healthy Egyptian Adolescents: Where Do We Stand?. *Medicine (Baltimore)*. 2021;100(29):e26661. <https://doi.org/10.1097/MD.00000000000026661>
 21. Moghadasi N, Alimohammadi I, Safari Variani A, et al. The Effect of Mobile Radiation on the Oxidative Stress Biomarkers in Pregnant Mice. *Journal of Family and Reproductive Health*. 2021 Sep;15(3):172-178. <https://doi.org/10.18502/jfrh.v15i3.7134>
 22. Nguyen CV, Le TT, Nguyen NH. The Impact of Cigarette Prices on Smoking Participation and Tobacco Expenditure in Vietnam. *PLoS One*. 2021;16(12):e0260415. <https://doi.org/10.1371/journal.pone.0260415>
 23. Septiani R. Hubungan Lama Merokok dan Frekuensi Merokok dengan Kadar Hemoglobin (HB) Pada Perokok Aktif. *Babul Ilmi: Jurnal Ilmiah Multi Science Kesehatan*. 2022;14(1):30-40. <https://jurnal.stikes-aisyiyah-palembang.ac.id/index.php/Kep/article/view/809>
 24. Weir CB, Jan A. *BMI Classification Percentile And Cut Off Points*. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2023.
 25. Salsabila NN, Indraswari N, Sujatmiko B. Gambaran Kebiasaan Merokok Di Indonesia Berdasarkan *Indonesia Family Life Survey 5* (IFLS 5). *Jurnal Ekonomi Kesehatan Indonesia*. 2022;7(1):13-22. <https://journal.fkm.ui.ac.id/jurnal-eki/article/view/5394>

26. Islam MS, Rashid M, Sizar MI, et al. Cigarette Smoking and Associated Factors Among Men in Five South Asian Countries: A Pooled Analysis of Nationally Representative Surveys. *PLoS One*. 2022;17(11):e0277758. <https://doi.org/10.1371/journal.pone.0277758>
27. Foulds J, Veldheer S, Pachas G, et al. The Effects of Reduced Nicotine Content Cigarettes on Biomarkers of Nicotine and Toxicant Exposure, Smoking Behavior and Psychiatric Symptoms in Smokers with Mood or Anxiety Disorders: A Double-Blind Randomized Trial. *PLoS One*. 2022;17(11):e0275522. <https://doi.org/10.1371/journal.pone.0275522>
28. Omare MO, Kibet JK, Cherutoi JK, et al. A Review of Tobacco Abuse And Its Epidemiological Consequences. *Zeitschrift Fur Gesundheitswissenschaften = Journal of Public Health*. 2022;30(6):1485-500. <https://doi.org/10.1007/s10389-020-01443-4>
29. Golestan YP, Kalan ME, Taleb ZB, et al. The Effect of Price on Cigarette Consumption, Distribution, and Sale In Tehran: A Qualitative Study. *BMC Public Health*. 2021;21(1):1720. <https://doi.org/10.1186/s12889-021-11733-5>
30. Lorensia A, Suryadinata RV, Arganitya GN. Relationship of Food Vitamin D Intake with Obesity in Adolescent in Surabaya. *Global Medical & Health Communication*. 2022;10(2):104-110. <https://ejournal.unisba.ac.id/index.php/gmhc/article/view/9024>
31. AlFaris NA, Alshammari GM, AlTamimi JZ, et al. Evaluating The Effects of Different Processing Methods on The Nutritional Composition of Shrimp and The Antioxidant Activity of Shrimp Powder. *Saudi Journal of Biological Sciences*. 2022;29(1):640-9. <https://doi.org/10.1016/j.sjbs.2021.09.029>
32. Sridonpai P, Judprasong K, Tirakompong N, et al. Effects of Different Cooking Methods on the Vitamin D Content of Commonly Consumed Fish in Thailand. *Foods*. 2022;11(6):819. <https://doi.org/10.3390/foods11060819>
33. Andrade JM, Grandoff PG, Schneider ST. Vitamin D Intake and Factors Associated With Self-Reported Vitamin D Deficiency Among US Adults: A 2021 Cross-Sectional Study. *Frontiers in Nutrition*. 2022;9:899300. <https://doi.org/10.3389/fnut.2022.899300>
34. Vičič V, Mikuš RP, Kugler S, et al. Vitamin D Fortification of Eggs Alone and in Combination with Milk in Women Aged 44-65 Years: Fortification Model and Economic Evaluation. *Slovenian Journal of Public Health*. 2022;62(1):30-33. <https://doi.org/10.2478/sjph-2023-0005>
35. Daly RM, De Ross B, Gianoudis J, et al. Dose-Response Effect of Consuming Commercially Available Eggs on Wintertime Serum 25-Hydroxyvitamin D Concentrations in Young Australian Adults: a 12-Week Randomized Controlled Trial. *The Journal of Nutrition*. 2022;152(7):1702-1710. <https://doi.org/10.1093/jn/nxac044>
36. Pellegrino L, Marangoni F, Muscogiuri G, et al. Vitamin D Fortification of Consumption Cow's Milk: Health, Nutritional and Technological Aspects. A Multidisciplinary Lecture of the Recent Scientific Evidence. *Molecules*. 2021;26(17):5289. <https://doi.org/10.3390/molecules26175289>
37. Weir RR, Johnston M, Lowis C, et al. Vitamin D₃ Content of Cows' Milk Produced in Northern Ireland and Its Efficacy As A Vehicle For Vitamin D Fortification: A UK Model. *International Journal of Food Sciences and Nutrition*. 2021;72(4):447-455. <https://doi.org/10.1080/09637486.2020.1837743>
38. Zhang Y, Lin S, Li J, et al. Interaction of Passive Smoking and Diet Habits on Vitamin D Deficiency among Women of Reproductive Age in Rural Central China. *Nutrients*. 2022;15(1):126. <https://doi.org/10.3390/nu15010126>

39. Tiwari RK, Sharma V, Pandey RK, et al. Nicotine Addiction: Neurobiology and Mechanism. *Journal of Pharmacopuncture*. 2020;23(1):1-7.
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7163392/>
40. Lorensia A, Suryadinata RV. Profil of Omega-3 Food Intake and Its Association with Socioeconomic Status on Online Motorcycle Driver Smokers. *Healthcare in Low-resource Settings*. 2023;11(s1):11164.
<https://www.pagepressjournals.org/index.php/hls/article/view/11164>
41. Carlberg C. Vitamin D and Pigmented Skin. *Nutrients*. 2022;14(2):325.
<https://doi.org/10.3390/nu14020325>
42. Raymond-Lezman JR, Riskin SI. Benefits and Risks of Sun Exposure to Maintain Adequate Vitamin D Levels. *Cureus*. 2023;15(5):e38578.
<https://doi.org/10.7759/cureus.38578>
43. Sultana N. Sun Awareness and Sun Protection Practices. *Clinical, Cosmetic and Investigational Dermatology*. 2020;13:717-730.
<https://doi.org/10.2147/CCID.S265477>
44. Shan L, Dong H, Wang T, et al. Screen Time, Age and Sunshine Duration Rather Than Outdoor Activity Time Are Related to Nutritional Vitamin D Status in Children With ASD. *Frontiers in Pediatrics*. 2022;9:806981.
<https://doi.org/10.3389/fped.2021.806981>
45. Rocka A, Jasielska F, Madras D, et al. The Impact of Digital Screen Time on Dietary Habits and Physical Activity in Children and Adolescents. *Nutrients*. 2022;14(14):2985.
<https://doi.org/10.3390/nu14142985>
46. Zhang J, Cao ZB. Exercise: A Possibly Effective Way to Improve Vitamin D Nutritional Status. *Nutrients*. 2022;14(13):2652.
<https://doi.org/10.3390/nu14132652>
47. Lorensia A, Suryadinata RV, Chandra NLMR. Profile of Vitamin D Status, Physical Activity, and Lung Health in Construction Workers. *Community Medicine and Public Health of Indonesia Journal*. 2020;1(2):117-24.
<http://comphi.sinergis.org/index.php/comphi/article/view/19>
48. Lorensia A, Muntu CM, Suryadinata RV, et al. Effect of lung function disorders and physical activity on smoking and non-smoking students. *Journal of Preventive Medicine and Hygiene*. 2021; 62(1):89-96.
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8283647>