

RESEARCH ARTICLE

Relationship of Physical Activity and Vitamin D Levels in Elderly Women

Amelia Lorensia,¹ Marthy Meliana Ariyanti Jalmav,² Rivan Virlando Suryadinata,³
Muhammad Reza Aprililo Akbar⁴

¹Department of Clinical-Community Pharmacy, Faculty of Pharmacy, Universitas Surabaya, Surabaya, Indonesia,

²Faculty of Pharmacy, Universitas Anwar Medika, Sidoarjo, Bandung, Indonesia,

³Department of Public Health, Faculty of Medicine, Universitas Surabaya, Surabaya, Indonesia,

⁴Faculty of Pharmacy, Universitas Surabaya, Surabaya, Indonesia

Abstract

Older adults are at risk of vitamin D deficiency, especially in older women, due to a decrease in the hormone estrogen, which causes decreased bone density and increased risk of fractures. There is a relationship between vitamin D deficiency and lack of physical activity due to the storage of vitamin D in adipose tissue. Physical activity was measured using the International Physical Activity Questionnaire (IPAQ). The aim was to determine the relationship between physical activity and vitamin D in elderly women. This cross-sectional analytic observational study was conducted at Nursing Home X in Surabaya city from May to August 2023. The variables were physical activity (measured by IPAQ to estimate physical activity levels) and vitamin D level (measured by VIDAS® instrument to determine 25(OH) levels). The subjects were all women aged ≥ 60 years. The sampling technique uses the purposive sampling method. For an analysis of the relationship between physical activity levels and vitamin D levels, the Pearson correlation test was used with a ratio data scale. This research received 49 respondents. The average physical activity level was 1316.82 ± 720.90 , and most respondents had a moderate physical activity level of 44.90%. Vitamin D levels were ≤ 30 ng/ml for 46 respondents (93.87%). The results of the Pearson correlation test were obtained with a $r_{\text{value}} = 0.089$ and a significance value (Sig.) of 0.542. There was no significant correlation between physical activity and vitamin D levels. Other factors such as age, health conditions, and vitamin D intake can affect the results. Further research can be developed into measurements to examine physical activity in old age.

Keywords: IPAQ, elderly woman, nursing home, physical activity, vitamin D

Introduction

Vitamin D deficiency is a common global health problem with high prevalence, especially in South Asia countries. Vitamin D mainly regulates calcium and phosphorus metabolism and promotes bone growth.¹ Older adults are at risk of vitamin D deficiency because vitamin D production and metabolism change with age due to various factors, such as reduced sun exposure and skin production capacity.² Physically, geriatrics are less active and have poor nutritional status. In addition, there are physiological processes that exacerbate the decrease in vitamin D levels in the body, such as decreased production of vitamin D in the skin after sun exposure caused by skin atrophy, eating foods that are low in vitamin D, impaired

gastrointestinal absorption, and decreased production $1,25(\text{OH})_2\text{D}$ in the kidney.³ Vitamin D deficiency is prevalent in postmenopausal women, and this undoubtedly exacerbates the risk of cardiovascular disease and menopause-related dyslipidemia. Estrogen levels drop significantly as women go through menopause, and a lack of estrogen weakens bones. Vitamin D helps the body absorb and use calcium to maintain bone structure. 25-hydroxyvitamin D [$25(\text{OH})\text{D}$] serves the purpose of calcium absorption and appears to be hormonally sensitive. Vitamin D deficiency is associated with low bone mass and increased fracture risks. Menopausal women are at risk of decreased bone density and increased risk of fractures due to decreased estrogen levels,¹ reduced vitamin D synthesis in the skin, or changes in body composition relevant to vitamin

Copyright ©2024 by authors. This is an open access article under a Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License (<https://creativecommons.org/licenses/by-nc-sa/4.0>).

Received: 20 February 2024; Revised: 10 October 2024; Accepted: 10 December 2024; Published: 20 December 2024

Correspondence: Amelia Lorensia. Department of Clinical-Community Pharmacy, Faculty of Pharmacy, Universitas Surabaya. Jln. Raya Kalirungkut, Surabaya 60292, East Java, Indonesia. E-mail: amelia.lorensia@gmail.com, amelia.lorensia@staff.ubaya.ac.id

D status and physiology.⁴ Vitamin D deficiency is defined as vitamin D levels below 20 ng/ml (50 nmol/liter), and vitamin D deficiency is 21–29 ng/ml (525–725 nmol/liter). Vitamin D deficiency can also be a risk factor for diseases such as immune system diseases and infectious diseases caused by coronavirus disease 2019 (COVID-19).⁵ Research by Pereira et al.⁶ reported that low vitamin D levels were associated with COVID-19.

Research by Cui et al.⁵ shows that 15.7% of the global adult population has vitamin D deficiency. Vitamin D deficiency can occur in all age groups, including menopausal women. Indonesia currently has 7.4% of menopausal women from the total population, and in 2020, it is estimated to reach 11.54% with an average age of menopause of 49 years. According to Statistics Indonesia, in 2015, the number of women in Indonesia who entered menopause reached 21.22 million. It is estimated that by 2025, there will be 60 million menopausal women.⁷ Another study on the incidence of vitamin D deficiency in Indonesian women showed that 95% of them (148 subjects out of a total of 156) had vitamin D deficiency.⁸

There is a relationship between vitamin D deficiency and lack of physical activity due to the storage of vitamin D in adipose tissue due to its high fat solubility and decreased exposure to sunlight in obese subjects due to limited physical activity and limited mobility. It is proven that physical activity increases glucose, calcium, and vitamin D metabolism and reduces body weight by increasing lipolysis.⁹ Someone who does high physical activity will be associated with increased vitamin D concentration. Suppose someone does less physical activity (low physical activity). In that case, it will cause a decrease in vitamin D concentration, while someone with high physical activity will most likely be associated with optimal vitamin D levels.¹⁰ Vitamin D can also help improve the body's performance in physical activities in populations at risk of vitamin D deficiency. Previous research by Nascimento et al.¹¹ found a significant increase in levels and improvements in the body, so there was an interaction between vitamin D levels and physical activity in maintaining body performance.

Measurement of physical activity in this study used a questionnaire, namely the extended version of the International Physical Activity Questionnaire (IPAQ), which is used to estimate

physical activity levels. Data on the relationship between physical activity and vitamin D levels in Indonesia have been collected by Suryadinata et al.,¹² conducted at Public Health Center Taman, Sidoarjo district, East Java, in March–July 2017. The results showed geriatrics with obesity and non-obesity had similar levels of physical activity, but vitamin D status in obesity tended to be lower than non-obese. However, similar studies have never been conducted on elderly women in nursing homes. These data would greatly assist health workers in improving the quality of life of these patients. This research aimed to determine the relationship between physical activity and vitamin D in elderly women.

Methods

This study used analytic observational research with a cross-sectional design. Data was collected from May 2023 to August 2023 at Nursing Home X in Surabaya city. The Institutional Ethical Committee of the University of Surabaya approved the study, 207/KE/VIII/2023.

This study's population was all women aged ≥ 60 who had experienced menopause in Nursing Home X in Surabaya city. The samples taken in this study were all populations that met the inclusion criteria. The sampling technique used is purposive sampling, which meets the requirement of not having physical limitations such as motor disorders.

Each respondent will be interviewed to fill out the IPAQ and take blood samples to check vitamin D levels. A phlebotomist from Pramita Laboratorium will take a blood sample and test it at an accredited laboratory.

Blood sampling was carried out simultaneously during the day by the phlebotomist, taking into account the time of blood sampling, which can affect vitamin D levels.¹³ After blood collection, the laboratory will process the blood by centrifuging to obtain serum, which will then be tested for vitamin D levels using VIDAS® 25-OH Vitamin D Total (VITD), which is an automatic quantitative test used on the VIDAS® instrument to determine 25(OH) levels. D total in human serum or plasma with the enzyme-linked fluorescence assay (ELFA) technique.

The method for measuring physical activity uses the modified IPAQ. The physical activity questionnaire contains 12 questions about light,

moderate, and vigorous physical activity activities. Respondents were interviewed, and then results were obtained, which categorized respondents into mild, moderate, and vigorous categories using the metabolic equivalent of task (MET). Physical activity is divided into three categories, namely the mild category (<600 MET minutes/week), the moderate category (600–1,500 MET minutes/week), and the vigorous category (>1,500 MET minutes/week). This questionnaire contains questions about the type of activity, duration, and frequency of a person doing physical activity in the last 7 days. Physical activity can be measured by measuring the amount of energy expended/needed in each minute of activity. According to the IPAQ assessment protocol, MET-minutes/week for a given activity (walking moderate-intensity activity or vigorous-intensity activity) is calculated by multiplying the MET value of the given activity (3.3 for walking, 4.0 for moderate-intensity activity, and 8.0 for high-intensity activities) according to hours spent in a particular activity, for example walking at work = $3.3 \times \text{minutes walking} \times \text{days walking at work}$. MET-minutes/week = (MET \times activity minutes \times days). Information: MET-minutes/week = METs produced in one week MET = MET value based on activity level minutes = duration of activity in one day days = number of days in one activity.^{14,15} Results are presented as estimates of energy expenditure in metabolic equivalent minutes per week (MET-hours/week).^{16,17} The metabolic equivalent of task, or MET, is valid for describing energy expenditure from certain activities. MET is the ratio of the rate of energy expended during activity to the rate of energy expended at rest.¹⁸

Before use, the questionnaire (IPAQ)^{11,16,19} must be used as a research instrument, and a validity test must first be carried out, consisting of content and construct validation. Content validity through rational analysis or expert judgment by a panel to ensure that measurements are representative. Meanwhile, construct validity refers to the extent to which a measuring instrument shows results following theory. Reliability shows that the questionnaire is reliable and consistently provides the same results when measured twice or more with the same instrument. Cronbach's alpha value was >0.6, so the research instrument would be reliable.

The relationship between physical activity levels and vitamin D levels was analyzed using the

Pearson correlation test, with the data scale being the ratio. The data was tested for normality using the Shapiro-Wilk test; if the p-value > 0.05, it was concluded that the data was normally distributed. The relationship test used the Pearson correlation test; if the p-value < 0.05, it can be concluded that there is a significant relationship between physical activity and vitamin D levels.

Results

This research involved 49 geriatric women as respondents. Data were collected using physical activity questionnaire interviews and vitamin D serum tests. The study involved elderly women aged 60 years and over. The highest number of high-risk elderly women >70 years was 43 respondents, while the rest were in the 60–70-year age group.

Regarding body mass index (BMI) characteristics, most respondents were of normal weight, and 34 respondents had an average standard deviation of $21.58 \pm 2.16 \text{ kg/m}^2$. Most of them had no disease history (Table 1).

Based on the data in Table 2, respondents' most common physical activity profile was exercise (44 respondents) and walking 100 m (39 respondents). At the same time, physical activities that were only done by some respondents were cooking and cleaning the garden or the like. Seven respondents carry out physical activity in the mild category, 22 in the moderate category,

Table 1 Frequency Distribution of Respondent Characteristics

Characteristics	n=49
Age (years)	
60–70	6
>70	43
BMI (kg/m ²)	
Very thin (<16)	6
Skinny (16–18.4)	6
Normal (18.5–24.9)	34
Overweight (25–29.9)	3
Obesity (>30)	0
Medical history	
Type 2 diabetes mellitus	5
Hypertension	10
Osteoarthritis	7
Osteoporosis	3
There isn't any	27

Table 2 Profile of Respondents' Physical Activity Questionnaire Answers

No.	Types of Physical Activity	Answer (Minutes)	n=49
1	Walk 100 m	5	1
		10	5
		15	6
		20	1
		30	23
		60	3
		Didn't do	10
		2	Walk more than 100 m
15	3		
60	12		
Didn't do	32		
3	Cook	Didn't do	49
4	Wash clothes by hand	10	4
		15	2
		20	2
		30	9
		Didn't do	32
5	Cleaning room	5	1
		10	5
		15	1
		30	2
		Didn't do	40
6	Exercise	10	7
		15	13
		20	4
		25	6
		30	4
		45	3
		60	8
		Didn't do	5
7	Cleaning the garden	Didn't do	49
8	Gardening/farming	30	2
		Didn't do	47
9	Sew with hand	10	2
		15	3
		20	1
		60	4
		Didn't do	39
10	Sewing by machine	60	1
		Didn't do	48
11	Read	10	4
		15	2
		20	1
		30	4
		60	4
		Didn't do	34
12	Watch TV	5	1
		10	3
		15	3
		20	1
		30	6
		60	2
Didn't do	33		

and 20 in the vigorous category. The average physical activity level was 1316.82 ± 720.90 , and most respondents had a moderate physical activity level of 44.90% (Table 3).

Vitamin D levels were measured during the research by taking blood samples from each respondent. In Table 4, the results showed that vitamin D levels^{16,17} were <20 ng/ml for 27 respondents, vitamin D levels were $20-30$ ng/ml for 19 respondents, and vitamin D levels were >30 ng/ml for as many as three respondents.

Based on Table 5, the results showed that deficiency (<20 ng/ml) for 27 respondents (mean \pm SD= 17.40 ± 2.07), insufficiency ($20-30$ ng/ml) for 19 respondents (mean \pm SD= 23.64 ± 2.42), and normal level (>30 ng/ml) for three respondents (mean \pm SD= 31.33 ± 1.15). The total mean \pm SD was 18.41 ± 6.60 . In the mild physical activity category, two respondents had vitamin D levels <20 ng/ml, and $20-30$ ng/ml for five respondents. In the category of moderate physical activity, 14 respondents had vitamin D levels <20 ng/ml, and $20-30$ ng/ml as many as eight respondents. In the category of vigorous physical activity, 11 respondents had vitamin D levels <20 ng/ml, six respondents had $20-30$ ng/ml, and three had >30 ng/ml.

The normality test results are as follows: In the Shapiro-Wilk normality test, a significance value (Sig.) was obtained, which was 0.133 for physical activity and 0.94 for vitamin D levels. Because both data have a significance value of >0.05 , it can be concluded that the physical activity data and vitamin D levels were usually distributed. The results of the Pearson correlation test are as follows: obtained an r_{value} (Pearson correlation) of 0.089 and a significance value (Sig.) of 0.542. Based on a significance value (α) >0.05 , it means there is no significant correlation between physical activity and vitamin D levels (Figure).

Discussion

This research involved 49 geriatric women as respondents. Since this study only involved 49 respondents, which may need to be more representative of a larger group, the results cannot be generalized widely and only to elderly women with similar characteristics.

Based on age, the highest number in the group of high-risk elderly women, namely 70 years and over, was 43 respondents, while in the 60-70

Table 3 Physical Activity Level of Respondents

Category Physical Activity	n=49	Mean±SD	Total Mean±SD
Mild (<600)	7	312.21±162.37	1,316.82±720.90
Moderate (600–1,500)	22	1,038.75±401.43	
Vigorous (>1,500)	20	2,074.17±581.39	

Table 4 Presentation of Data on Vitamin D Level Values

Vitamin D Level (ng/ml)	n=49	Mean±SD	Total Mean±SD
Deficiency (<20)	27	17.40±2.07	18.41±6.60
Insufficiency (20–30)	19	23.64±2.42	
Normal (>30)	3	31.33±1.15	

Table 5 Cross Tabulation of Vitamin D Levels with Physical Activity

Physical Activity	Number of Respondents based on Vitamin D Level Value			Total n=49
	Deficiency n=27	Insufficiency n=19	Normal n=3	
Mild (<600)	2	5	0	7
Moderate (600–1,500)	14	8	0	22
Vigorous (>1,500)	11	6	3	20

Note: deficiency (<20 ng/ml), insufficiency (20–30 ng/ml), normal (>30 ng/ml)




Figure Distribution of Physical Activity Data and Vitamin D Levels

year age group, there were only six respondents. Uneven age distribution may affect the results, especially physiological changes in different age groups. Age can affect vitamin D levels, such as a decreased vitamin D metabolism in old age. A decrease in calcium absorption causes this, the presence of intestinal resistance to absorption of circulating calcium $1,25(\text{OH})_2\text{D}$, a decrease in kidney function resulting in a reduction of $1,25(\text{OH})_2\text{D}$ production in the kidneys, and a decrease in vitamin D production in the skin, nine caused by decreasing the concentration of 7-dehydrocholesterol in the epidermis which can reduce the formation of vitamin D_3 . So aging can also cause calcium deficiency, hyperparathyroidism, increased bone loss, and osteoporosis.^{20,21}

The majority of respondents (27 people) had no history of illness, while only several suffered from diabetes, hypertension, or osteoarthritis. This imbalance in health characteristics can affect physical activity and vitamin D levels. History of disease can affect vitamin D levels, including type 2 diabetes mellitus. The relationship between vitamin D deficiency and insulin resistance can develop through inflammation because vitamin D deficiency is associated with increased inflammatory markers. In addition, genetic polymorphisms of vitamin D-related genes may predispose to impaired glycemic control and type 2 diabetes. Epidemiological studies suggest an association between low serum 25-hydroxyvitamin D_3 ($25(\text{OH})\text{D}_3$) concentrations and an increased risk of metabolic syndrome and type 2 diabetes. The increase in fat mass partly explains it.^{20,21}

Most respondents had low vitamin D levels (<20 ng/ml), and only a few had levels above 30 ng/ml. This uneven distribution may limit statistical power for analysis and comparison between groups. The research showed no significant correlation between physical activity and vitamin D levels. It differs from previous research by Trovato et al.,²² which stated that higher dietary intake of vitamin D and sunlight exposure are associated with a lower likelihood of having high perceived stress among physically active individuals. These results may be related to previous research by Zhang and Cao,²³ that endurance exercise can significantly increase serum $25(\text{OH})\text{D}$ levels in vitamin D-deficient people but has no significant effect on vitamin

D-sufficient people. The effects of exercise on $25(\text{OH})\text{D}$ levels in the circulation may depend on factors such as vitamin D nutritional status, exercise type and intensity, and sex.

Physical activity can be measured with a tool, namely an accelerometer, pedometer, and self-report questionnaire (IPAQ) instrument. This research uses a self-report questionnaire because it has the advantage of cheaper costs, does not burden respondents, and obtains the flow of daily physical activity from respondents. Still, the self-report questionnaire has areas for improvement, such as less accuracy and reliability, because the data is based only on respondents' memories.²⁴

This study used a modified IPAQ to measure the intensity of physical activity. The IPAQ is the most frequently and commonly used measurement tool in measuring physical activity levels. However, the questionnaire has limitations because it is based on respondents' memories. As in previous research by Cleland et al.,¹⁶ results showed the majority of older adults under-report their level of moderate-to-vigorous physical activity and sedentary behavior when completing the IPAQ, and the linear relationship above the mean shows an error from under to over-reporting as the mean increases.

Suggestions for future research include expanding research on older-specific physical activity measures, such as previous research by VandeBunte et al.,²⁵ where actigraphy metrics are not available, standard self-report measures of PA are used in older adults to capture more structured activity and exercise routines (e.g., duration, type of exercise). The Community Healthy Activities Model Program for Seniors (CHAMPS) and the Physical Activity Scale for the Elderly (PASE) are two widely used self-report PA questionnaires in older adults. Fitbit monitors have demonstrated inter-device reliability with other actigraphy monitors. Results showed that Fitbit's step counts showed a stronger association with PASE, while Fitbit's burned calories were more strongly associated with CHAMPS-MET. Fitbit results had more consistent convergence with relevant outcomes of interest (e.g., brain and cardiometabolic health indices) compared to subjective measures; however, considerable heterogeneity in these associations was observed.

Most respondents had moderate physical activity levels, while only a few were categorized as mild or vigorous. This imbalance may limit

the broad analysis of the relationship between physical activity and vitamin D. Based on the distribution of physical activity profiles; two physical activities are often done: gymnastics and walking 100 meters. Regarding walking activity alone, many age-related changes contribute to the increased energy requirements for walking, which can be two to four times that of a healthy adult. Age-related biomechanical factors, such as a flexed trunk, limited hip extension, and reduced ankle movement in gait, result in less use of stored passive energy and greater demands on muscle activity. Age-related neuromuscular factors alter efficient muscle recruitment patterns and the timing of limb movements, resulting in inefficient walking due to compensating for body changes.²⁰ Several factors can influence a person in carrying out physical activity, including:^{14,15}

(1) Intrinsic factors. This factor is more directed at a person's inner motivation for activities. A person with good feelings will do more activities, while a person with bad feelings will do less. Even though the questionnaire has been validated, a person's motivation to move can also influence a person's physical activity.

(2) Environmental factors. The environment can influence physical activity, and weather that is too hot or rainy can be a barrier. Changing environmental weather conditions can also affect a person's ability to carry out physical activities, mainly because most respondents' physical activities are carried out in open areas, which the weather and environment will significantly influence.

(3) Physical consideration factors. A person's body condition can influence their physical activity. They will do physical activity if they feel their body is in good condition, but they will not if they feel tired or injured. In this study, activity restrictions are one of the requirements for research subjects so that the influence of this factor can be minimized.

(4) Routine factor. In the routine factor, a person carries out daily physical activity and/or other demands. External factors such as sun exposure, dietary intake, or, although these factors can significantly affect vitamin D levels.

During the research, vitamin D levels were measured to determine vitamin D levels in elderly women by taking blood samples from each research respondent. From a total of 49 elderly women, the majority of vitamin D levels <20 ng/

ml were obtained by 27 respondents (55%) with an average of 17.40 ng/ml.

Blood samples were collected only once. This may need to be revised to adequately reflect the variation in vitamin D levels, which can change depending on daily conditions, such as sun exposure and diet. Daily variation was minimized by collecting samples at the same time of day. Blood sampling was carried out simultaneously during the noonday by the phlebotomist, taking into account the time of blood sampling, which can affect vitamin D levels.¹³

This technique can cause bias because the sample is selected based on specific criteria and does not represent the entire population of elderly women. As a result, the study's results may be challenging to apply to all older adults. To minimize the influence of vitamin D variations, several factors that can affect vitamin D variations in the respondent's body condition are minimized by not having significant changes in these factors from one month earlier (such as medication, health conditions, and lifestyle). Several factors can influence vitamin D levels, including the following:^{20,21}

(1) Body mass index (BMI). Vitamin D is a fat-soluble vitamin, and people with a BMI >30 kg/m² tend to have higher fat than those with a normal BMI. The association between vitamin D deficiency and overweight and obese patients may be explained by the storage of vitamin D in adipose tissue due to its high-fat solubility and decreased exposure to sunlight in obese subjects due to limited physical activity and mobility.

(2) Lack of intake. All age-related changes in vitamin D metabolism are magnified if there is concomitant vitamin D deficiency, as it limits the supply of substrates for 25OHD and, ultimately, 1,25(OH)₂D. Substrate deficiency is a common problem in older people and is essential to recognize because it can be prevented and treated. There may be a vitamin D deficiency either from diet or from lack of sunlight, and a subsequent decrease in serum 25OHD limits 1,25(OH)₂D production, especially if there is also renal dysfunction. Serum 1,25(OH)₂D levels decrease when serum 25OHD levels fall below 10 ng/mL in young and older adults.

Conclusions

The correlation test results between physical

activity and vitamin D levels in elderly women showed no significant correlation. Other factors affecting the research results include age, which has a wide range of variations, health conditions, and vitamin D intake. Further research can develop measurements specifically for examining physical activity in old age.

Conflict of Interest

The authors declare no conflict of interest.

Acknowledgment

This research was funded by the Institute of Research and Community Service at Universitas Surabaya.

References

1. Mei Z, Hu H, Zou Y, Li D. The role of vitamin D in menopausal women's health. *Front Physiol.* 2023;14:1211896.
2. Giustina A, Bouillon R, Dawson-Hughes B, Ebeling PR, Lazaretti-Castro M, Lips P, et al. Vitamin D in the older population: a consensus statement. *Endocrine.* 2023;79(1):31–44.
3. Borowicz W, Ptaszkowski K, Ptaszowska L, Rosińczuk J, Murawska-Ciałowicz E. Association between serum vitamin D levels and physical outcomes of patients who underwent rehabilitation following ischemic stroke. *Med Sci Monit.* 2023;29:e940115.
4. Liu W, Wu Z, Zhu D, Chen G, Yan G, Zhang S, et al. Vitamin D and lipid profiles in postmenopausal women: a meta-analysis and systematic review of randomized controlled trials. *Front Mol Biosci.* 2021;8:799934.
5. Cui A, Zhang T, Xiao P, Fan Z, Wang H, Zhuang Y. Global and regional prevalence of vitamin D deficiency in population-based studies from 2000 to 2022: a pooled analysis of 7.9 million participants. *Front Nutr.* 2023;10:1070808.
6. Pereira M, Dantas Damascena A, Galvão Azevedo LM, de Almeida Oliveira T, da Mota Santana J. Vitamin D deficiency aggravates COVID-19: systematic review and meta-analysis. *Crit Rev Food Sci Nutr.* 2020;62(5):1308–16.
7. Widyantari NPS, Wijaya IPA, Susila IMDP. Hubungan tingkat pengetahuan tentang menopause dengan kecemasan menghadapi menopause pada ibu Pembinaan Kesejahteraan Keluarga. *Caring.* 2019;3(2):56–9.
8. Sari DK, Alrasyid DH, Nurlndrawaty L, Zulkifli L. Occurrence of vitamin D deficiency among women in North Sumatera, Indonesia. *Malaysian Journal of Nutrition.* 2014;20(1):63–70.
9. Dominguez LJ, Farruggia M, Veronese N, Barbagallo M. Vitamin D sources, metabolism, and deficiency: available compounds and guidelines for its treatment. *Metabolites.* 2021;11(4):255.
10. Wiciński M, Adamkiewicz D, Adamkiewicz M, Śniegocki M, Podhorecka M, Szychta P, et al. Impact of vitamin D on physical efficiency and exercise performance—a review. *Nutrients.* 2019;11(11):2826.
11. Nascimento NAP, Moreira PFP, Carvalho VA, Aragão L, Marin-Mio RV, Lazaretti-Castro M, et al. Effect of vitamin D Level and physical exercise on the physical performance and functional test results in elderly women. *J Geriatr Med Gerontol.* 2019;5(1):061.
12. Suryadinata RV, Lorensia A, Tangkilisan EC. Effect of physical activity and vitamin D status on geriatrics obesity. *GMHC.* 2019;7(1):1–6.
13. Abu Jadayil S, Abu Jadayel B, Takruri H, Muwalla M, McGrattan AM. Study of the fluctuation of serum vitamin D concentration with time during the same day and night on a random sample of healthy adults. *Clin Nutr ESPEN.* 2021;46:499–504.
14. Li J, Huang Z, Si W, Shao T. The effects of physical activity on positive emotions in children and adolescents: a systematic review and meta-analysis. *Int J Environ Res Public Health.* 2022;19(21):14185.
15. Ho JY, Lam HYC, Huang Z, Liu S, Goggins WB, Mo PKH, et al. Factors affecting outdoor physical activity in extreme temperatures in a sub-tropical Chinese urban population: an exploratory telephone survey. *BMC Public Health.* 2023;23(1):101.
16. Cleland C, Ferguson S, Ellis G, Hunter RF. Validity of the International Physical Activity Questionnaire (IPAQ) for assessing moderate-to-vigorous physical activity and sedentary behaviour of older adults in the United Kingdom. *BMC Med Res Methodol.* 2018;18(1):176.

17. Sember V, Meh K, Sorić M, Starc G, Rocha P, Jurak G. Validity and reliability of International Physical Activity Questionnaires for adults across EU countries: systematic review and meta analysis. *Int J Environ Res Public Health*. 2020;17(19):7161.
18. Khan SR, Claeson M, Khan A, Neale RE. The effect of physical activity on vitamin D: a systematic review and meta-analysis of intervention studies in humans. *Public Health Pract (Oxf)*. 2024;7:100495.
19. Heydenreich J, Schutz Y, Melzer K, Kayser B. Comparison of conventional and individualized 1-MET values for expressing maximum aerobic metabolic rate and habitual activity related energy expenditure. *Nutrients*. 2019;11(2):458.
20. Wang LK, Hung KC, Lin YT, Chang YJ, Wu ZF, Ho CH, et al. Age, Gender and season are good predictors of vitamin D status independent of body mass index in office workers in a subtropical region. *Nutrients*. 2020;12(9):2719.
21. Lorensia A, Suryadinata RV, Arganitya GN. Relationship of vitamin D intake with obesity in adolescents. *GMHC*. 2022;10(2):104–10.
22. Trovato B, Godos J, Varrasi S, Roggio F, Castellano S, Musumeci G. Physical activity, sun exposure, vitamin D intake and perceived stress in Italian adults. *Nutrients*. 2023;15(10):2301.
23. Zhang J, Cao ZB. Exercise: a possibly effective way to improve vitamin D nutritional status. *Nutrients*. 2022;14(13):2652.
24. Lorensia A, Suryadinata RV, Inu IA. Comparison of vitamin D status and physical activity related with obesity in student. *J Appl Pharm Sci*. 2022;12(4):108–18.
25. VandeBunte A, Gontrum E, Goldberger L, Fonseca C, Djukic N, You M, et al. Physical activity measurement in older adults: wearables versus self-report. *Front Digit Health*. 2022;4:869790.

Home > Archives > **Vol 12, No 3 (2024)**

USER

Username

Password

Remember me

VOL 12, NO 3 (2024)

DOI: <https://doi.org/10.29313/gmhc.v12i3>

DECEMBER 2024
FULL ISSUE

View or download the full issue

VOL. 12 NO. 3, DECEMBER 2024

TABLE OF CONTENTS

Articles

Promotion of Crypt-like Structures in Intestinal Organoid Development through the Addition of Graphene Oxide in Cell-based Assays Haura Labibah Salsabil Sulaksono, Mutiara Mila Kamilah, Lia Faridah, I Made Joni, Kozo Watanabe, Savira Ekawardhani	PDF 169-179
Polymorphisms of rs7055763 and rs41307258 in TBX22 Gene Haplotype as Risk Factors for Non-syndromic Cleft Palate Indonesian Deutero-Malay Population Nita Susanti, Irfan Ullah, Haura Labibah Salsabil Sulaksono, Saskia Lenggogeni Nasroen, Ani Melani Maskoen	PDF 180-189
Association between Clinical Characteristics of Pregnant Women in the Third Trimester and Low Back Pain Mira Dyani Dewi, Meike Rachmawati	PDF 190-194
Relationship of Physical Activity and Vitamin D Levels in Elderly Women Amelia Lorensia, Marthy Meliana Ariyanti Jalmav, Rivan Virlando Suryadinata, Muhammad Reza Aprililo Akbar	PDF 195-203
Ergonomic Risk Level of Work Posture in Leather Shoes Industry Ekawati Ekawati, Yuliani Setyaningsih, Bina Kurniawan, Ida Wahyuni	PDF 204-211
Effect of Ketapang Leaf Extract on HDL Levels of Rattus norvegicus Induced by High-fat Foods Nurfadly Nurfadly, Muhammad Fajar, Isra Thristy, Yenita Yenita	PDF 212-217
COVID-19 Treatment Patterns in Patients with Acute Respiratory Failure at Dr. Hasan Sadikin General Hospital Bandung in 2021-2022 Enny Rohmawaty, Aklila Qurrota Aini Sumardi, Iceu Dimas Kulsum	PDF 218-226
Analysis of FOXE1 rs3758249, IRF6 rs2235375, MTRR A66G in Non-syndromic Cleft Lip and Palate among Indonesian Deutero-Malay Population Saskia Lenggogeni Nasroen, Ani Melani Maskoen, Asih Rahaju, Rudi Satria Darwis, Maulia Putri, Levina Azzahra Rachman, Putri Nurul Zannah	PDF 227-233
Associated Tinnitus Risk Factors in Patients with Type 2 Diabetes Mellitus Noorma Zuliana Puryani, Yenny Yenny	PDF 234-241
The Impact of Social Value Orientation on Pro-environmental Behavior Milda Yanuvianti, Wilis Srisayekti, Efi Fitriana	PDF 242-248
Autism Spectrum Disorder: a Two-center Evaluation of Pharmacological Intervention and Behavioral Therapies on Core Symptoms Purboyo Solek, Uni Gamayani, Kusnandi Rusmil, Irvan Afriandi, Taufan Prasetya, Anggia Farrah Rizqiamuti, Eka Nurfitri, Burhan Burhan, Indra Sahril, Kevin Gunawan	PDF 249-255
Correlation between Length-to-width Ratio of Gallbladder and Gamma-glutamyl Transferase Value in Biliary Atresia Ryan Elmanar, Lenny Violetta, Rosy Setiawati	PDF 256-262

ACCREDITATION



Number: 30/E/KPT/2019



AUTHOR GUIDELINES

ONLINE SUBMISSION

FOCUS AND SCOPE

EDITORIAL TEAM

PEER REVIEWERS

PUBLICATION ETHICS

ABSTRACT AND INDEXING

AUTHOR FEES

CITATIONS

NOTIFICATIONS

- View
- Subscribe

CURRENT ISSUE

ATOM	1.0
RSS	2.0
RSS	1.0

Global Medical and Health Communication In Collaboration with:



OPEN JOURNAL SYSTEMS

Journal Help

Global Medical and Health Communication Indexed by:



Global Medical and Health Communication Submission Template



Ethical Statement Template



Copyright Transfer Statement



RECOMMENDED TOOLS

EndNote



pISSN 2301-9123 | eISSN 2460-5441

Visitor since 19 October 2016:

0000210834



Visitors				See more
164.540	5.110	980	752	
15.116	1.489	843	613	



Global Medical and Health Communication is licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License.

GLOBAL MEDICAL AND HEALTH COMMUNICATION

Published by: UPT Publikasi Ilmiah Universitas Islam Bandung

- RUMAH
- TENTANG
- LOGIN
- DAFTAR
- KATEGORI
- MENCARI
- SAAT INI
- ARSIP
- PENGUMUMAN

Beranda > Vol 12, No 3 (2024)


PENGGUNA

Nama belakang

Kata sandi

Ingat aku

Komunikasi Medis dan Kesehatan Global Diindeks oleh:



KOMUNIKASI MEDIS DAN KESEHATAN GLOBAL

Global Medical and Health Communication merupakan jurnal yang memuat artikel-artikel kedokteran dan kesehatan yang terbit sejak tahun 2013. Artikel-artikel merupakan hasil penelitian asli yang perlu disebarluaskan dan ditulis dalam **bahasa Inggris**.

Dalam kurun waktu yang tidak begitu lama, **Global Medical and Health Communication** yang dikelola oleh **Fakultas Kedokteran Universitas Islam Bandung (Unisba)** dan diterbitkan oleh **UPT Publikasi Ilmiah Unisba** telah terakreditasi oleh Akreditasi Jurnal Nasional (**Arjuna**) yang dikelola oleh Direktorat Jenderal Pendidikan Tinggi, Riset, dan Teknologi, Kementerian Pendidikan, Kebudayaan, Riset, dan Teknologi Republik Indonesia

dengan Peringkat 2 (**Sinta 2**) mulai Volume 7 Nomor 2 Tahun 2019 sampai dengan Volume 12 Nomor 2 Tahun 2024. Jurnal ini juga telah terindeks di **Directory of Open Access Journals (DOAJ)** pada tanggal 9 Mei 2017 dan

Crossref pada tanggal 2 Januari 2018. Dengan terindeksnya DOAJ dan Crossref, jurnal ini dapat menjangkau khalayak internasional.

Capaian ini mendapat respon positif dari para peneliti, dosen, dan pemerhati kesehatan yang ditunjukkan dengan jumlah artikel yang masuk tiga kali lipat dari jumlah artikel yang diterima pada awalnya. Kualitas artikel juga menunjukkan peningkatan dari segi metodologi dan penulisan yang bermanfaat bagi khalayak. Hasil penelitian sebaiknya disebarluaskan sedini mungkin agar dapat dimanfaatkan dengan baik. Penerbitan **Global Medical and Health Communication** setiap **6 (enam) bulan** dalam setahun akan diterbitkan setiap **4 (empat) bulan** mulai tahun 2017.

Berita dari bidang sains dan kesehatan tentang tantangan dan peluang di Indonesia masih sangat jarang dan **Global Medical and Health Communication** berharap dapat menjadi sarana terbaik untuk mendukung para peneliti, dosen, dan praktisi kesehatan untuk menjadi suara Indonesia, khususnya di bidang kesehatan.

pISSN 2301-9123 | eISSN 2460-5441



JIL. 12, NOMOR 3 (2024)

DESEMBER 2024 (TERBIT SEDANG DALAM PROSES)
DAFTAR ISI

Artikel

Peningkatan Struktur Mirip Kripta dalam Perkembangan Organoid Usus melalui Penambahan Oksida Grafena dalam Pengujian Berbasis Sel Haura Labibah Salsabil Sulaksono, Mutiara Mila Kamilah, Lia Faridah, I Made Joni, Kozo Watanabe, Savira Ekawardhani	BAHASA INDONESIA: 169-179
Polimorfisme rs7055763 dan rs41307258 pada Haplotipe Gen TBX22 sebagai Faktor Risiko Sumbing Langit-langit Non-Sindromik Populasi Deutero-Melayu Indonesia Nita Susanti, Irfan Ullah, Haura Labibah Salsabil Sulaksono, Saskia Lenggogeni Nasroen, Ani Melani Maskoen	BAHASA INDONESIA: 180-189
Hubungan Karakteristik Klinis Ibu Hamil Trimester III dengan Nyeri Punggung Bawah Mira Dyani Dewi, Meike Rachmawati	BAHASA INDONESIA: 190-194
Hubungan Aktivitas Fisik dan Kadar Vitamin D pada Wanita Lansia Amelia Lorensia, Marthy Meliana Ariyanti Jalmav, Rivan Virlando Suryadinata, Muhammad Reza Aprililo Akbar	BAHASA INDONESIA: 195-203

pISSN 2301-9123 | eISSN 2460-5441

Pengunjung sejak 19 Oktober 2016:
0000208907



ACCREDITATION



- AUTHOR GUIDELINES
- ONLINE SUBMISSION
- FOCUS AND SCOPE
- EDITORIAL TEAM
- PEER REVIEWERS
- PUBLICATION ETHICS
- ABSTRACT AND INDEXING
- AUTHOR FEES
- CITATIONS

PEMBERITAHUAN

- Melihat
- Berlangganan

EDISI SAAT INI

RIQM	1.0
RSS	2.0
RSS	1.0

Komunikasi Medis dan Kesehatan Global Bekerja Sama dengan:



SISTEM JURNAL TERBUKA
Bantuan Jurnal

Visitors [See more](#)

163.768	5.005	977	751
15.020	1.481	828	609

FLAG counter



Komunikasi Medis dan Kesehatan Global dilisensikan di bawah [Lisensi Internasional Creative Commons Atribusi-NonKomersial-BerbagiSerupa 4.0](#) .

GLOBAL MEDICAL AND HEALTH COMMUNICATION

Published by: UPT Publikasi Ilmiah Universitas Islam Bandung

- RUMAH
- TENTANG
- LOGIN
- DAFTAR
- KATEGORI
- MENCARI
- SAAT INI
- ARSIP
- PENGUMUMAN

Beranda > Tentang Jurnal > **Orang-orang**


PENGGUNA

Nama belakang

Kata sandi

Ingat aku

Komunikasi Medis dan Kesehatan Global Diindeks oleh:



RAKYAT

Peninjau

Agung Firmansyah Sumantri, Departemen Penyakit Dalam, Fakultas Kedokteran, Universitas Islam Bandung, Bandung, Indonesia

Arief Budi Yulianti, Departemen Biologi Medis, Fakultas Kedokteran, Universitas Islam Bandung, Bandung, Indonesia

Arif Susanto, Kesehatan, Keselamatan & Lingkungan (HSE), PT Freeport Indonesia, Tembaga, Indonesia

Budi Setiabudiawan, Departemen Kesehatan Anak Fakultas Kedokteran Universitas Padjadjaran/Dr. Rumah Sakit Umum Hasan Sadikin Bandung, Indonesia

Cordia Ming Yeuk Chu, Pusat Lingkungan dan Kesehatan Populasi, Universitas Griffith, Brisbane, Australia

Dicky Santosa, Departemen Kesehatan Anak, Fakultas Kedokteran, Universitas Islam Bandung, Bandung, Indonesia

Djatinika Setiabudi, Departemen Kesehatan Anak, Fakultas Kedokteran, Universitas Padjadjaran, Bandung, Indonesia

Ermi Ndoen, UNICEF Indonesia, Jakarta, Indonesia

Ferry Efendi, Fakultas Keperawatan, Universitas Airlangga, Surabaya, Indonesia

Hendro Sudjono Yuwono, Departemen Bedah, Fakultas Kedokteran, Universitas Islam Bandung, Bandung, Indonesia

Henri Muflihah, Departemen Farmakologi, Fakultas Kedokteran, Universitas Islam Bandung, Bandung, Indonesia

Herri S Sastramihardja, Departemen Farmakologi, Fakultas Kedokteran, Universitas Islam Bandung, Bandung, Indonesia

Hidayat Wijayanegara, Departemen Obstetri dan Ginekologi, Fakultas Kedokteran, Universitas Islam Bandung, Bandung, Indonesia

Ieva B Akbar, Departemen Fisiologi, Fakultas Kedokteran, Universitas Islam Bandung, Bandung, Indonesia

Johanes Cornelius Mose, Departemen Obstetri dan Ginekologi, Fakultas Kedokteran, Universitas Padjadjaran, Bandung, Indonesia

Lelly Yuniarti, Departemen Biokimia, Nutrisi, dan Biomolekuler, Fakultas Kedokteran, Universitas Islam Bandung, Bandung, Indonesia

Leri Septiani, STIKes Dharma Husada/RSIA Grha Bunda, Bandung, Indonesia

Lia Faridah, Departemen Ilmu Biomedis, Fakultas Kedokteran, Universitas Padjadjaran, Bandung, Indonesia

M Ahmad Djojogugito, Departemen Ortopedi dan Traumatologi, Fakultas Kedokteran, Universitas Islam Bandung, Bandung, Indonesia

Mas Rizky Anggun Adipurna Syamsunarno, Departemen Biokimia dan Biologi Molekuler, Fakultas Kedokteran, Universitas Padjadjaran, Jatinangor, Indonesia

Maya Tejasari, Departemen Histologi, Fakultas Kedokteran, Universitas Islam Bandung, Bandung, Indonesia

Mia Kusmiati, Departemen Pendidikan Kedokteran, Bioetika dan Humaniora, Fakultas Kedokteran, Universitas Islam Bandung, Bandung, Indonesia

Nuzirwan Acang, Departemen Penyakit Dalam, Fakultas Kedokteran, Universitas Islam Bandung, Bandung, Indonesia

Riyadi Adrizain, Departemen Kesehatan Anak, Fakultas Kedokteran, Universitas Padjadjaran, Bandung, Indonesia

Rizky Suganda Prawiradilaga, Departemen Biokimia, Gizi dan Biomolekuler, Fakultas Kedokteran, Universitas Islam Bandung, Bandung, Indonesia

Titik Respati, Departemen Kesehatan Masyarakat, Fakultas Kedokteran, Universitas Islam Bandung, Bandung, Indonesia

Umar Fahmi Achmadi, Departemen Kesehatan Lingkungan, Fakultas Kesehatan Masyarakat, Universitas Indonesia, Depok, Indonesia

Wahyu Widowati, Departemen Biomolekuler, Fakultas Kedokteran, Universitas Kristen Maranatha, Bandung, Indonesia

Wawang Setiawan Sukarya, Departemen Obstetri dan Ginekologi, Fakultas Kedokteran, Universitas Muhammadiyah Prof. Dr. HAMKA, Jakarta Selatan, Indonesia

Wida Purbaningsih, Departemen Histologi, Fakultas Kedokteran, Universitas Islam Bandung, Bandung, Indonesia

ACCREDITATION



- AUTHOR GUIDELINES**
- ONLINE SUBMISSION**
- FOCUS AND SCOPE**
- EDITORIAL TEAM**
- PEER REVIEWERS**
- PUBLICATION ETHICS**
- ABSTRACT AND INDEXING**
- AUTHOR FEES**
- CITATIONS**

PEMBERITAHAUAN

- Melihat
- Berlangganan

Komunikasi Medis dan Kesehatan Global Bekerja Sama dengan:



SISTEM JURNAL TERBUKA
Bantuan Jurnal


Yani Triyani , Departemen Patologi Klinik, Fakultas Kedokteran, Universitas Islam Bandung, Bandung, Indonesia

Yulia Sofiatin , Divisi Epidemiologi dan Biostatistik, Departemen Kesehatan Masyarakat, Universitas Padjadjaran, Bandung, Indonesia

pISSN 2301-9123 | eISSN 2460-5441

Pengunjung sejak 19 Oktober 2016:

0000208916



Komunikasi Medis dan Kesehatan Global dilisensikan di bawah [Lisensi Internasional Creative Commons Atribusi-NonKomersial-BerbagiSerupa 4.0](#) .

GLOBAL MEDICAL AND HEALTH COMMUNICATION

Published by: UPT Publikasi Ilmiah Universitas Islam Bandung

- RUMAH
- TENTANG
- LOGIN
- DAFTAR
- KATEGORI
- MENCARI
- SAAT INI
- ARSIP
- PENGUMUMAN

Beranda > Tentang Jurnal > **Tim Editorial**

PENGUNA

Nama belakang

Kata sandi

Ingat aku

Komunikasi Medis dan Kesehatan Global Diindeks oleh:



Template Pernyataan Etika



Pernyataan Pengalihan Hak Cipta



RECOMMENDED TOOLS



TIM REDAKSI

Pemimpin redaksi

Santun Bhkti Rahimah, Departemen Farmakologi, Fakultas Kedokteran, Universitas Islam Bandung, Bandung, Indonesia

Redaktur Pelaksana

Winni Maharani, Departemen Mikrobiologi, Fakultas Kedokteran, Universitas Islam Bandung, Bandung, Indonesia

Dewan Redaksi Eksternal

Badrul Hisham Yahaya, Institut Kedokteran dan Gigi Tingkat Lanjut, Universiti Sains Malaysia, Kepala Batas, Malaysia

Jerico Franciscus Padosi, Fakultas Kesehatan Masyarakat dan Pekerjaan Sosial, Universitas Teknologi Queensland, Kelvin Grove, Queensland, Australia

Roy Rillera Marzo, Universitas Asia Metropolitan, Johor, Malaysia

Dewan Redaksi Internal

Ike Rahmawaty Alie, Departemen Fisiologi, Fakultas Kedokteran, Universitas Islam Bandung, Bandung, Indonesia

Indri Budiarti, Departemen Obstetri dan Ginekologi, Fakultas Kedokteran, Universitas Islam Bandung, Bandung, Indonesia

Lisa Adhia Garina, Departemen Kesehatan Anak, Fakultas Kedokteran, Universitas Islam Bandung, Bandung, Indonesia

Mirasari Putri, Departemen Biokimia, Nutrisi, dan Biomolekuler, Fakultas Kedokteran, Universitas Islam Bandung, Bandung, Indonesia

Yuktiana Kharisma, Departemen Patologi Anatomi, Fakultas Kedokteran, Universitas Islam Bandung, Bandung, Indonesia

Korektor

Herry Garna, Departemen Kesehatan Anak, Fakultas Kedokteran, Universitas Islam Bandung, Bandung, Indonesia

Editor naskah

Yudi Feriandi, Departemen Pendidikan Kedokteran, Bioetika, Humaniora dan Sisipan Islam, Fakultas Kedokteran, Universitas Islam Bandung, Bandung, Indonesia

Editor Tata Letak

Agus Chalid, Fakultas Kedokteran Universitas Islam Bandung, Bandung, Indonesia

Asisten Editorial


Agus Chalid, Fakultas Kedokteran Universitas Islam Bandung, Bandung, Indonesia

Ihsan Mulyadi Kurniawan, Fakultas Kedokteran Universitas Islam Bandung, Bandung, Indonesia

pISSN 2301-9123 | eISSN 2460-5441

Pengunjung sejak 19 Oktober 2016:

0000208914



Komunikasi Medis dan Kesehatan Global dilisensikan di bawah [Lisensi Internasional Creative Commons Atribusi-NonKomersial-BerbagiSerupa 4.0](#) .

ACCREDITATION



Nomor: 30/E/KPT/2019



AUTHOR GUIDELINES

ONLINE SUBMISSION

FOCUS AND SCOPE

EDITORIAL TEAM

PEER REVIEWERS

PUBLICATION ETHICS

ABSTRACT AND INDEXING

AUTHOR FEES

CITATIONS

PEMBERITAHAUAN

- Melihat
- Berlangganan

Komunikasi Medis dan Kesehatan Global Bekerja Sama dengan:



SISTEM JURNAL TERBUKA

[Bantuan Jurnal](#)

SERTIFIKAT

Direktorat Jenderal Penguatan Riset dan Pengembangan,
Kementerian Riset, Teknologi, dan Pendidikan Tinggi



Kutipan dari Keputusan Direktur Jenderal Penguatan Riset dan Pengembangan,
Kementerian Riset, Teknologi, dan Pendidikan Tinggi Republik Indonesia
Nomor: 30/E/KPT/2019
Tentang Hasil Akreditasi Jurnal Ilmiah Periode 6 Tahun 2019

Global Medical & Health Communication

E-ISSN: 24605441

Penerbit: Pusat Penerbitan Universitas-Lembaga Penelitian dan Pengembangan Masyarakat
(P2U-LPPM), Universitas Islam Bandung


Ditetapkan sebagai Jurnal Ilmiah

TERAKREDITASI PERINGKAT 2

Akreditasi berlaku selama 5 (lima) tahun, yaitu
Volume 7 Nomor 2 Tahun 2019 sampai Volume 11 Nomor 2 Tahun 2024

Jakarta, 11 November 2019

Direktur Jenderal Penguatan Riset dan Pengembangan



Dr. Muhammad Dimiyati
NIP. 195912171984021001



Relationship of Physical Activity and Vitamin D Levels in Elderly Women

by Amelia Lorensia

Submission date: 23-Dec-2024 09:42PM (UTC+0700)

Submission ID: 2557741079

File name: 13499-48487-1-PB_Amelia.pdf (705.33K)

Word count: 5465

Character count: 28590

Relationship of Physical Activity and Vitamin D Levels in Elderly Women

ORIGINALITY REPORT

10%

SIMILARITY INDEX

9%

INTERNET SOURCES

11%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

1	www.researchsquare.com Internet Source	1%
2	1library.net Internet Source	1%
3	biospraysehatalami.com Internet Source	1%
4	fitabase.com Internet Source	1%
5	ejournal.unsap.ac.id Internet Source	1%
6	bmcmmedresmethodol.biomedcentral.com Internet Source	1%
7	Achmad Gatot Sunarianto, Ning Arti Wulandari, Andri Darmawan. "Penurunan Hemoglobin pada Penyakit Ginjal Kronik Setelah Hemodialisis di RSUD "KH" Batu", Jurnal Ners dan Kebidanan (Journal of Ners and Midwifery), 2019 Publication	1%

8 Zhaojun Mei, Hong Hu, Yi Zou, Dandan Li. "The role of vitamin D in menopausal women's health", *Frontiers in Physiology*, 2023
Publication 1%

9 mdpi-res.com
Internet Source 1%

10 www.selcukmedj.org
Internet Source 1%

11 Anna M. VandeBunte, Nina Djukic, Michelle You, Corrina Fonseca, Joel H Kramer, Kaitlin B Casaletto. "Defining physical activity: A comparison of actigraphy monitoring and self-report measures", *Alzheimer's & Dementia*, 2021
Publication 1%

Exclude quotes On

Exclude matches < 1%

Exclude bibliography On

RESEARCH ARTICLE

Relationship of Physical Activity and Vitamin D Levels in Elderly Women**Amelia Lorensia,¹ Marthy Meliana Ariyanti Jalmav,² Rivan Virlando Suryadinata,³
Muhammad Reza Aprililo Akbar⁴**¹Department of Clinical-Community Pharmacy, Faculty of Pharmacy, Universitas Surabaya, Surabaya, Indonesia,²Faculty of Pharmacy, Universitas Anwar Medika, Sidoarjo, Bandung, Indonesia,³Department of Public Health, Faculty of Medicine, Universitas Surabaya, Surabaya, Indonesia,⁴Faculty of Pharmacy, Universitas Surabaya, Surabaya, Indonesia**Abstract**

Older adults are at risk of vitamin D deficiency, especially in older women, due to a decrease in the hormone estrogen, which causes decreased bone density and increased risk of fractures. There is a relationship between vitamin D deficiency and lack of physical activity due to the storage of vitamin D in adipose tissue. Physical activity was measured using the International Physical Activity Questionnaire (IPAQ). The aim was to determine the relationship between physical activity and vitamin D in elderly women. This cross-sectional analytic observational study was conducted at Nursing Home X in Surabaya city from May to August 2023. The variables were physical activity (measured by IPAQ to estimate physical activity levels) and vitamin D level (measured by VIDAS® instrument to determine 25(OH) levels). The subjects were all women aged ≥60 years. The sampling technique uses the purposive sampling method. For an analysis of the relationship between physical activity levels and vitamin D levels, the Pearson correlation test was used with a ratio data scale. This research received 49 respondents. The average physical activity level was 1316.82±720.90, and most respondents had a moderate physical activity level of 44.90%. Vitamin D levels were ≤30 ng/ml for 46 respondents (93.87%). The results of the Pearson correlation test were obtained with a $r_{\text{value}}=0.089$ and a significance value (Sig.) of 0.542. There was no significant correlation between physical activity and vitamin D levels. Other factors such as age, health conditions, and vitamin D intake can affect the results. Further research can be developed into measurements to examine physical activity in old age.

Keywords: IPAQ, elderly woman, nursing home, physical activity, vitamin D**Introduction**

Vitamin D deficiency is a common global health problem with high prevalence, especially in South Asia countries. Vitamin D mainly regulates calcium and phosphorus metabolism and promotes bone growth.¹ Older adults are at risk of vitamin D deficiency because vitamin D production and metabolism change with age due to various factors, such as reduced sun exposure and skin production capacity.² Physically, geriatrics are less active and have poor nutritional status. In addition, there are physiological processes that exacerbate the decrease in vitamin D levels in the body, such as decreased production of vitamin D in the skin after sun exposure caused by skin atrophy, eating foods that are low in vitamin D, impaired

gastrointestinal absorption, and decreased production 1,25(OH)₂D in the kidney.³ Vitamin D deficiency is prevalent in postmenopausal women, and this undoubtedly exacerbates the risk of cardiovascular disease and menopause-related dyslipidemia. Estrogen levels drop significantly as women go through menopause, and a lack of estrogen weakens bones. Vitamin D helps the body absorb and use calcium to maintain bone structure. 25-hydroxyvitamin D [25(OH)D] serves the purpose of calcium absorption and appears to be hormonally sensitive. Vitamin D deficiency is associated with low bone mass and increased fracture risks. Menopausal women are at risk of decreased bone density and increased risk of fractures due to decreased estrogen levels,¹ reduced vitamin D synthesis in the skin, or changes in body composition relevant to vitamin

Copyright ©2024 by authors. This is an open access article under a Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License (<https://creativecommons.org/licenses/by-nc-sa/4.0>).

Received: 20 February 2024; Revised: 10 October 2024; Accepted: 10 December 2024; Published: 20 December 2024

Correspondence: Amelia Lorensia. Department of Clinical-Community Pharmacy, Faculty of Pharmacy, Universitas Surabaya. Jln. Raya Kalirungkut, Surabaya 60292, East Java, Indonesia. E-mail: amelia.lorensia@gmail.com, amelia.lorensia@staff.ubaya.ac.id

D status and physiology.⁴ Vitamin D deficiency is defined as vitamin D levels below 20 ng/ml (50 nmol/liter), and vitamin D deficiency is 21–29 ng/ml (525–725 nmol/liter). Vitamin D deficiency can also be a risk factor for diseases such as immune system diseases and infectious diseases caused by coronavirus disease 2019 (COVID-19).⁵ Research by Pereira et al.⁶ reported that low vitamin D levels were associated with COVID-19.

Research by Cui et al.⁵ shows that 15.7% of the global adult population has vitamin D deficiency. Vitamin D deficiency can occur in all age groups, including menopausal women. Indonesia currently has 7.4% of menopausal women from the total population, and in 2020, it is estimated to reach 11.54% with an average age of menopause of 49 years. According to Statistics Indonesia, in 2015, the number of women in Indonesia who entered menopause reached 21.22 million. It is estimated that by 2025, there will be 60 million menopausal women.⁷ Another study on the incidence of vitamin D deficiency in Indonesian women showed that 95% of them (148 subjects out of a total of 156) had vitamin D deficiency.⁸

There is a relationship between vitamin D deficiency and lack of physical activity due to the storage of vitamin D in adipose tissue due to its high fat solubility and decreased exposure to sunlight in obese subjects due to limited physical activity and limited mobility. It is proven that physical activity increases glucose, calcium, and vitamin D metabolism and reduces body weight by increasing lipolysis.⁹ Someone who does high physical activity will be associated with increased vitamin D concentration. Suppose someone does less physical activity (low physical activity). In that case, it will cause a decrease in vitamin D concentration, while someone with high physical activity will most likely be associated with optimal vitamin D levels.¹⁰ Vitamin D can also help improve the body's performance in physical activities in populations at risk of vitamin D deficiency. Previous research by Nascimento et al.¹¹ found a significant increase in levels and improvements in the body, so there was an interaction between vitamin D levels and physical activity in maintaining body performance.

Measurement of physical activity in this study used a questionnaire, namely the extended version of the International Physical Activity Questionnaire (IPAQ), which is used to estimate

physical activity levels. Data on the relationship between physical activity and vitamin D levels in Indonesia have been collected by Suryadinata et al.,¹² conducted at Public Health Center Taman, Sidoarjo district, East Java, in March–July 2017. The results showed geriatrics with obesity and non-obesity had similar levels of physical activity, but vitamin D status in obesity tended to be lower than non-obese. However, similar studies have never been conducted on elderly women in nursing homes. These data could greatly assist health workers in improving the quality of life of these patients. This research aimed to determine the relationship between physical activity and vitamin D in elderly women.

Methods

This study used analytic observational research with a cross-sectional design. Data was collected from May 2023 to August 2023 at Nursing Home X in Surabaya city. The Institutional Ethical Committee of the University of Surabaya approved the study, 207/KE/VIII/2023.

This study's population was all women aged ≥ 60 who had experienced menopause in Nursing Home X in Surabaya city. The samples taken in this study were all populations that met the inclusion criteria. The sampling technique used is purposive sampling, which meets the requirement of not having physical limitations such as motor disorders.

Each respondent will be interviewed to fill out the IPAQ and take blood samples to check vitamin D levels. A phlebotomist from Pramita Laboratorium will take a blood sample and test it at an accredited laboratory.

Blood sampling was carried out simultaneously during the day by the phlebotomist, taking into account the time of blood sampling, which can affect vitamin D levels.¹³ After blood collection, the laboratory will process the blood by centrifuging to obtain serum, which will then be tested for vitamin D levels using VIDAS® 25-OH Vitamin D Total (VITD), which is an automatic quantitative test used on the VIDAS® instrument to determine 25(OH) levels. D total in human serum or plasma with the enzyme-linked fluorescence assay (ELFA) technique.

The method for measuring physical activity uses the modified IPAQ. The physical activity questionnaire contains 12 questions about light,

moderate, and vigorous physical activity activities. Respondents were interviewed, and then results were obtained, which categorized respondents into mild, moderate, and vigorous categories using the metabolic equivalent of task (MET). Physical activity is divided into three categories, namely the mild category (<600 MET minutes/week), the moderate category (600–1,500 MET minutes/week), and the vigorous category (>1,500 MET minutes/week). This questionnaire contains questions about the type of activity, duration, and frequency of a person doing physical activity in the last 7 days. Physical activity can be measured by measuring the amount of energy expended/needed in each minute of activity. According to the IPAQ assessment protocol, MET-minutes/week for a given activity (walking moderate-intensity activity or vigorous-intensity activity) is calculated by multiplying the MET value of the given activity (3.3 for walking, 4.0 for moderate-intensity activity, and 8.0 for high-intensity activities) according to hours spent in a particular activity, for example walking at work = 3.3 × minutes walking × days walking at work. MET-minutes/week = (MET × activity minutes × days). Information: MET-minutes/week = METs produced in one week MET = MET value based on activity level minutes = duration of activity in one day days = number of days in one activity.¹⁴⁻¹⁵ Results are presented as estimates of energy expenditure in metabolic equivalent minutes per week (MET-hours/week).^{16,17} The metabolic equivalent of task, or MET, is valid for describing energy expenditure from certain activities. MET is the ratio of the rate of energy expended during activity to the rate of energy expended at rest.¹⁸

Before use, the questionnaire (IPAQ)^{11,16,19} must be used as a research instrument, and a validity test must first be carried out, consisting of content and construct validation. Content validity through rational analysis or expert judgment by a panel to ensure that measurements are representative. Meanwhile, construct validity refers to the extent to which a measuring instrument shows results following theory. Reliability shows that the questionnaire is reliable and consistently provides the same results when measured twice or more with the same instrument. Cronbach's alpha value was >0.6, so the research instrument would be reliable.

The relationship between physical activity levels and vitamin D levels was analyzed using the

Pearson correlation test, with the data scale being the ratio. The data was tested for normality using the Shapiro-Wilk test; if the p-value > 0.05, it was concluded that the data was normally distributed. The relationship test used the Pearson correlation test; if the p-value < 0.05, it can be concluded that there is a significant relationship between physical activity and vitamin D levels.

Results

This research involved 49 geriatric women as respondents. Data were collected using physical activity questionnaire interviews and vitamin D serum tests. The study involved elderly women aged 60 years and over. The highest number of high-risk elderly women >70 years was 43 respondents, while the rest were in the 60–70-year age group.

Regarding body mass index (BMI) characteristics, most respondents were of normal weight, and 34 respondents had an average standard deviation of 21.58 ± 2.16 kg/m². Most of them had no disease history (Table 1).

Based on the data in Table 2, respondents' most common physical activity profile was exercise (44 respondents) and walking 100 m (39 respondents). At the same time, physical activities that were only done by some respondents were cooking and cleaning the garden or the like. Seven respondents carry out physical activity in the mild category, 22 in the moderate category,

Table 1 Frequency Distribution of Respondent Characteristics

Characteristics	n=49
Age (years)	
60–70	6
>70	43
BMI (kg/m ²)	
Very thin (<16)	6
Skinny (16–18.4)	6
Normal (18.5–24.9)	34
Overweight (25–29.9)	3
Obesity (>30)	0
Medical history	
Type 2 diabetes mellitus	5
Hypertension	10
Osteoarthritis	7
Osteoporosis	3
There isn't any	27

Table 2 Profile of Respondents' Physical Activity Questionnaire Answers

No.	Types of Physical Activity	Answer (Minutes)	n=49
1	Walk 100 m	5	1
		10	5
		15	6
		20	1
		30	23
		60	3
		Didn't do	10
2	Walk more than 100 m	10	2
		15	3
		60	12
		Didn't do	32
3	Cook	Didn't do	49
4	Wash clothes by hand	10	4
		15	2
		20	2
		30	9
5	Cleaning room	Didn't do	32
		5	1
		10	5
		15	1
6	Exercise	30	2
		Didn't do	40
		10	7
		15	13
		20	4
		25	6
		30	4
7	Cleaning the garden	45	3
		60	8
		Didn't do	5
		Didn't do	49
		8	Gardening/farming
9	Sew with hand	Didn't do	47
		10	2
		15	3
		20	1
		60	4
10	Sewing by machine	Didn't do	39
		60	1
		Didn't do	48
11	Read	10	4
		15	2
		20	1
		30	4
		60	4
		Didn't do	34
12	Watch TV	5	1
		10	3
		15	3
		20	1
		30	6
		60	2
Didn't do	33		

and 20 in the vigorous category. The average physical activity level was 1316.82 ± 720.90 , and most respondents had a moderate physical activity level of 44.90% (Table 3).

Vitamin D levels were measured during the research by taking blood samples from each respondent. In Table 4, the results showed that vitamin D levels^{16,17} were <20 ng/ml for 27 respondents, vitamin D levels were 20–30 ng/ml for 19 respondents, and vitamin D levels were >30 ng/ml for as many as three respondents.

Based on Table 5, the results showed that deficiency (<20 ng/ml) for 27 respondents (mean \pm SD= 17.40 ± 2.07), insufficiency (20–30 ng/ml) for 19 respondents (mean \pm SD= 23.64 ± 2.42), and normal level (>30 ng/ml) for three respondents (mean \pm SD= 31.33 ± 1.15). The total mean \pm SD was 18.41 ± 6.60 . In the mild physical activity category, two respondents had vitamin D levels <20 ng/ml, and 20–30 ng/ml for five respondents. In the category of moderate physical activity, 14 respondents had vitamin D levels <20 ng/ml, and 20–30 ng/ml as many as eight respondents. In the category of vigorous physical activity, 11 respondents had vitamin D levels <20 ng/ml, six respondents had 20–30 ng/ml, and three had >30 ng/ml.

The normality test results are as follows: In the Shapiro-Wilk normality test, a significance value (Sig.) was obtained, which was 0.133 for physical activity and 0.94 for vitamin D levels. Because both data have a significance value of >0.05 , it can be concluded that the physical activity data and vitamin D levels were usually distributed. The results of the Pearson correlation test are as follows: obtained an r_{value} (Pearson correlation) of 0.089 and a significance value (Sig.) of 0.542. Based on a significance value (α) >0.05 , it means there is no significant correlation between physical activity and vitamin D levels (Figure).

Discussion

This research involved 49 geriatric women as respondents. Since this study only involved 49 respondents, which may need to be more representative of a larger group, the results cannot be generalized widely and only to elderly women with similar characteristics.

Based on age, the highest number in the group of high-risk elderly women, namely 70 years and over, was 43 respondents, while in the 60–70

Table 3 Physical Activity Level of Respondents

Category Physical Activity	n=49	Mean±SD	Total Mean±SD
Mild (<600)	7	312.21±162.37	1,316.82±720.90
Moderate (600–1,500)	22	1,038.75±401.43	
Vigorous (>1,500)	20	2,074.17±581.39	

Table 4 Presentation of Data on Vitamin D Level Values

Vitamin D Level (ng/ml)	n=49	Mean±SD	Total Mean±SD
Deficiency (<20)	27	17.40±2.07	18.41±6.60
Insufficiency (20–30)	19	23.64±2.42	
Normal (>30)	3	31.33±1.15	

Table 5 Cross Tabulation of Vitamin D Levels with Physical Activity

Physical Activity	Number of Respondents based on Vitamin D Level Value			Total n=49
	Deficiency n=27	Insufficiency n=19	Normal n=3	
Mild (<600)	2	5	0	7
Moderate (600–1,500)	14	8	0	22
Vigorous (>1,500)	11	6	3	20

Note: deficiency (<20 ng/ml), insufficiency (20–30 ng/ml), normal (>30 ng/ml)




Figure Distribution of Physical Activity Data and Vitamin D Levels

year age group, there were only six respondents. Uneven age distribution may affect the results, especially physiological changes in different age groups. Age can affect vitamin D levels, such as a decreased vitamin D metabolism in old age. A decrease in calcium absorption causes this, the presence of intestinal resistance to absorption of circulating calcium $1,25(\text{OH})_2\text{D}$, a decrease in kidney function resulting in a reduction of $1,25(\text{OH})_2\text{D}$ production in the kidneys, and a decrease in vitamin D production in the skin, nine caused by decreasing the concentration of 7-dehydrocholesterol in the epidermis which can reduce the formation of vitamin D₃. So aging can also cause calcium deficiency, hyperparathyroidism, increased bone loss, and osteoporosis.^{20,21}

The majority of respondents (27 people) had no history of illness, while only several suffered from diabetes, hypertension, or osteoarthritis. This imbalance in health characteristics can affect physical activity and vitamin D levels. History of disease can affect vitamin D levels, including type 2 diabetes mellitus. The relationship between vitamin D deficiency and insulin resistance can develop through inflammation because vitamin D deficiency is associated with increased inflammatory markers. In addition, genetic polymorphisms of vitamin D-related genes may predispose to impaired glycemic control and type 2 diabetes. Epidemiological studies suggest an association between low serum 25-hydroxyvitamin D₃ ($25(\text{OH})\text{D}_3$) concentrations and an increased risk of metabolic syndrome and type 2 diabetes. The increase in fat mass partly explains it.^{20,21}

Most respondents had low vitamin D levels (<20 ng/ml), and only a few had levels above 30 ng/ml. This uneven distribution may limit statistical power for analysis and comparison between groups. The research showed no significant correlation between physical activity and vitamin D levels. It differs from previous research by Trovato et al.,²² which stated that higher dietary intake of vitamin D and sunlight exposure are associated with a lower likelihood of having high perceived stress among physically active individuals. These results may be related to previous research by Zhang and Cao,²³ that endurance exercise can significantly increase serum $25(\text{OH})\text{D}$ levels in vitamin D-deficient people but has no significant effect on vitamin

D-sufficient people. The effects of exercise on $25(\text{OH})\text{D}$ levels in the circulation may depend on factors such as vitamin D nutritional status, exercise type and intensity, and sex.

Physical activity can be measured with a tool, namely an accelerometer, pedometer, and self-report questionnaire (IPAQ) instrument. This research uses a self-report questionnaire because it has the advantage of cheaper costs, does not burden respondents, and obtains the flow of daily physical activity from respondents. Still, the self-report questionnaire has areas for improvement, such as less accuracy and reliability, because the data is based only on respondents' memories.²⁴

This study used a modified IPAQ to measure the intensity of physical activity. The IPAQ is the most frequently and commonly used measurement tool in measuring physical activity levels. However, the questionnaire has limitations because it is based on respondents' memories. As in previous research by Cleland et al.,¹⁶ results showed the majority of older adults under-report their level of moderate-to-vigorous physical activity and sedentary behavior when completing the IPAQ, and the linear relationship above the mean shows an error from under to over-reporting as the mean increases.

Suggestions for future research include expanding research on older-specific physical activity measures, such as previous research by VandeBunte et al.,²⁵ where actigraphy metrics are not available, standard self-report measures of PA are used in older adults to capture more structured activity and exercise routines (e.g., duration, type of exercise). The Community Healthy Activities Model Program for Seniors (CHAMPS) and the Physical Activity Scale for the Elderly (PASE) are two widely used self-report PA questionnaires in older adults. Fitbit monitors have demonstrated inter-device reliability with other actigraphy monitors. Results showed that Fitbit's step counts showed a stronger association with PASE, while Fitbit's burned calories were more strongly associated with CHAMPS-MET. Fitbit results had more consistent convergence with relevant outcomes of interest (e.g., brain and cardiometabolic health indices) compared to subjective measures; however, considerable heterogeneity in these associations was observed.

Most respondents had moderate physical activity levels, while only a few were categorized as mild or vigorous. This imbalance may limit

the broad analysis of the relationship between physical activity and vitamin D. Based on the distribution of physical activity profiles; two physical activities are often done: gymnastics and walking 100 meters. Regarding walking activity alone, many age-related changes contribute to the increased energy requirements for walking, which can be two to four times that of a healthy adult. Age-related biomechanical factors, such as a flexed trunk, limited hip extension, and reduced ankle movement in gait, result in less use of stored passive energy and greater demands on muscle activity. Age-related neuromuscular factors alter efficient muscle recruitment patterns and the timing of limb movements, resulting in inefficient walking due to compensating for body changes.²⁰ Several factors can influence a person in carrying out physical activity, including:^{14,15}

(1) Intrinsic factors. This factor is more directed at a person's inner motivation for activities. A person with good feelings will do more activities, while a person with bad feelings will do less. Even though the questionnaire has been validated, a person's motivation to move can also influence a person's physical activity.

(2) Environmental factors. The environment can influence physical activity, and weather that is too hot or rainy can be a barrier. Changing environmental weather conditions can also affect a person's ability to carry out physical activities, mainly because most respondents' physical activities are carried out in open areas, which the weather and environment will significantly influence.

(3) Physical consideration factors. A person's body condition can influence their physical activity. They will do physical activity if they feel their body is in good condition, but they will not if they feel tired or injured. In this study, activity restrictions are one of the requirements for research subjects so that the influence of this factor can be minimized.

(4) Routine factor. In the routine factor, a person carries out daily physical activity and/or other demands. External factors such as sun exposure, dietary intake, or, although these factors can significantly affect vitamin D levels.

During the research, vitamin D levels were measured to determine vitamin D levels in elderly women by taking blood samples from each research respondent. From a total of 49 elderly women, the majority of vitamin D levels <20 ng/

ml were obtained by 27 respondents (55%) with an average of 17.40 ng/ml.

Blood samples were collected only once. This may need to be revised to adequately reflect the variation in vitamin D levels, which can change depending on daily conditions, such as sun exposure and diet. Daily variation was minimized by collecting samples at the same time of day. Blood sampling was carried out simultaneously during the noonday by the phlebotomist, taking into account the time of blood sampling, which can affect vitamin D levels.¹³

This technique can cause bias because the sample is selected based on specific criteria and does not represent the entire population of elderly women. As a result, the study's results may be challenging to apply to all older adults. To minimize the influence of vitamin D variations, several factors that can affect vitamin D variations in the respondent's body condition are minimized by not having significant changes in these factors from one month earlier (such as medication, health conditions, and lifestyle). Several factors can influence vitamin D levels, including the following:^{20,21}

(1) Body mass index (BMI). Vitamin D is a fat-soluble vitamin, and people with a BMI >30 kg/m² tend to have higher fat than those with a normal BMI. The association between vitamin D deficiency and overweight and obese patients may be explained by the storage of vitamin D in adipose tissue due to its high-fat solubility and decreased exposure to sunlight in obese subjects due to limited physical activity and mobility.

(2) Lack of intake. All age-related changes in vitamin D metabolism are magnified if there is concomitant vitamin D deficiency, as it limits the supply of substrates for 25OHD and, ultimately, 1,25(OH)₂D. Substrate deficiency is a common problem in older people and is essential to recognize because it can be prevented and treated. There may be a vitamin D deficiency either from diet or from lack of sunlight, and a subsequent decrease in serum 25OHD limits 1,25(OH)₂D production, especially if there is also renal dysfunction. Serum 1,25(OH)₂D levels decrease when serum 25OHD levels fall below 10 ng/mL in young and older adults.

Conclusions

The correlation test results between physical

activity and vitamin D levels in elderly women showed no significant correlation. Other factors affecting the research results include age, which has a wide range of variations, health conditions, and vitamin D intake. Further research can develop measurements specifically for examining physical activity in old age.

Conflict of Interest

The authors declare no conflict of interest.

Acknowledgment

This research was funded by the Institute of Research and Community Service at Universitas Surabaya.

References

- Mei Z, Hu H, Zou Y, Li D. The role of vitamin D in menopausal women's health. *Front Physiol.* 2023;14:1211896.
- Giustina A, Bouillon R, Dawson-Hughes B, Ebeling PR, Lazaretti-Castro M, Lips P, et al. Vitamin D in the older population: a consensus statement. *Endocrine.* 2023;79(1):31–44.
- Borowicz W, Ptaszkowski K, Ptaszkowska L, Rosińczuk J, Murawska-Ciałowicz E. Association between serum vitamin D levels and physical outcomes of patients who underwent rehabilitation following ischemic stroke. *Med Sci Monit.* 2023;29:e940115.
- Liu W, Wu Z, Zhu D, Chen G, Yan G, Zhang S, et al. Vitamin D and lipid profiles in postmenopausal women: a meta-analysis and systematic review of randomized controlled trials. *Front Mol Biosci.* 2021;8:799934.
- Cui A, Zhang T, Xiao P, Fan Z, Wang H, Zhuang Y. Global and regional prevalence of vitamin D deficiency in population-based studies from 2000 to 2022: a pooled analysis of 7.9 million participants. *Front Nutr.* 2023;10:1070808.
- Pereira M, Dantas Damascena A, Galvão Azevedo LM, de Almeida Oliveira T, da Mota Santana J. Vitamin D deficiency aggravates COVID-19: systematic review and meta-analysis. *Crit Rev Food Sci Nutr.* 2020;62(5):1308–16.
- Widyantari NPS, Wijaya IPA, Susila IMDP. Hubungan tingkat pengetahuan tentang menopause dengan kecemasan menghadapi menopause pada ibu Pembinaan Kesejahteraan Keluarga. *Caring.* 2019;3(2):56–9.
- Sari DK, Alrasyid DH, Nurlndrawaty L, Zulkiffi L. Occurrence of vitamin D deficiency among women in North Sumatera, Indonesia. *Malaysian Journal of Nutrition.* 2014;20(1):63–70.
- Dominguez LJ, Farruggia M, Veronese N, Barbagallo M. Vitamin D sources, metabolism, and deficiency: available compounds and guidelines for its treatment. *Metabolites.* 2021;11(4):255.
- Wiciński M, Adamkiewicz D, Adamkiewicz M, Śniegocki M, Podhorecka M, Szycha P, et al. Impact of vitamin D on physical efficiency and exercise performance—a review. *Nutrients.* 2019;11(11):2826.
- Nascimento NAP, Moreira PFP, Carvalho VA, Aragão L, Marin-Mio RV, Lazaretti-Castro M, et al. Effect of vitamin D Level and physical exercise on the physical performance and functional test results in elderly women. *J Geriatr Med Gerontol.* 2019;5(1):061.
- Suryadinata RV, Lorensia A, Tangkilisan EC. Effect of physical activity and vitamin D status on geriatrics obesity. *GMHC.* 2019;7(1):1–6.
- Abu Jadayil S, Abu Jadayel B, Takruri H, Muwalla M, McGrattan AM. Study of the fluctuation of serum vitamin D concentration with time during the same day and night on a random sample of healthy adults. *Clin Nutr ESPEN.* 2021;46:499–504.
- Li J, Huang Z, Si W, Shao T. The effects of physical activity on positive emotions in children and adolescents: a systematic review and meta-analysis. *Int J Environ Res Public Health.* 2022;19(21):14185.
- Ho JY, Lam HYC, Huang Z, Liu S, Goggins WB, Mo PKH, et al. Factors affecting outdoor physical activity in extreme temperatures in a sub-tropical Chinese urban population: an exploratory telephone survey. *BMC Public Health.* 2023;23(1):101.
- Cleland C, Ferguson S, Ellis G, Hunter RF. Validity of the International Physical Activity Questionnaire (IPAQ) for assessing moderate-to-vigorous physical activity and sedentary behaviour of older adults in the United Kingdom. *BMC Med Res Methodol.* 2018;18(1):176.

17. Sember V, Meh K, Sorić M, Starc G, Rocha P, Jurak G. Validity and reliability of International Physical Activity Questionnaires for adults across EU countries: systematic review and meta analysis. *Int J Environ Res Public Health*. 2020;17(19):7161.
18. Khan SR, Claeson M, Khan A, Neale RE. The effect of physical activity on vitamin D: a systematic review and meta-analysis of intervention studies in humans. *Public Health Pract (Oxf)*. 2024;7:100495.
19. Heydenreich J, Schutz Y, Melzer K, Kayser B. Comparison of conventional and individualized 1-MET values for expressing maximum aerobic metabolic rate and habitual activity related energy expenditure. *Nutrients*. 2019;11(2):458.
20. Wang LK, Hung KC, Lin YT, Chang YJ, Wu ZF, Ho CH, et al. Age, Gender and season are good predictors of vitamin D status independent of body mass index in office workers in a subtropical region. *Nutrients*. 2020;12(9):2719.
21. Lorensia A, Suryadinata RV, Arganitya GN. Relationship of vitamin D intake with obesity in adolescents. *GMHC*. 2022;10(2):104–10.
22. Trovato B, Godos J, Varrasi S, Roggio F, Castellano S, Musumeci G. Physical activity, sun exposure, vitamin D intake and perceived stress in Italian adults. *Nutrients*. 2023;15(10):2301.
23. Zhang J, Cao ZB. Exercise: a possibly effective way to improve vitamin D nutritional status. *Nutrients*. 2022;14(13):2652.
24. Lorensia A, Suryadinata RV, Inu IA. Comparison of vitamin D status and physical activity related with obesity in student. *J Appl Pharm Sci*. 2022;12(4):108–18.
25. VandeBunte A, Gontrum E, Goldberger L, Fonseca C, Djukic N, You M, et al. Physical activity measurement in older adults: wearables versus self-report. *Front Digit Health*. 2022;4:869790.