



Antibiotics consumption in neurosurgery versus appendectomy: a call for antibiotic stewardship initiatives

Rika Yulia¹, Steven Victoria Halim¹, Fauna Herawati¹,
Dian Natasya Raharjo¹, Eko Setiawan¹, Heru Wijono²

1) Departement of Clinical and Community Pharmacy, Faculty of Pharmacy, University of Surabaya, East Java, Indonesia

2) Faculty of Medicine, University of Surabaya, East Java, Indonesia

Abstract

Background and aims. Surgical site infection (SSI) remains a significant global health concern, including in Indonesia. The administration of prophylactic antibiotics plays a pivotal role in the prevention and the reduction in mortality rates associated with SSIs. Prophylactic antibiotic use is recommended in surgical procedures that are at risk of infection, including in clean surgical operations that last more than three hours and clean-contaminated surgeries. The objective of this study was to analyze the quantity of prophylactic antibiotic consumption and to compare the consumption of antibiotics between neurosurgery (clean surgery) and appendectomy (clean-contaminated surgery).

Methods. Data in this observational study were obtained from the medical records of patients who underwent neurosurgery and appendectomy at two hospitals in Surabaya: one private and one public hospital. The quantity of antibiotic consumption will be analyzed descriptively and presented as a defined daily dose (DDD) per 100 bed days.

Results. Research findings revealed a higher quantity of antibiotic consumption in elective neurosurgery, representing 47.43 DDD/100 bed-days, in comparison to 21.26 DDD/100 bed-days and 76.34 DDD/100 bed-days in elective and emergency appendectomy, respectively. The most frequently used antibiotics were broad-spectrum antibiotics, which included cefixime (36.91 DDD/100 bed-days) and ceftriaxone (5.45 DDD/100 bed-days) in elective neurosurgery; and ceftriaxone (14.94 vs 50.86 DDD/100 bed-days) and metronidazole (5.75 vs 19.16 DDD/100 bed-days) in elective and emergency appendectomy, respectively.

Conclusion. The consumption of prophylaxis antibiotics remains a significant concern. In order to develop strategies to prevent bacterial resistance through an antibiotic stewardship program, it is essential to evaluate antibiotic consumption and monitor trends over time.

Keywords: antibiotic prophylaxis, neurosurgery, appendectomy, antimicrobial stewardship, drug evaluation

DOI: 10.15386/mpr-2823

Manuscript received: 30.09.2024

Received in revised form: 15.01.2025

Accepted: 15.02.2025

Address for correspondence:

Steven Victoria Halim

stevenvictoria@staff.ubaya.ac.id

This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License <https://creativecommons.org/licenses/by-nc-nd/4.0/>

Background and aims

The prevalence of surgical site infections (SSIs) represents a significant global public health concern in the context of surgical procedures. The incidence of SSIs varies among countries, with a global incidence rate of 5-10% [1]. Additionally, SSIs are becoming more prevalent in low- and middle-income countries, with an incidence rate that could exceed 15% [2-4]. Indonesia is one

of the low-middle income countries, with a reported SSI incidence ranging from 13.10% to 56.7% [5-8].

SSIs have been identified as one of major risk factors of 30-day postoperative mortality, which further make SSI listed as the third leading cause of death globally [9-11]. Moreover, SSIs have been linked to an increased length of stay, higher healthcare costs, increased morbidity, poorer therapeutic outcomes, and a longer

course of antibiotic therapy [3,12,13]. Hence, in order to prevent the incidence of SSI, antibiotic prophylaxis is required in surgical procedures, especially clean surgery and clean-contaminated surgery [14]. Clean surgery is defined as a surgical procedure that is planned and performed in uninfected anatomical areas, without opening the gastrointestinal, biliary, urinary, or respiratory tracts. These procedures typically involve primary skin closure, with or without the application of closed drains. On the other hand, clean-contaminated surgery refers to operations involving hollow organs, such as the gastrointestinal, biliary, urinary, respiratory, or reproductive systems (excluding the ovaries), or surgeries where contamination is not evident. In general, the administration of prophylactic antibiotics is not required for clean surgery, unless the procedure is performed within a body cavity which, if infected, would pose a significant risk, such as neurosurgery or clean surgery lasting more than three hours [14].

Neurosurgery is classified as a clean surgery, with a relatively high incidence of surgical site infections [14-17]. A systematic review and meta-analysis conducted by Lee et al. (2024), revealed that the incidence of SSIs at one and three months post-neurosurgery was 4.03% and 6.17%, respectively. Furthermore, the study indicated that 13.9% of patients were readmitted and 16.3% required reoperation due to SSI post-neurosurgery [18]. It is worth emphasizing that SSIs in neurosurgery could result in significant complications, including meningitis, encephalitis, epidural abscess, subdural empyema, and brain abscess, which can be life-threatening [19-21].

Another type of surgical procedure is the appendectomy, which is classified as clean-contaminated surgery [14]. The global incidence of surgical site infection (SSI) in patients undergoing appendectomy is comparatively high, with a range of 6.4% to 7.7% [22-25]. Notably, this incidence is even higher in low-middle income countries, including Indonesia, reaching rates up to 11% [22]. The SSI post-appendectomy can be caused by the presence of a large number of normal flora within the intestine, which may enter and infect the surgical wound [22,23].

To reduce the prevalence of SSI in surgical procedures, including neurosurgery and appendectomy, antibiotic stewardship programs (ASPs) should be implemented [26-28]. These programs are systematic initiatives designed to promote the responsible use of antimicrobials, enhance treatment efficacy, limit the emergence of resistance, and control the spread of infections caused by resistant microorganisms [29,30]. A commonly used metric in ASPs is the defined daily dose (DDD), which was developed by the World Health Organization (WHO). Within the framework of ASPs, the DDD is used to quantify the antibiotic consumption, monitor and evaluate patterns of antibiotic use, identify trends, and detect potential overuse or misuse in clinical practice [29-31].

Prior to the present study, the available published research on the quantity of prophylactic antibiotic use in patients undergoing neurosurgery and appendectomy was relatively limited. The study by Nasare et al. (2023) revealed an increase in the consumption of antimicrobial agents in neurosurgery from 125.7 to 155.5 DDDs/100 days within a five-year observation. The consumption of Watch group antibiotics, as defined by the WHO AWaRe classification, was higher than that of Access group antibiotics in both the intensive care unit (ICU) and in patient department settings, representing over 75% of the total antibiotic consumption [32]. Effort to prevent SSI by relying heavily on the use of antibiotic should be taken with precaution because antibiotic consumption was associated with the developmental of antibiotic resistant pathogens [28,33].

A relatively limited number of studies aiming to identify the use of antibiotic prophylaxis in an Indonesian setting have been documented in the published literature [34,35]. Majority of the studies were specifically conducted among adult patients undergoing surgery and ceftriaxone was found as the most common used antibiotic for prophylaxis in both neurosurgery and appendectomy surgery [34-36]. It should be mentioned that the profile of antibiotic use in pediatric patients in an Indonesian setting is yet to be explored.

In light of the potential risk of SSIs and the pivotal role of responsible antibiotic use, it is essential to assess the utilization of antibiotics. The most common method of evaluating antibiotic use is to assess the quantity of antibiotics using the defined daily dose (DDD) [28,33,37,38]. Based on the classification of the surgical procedure, the quantity of antibiotics administered during clean surgery (neurosurgery) is expected to be lower than in clean-contaminated surgery (appendectomy). However, no studies have been conducted to compare the antibiotic consumption in both types of surgery, especially in Indonesia. Thus, this study aims to analyze the quantity of prophylactic antibiotic consumption and to compare the consumption of antibiotics in patients undergoing neurosurgery (clean surgery) and appendectomy (clean-contaminated surgery).

Methods

Study design and setting

This research is a descriptive observational study with retrospective data collection. Data were collected from medical records in two tertiary care hospitals in Surabaya, i.e one private hospital and one public hospital. The private hospital has 205 beds, whereas public hospital has 293 beds. This research has been reviewed by the Institutional Ethical Committee of the University of Surabaya and obtained a research ethics permit (letter number 013/KE/KEPK/2020 and 142/KE/XI/2020).

Population and sample

The population in this research consists of all patients who underwent neurosurgery or appendectomy

from January 2019 to November 2020. The research sample was selected using a purposive sampling technique. Patients who met the research criteria were included in this research. The inclusion criteria in this research were: (1) patients aged over 5 years; (2) patients who received prophylactic antibiotics before surgery; and (3) patients with complete medical records.

Data collection

Data were collected using a standardized data collection form consisting of two sections: patient characteristics and antibiotic profile. Patient characteristics included sex, age, diagnosis and length of stay (LOS), type of surgical procedure. The type surgical procedure may be categorized as either elective or emergency surgery. Elective surgery is defined as a surgical procedure that is not an emergency and can be scheduled in advance. In contrast, emergency surgery is defined as a surgical procedure that must be performed without delay in order to save the life of a patient. Antibiotic profile in this study included type of antibiotics, dosage, route of administration, and Anatomical Therapeutic Chemical (ATC) code.

Data analysis

Data were analyzed descriptively. Patient characteristics were collected and presented in tables and graphs, followed by descriptive explanations. Antibiotic quantity profiles were classified using the ATC system to obtain DDD values for each antibiotic and analyzed using the Defined Daily Dose (DDD)/100 days, according to the following formula [40]:

$$\text{DDD}/100 \text{ bed-days} = \frac{\text{Antibiotic consumption (g)}}{\text{WHO DDD antibiotic (g)}} \times \frac{100}{\text{total patients' length of stay}}$$

In addition, antibiotics prophylaxis in this study were also classified according to the 2021 WHO AWaRe classification [40].

Results

A total of 180 medical records were reviewed in this research, consisting of 142 records from patients who had neurosurgery and 38 records from patients who had an appendectomy. Of the 38 patients who underwent appendectomy, 23 patients had uncomplicated appendicitis and 15 patients had complicated appendicitis. The majority of patients undergoing neurosurgery and appendectomy with uncomplicated appendicitis were female (more than 60%), whereas the majority of patients with complicated appendicitis were male (73%). Most neurosurgery patients (81.69%) were over 46 years old. Out of 180 patients, 18.33% were patients under 17 years old (Table I).

All patients included in this study underwent elective neurosurgery. The antibiotic with the highest defined daily dose (DDD) per 100 bed-days in patients undergoing neurosurgery is cefixime (36.91 DDD/100 bed-days), which implies that, on average, each patient receives 0.36 DDD of cefixime per inpatient day. According to the 2021 WHO AWaRe classification, most antibiotics used in neurosurgery were categorized as 'Watch' and 'Access' antibiotics, each accounting for 37.5%. In addition, two reserve antibiotics were found to be administered in patient underwent neurosurgery: tigecycline (2.58 DDD/100 bed-days) and meropenem (0.16 DDD/100 bed-days). A detailed description of the antibiotics used in neurosurgery could be found in table II.

Table I. Patients characteristics (N=180).

Characteristics	Clean surgery (N=142)	Clean-contaminated surgery (N=38)	
	Neurosurgery (n, %)	Uncomplicated appendicitis ¹ (n, %)	Complicated appendicitis ² (n, %)
Gender			
• Male	48 (33.80)	9 (39.13)	11 (73.33)
• Female	94 (66.20)	14 (60.87)	4 (26.67)
Age (years)			
• 6 – <12	2 (1.41)	3 (13.04)	1 (6.67)
• 12 – <17	3 (2.11)	4 (17.39)	2 (13.33)
• 17 – <26	4 (2.82)	2 (8.70)	2 (13.33)
• 26 – <36	3 (2.11)	3 (13.04)	4 (26.67)
• 36 – <46	14 (9.86)	6 (26.09)	2 (13.33)
• 46 – <56	35 (24.65)	4 (17.39)	1 (6.67)
• 56 – <65	36 (25.35)	1 (4.35)	2 (13.33)
• 65<	45 (31.69)	0	1 (6.67)
Total LOS (days)	4,072	87	86

¹ – acute appendicitis (K35), acute appendicitis unspecified (K35.8); ² – generalized peritonitis (K35.2), perforation (K35.3), peritoneal abscess (K35.3); LOS: length of stay

Table II. Prescribed prophylactic antibiotics for neurosurgery in DDD/100 bed-days.

Variable	ATC code	WHO AWaRe classification	DDD/100 bed-days
Elective			
Tigecycline	J01AA12	Reserve	2.58
Meropenem	J01DH02	Reserve	0.16
Ceftriaxone	J01DD04	Watch	5.45
Cefixime	J01DD08	Watch	36.91
Moxifloxacin	J01MA14	Watch	0.80
Gentamicin	J01GB03	Access	1.06
Amikacin	J01GB06	Access	0.25
Metronidazole	J01XD01	Access	0.22
Total			47.43
Emergency	-	-	-

Table III. Prescribed prophylactic antibiotics for appendectomy in DDD/100 bed-days.

Variable	ATC code	WHO AWaRe classification	Type of appendicitis		
			Uncomplicated appendicitis ¹	Complicated appendicitis ²	Total
Elective					
Cefuroxime	J01DC02	Watch	0.57	-	0.57
Ceftriaxone	J01DD04	Watch	14.94	4.65	19.59
Metronidazole	J01XD01	Access	5.75	4.65	10.40
Total			21.26	9.30	30.56
Emergency					
Cefotaxime	J01DD01	Watch	-	0.58	0.58
Ceftriaxone	J01DD04	Watch	50.86	61.63	112.49
Cefixime	J01DD08	Watch	4.02	0.58	4.60
Ciprofloxacin	J01MA02	Watch	-	4.36	4.36
Cefazolin	J01DB04	Access	2.30	1.55	3.85
Metronidazole	J01XD01	Access	19.16	54.65	73.81
Total			76.34	123.35	199.69

¹ – acute appendicitis (K35), acute appendicitis unspecified (K35.8); ² – generalized peritonitis (K35.2), perforation (K35.3), peritoneal abscess (K35.3); A – WHO Access classification; Wa – WHO Watch classification; Re – WHO Reserve classification.

The patients with appendicitis included in this study underwent either elective or emergency appendectomy. Ceftriaxone is the most commonly used as a prophylactic antibiotic both in elective and emergency appendectomy with 19.59 DDD/100 bed-days and 112.49 DDD/100 bed-days, respectively. The antibiotic recommended for appendectomy prophylaxis by American Society of Health System Pharmacist (ASHP) guideline and the Ministry of Health Republic Indonesia guideline published in 2021 is cefazolin, either alone or in combination with metronidazole. Metronidazole was identified as the second most commonly used antibiotic in both elective (10.40 DDD/100 bed-days) and emergency appendectomy procedures (73.81 DDD/100 bed-days). In contrast, the use of cefazolin in emergency appendectomy cases accounted for only approximately 2% of the total DDD (3.85 out of 199.69 DDD/100 bed-days; table III). According to the 2021 WHO AWaRe classification, antibiotics for both elective and emergency appendectomies are mainly categorized as ‘Watch’.

The use of prophylactic antibiotic in clean surgery (elective neurosurgery) was found to be two times

higher than in clean-contaminated surgery (elective appendectomy), with 47.43 DDD/100 bed-days compared to 21.26 DDD/100 bed-days. A detailed description of the antibiotics used in neurosurgery, compared with those used in appendectomy, can be found in table II and table III.

Antibiotics for patients with appendicitis were administered at a relatively various points of time in the surgical process, including before, during, and after surgery. The quantity of antibiotic use in patients with uncomplicated appendicitis before, during, and after surgery was 12.64 DDD/100 bed-days, 27.30 DDD/100 bed-days, and 57.66 DDD/100 bed-days, respectively. In contrast, patients with complicated appendicitis showed a markedly increased quantity of antibiotic use before, during, and after surgery, reaching 13.28 DDD/100 bed-days, 23.45 DDD/100 bed-days, and 95.93 DDD/100 bed-days, respectively. The use of antibiotics after elective appendectomy surgery (beyond 24 hours after the procedure) is 1-4 times higher than during surgery. Furthermore, antibiotic use for emergency appendectomy after surgery is 2-5 times higher compared to during surgery (Table IV).

Table IV. Prescribed antibiotic for appendectomy.

Variable	ATC code	Uncomplicated ¹ – Complicated ²		
		Before	During	After
Elective				
Cefuroxime (Wa)	J01DC02	0.00 – 0.00	0.57 – 0.00	0.00 – 0.00
Ceftriaxone (Wa)	J01DD04	2.30 – 1.16	3.45 – 1.16	9.20 – 2.33
Metronidazole (A)	J01XD01	0.00 – 1.16	0.00 – 1.16	5.75 – 2.33
Total		2.30 – 2.32	4.02 – 2.32	14.95 – 4.66
Emergency				
Cefazolin (A)	J01DB04	0.00 – 0.78	2.30 – 0.78	0.00 – 0.00
Cefotaxime (Wa)	J01DD01	0.00 – 0.00	0.00 – 0.58	0.00 – 0.00
Ceftriaxone (Wa)	J01DD04	6.90 – 7.56	17.53 – 11.63	26.44 – 42.44
Cefixime (Wa)	J01DD08	0.00 – 0.00	0.00 – 0.00	4.02 – 0.58
Ciprofloxacin (Wa)	J01MA02	0.00 – 0.87	0.00 – 0.58	0.00 – 2.91
Metronidazole (A)	J01XD01	3.45 – 1.74	3.45 – 7.56	12.26 – 45.35
Total		10.35 – 10.95	23.28 – 21.13	42.72 – 91.28

¹ – acute appendicitis (K35), acute appendicitis unspecified (K35.8); ² – generalized peritonitis (K35.2), perforation (K35.3), peritoneal abscess (K35.3); A – WHO Access classification; Wa – WHO Watch classification; Re – WHO Reserve classification.

Discussion

A number of different types of antibiotics are used in neurosurgery for prophylaxis and cefixime is the most frequently used antibiotic in patients undergoing neurosurgery. Furthermore, ceftriaxone is the second most commonly used antibiotic in neurosurgery. Cefixime and ceftriaxone are classified as the third-generation cephalosporin antibiotics with a broad spectrum of activity [41]. In accordance with the national antibiotic guidelines provided by ASHP guideline and the Ministry of Health Republic Indonesia guideline, cefazolin at a dose of 2 grams intravenously is the recommended prophylactic antibiotic for patients undergoing neurosurgery, including craniotomy, spinal cord surgery, and cerebrospinal fluid shunting [14,42]. The published evidence indicated that the use of third-generation cephalosporins was not superior compared to conventional antibiotics (cefazoline, trimethoprim/sulfamethoxazole, ampicillin/sulbactam, vancomycin and gentamicin) as a prophylaxis in neurosurgery. It was documented that the overall incidence of surgical site infections (SSIs) following neurosurgery was not statistically different between the use of third generation cephalosporins and conventional antibiotics (OR 0.94; 95% CI 0.59-1.52). The results of the study regarding the incidence of SSIs, such as osteomyelitis, meningitis, intracranial infection and disc space also demonstrated a non-significant difference between third generation cephalosporins and conventional antibiotics (OR 0.88; 95% CI 0.45-1.74) [43]. In addition to the lack of superiority in efficacy, it should be emphasized that the overuse of third-generation cephalosporins can result in the emergence of third-generation cephalosporin-resistant Enterobacterales (3GCRE) due to the production of beta-lactamases, particularly extended-spectrum beta-

lactamases (ESBL). This phenomenon could potentially lead to the incremental of last-resort antibiotics utilisation, including carbapenems, which could exacerbate the global resistance problem [44,45]. The World Health Organization (WHO) has previously identified third-generation cephalosporin-resistant Enterobacterales (3GCRE) as a critical priority with the highest priority for monitoring in terms of resistance issues globally [45].

Ceftriaxone and metronidazole are two types of prophylactic antibiotics that are commonly employed in patients undergoing appendectomy, both in elective and emergency surgery. The recommended antibiotic for prophylaxis in patients undergoing appendectomy is cefazoline, administered intravenously at a dose of 2 grams alone [14,42] or in combination with metronidazole, administered intravenously at a dose of 500 mg [46-48]. A review of the literature reveals no studies that have directly compared the efficacy and safety of cefazoline-metronidazole and ceftriaxone-metronidazole as prophylactic antibiotics for appendectomy. A number of published studies have demonstrated that the combination of ceftriaxone with metronidazole represents the optimal antibiotic choice for patients presenting with complicated or perforated appendicitis [49-52]. As previously described, the inappropriate use of ceftriaxone as a prophylactic antibiotic has the potential to result in the emergence of 3GCRE bacteria [44,45].

The consumption of antibiotics in elective neurosurgery (clean surgery) demonstrated a higher quantity than that observed in appendectomy (clean-contaminated surgery) which 47.43 DDD/100 bed-days compared to 21.26 DDD/100 bed-days. In general, prophylactic antibiotics are not indicated for clean surgeries, except in procedures with higher risk of infection, including

neurosurgery and surgeries that exceed 3 hours in duration. In the national antibiotic guidelines provided by the Ministry of Health in 2021, the recommended antibiotics for patients undergoing neurosurgery and appendectomy are Cefazoline 2 grams intravenously as a single dose, which may be re-administered if the surgical procedure exceeds 3 hours [14,42]. The increased consumption of antibiotics in neurosurgery that classified as clean surgery can be attributed to a number of factors. Primarily, the duration of neurosurgery is longer than that of appendectomy, which increases the likelihood of re-administration of antibiotics (repeat antibiotic treatment). Secondly, neurosurgeons tend to administer antibiotics following surgery to prevent the occurrence of surgical site infections, which can lead to an increased risk of postoperative mortality [53].

The quantity of antibiotics administered after surgery was found to be greater than that administered during surgery, in both elective and emergency surgical procedures. This suggests that the practice of administering antibiotics for an extended period following the patient's discharge from the hospital is common. The practice of administering postoperative antibiotics has also been demonstrated in several published studies [54-56]. A number of published studies found that patients who received postoperative prophylactic antibiotics did not show a significantly lower incidence of SSIs compared to patients who did not receive postoperative antibiotics [55,57-59]. A summary of systematic review of 44 trials comparing the use of prolonged antibiotics compared to no postoperative antibiotics in different types of surgery revealed that only three trials demonstrated a reduced risk of surgical site infection (SSI). In the context of appendectomies, a review of five trials revealed a non-significant difference (OR 0.71; 95% CI 0.41-1.23) in the incidence of SSI between prolonged antibiotic administration and single-dose prophylactic antibiotic administration [60].

The findings of this research are crucial for advancing both future research and clinical practice. The results revealed that the types of prophylactic antibiotics used in neurosurgery and appendectomy varied and did not fully comply with the guidelines established by the ASHP and the Ministry of Health of the Republic of Indonesia. These findings should prompt further research involving interprofessional collaboration among all healthcare workers in hospitals to identify effective strategies for enhancing adherence to prophylactic antibiotic guidelines in surgical procedures [26-28]. Additionally, the findings of this study could provide a basis for the development of hospital policies regarding the rational use of prophylactic antibiotics.

The limitations of this study are inherent to its cross-sectional research design, which provides a description of antibiotic consumption only for a specific period. Consequently, further research is required to quantify antibiotic consumption on a continuous basis, thus

providing data that can be used to predict the incidence of bacterial resistance, assist in the selection of optimal antibiotics for patients and effective antibiotic stewardship program. In addition, we included all patients, those with out of pocket expenses and patients covered by health insurance. The different payment methods of medical services could impact on the prescribing behavior of intra-hospital physicians [61,62]. Lastly, antibiotic consumption is influenced by the patient visiting the hospital during a specific period, which may result in differences in antibiotic consumption patterns between these two tertiary care hospitals and other hospitals.

Conclusion

The use of broad-spectrum antibiotics, including the third-generation cephalosporins and metronidazole, as a prophylaxis in neurosurgery and appendectomy were a common practice in our research sites. Moreover, some of restricted antibiotics were also found as the prophylaxis antibiotics for neurosurgery and appendectomy. Owing to the further negative consequences of these prescribing practices, effort should be made to facilitate the successful implementation of an antibiotic stewardship program.

Acknowledgements

The authors would like to thank the Directorate of Research, Technology and Community Service (DRTPM) Directorate General of Higher Education, Research and Technology (Ditjen Diktilistik) Ministry of Education, Culture, Research and Technology (Kemendikbudristek) Republic of Indonesia for providing research funding for this project, as outlined in the contract number 051/ST-Lit/LPPM-01/FF/VII/2024. We would also like to acknowledge the director of the hospital for granting permission for this research to be conducted and all hospital staff who provided assistance during the data collection process.

References

1. Mengistu DA, Alemu A, Abdukadir AA, Mohammed Husen A, Ahmed F, Mohammed B, et al. Global incidence of surgical site infection among patients: systematic review and meta-analysis. *Inquiry*. 2023;60:469580231162549.
2. Mehtar S, Wanyoro A, Ogunola F, Ameh EA, Nthumba P, Kilpatrick C, et al. Implementation of surgical site infection surveillance in low- and middle-income countries: A position statement for the International Society for Infectious Diseases. *Int J Infect Dis*. 2020;100:123-131.
3. Monahan M, Jowett S, Pinkney T, Brocklehurst P, Morton DG, Abdali Z, et al. Surgical site infection and costs in low- and middle-income countries: A systematic review of the economic burden. *PLoS One*. 2020;15:e0232960.
4. Rickard J, Beilman G, Forrester J, Sawyer R, Stephen A, Weiser TG, et al. Surgical infections in low- and middle-

- income countries: a global assessment of the burden and management needs. *Surg Infect (Larchmt)*. 2020;21:478-494.
5. Atira A, Salmiyah E, Purwandi DP. Surgical site infection incidence in patients post appendectomy. *Global Health Science*. 2021;6:101-104.
 6. Aulya Y, Novelia S, Isnaeni A. Difference in surgical site infection incidence between elective and emergency cesarean section at Harapan Jayakarta Hospital in 2019. *Journal for Quality in Women's Health*. 2021:115-122.
 7. Chairani F, Puspitasari I, Asdie RH. Incidence and risk factors of surgical site infection in obstetric and gynecologic surgery at the hospital. *J Manag Pharm Pract*. 2019; 9:274.
 8. Marsanto AE, Paraton H, Budi Prasetyo B. Pattern of disease and type of operation of surgical site infection in obstetrics and gynecology at Dr Soetomo Hospital, Surabaya, Indonesia. *Majalah Obstetri dan Ginekologi*. 2019;27:49-55.
 9. Nepogodiev D, Martin J, Biccard B, Makupe A, Bhangu A. National Institute for Health Research Global Health Research Unit on Global Surgery. Global burden of postoperative death. *Lancet*. 2019;393:401.
 10. O'Brien WJ, Gupta K, Itani KMF. Association of Postoperative Infection With Risk of Long-term Infection and Mortality. *JAMA Surg*. 2020;155:61-68.. Erratum in: *JAMA Surg*. 2020;155:94.
 11. Wan YI, Patel A, Achary C, Hewson R, Phull M, Pearse RM; International Surgical Outcomes Study (ISOS) Group. Postoperative infection and mortality following elective surgery in the International Surgical Outcomes Study (ISOS). *Br J Surg*. 2021;108:220-227.
 12. HirHirani S, Trivedi NA, Chauhan J, Chauhan Y. A study of clinical and economic burden of surgical site infection in patients undergoing caesarian section at a tertiary care teaching hospital in India. *PLoSOne*. 2022;17:e0269530.
 13. Totty JP, Moss JWE, Barker E, Mealing SJ, Posnett JW, Chetter IC, et al. The impact of surgical site infection on hospitalisation, treatment costs, and health-related quality of life after vascular surgery. *Int Wound J*. 2021;18:261-268.
 14. Ministry of Health Republic Indonesia (2021) Regulation of the Ministry of Health Republic Indonesia Number 28 of 2021 on Guidelines for Antibiotic Use. 1-98.
 15. Stienen MN, Moser N, Krauss P, Regli L, Sarnthein J. Incidence, depth, and severity of surgical site infections after neurosurgical interventions. *Acta Neurochir (Wien)*. 2019;161:17-24.
 16. Abdel-Latif AM, Moharram AA, Higazy A, Ghoneim NI, Shafei O, Abdelhady SG, et al. Incidence and perioperative risk factors for surgical site infections in neurosurgery: prospective observational study. *Egyptian J Neurosurg*. 2022;37:2-5.
 17. López Pereira P, Díaz-Agero Pérez C, López Fresneña N, Las Heras Mosteiro J, Palancar Cabrera A, Rincón Carlavilla ÁL, et al. 'Epidemiology of surgical site infection in a neurosurgery department'. *Br J Neurosurg*. 2017;31:10-15.
 18. Lee KS, Borbas B, Plaha P, Ashkan K, Jenkinson MD, Price SJ. Incidence and Risk Factors of Surgical Site Infection After Cranial Surgery for Patients with Brain Tumors: A Systematic Review and Meta-analysis. *World Neurosurg*. 2024;185:e800-e819.
 19. Nanda A. *Complications in neurosurgery*: Elsevier Health Sciences: 2018: 217-224
 20. Rolston JD, Han SJ, Lau CY, Berger MS, Parsa AT. Frequency and predictors of complications in neurological surgery: national trends from 2006 to 2011. *J Neurosurg*. 2014;120:736-745.
 21. Venkatapura RJ, Dubey SK, Panda N, Chakrabarti D, Venkataramaiah S, Rath GP et al. Postoperative Neurological Complications after a Cranial Surgery: A Multicentre Prospective Observational Study. *J Neuroanaesthesiol Crit Care*. 2021;8:034-039.
 22. Danwang C, Bigna JJ, Tochie JN, Mbonda A, Mbanga CM, Nzalie RNT, et al. Global incidence of surgical site infection after appendectomy: a systematic review and meta-analysis. *BMJ Open*. 2020;10:e034266.
 23. Singh R, Singla P, Chaudhary U. Surgical site infections: classification, risk factors, pathogenesis and preventive management. *Int J Pharm Res Health Sci*. 2014;2:203-214.
 24. Gillespie BM, Harbeck E, Rattray M, Liang R, Walker R, Latimer S, et al. Worldwide incidence of surgical site infections in general surgical patients: A systematic review and meta-analysis of 488,594 patients. *Int J Surg*. 2021;95:106136.
 25. Yang L, Zheng R, Li H, Ren Y, Chen H. The burden of appendicitis and surgical site infection of appendectomy worldwide. *J Infect Dev Ctries*. 2023;17:367-373.
 26. Sarang B, Tiwary A, Gadgil A, Roy N. Implementing antimicrobial stewardship to reduce surgical site infections: Experience and challenges from two tertiary-care hospitals in Mumbai, India. *J Glob Antimicrob Resist*. 2020;20:105-109.
 27. Martinez-Sobalvarro JV, Júnior AAP, Pereira LB, Baldoni AO, Ceron CS, Dos Reis TM. Antimicrobial stewardship for surgical antibiotic prophylaxis and surgical site infections: a systematic review. *Int J Clin Pharm*. 2022;44:301-319.
 28. Kourbeti I, Kamiliou A, Samarkos M. Antibiotic Stewardship in Surgical Departments. *Antibiotics (Basel)*. 2024;13:329.
 29. Center for Disease Control and Prevention. The core elements of hospital antibiotic stewardship programs: 2019. CDC. 2019. . Available from: <https://www.cdc.gov/antibiotic-use/healthcare/pdfs/hospital-core-elements-H.pdf>
 30. Center for Disease Control and Prevention. Core elements of antibiotic stewardship for health departments. CDC. 2023. Available from: <https://www.cdc.gov/antibiotic-use/media/pdfs/Core-Elements-ASHD-508.pdf>
 31. Misra A, Sharma S, Rai S, Rangari G, Katiboina SR, Kutikuppala LVS, et al. Defined Daily Dose (DDD): An Essential Metric in the Antimicrobial Stewardship Programmes (AMSPs) in the Healthcare Sector. *Journal of Antimicrobial Stewardship Practices and Infectious*

- Diseases, 2023;1:27-33.
32. Nasare NV, Bhardwaj A, Bablani V, Gupta R, Kushwaha S, Sharma S. Antimicrobial Utilization Study in a Neurology Setting. *J Indian Med Assoc.* 2023;121:41-44.
33. Birgand G, Dhar P, Holmes A. The threat of antimicrobial resistance in surgical care: the surgeon's role and ownership of antimicrobial stewardship. *Br J Surg.* 2023;110:1567-1569.
34. Azrifitria A, Mudahar H, Fazriyah N, editors. A retrospective study of the antibiotics prophylactic use in appendectomy patient with ATC/DDD and DU 90% method in Regional General Hospital in Jakarta. Proceedings of the 1st International Conference on Health Science, ICHS 2020, 26-27 October 2020, Jakarta, Indonesia; 2021.
35. Anggraini W, Wiraningtias NB, Inayatilah FR, Indrawijaya YYA. Evaluation of antibiotics use in post-surgical acute appendicitis patients at the Pasuruan Regency General Hospital in 2018. *Pharmaceutical Journal of Indonesia.* 2020;6:15-20.
36. Noer SS, Yulia R, Herawati F, Zamroni A. Analysis of Antibiotic Use in Neurosurgery Patients in the ICU of Dr. R. Sosodoro Djatikoesoemo Regional Hospital, Bojonegoro. *Syntax Literate: Jurnal Ilmiah Indonesia,* 2022;7:9037-9052.
37. Pinchera B, Buonomo AR, Schiano Moriello N, Scotto R, Villari R, Gentile I. Update on the Management of Surgical Site Infections. *Antibiotics (Basel).* 2022;11:1608.
38. World Health Organization. Defined Daily Dose (DDD). 2024, August 10 Available from: <https://www.who.int/tools/atc-ddd-toolkit/about-ddd>.
39. Ministry of Health Republic Indonesia. Regulation of the Ministry of Health Republic Indonesia Number 8 of 2015 on the Antimicrobial Resistance Control Program in Hospitals. 2015; 1-32.
40. World Health Organization. 2021 AWaRe classification. 2021, September 9 Available from: <https://www.who.int/publications/i/item/2021-aware-classification>
41. Karaali C, Emiroglu M, Esin H, Sert I, Aydın C, Atalay S, et al. Assessment of prophylactic antibiotic usage habits of the general surgeons in Turkey. *J Infect Dev Ctries.* 2020;14:758-764.
42. American Society of Health System Pharmacist. Clinical practice guidelines for antimicrobial prophylaxis in surgery. ASHP therapeutic guidelines. 2022. Available from: <https://www.ashp.org/surgical-guidelines>
43. Gallagher J, Macdougall C. *Antibiotics simplified.* 5th ed. United States: Jones & Barlett Learning; 2022, pp. 40-85.
44. Liu W, Neidert MC, Groen RJ, Woernle CM, Grundmann H. Third-generation cephalosporins as antibiotic prophylaxis in neurosurgery: what's the evidence? *Clin Neurol Neurosurg.* 2014;116:13-19.
45. Belley A, Morrissey I, Hawser S, Kothari N, Knechtel P. Third-generation cephalosporin resistance in clinical isolates of Enterobacterales collected between 2016-2018 from USA and Europe: genotypic analysis of β -lactamases and comparative in vitro activity of cefepime/enmetazobactam. *J Glob Antimicrob Resist.* 2021;25:93-101.
46. World Health Organization. WHO Bacterial Priority Pathogens List, 2024: bacterial pathogens of public health importance to guide research, development and strategies to prevent and control antimicrobial resistance. 2024, September 15 Available from: <https://iris.who.int/bitstream/handle/10665/376776/9789240093461-eng.pdf?sequence=1>.
47. Bratzler DW, Dellinger EP, Olsen KM, Perl TM, Auwaerter PG, Bolon MK, et al. Clinical practice guidelines for antimicrobial prophylaxis in surgery. *Am J Health Syst Pharm.* 2013;70:195-283.
48. Central administration of pharmaceutical care, General administration for drug utilization and pharmacy practice . National Guidelines for Antimicrobial Prophylaxis in Surgery. 2022, pp. 1-36.
49. Stanford Health Care. SHC Surgical Antimicrobial Prophylaxis Guidelines. Stanford Antimicrobial Safety and Sustainability Program. 2019, pp. 1-6.
50. Hurst AL, Olson D, Somme S, Child J, Pyle L, Ranade D, et al. Once-Daily Ceftriaxone Plus Metronidazole Versus Ertapenem and/or Cefoxitin for Pediatric Appendicitis. *J Pediatric Infect Dis Soc.* 2017;6:57-64.
51. Kashtan MA, Graham DA, Melvin P, Hills-Dunlap JL, Anandalwar SP, Rangel SJ. Ceftriaxone with Metronidazole versus Piperacillin/Tazobactam in the management of complicated appendicitis in children: Results from a multicenter pediatric NSQIP analysis. *J Pediatr Surg.* 2022;57:365-372.
52. Lee JY, Ally S, Kelly B, Kays D, Thames L. Once Daily Dosing of Ceftriaxone and Metronidazole in Children With Perforated Appendicitis. *J Pediatr Pharmacol Ther.* 2016;21:140-145.
53. Zeineddin S, Pitt JB, Linton S, De Boer C, Hu A, Carter M, et al. Postoperative Antibiotics for Complicated Appendicitis in Children: Piperacillin/Tazobactam Versus Ceftriaxone with Metronidazole. *J Pediatr Surg.* 2023;58:1128-1132.
54. Bialek Ł, Rydzińska M, Vetterlein MW, Dobruch J, Skrzypczyk MA. Trauma and Reconstructive Urology Working Party of the European Association of Urology Young Academic Urologists. A Systematic Review on Postoperative Antibiotic Prophylaxis after Pediatric and Adult Male Urethral Reconstruction. *J Clin Med.* 2023;12:6162.
55. Clean Cut Investigators Group. An observational cohort study on the effects of extended postoperative antibiotic prophylaxis on surgical-site infections in low- and middle-income countries. *Br J Surg.* 2024;111:znad438.
56. Muliani N, Herawati F, Yulia R, Wijono H. Quantity and quality profiles of antibiotics pre, on, and post surgery in a hospital setting. *Int J Clin Pharm.* 2021;43:1302-1310.
57. de Jonge SW, Boldingh QJJ, Solomkin JS, Dellinger EP, Egger M, Salanti G, et al. Effect of postoperative continuation of antibiotic prophylaxis on the incidence of surgical site infection: a systematic review and meta-analysis. *Lancet Infect Dis.* 2020;20:1182-1192.
58. Dhole S, Mahakalkar C, Kshirsagar S, Bhargava A. Antibiotic Prophylaxis in Surgery: Current Insights and

- Future Directions for Surgical Site Infection Prevention. *Cureus*. 2023;15:e47858.
59. Wang F, Chin R, Piper M, Esserman L, Sbitany H. Do Prolonged Prophylactic Antibiotics Reduce the Incidence of Surgical-Site Infections in Immediate Prosthetic Breast Reconstruction? *Plast Reconstr Surg*. 2016;138:1141-1149.
60. World Health Organization. Summary of a systematic review on surgical antibiotic prophylaxis prolongation. Global guidelines for the prevention of surgical site infection. 2018.
61. Bahari NI, Baharom M, Zahid SNA, Daud F. Behavioral Impact on Clinical Specialist Payment Method: A Systematic Review. *Iran J Public Health*. 2022;51:1469-1480.
62. Quinn AE, Trachtenberg AJ, McBrien KA, Ogundeji Y, Souri S, Manns L, et al. Impact of payment model on the behaviour of specialist physicians: A systematic review. *Health Policy*. 2020;124:345-358.

MPR

Vol. 98 / no. 2 / 2025

MEDICINE and PHARMACY REPORTS

established in 1920 as CLUJUL MEDICAL



IULIU
HATIEGANU
MEDICAL
UNIVERSITY
PRESS



THE ROLE OF PHARMACISTS IN ENSURING RATIONAL ANTIBIOTIC THERAPY IN THE INTERDISCIPLINARY TEAM • COLONIC POLYPS IN NORTH-WESTERN ROMANIA: A COMPREHENSIVE SINGLE-CENTER ANALYSIS OF ENDOSCOPIC AND PATHOLOGICAL FEATURES • RETURN TO WORK OF HEAD AND NECK CANCER PATIENTS AFTER CURATIVE RADIOTHERAPY • SKULL BASE CHORDOMAS - EMPHASIS ON SURGICAL STRATEGY AND RECURRENCE-FREE SURVIVAL • LAPAROSCOPIC SEALING DEVICES ON ANIMAL MODELS: SEARCHING THE LEAST HARMFUL FOR THE SURROUNDING TISSUE • PHYTOCHEMICAL PROFILE AND A PRELIMINARY *IN OVO* SCREENING OF THE ETHANOLIC EXTRACT OF *COFFEA ARABICA* GREEN SEEDS



Search

[Home](#) / Editorial Team

Editor in Chief

Radu Badea

Associate Chief Editor

Petru-Adrian Mircea

Senior Editor

Dan L. Dumitrașcu

Honorary Board Adviser

Monica Acalovschi

Associate Editors

Ioana Berindean Neagoe

Radu Nicolaie Oprean

Editorial Assistant

Monica Platon Lupșor

Editorial Office

Mîndra Eugenia Badea

Mihaela Băciuț

Cristian Bârsu

Cosmin-Nicolae Caraiani

Simona Clichici

Horațiu Colosi

Doinița Crișan

Ofelia Crișan

Diana Dudea

Sorin Marian Dudea

Daniela Fodor

Călin Lazăr

Felicia Loghin

Valentin Muntean

Andrada Seicean

Copy Editor

Ioana Robu

Website Manager

Daniel-Corneliu Leucuța

Art and GraphicDesign

Adrian Grecu

Desktop Publishing

Anne-Marie Constantin

Editorial Secretary

Dorina Sorcoi

Editorial board


Ludovico Abenavoli (Italy)

Andrei Achimaș (Romania)

Istvan Altorjay (Hungary)

Dinu Antonescu (Romania)

Adriana Băban (Romania)



Grigore Băciuț (Romania)
Marius Bojiță (Romania)
Ion I. Bruckner (Romania)
Anca Buzoianu (Romania)
Radu Câmpian (Romania)
Constantin Ciuce (Romania)
Ștefan Ciurea (TX, USA)
Douglas Drossman (NC, USA)
Dorin Farcău (Romania)
Ioan Ștefan Florian (Romania)
Shigeo Fuji (Japan)
Ion Fulga (Romania)
Alexandru Georgescu (Romania)
Liana Gheorghe (Romania)
Gabriel Ghiaur (MD, USA)
Ravnit Grewal (South Africa)
Mircea Grigorescu (Romania)
Waseem TY Hamoudy (Jordan)
Nicolae Hâncu (Romania)
Laszlo Herszenyi (Hungary)
Alexandru Irimie (Romania)
Cornel Iancu (Romania)
Vladimir Lazar (France)
Sorin Man (Romania)
Peter Manu (NY, USA)
Traian Mihăescu (Romania)
Adriana Mureșan (Romania)
Dafin Mureșanu (Romania)
Laurențiu Nedelcu (Romania)
Manuela G. Neuman (Canada)
Angelo Paradiso (Italy)
Alina Pârvu (Romania)
Virgil Păunescu (Romania)
Aurel Popa-Wagner (Germany)
Mihai Popescu (Romania)
Paul J. Porr (Romania)
Piero Portincasa (Italy)
Liren Qian (China)
Liliana Rogozea (Romania)
Vlaicu Sandor (Romania)
Adrian Săftoiu (Romania)
Florin Stamatian (Romania)
Luminița Stanciu (United Kingdom)
Ciprian Tomuleasa (Romania)
Radu Tutuian (Switzerland)




[Contact](#)



Search

[Home](#) / [Archives](#) / Vol 98 No 2 (2025)

Full Issue

-  [Table of content](#)
-  [Complete issue](#)
-  [Cover - Marcel LUPȘE](#)

Reviews

[The importance of multimodal diagnostic methods for therapeutic decision making for overactive bladder in women](#)

Mihaela Ivanov, Emil Ceban
165-175

[The role of pharmacists in ensuring rational antibiotic therapy within the interdisciplinary team](#)

Aneta Królak-Ulińska, Urszula Religioni, Beata Chełstowska, Edwin Panford-Quainoo, Zbigniew Doniec, Regis Vaillancourt, Piotr Merks
176-182

Original Research

[Colonic polyps in North-Western Romania: a comprehensive single-center analysis of endoscopic and pathological features](#)

Andrada Albu, Cosmin I. Baroana, Radu A. Farcas, Eduard Freidrich, Abdulrahman Ismaiel, Dan L. Dumitrascu, Simona Grad
183-189

[Evolution of metabolic disorders after resection of pheochromocytomas and paragangliomas: a single-center study](#)

Fatim Zahra Bentebbaa, Imane Rami, Imane Assarrar, Rania Elamel, Latifa Boutaybi, Siham Rouf, Hanane Latrech
190-195

[Return to work of head and neck cancer patients after curative radiotherapy](#)

Renata Zahu, Daniela Urian, Monica Chirila, Cristina Tiple, Magdalena Chirila, Stefan Cristian Vesa, Gabriel Kacso
196-202

[Physician migration in Romania: a study on the emigration preferences within the Cluj County healthcare system](#)

Bogdan Stancu, Stefan Lucian Popa, Abdulrahman Ismaiel, Daria Claudia Turtoi, Anca Monica Brata, Traian Adrian Duse, Cristina Pop, Maria Barsan, Alexandru Marius Padureanu, Miruna Oana Dita, Andrei Pop, Dinu Iuliu Dumitrascu, Vlad Dumitru Brata, Florin Vasile Mihaileanu, Razvan Alexandru Ciocan, Claudia Diana Gherman, Daniel Corneliu Leucuta, Aida Puia, Ion Cosmin Puia
203-209

[Skull base chordomas - emphasis on surgical strategy and recurrence-free survival](#)

[Laparoscopic sealing devices on animal models: searching the least harmful for the surrounding tissue](#)

Bogdan Petrut, Cristina Eliza Bujoreanu, Vasile Vlad Hardo, Cristian Vasile Maris, Adrian Grivei, Bogdan Fetica
217-222

[The influence of different types of bevels on the fracture resistance of directly restored lower anterior teeth](#)

Mădălina Balaj, Simona Iacob, Marius Bud, Ada Delean, Mircea Cristian Dudescu, Smaranda Buduru
223-229

[Antibiotics consumption in neurosurgery versus appendectomy: a call for antibiotic stewardship initiatives](#)

Rika Yulia, Steven Victoria Halim, Fauna Herawati, Dian Natasya Raharjo, Eko Setiawan, Heru Wijono
230-238

[Phytochemical profile and a preliminary in ovo screening of the ethanolic extract of Coffea arabica green seeds](#)

Grațiana Ruse, Ștefana Avram, Alex-Robert Jîjîe, Ana-Maria Vlase, Dalia Pătrașcu, Raluca Dumitru, George Drăghici, Alina-Arabela Jojic, Florina Ardelean, Diana-Simona Tchiakpe-Antal
239-251

Case Report

[Primary ciliary dyskinesia: a case report of double DNAH11 mutant alleles](#)

Lorin-Manuel Pîrlog, Andrada-Adelaida Pătrășcanu, Eniko Kutasi, Irina Iordănescu, Mariela Sanda Militaru
252-256

[A rare case of Wilkie`s syndrome with infantile hypertrophic pyloric stenosis](#)

Petar Stamov, Sylvia Stamova
257-261

Medicine and Pharmacy Reports

<div>COUNTRY</div> <div>Romania</div> <div><div>Universities and research institutions in Romania</div><div>Media Ranking in Romania</div></div>	<div>SUBJECT AREA AND CATEGORY</div> <div>Medicine</div> <div>Medicine (miscellaneous)</div>	<div>PUBLISHER</div> <div>Iuliu Hatieganu Medical University Press</div>	<div>SJR 2024</div> <div>0.442Q3</div> <div>H-INDEX</div> <div>33</div>
<div>PUBLICATION TYPE</div> <div>Journals</div>	<div>ISSN</div> <div>26020807, 26680572</div>	<div>COVERAGE</div> <div>2019-2025</div>	<div>INFORMATION</div> <div>Homepage</div> <div>How to publish in this journal</div> <div>info@medpharmareports.com</div>

SCOPE

MOTTO: Serving the medical community for the benefit of the society.. Medicine and Pharmacy Reports is a quarterly journal dedicated to the dissemination of progress in health sciences. The journal publishes original research, reviews, case reports, letters, and occasionally historical material, all peer reviewed. Medicine and Pharmacy Reports was founded in February 1920 under the name “Clujul Medical”, as the journal of the Medical School in Cluj, and appeared almost without interruption. Starting with 2019 it is entitled Medicine and Pharmacy Reports Medicine and Pharmacy Reports is published by one of the most prestigious medical schools in Romania and aims at representing a scientific forum in health sciences. It is open access and it promotes original papers of researchers worldwide, along with contributions of top specialists in the medical and pharmaceutical fields.

Join the conversation about this journal

FIND SIMILAR JOURNALS ?

options

1

World Journal of Clinical Cases

CHN

54%

2

Medicina (Lithuania)

CHE

52%

3

Nigerian Journal of Clinical Practice

NGA

52%

4

Medical Science Monitor

USA


50%

5

Medicine (United States)

USA

50%



Email

(will not be published)

Submit

The users of Scimago Journal & Country Rank have the possibility to dialogue through comments linked to a specific journal. The purpose is to have a forum in which general doubts about the processes of publication in the journal, experiences and other issues derived from the publication of papers are resolved. For topics on particular articles, maintain the dialogue through the usual channels with your editor.

Developed by:



Powered by:



Follow us on @ScimagoJR

Scimago Lab, Copyright 2007-2025. Data Source: Scopus®

EST MODUS IN REBUS

Horatio (Satire 1,1,106)

[Legal Notice](#)

[Privacy Policy](#)

Source details

Medicine and Pharmacy Reports

Formerly known as: Clujul Medical

Years currently covered by Scopus: from 2019 to 2025

Publisher: Iuliu Hatieganu Medical University Press


ISSN: 2602-0807 E-ISSN: 2668-0572

Subject area: Medicine: General Medicine

Source type: Journal


[View all documents >](#)

[Set document alert](#)

 [Save to source list](#)

CiteScore 2023

3.1




SJR 2023

0.373



SNIP 2023

0.777



CiteScore

CiteScore rank & trend

Scopus content coverage

CiteScore

2023

▼

3.1


=

839 Citations 2020 - 2023

267 Documents 2020 - 2023

Calculated on 05 May, 2024

CiteScoreTracker 2024



2.4

=

620 Citations to date

256 Documents to date

Last updated on 05 April, 2025 • Updated monthly

CiteScore rank 2023

Category	Rank	Percentile
Medicine		
General Medicine	#167/636	73rd

[View CiteScore methodology >](#)

[CiteScore FAQ >](#)

[Add CiteScore to your site !\[\]\(291e070cef6c4d5e78fefe4696ef53be_img.jpg\)](#)

About Scopus

What is Scopus

Content coverage

Scopus blog

Scopus API

Privacy matters

Language

日本語版を表示する

查看简体中文版本

查看繁體中文版本

Просмотр версии на русском языке

Customer Service

Help

Tutorials

Contact us

ELSEVIER

Terms and conditions ↗ Privacy policy ↗ Cookies settings

All content on this site: Copyright © 2025 Elsevier B.V. ↗, its licensors, and contributors. All rights are reserved, including those for text and data mining, AI training, and similar technologies. For all open access content, the relevant licensing terms apply.
We use cookies to help provide and enhance our service and tailor content.By continuing, you agree to the use of cookies ↗.

