



## Demographic Factors and BMI on Declined Lung Function and Vitamin D Levels in Active Smokers

Pandu Laksono<sup>1</sup>, Amelia Lorensia<sup>1</sup>✉, Heru Wijono<sup>2</sup>

<sup>1</sup>Faculty of Pharmacy, University of Surabaya (Universitas Surabaya (UBAYA)) – Indonesia

<sup>2</sup>Faculty of Medicine, University of Surabaya (Universitas Surabaya (UBAYA)) – Indonesia

### Article Info

#### Article History:

Submitted May 2024

Accepted November 2024

Published July 2025

#### Keywords:

vitamin D; lung function; BMI; smoking; demographics

#### DOI

<https://doi.org/10.15294/kemas.v21i1.4547>

### Abstract

Cigarettes are one of the causes of health problems in the world. Smoking has been known to have a direct impact on reducing lung function. Smoking increases the risk of vitamin D deficiency. Vitamin D is a vital nutrient as a lung immunomodulator. Deficient levels will cause problems with lung health, especially in active smokers. This cross-sectional research using multivariate path analysis and the SEM-PLS method has three objectives. The direct influence of three independent variables, including obesity, smoking, and demographics. Regarding vitamin D status and lung function, analyzing the direct effect of vitamin D on lung function, and thirdly, analyzing the indirect one of the three independent variables on lung function through vitamin D levels. This research took time from October 2023 to January 2024 and involved 47 active smoker respondents whose vitamin D level status and lung function were measured. There was a significant direct effect of obesity level on vitamin D levels in the active smoker population ( $p < 0.05$ ;  $f\text{-square} = 2.889$ ). While demographic factors ( $p > 0.05$ ;  $f\text{-square} = 0.030$ ) and smoking frequency ( $p > 0.05$ ;  $f\text{-square} = 0.003$ ) did not have a direct significant effect. Demographic factors, obesity, and smoking frequency don't have significant direct effect on lung function in active smokers ( $p > 0.05$ ).

### Introduction

Indonesia is one of the countries with the highest number of smokers in the world. Based on data from the 2021 Global Adult Tobacco Survey (GATS) released by the Ministry of Health of the Republic of Indonesia (KEMENKES RI), the smoking prevalence of the adult population in Indonesia reached 33.5% in 2021 with an addition of 8 million people over the last 10 years (CDC, 2021). Exposure to cigarette smoke can cause inflammation of the airways and accumulation of mucus in the lungs, resulting in symptoms of shortness of breath and accelerating the decline in lung function. It is often associated with the emergence of lung diseases such as lung cancer, chronic obstructive pulmonary disease (COPD), asthma, and tuberculosis (Chung *et al.*, 2023).

Smokers experience decreased lung function compared to non-smokers, which can be measured using spirometry through a decrease in forced expiratory volume in 1 second (FEV1) values reaching  $>50$  mL per year (Lorensia *et al.*, 2021). The higher the intensity of smoking, the lower the rate of decline in FEV1 and FVC. Greater than non-smokers.8 Further reduction in lung function due to smoking will have an impact on various lung health problems, such as COPD, including emphysema, chronic bronchitis, and asthma (Tian *et al.*, 2023).

Another mechanism that also plays a role in causing damage or decreased lung function is the involvement of levels of a vitamin in the body, namely vitamin D. Vitamin D has a protective mechanism for lung function

✉ Correspondence Address:  
Faculty of Pharmacy, University of Surabaya (Universitas Surabaya (UBAYA)) –  
Indonesia  
Email: [amelia.lorensia@gmail.com](mailto:amelia.lorensia@gmail.com)

through increasing the secretion of the antimicrobial peptide cathelicidin, decreasing chemokine production, inhibiting dendritic cell activation and changing cell activation. T. This cellular mechanism is vital for the response of the lung organ to the threat of infection and the development of allergic lung diseases such as asthma (Bishop *et al.*, 2020). A decrease in the production of the inactive form of vitamin D (25(OH)D) in lung epithelial cells is thought to be caused by exposure to smoke due to smoking activities (Lorensia *et al.*, 2024). A meta-analysis result by Yang *et al.* (2021) of 24 studies with 11,340 participants showed that levels of vitamin D in the inactive form 25(OH)D were lower in smokers than non-smokers (Yang *et al.*, 2021). In addition, the expression level of vitamin D receptors can also be influenced by exposure to cigarette smoke (Ahn *et al.*, 2021). A study by Ghosh *et al.* (2020) also stated that vitamin D deficiency plays a role in changes in lung structure and decreased lung function.

There are other influencing factors that can be part of the impact of decreased lung function besides smoking, namely, obesity. Individuals who are obese show reduced lung volume and capacity when compared to people of normal weight. Larger fat deposits in the abdominal area produce greater resistance to diaphragm contraction, thereby inhibiting respiratory ventilation mechanisms (Cao *et al.*, 2022). Based on the explanation above, several previous studies have been conducted that examined the effect of vitamin D on lung function. However, there have been no studies that have examined this by involving factors such as obesity levels, smoking, and demographics in Indonesia, which is one of the countries with the highest prevalence of smoking, accompanied by consequences in the form of increased death rates and lung disease sufferers due to smoking. Therefore, research will be carried out that will examine the influence of demographic factors, BMI, and smoking intensity, on the decline in lung function and vitamin D levels in active smokers. There is a study by Abi-Ayad *et al.* (2023) who measured smokers' vitamin D levels using blood plasma samples, found that vitamin D deficiency was associated with lower lung function conditions, lung function in this case FEV1, FVC, and FEV1/FVC measured

using a spirometer experienced faster decrease in smoking subjects. It shows that adequate serum vitamin D levels are associated with a protective effect against the detrimental effects of smoking on lung function.

## Method

This study used an observational clinical trial with a cross-sectional design where the data collection stage was carried out once at a time. The independent variables in this study were demographic factors, BMI, and smoking intensity. The dependent variable in this study was the lung function value (percentage of FEV1/FVC ratio). The mediating variable in this study was vitamin D levels. Demographic factors consist of age and education level. Age is a measure of the patient's length of life, which is calculated based on the patient's date of birth until they become a research respondent. Educational level is a measure of the respondent's level of education as evidenced by possession of the latest educational certificate. Body Mass Index, or BMI, is a value obtained from the mass and height of each sample individual. BMI can be calculated using a formula by dividing the individual's body weight in kilograms (kg) by the square of body height in meters squared ( $m^2$ ) (Weir & Jan, 2023). Smoking intensity is the habit of smoking tobacco cigarettes. Smoker classification can be calculated using the Brinkman Index (IB)=number of cigarettes smoked per day x length of smoking (years) (Herath *et al.*, 2022). Lung function measurements using a handheld spirometer. The level of lung function impairment based on the percentage of the FEV1/FVC ratio is divided into 4 categories, namely normal-mild obstruction, moderate obstruction, severe obstruction, and very severe obstruction (Stanojevic, 2021).

The vitamin D levels that will be measured are the most abundant metabolite form in serum, namely 25(OH)D, 25(OH) levels reflect skin production of vitamin D3 and vitamin D (D2 and D3) from food, 25(OH) D has a half-life (The long  $t_{1/2}$ ) in the blood circulation is 3-4 weeks compared to the active vitamin D metabolite 1,25(OH) $_2$ D which only has a short  $t_{1/2}$  of around 4-6 hours (Tuckey *et al.*, 2019). Methods for observing vitamin

D levels (25(OH)D), which are the ELFA (Enzyme Linked Fluorescent Assay) method used is the VIDAS tool from bioMerieux. The type of specimen that can be used in testing can be serum or blood plasma. In this study, serum 25(OH)D levels were classified as deficient (<20 ng/mL), insufficiency (20-29 ng/mL), and normal (30-100 ng/mL) (Ahn *et al.*, 2021; Amrein *et al.*, 2020).

The population is active smokers located in the Mejoyo area, Rungkut District, Surabaya City. The accessible population is active smoking respondents in the Mejoyo 2 RT 6 Surabaya area who have filled out the questionnaire, can be found, and are not included in the exclusion criteria. The sample in this study was part of the affordable population who met the criterias: (1) aged 18-60 years; (2) didn't use vape; (3) no history of COVID-19 infection; (4) no history of diseases or conditions that can affect serum vitamin D levels; (5) not taking supplements containing vitamin D during the last 1 month before checking 25(OH)D levels; and (6) no history of diseases or conditions that can affect lung function. The minimum number of samples was calculated using the Slovin method. So the number of samples required in this research was 45 people. The sampling technique was purposive sampling.

The first meeting involved a recording of demographic data for the entire sample through a direct interview process with research respondents, and then continued with measuring height and weight. The researcher measured the height of the sample using a height

measuring device with the Onemed brand type HT701 Wireless. Body weight measurements were carried out by researchers using a SPEEDS brand digital weight scale, type LX040-8 USB, which has a sensitivity of one digit after the comma. Lung function measurements were carried out by researchers who had received special training from clinicians using a Contec SP10 handheld spirometer to obtain FEV1 and FVC values. The lung function measurement procedure for each respondent was repeated 3 times. The results in milliliters (mL) are recorded to calculate the percentage of the FEV1/FVC ratio. Measurement of Vitamin D or 25(OH)D levels using the VIDAS® 25 OH Vitamin D TOTAL (VITD) tool with the ELFA technique carried out by a standardized laboratory in Surabaya.

The aim of carrying out multivariate analysis is to determine the magnitude of the influence between variables determined based on the  $P_{\text{value}}$  and  $t_{\text{statistic}}$  value using the Structural Equation Modeling with Partial Least Squares (SEM-PLS) method with the help of the SmartPLS application. SEM-PLS is a powerful analysis method and is often referred to as soft modeling, because it eliminates the assumptions of Ordinary Least Square (OLS) regression such as data must be normally distributed in a multivariate manner and there is no problem of multicollinearity between independent variables, SEM-PLS can be used to testing weak theories and weak data (small samples and data normality problems). The analysis of the magnitude




Figure 1. Path Analysis

#### Information :

- i. Direct influence of obesity level on lung function
- ii. Direct influence of obesity level on vitamin D levels
- iii. Direct effect of smoking on vitamin D levels
- iv. Direct effect of smoking on lung function
- v. Direct influence of demographics on vitamin D levels
- vi. Direct influence of demographics on lung function
- vii. Direct influence of vitamin D levels on lung function

of the effect in this research will estimate the value of the  $P_{\text{value}}$  and also the value of the  $t_{\text{statistic}}$ . If the test results show a  $P_{\text{value}} < 0.05$  and a  $t_{\text{statistic}} \geq t_{\text{table}}$  value, then the influence of the two variables is said to be significant. It comes along with the conclusion that there is an influence of the independent variable on the dependent variable of active smoking. However, if the  $P_{\text{value}} > 0.05$  and the  $t_{\text{statistic}} < t_{\text{table}}$  value, then the influence of the two variables is said to be insignificant, with the conclusion that there is no influence of the independent variable on the dependent variable in active smokers. An overview of the path analysis model used in the research can be seen in **Figure 1**.

## Result and Discussion

This research was conducted from October 2023 to March 2024, located in the Mejoyo 2 RT 6 Surabaya, Kalirungkut, East Surabaya. This research has received an ethical certificate number 232/KE/IX/2023 from the University of Surabaya. Based on the preliminary study, the affordable population was 54 smokers in the area, and 47 people were willing and met the research subject criteria. Most of the respondents were the largest early elderly, 23 respondents (48.93%). Based on BMI measurements, the normal BMI group was the largest, namely 29 respondents (61.70%).

Based on the measurement of education level, the Secondary Education group was the largest, namely 20 respondents (42.55%). Based on smoking frequency, the moderate smoker group was the largest, namely 27 respondents (57.44%). Data on the distribution of respondents' characteristics can be seen in Table 1.

In this study, vitamin D (25(OH)D) levels were measured in 47 respondents using the ELFA method with the VIDAS® tool from Biomerieux. Based on the classification of vitamin D levels of the total respondents measured, the insufficiency group was the largest group with 23 respondents (48.93%). The results of the percentage value of the FEV1/FVC ratio, of the total respondents who were measured, the moderate obstruction group had the largest, namely 21 respondents (44.68%) (**Table 2**).

The implementation of multivariate statistical analysis in this research was carried out using a quantitative technical approach, where there were two analyzes used, namely descriptive analysis and hypothesis testing or statistical analysis using the Partial Least Square (PLS) method using the SmartPLS program. There are 5 (five) variables involved in this research, which include obesity levels, smoking, demographics, vitamin D, and lung function.

Table 1. Demographic Profile of Respondents

		Frequency (n=47)	Percentage (%)	$\bar{x} \pm SD$
Age (years)	Late Adulthood (36-45)	10	21.27	51.20±6.43
	Early Seniors (46-55)	23	48.93	
	Late Seniors (56-60)	14	29.78	
Level of education	Basic Education (elementary school-junior high school)	19	40.42	
	Secondary Education (senior high school)	20	42.55	
	higher education	8	17.02	
BMI (kg/m <sup>2</sup> )	Normal (18.5-25.0)	29	61.70	
	Fat (Overweight) (25.1-27.0)	14	29.78	
	Obesity (>27)	4	8.51	
Brinkman Index (BI)	Light (<200)	14	29.78	
	Moderate (200-599)	27	57.44	
	High (>600)	6	12.76	

$\bar{x}$  = average; SD= standard deviation

Source: Primary Data, 2024

Table 2. Distribution of Vitamin D and Lung Function Examination

		Frequency (n=47)	Percentage (%)	$\bar{x} \pm SD$
Vitamin D Classification -25(OH) D Content (ng/mL)	Deficiency (<20)	15	31.91	23.52±8.75
	Insufficiency (20-29)	23	48.93	
	Normal (30-100)	9	19.14	
Lung Function Classification (FEV1/ FVC ratio) (%)	Normal-Mild Obstruction (≥ 80)	7	14.89	95.43±5.72
	Moderate Obstruction (50- 79)	21	44.68	
	Severe Obstruction (30-49)	10	21.27	
	Very Severe Obstruction (<30)	9	19.14	

$\bar{x}$  = average; SD= standard deviation

Source: Primary Data, 2024

The research model evaluation stage in PLS consists of measuring model evaluation and structural model evaluation. The measurement model in this research consists of a reflective measurement model for the variables Levels of Obesity, Smoking, Demography, Vitamin D, and Lung Function by examining loading factor values  $\geq 0.50$ , composite reliability  $\geq 0.60$ , and average variance extracted (AVE)  $\geq 0.50$ . The level of obesity is measured by BMI (X1.2), where the outer loading value is between 0.688-0.936, which shows that BMI is strongly correlated in explaining the obesity level variable. Every change in BMI on obesity levels has a more significant effect. The smoking variable is measured from the Brinkman index (IB) of smoking frequency (X2.1), where the outer loading value is between 0.243-0.996, which shows that IB is correlated in explaining the smoking variable. The level of reliability of the motivation variable is well accepted, with a composite reliability value (0.618)  $> 0.60$ . Any change in smoking frequency on smoking activity has a more significant effect.

Demographic variables are measured by 2 (two) indicators, namely age (X3.1) and education level (X3.2), where the outer loading value is between 0.243-0.996, which shows that these two indicators correlate in explaining the obesity level variable. The level of reliability of the motivation variable is acceptable, even though the composite reliability value (0.406) is  $< 0.60$ , but has an AVE value close to 5.0.

Between the two measurement indicators, the validity of demographic variables appears to be more strongly reflected by the education level indicator (X3.3). It means that every change in age demographics has a more significant effect. The vitamin D variable is measured directly with 1 (one) indicator, namely the 25(OH)D level, where the outer loading value is 1,000, which shows that this indicator has a strong correlation in explaining the vitamin D variable. The level of reliability of the vitamin D variable is acceptable, with the composite reliability value (1,000)  $> 0.60$ . The validity measurement of the vitamin D variable looks strong, as reflected by the 25(OH)D level indicator (Y). The lung function variable is measured directly with 1 (one) indicator, namely the  $P_{\text{value}}$  of the percentage of the FEV1/FVC ratio, where the outer loading value is 1,000, which shows that this indicator has a strong correlation in explaining the lung function variable. The level of reliability of the vitamin D variable is acceptable, with a composite reliability value (1,000)  $> 0.60$ . The validity measurement of lung function variables appears to be strong, as reflected by the percentage indicator of the FEV1/FVC ratio (Z).

Structural model evaluation is a form of evaluation to determine the influence of various endogenous variables on exogenous ones by observing the R-square ( $R^2$ ) value (coefficient determination), if the  $R^2$  value = 0.19 then the exogenous influence on the endogenous is weak,



Table 3. Evaluation Results of Reflective Measurement Model, Inner VIF Measurement

Variables	Measurement Item Code	Indicators	<i>O u t e r Loading</i>	<i>Composite Reliability</i>	AVE	VIF
Obesity Rate	X1.2	BMI	0.936	0.802	0.675	1.178
Cigarette Frequency	X2.1	Smoking Frequency	0.996	0.618	0.525	1.024
Demographics	X3.1	Age	0.137	0.406	0.485	1.007
	X3.2	Level of education	0.975			1.007
Vitamin D	Y	25(OH)D levels	1.000	1.000	1.000	1.000
Lung Function	Z	FEV1/FVC ratio	1.000	1.000	1.000	1.000

AVE=average variance extracted; BMI=Body Mass Index; FEV<sub>1</sub>=Forced Expiratory Volume in 1 second; VIF = Variance Inflated Factor

Source: Primary Data, 2024

if the  $R^2$  value=0.33 the exogenous influence on the endogenous is moderate, if the  $R^2$  value=0.67 the exogenous influence on the endogenous is strong. Next, hypothesis testing is carried out to find out the significance of the influence of the observed research variables by observing the path coefficient value. The influence of the level of obesity, smoking, and demographics on vitamin D, with an  $R^2$  value of 0.782, means that the endogenous variable vitamin D is influenced by 78.2% by the exogenous variables the level of obesity, smoking, and demographics, while 21.8% is influenced by other factors outside the variable. It can be concluded that the influence of exogenous variables on endogenous variables is strong. The influence of the level of obesity, smoking, demographics, vitamin D on lung function with an  $R^2$  value of 0.190, meaning that the endogenous variable lung function is influenced by 19% by the exogenous variable the level of obesity, smoking, demographics, and vitamin D, while 81% is influenced by other factors in outside the variables studied. It can be concluded that the influence of exogenous variables on endogenous variables is weak.

Next, the evaluation of the structural model with path coefficients was carried out in three stages: The first stage, namely checking the absence of multicollinearity between variables and the inner VIF (Variance Inflated Factor). If the estimation results show an inner VIF value <5, then the level of multicollinearity between variables is low. All indicators have an inner VI value <5, so the estimates of all variables and indicators in SEM-PLS are not robust (not biased). The second stage, hypothesis testing, is carried out between variables by looking at the

$t_{\text{statistic}}$  value and  $P_{\text{value}}$ . If the  $t_{\text{statistic}}$  = calculation result is greater than the  $t_{\text{table}}$  (2.0166) and the  $P_{\text{value}}$  of the test results is <0.05, then there is a significant influence between the variables. The third stage, analysis of the results of the f-square value is carried out, namely the influence of variables at the structural level with the criteria f-square  $0.02 \leq$  no effect,  $0.02 \leq$  f-square  $\leq 0.14$  small effect,  $0.15 \leq$  f-square  $\leq 0.35$  has a medium effect, and  $>0.35$  has a high effect. The patch coefficient assesses the magnitude of the direct influence of exogenous variables on endogenous variables; the magnitude of the influence ranges from -1 (negative influence) to +1 (positive influence).


Hypothesis testing is not only direct observation between exogenous variables and endogenous variables, but also tested indirectly through mediating variables (vitamin D). The SEM-PLS model can also be useful as a predictor in developing strategies to improve health promotion related to the influence of obesity, smoking, and demographic factors on lung function and vitamin D levels directly, as well as on lung function indirectly through vitamin D levels in active smokers. Education provided in terms of increasing vitamin D (25(OH)D) levels is by reducing body weight through BMI indicators and reducing the frequency of smoking. It also applies to implementing education to improve lung function by improving or increasing vitamin D levels (Figure 2).

Body mass index (BMI) is the ratio of body weight to height squared. Most of the respondents had normal BMI values, namely 29 respondents (61.70%) and safe limit abdominal

Table 4. Structural Model Evaluation Results, consisting of Research Hypothesis Testing and Effect Size (f-square)

	Hypothesis	<i>Path coefficient</i>	$t_{\text{statistic}}$	$P_{\text{value}}$	<i>f-square</i>	Interpretations
Direct Hypothesis	Is there a direct effect of obesity level on vitamin D in active smokers?	0.908	9.497	0.000	2.889	There is a direct influence of the level of obesity on vitamin D levels in active smokers. The significant influence of the level of obesity on vitamin D has a high influence value.
	Is there a direct effect of the level of obesity on lung function in active smokers?	-0.090	1.076	0.283	0.030	There is no direct effect of the level of obesity on lung function in active smokers. The obesity level variable on vitamin D has a moderate influence value.
	Is there a direct effect of smoking on vitamin D in active smokers?	0.030	0.265	0.791	0.003	There is no direct effect of smoking on vitamin D in active smokers. The cigarette variable on vitamin D has a small influence value.
	Is there an effect of smoking on lung function in active smokers?	0.815	0.512	0.609	0.161	There is no direct effect of smoking on lung function in active smokers. The smoking variable on lung function has a small influence value.
	Is there a direct influence of demographics on vitamin D in active smokers?	-0.319	1.229	0.220	0.098	There is no direct influence of demographics on vitamin D in active smokers. Demographic variables on vitamin D have no influence.
	Is there a direct influence of demographics on lung function in active smokers?	0.122	0.654	0.513	0.016	There is no direct influence of demographics on lung function in active smokers. Demographic variables on lung function have no influence value.
	Is there a direct effect of vitamin D on lung function in active smokers?	-0.791	0.500	0.617	0.168	There is no direct effect of vitamin D on lung function in active smokers. The vitamin D variable on lung function has a moderate influence value.
Indirect Hypothesis	Is there an effect of obesity level on lung function through vitamin D levels in active smokers?	-	0.256	0.798	-	There is no influence of obesity level on lung function through vitamin D levels in active smokers. Vitamin D does not mediate the indirect relationship between obesity level and lung function.
	Is there an effect of smoking on lung function through vitamin D levels in active smokers?	-	1.087	0.277	-	There is no effect of smoking on lung function through vitamin D levels in active smokers. Vitamin D does not mediate the indirect relationship between smoking and lung function.
	Is there a demographic influence on lung function through vitamin D levels in active smokers?	-	0.454	0.650	-	There is no demographic influence on lung function through vitamin D levels in active smokers. Vitamin D did not mediate the indirect relationship between demographic variables and lung function.

Source: Primary Data, 2024



Source: Primary Data, 2024

Figure 2. SEM-PLS Model After Bootstrapping with  $P_{\text{value}}$  = between Variables Based on SmartPLS analysis

circumference of 38 respondents (80.85%). It is supported by the results of research in Indonesia which analyzed the BMI picture and found that the majority of respondents had a normal BMI of 45%. 75 Another study in Indonesia which aimed to find out the factors related to the incidence of central obesity in adults found that there was a prevalence Central obesity with excessive abdominal circumference based on age 25-34 years (22.9%) and 35-44 years (33.5%), indicates that there are more respondents with abdominal circumference within safe limits. The results of this study found 4 respondents (8.51%) in the peripheral obesity category ( $\text{BMI} > 27 \text{ kg/m}^2$ ) and 9 people (19.14%) with central obesity (abdominal circumference  $> 90 \text{ cm}$ ). Based on the evaluation of the structural model (inner model) using SmartPLS version 3, there was a significant direct influence between the obesity level variable on vitamin D levels ( $P_{\text{value}} < 0.05$  and  $t_{\text{statistic}} > t_{\text{table}}$ ) in active smokers. It is supported by a theory that states the relationship between obesity mechanisms in causing a decrease in vitamin D levels, including three mechanisms that can explain the relationship between deficiency in vitamin D levels in obese individuals. Individuals who are obese experience decreased exposure to sunlight compared to non-obese individuals (Mirza *et al.*, 2022). The release of adiponectin from fatty tissues was inversely correlated with

body weight and BMI suggesting a link between vitamin D deficiency and insulin resistance (Kausar *et al.*, 2022).

The evaluation of the control model (inner model) found no direct significant influence between the obesity level variable on lung function ( $P_{\text{value}} > 0.05$  and  $t_{\text{statistic}} < t_{\text{table}}$ ) in active smokers. These results have conclusions that are the opposite of several theories which state that obesity harms the lung organs, one of which is the development of OHS which is defined as a combination of obesity ( $\text{BMI} \geq 30 \text{ kg/m}^2$ ), hypercapnia (arterial  $\text{CO}_2 \geq 45 \text{ mmHg}$ ) and the presence of breathing disorders during sleep which causes alveolar hypoventilation (Masa *et al.*, 2019; Palma *et al.*, 2022). Research results that are not in accordance with theory or previous research can be caused by only a small portion of respondents being classified as obese, whereas based on lung function examinations also the majority of respondents (44.68%) included in the moderate obstruction category. Individuals who are obese show a decrease in lung volume and capacity when compared to people with a normal body weight, this is because in people with a normal BMI the diaphragm contracts to push the contents of the stomach down and forward without any obstacles from excess fat deposits, when Likewise, contraction of the external intercostal muscles (muscles that play a role in



the inspiration process) can push the ribs up and forward without resistance (Svartengren *et al.*, 2020). Therefore, in this study, the majority of respondents had a normal BMI and the majority were in the obstruction category. moderate (27.65%), because the majority of respondents had breathing patterns that were not influenced by obesity.

Most of the smokers were moderate smokers (57.44%). Evaluation of the structural model (inner model) found that there was no direct significant influence between smoking variables on vitamin D levels ( $P_{\text{value}} > 0.05$  and  $t_{\text{statistic}} < t_{\text{table}}$ ) in active smokers. Smoking, ultraviolet radiation, and age are considered important factors that contribute to the skin aging process in humans by increasing wrinkles on the skin surface. In addition, tobacco smoke can affect the expression level of vitamin D receptors. If the expression of vitamin D receptors increases, the physiological effects of vitamin D will be better, one of which is the effect on lung function (Yang *et al.*, 2021). Most of the respondents fell into the moderate level with IB. Based on the evaluation of the inner control model using SmartPLS, it was found that there was no direct significant influence between smoking variables on lung function ( $P_{\text{value}} > 0.05$  and  $t_{\text{statistic}} < t_{\text{table}}$ ) in active smokers. These results have conclusions that are the opposite of several theories, which state that cigarettes contain various dangerous substances and their pathophysiological mechanisms, which can disrupt lung function. Nicotine was found to be chemotactic for human neutrophils, neutrophils being the first cells recruited in the process of lung inflammation due to cigarette smoke, which ultimately causes airway obstruction (Ham *et al.*, 2022). Cigarettes cause airway inflammation, which occurs more precisely in the bronchioles, causing lung remodeling. - The lungs lose their elasticity during the air exchange process, which then results in chronic obstructive airway limitations (Karnati *et al.*, 2021). Inappropriate research results can be caused by other factors or variables that are not observed but can influence, among others, genetics, physical activity, food intake, knowledge, attitudes, and behavior, sun exposure, skin color, and air pollution.

Patient demographics, which include age and level of education, are one of the factors that have been widely studied and influence the reduction in vitamin D levels. Based on the evaluation of the structural model (inner model), the results found that there is no direct significant influence between demographic variables on vitamin D levels ( $P_{\text{value}} > 0.05$  and  $t_{\text{statistic}} < t_{\text{table}}$ ) in active smokers. These results are aligned with the theory regarding demographic relationships, whether studied in terms of age or education level, which can influence vitamin D levels. In terms of age, physiologically an individual can experience a decrease in kidney function, resulting in reducing the production of the active metabolite 1,25(OH)D with increasing age, due to a decline in the activity of the kidney enzyme 1 $\alpha$ -hydroxylase which converts 25(OH)D to 1,25(OH)D. Serum 1,25(OH)D levels are inversely proportional to serum creatinine levels and proportional to glomerular function rate (GFR). Aging is not only associated with a decrease in kidney function, but is also associated with a decrease in vitamin D production in the skin, in the form of a decline in the concentration of 7-dehydrocholesterol in the epidermis and a reduced response to UVB light exposure in elderly individuals compared to young adults, resulting in a decrease in the formation of previtamin D3 by 50% (Turner *et al.*, 2022; Huish *et al.*, 2021). The higher the level of education, the greater the knowledge and awareness of the importance of adequate vitamin D nutrition (Hamhoum & Aljefree, 2022). Therefore, this is a limitation of this research because there are several factors that were not observed but tend to influence vitamin D levels, including the influence of physical activity, knowledge and attitudes towards vitamin D, and genetics.

Age and education level are also considered factors that influence lung function. Based on the evaluation of the inner control model using SmartPLS, there was no direct significant influence between demographic variables on lung function ( $P_{\text{value}} > 0.05$  and  $t_{\text{statistic}} < t_{\text{table}}$ ) in active smokers. These results have conclusions that are the opposite of several theories, which state that demographics, both in terms of age and level of education, can influence lung function, especially in active

smokers. Other research results show that the older a person is, the lower the FEV1 value will be. In particular, the smaller the FVC value due to advanced age, the more significant the decrease in FEV1/FVC (Thomas *et al.*, 2019). It is different from the results of this study, where there was no influence of demographics on lung function. This result can be caused by factors that influence these two variables (demography and lung function), each of these two variables can be influenced by smoking factors and the level of obesity studied in this study to factors that were not examined in this study but can contribute to changes in lung function such as physical activity, knowledge, attitudes and behavior, etc. as previously explained.

The largest number of respondents belonged to the vitamin D insufficiency and moderate obstruction group (25.53%). Based on the evaluation of the inner control model, it was found that there was no direct significant influence of the vitamin D variable on lung function ( $P_{\text{value}} > 0.05$  and  $t_{\text{statistic}} < t_{\text{table}}$ ) in active smokers. These results have conclusions that are the opposite of several theories, which state that vitamin D can affect lung function, especially in active smokers. It is not aligned with previous theory and research, which states that there is an inverse relationship between vitamin D levels and lung function. Previous research by Ganji *et al.* (2020) involved 11,983 respondents aged  $\geq 20$  years. This study combines three NHANES data from 2007–2008, 2009–2010, and 2011–2012. The results of the study concluded that serum 25(OH)D levels were directly related to FVC and FEV1; in other words, serum 25(OH)D levels were associated with improved lung function values in healthy people, but not with the prevalence of asthma, emphysema, and chronic bronchitis. Other research also supports previous research, such as Wannamethee *et al.* (2021), who used a prospective cohort design involving 3575 male respondents (60–79 years), concluded that male respondents with COPD tend to experience vitamin D deficiency when compared to male respondents with normal lung function. In contrast to the results of this study, there was no effect of vitamin D on lung function. This result can be caused by factors that influence these two variables (vitamin D and lung function),

each of these two variables can be influenced by obesity, smoking and demographic factors studied in this study to factors that were not studied such as genetics, physical activity, food intake, knowledge, attitudes and behavior, sun exposure, skin color, air pollution.

The results of the evaluation of the structural model (inner model) found that there was no influence of the level of obesity on lung function through vitamin D levels in active smokers. Therefore, vitamin D does not mediate the indirect relationship between obesity levels and lung function. This result could be caused by factors that influence these two variables (vitamin D and lung function), each of these three variables can be influenced by smoking and demographic factors studied in this study to factors not studied such as genetics, physical activity, food intake, knowledge, attitudes and behavior, sun exposure, skin color, air pollution and so on, as previously explained.

This research still has several limitations, including other factors that cannot be controlled such as genetic factors, physical activity, knowledge, attitudes and behavior, intensity of exposure to sunlight, use of sunscreen, air pollution, food intake cannot be controlled in this research so that it can influence the research results. Current knowledge finds that a lack of physical activity is an important risk factor for vitamin D deficiency. Various observational studies show that maintaining vitamin D nutritional status is related to physical activity/exercise habits, where the level of physical activity is significantly positively correlated with 25(OH)D levels (Song *et al.*, 2020; Lorensia *et al.*, 2022). Variations in sun exposure can cause vitamin D levels to fluctuate. Indoor activity may be a factor in these fluctuations and may cause a decrease in vitamin D levels. A systematic review and meta-analysis aimed at identifying whether physical activity or exercise indoors compared to outdoors has a significant effect on vitamin D levels through subgroup analysis and multivariate meta-regression (Bârsan *et al.*, 2023).

## Conclusion

Obesity level factors (BMI) have a significant direct effect on vitamin D levels in active smokers. The demographic factors and smoking frequency do

not have a direct significant effect on vitamin D in active smokers. Demographic factors, obesity level, and smoking frequency do not have a significant direct effect on lung function in active smokers. Vitamin D has no direct significant effect on lung function in active smokers. Demographic factors, obesity level, and smoking frequency do not have a significant effect on lung function via vitamin D in active smokers. Therefore, it is recommended to maintain body weight and lose weight, especially in obese populations, to reduce the incidence of vitamin D deficiency. So it is necessary to carry out further research on the influence of other factors on vitamin D levels and lung function, such as genetic factors, physical activity, knowledge and attitudes, and behavior, intensity of exposure to sunlight, use of sunscreen, air pollution, and food intake that were not observed in this study to develop theories about the influence of broader factors.

### Acknowledgment

This research was funded by the Research and Community Service Institute (LPPM) of Universitas Surabaya.

### References

- Abi-Ayad, M., Nedjar, I., & Chabni, N., 2023. Association between 25-Hydroxy Vitamin D and Lung Function (FEV1, FVC, FEV1/FVC) in Children and Adults with Asthma: A Systematic Review. *Lung India*, 40(5), pp.449–56.
- Ahn, K.M., Kim, S.S., Lee, S.Y., Lee, S.H., & Park, H.W., 2021. Vitamin D Deficiency and Lung Function Decline in Healthy Individuals: A Large Longitudinal Observation Study. *Respir Med.*, 182, pp.106395.
- Amrein, K., Scherkl, M., Hoffmann, M., Neuwersch-Sommeregger, S., Köstenberger, M., Tmava, B.A., Martucci, G., Pilz, S., & Malle, O., 2020. Vitamin D Deficiency 2.0: An Update on the Current Status Worldwide. *Eur J Clin Nutr.*, 74(11), pp.1498–513.
- Bârsan, M., Chelaru, V.F., Râjnoveanu, A.G., Popa, Ș.L., Socaciu, A.I., & Bădulescu, A.V., 2023. Difference in Levels of Vitamin D between Indoor and Outdoor Athletes: A Systematic Review and Meta-Analysis. *Int J Mol Sci.*, 24(8), pp.7584.
- Bishop, E.L., Ismailova, A., Dimeloe, S., Hewison, M., & White, J.H., 2020. Vitamin D and Immune Regulation: Antibacterial, Antiviral, Anti-Inflammatory. *JBMR Plus*, 5(1), pp.e10405.
- Cao, Y., Li, P., Wang, Y., Liu, X., & Wu, W., 2022. Diaphragm Dysfunction and Rehabilitation Strategy in Patients with Chronic Obstructive Pulmonary Disease. *Front Physiol.*, 13, pp.872277.
- CDC., 2021. GATS (Global Adult Tobacco Survey) Comparison Fact Sheet Indonesia 2011 & 2021. *Global Adult Tobacco Survey*, pp.1–2.
- Chung, C., Lee, K.N., Han, K., Shin, D.W., & Lee, S.W., 2023. Effect of Smoking on the Development of Chronic Obstructive Pulmonary Disease in Young Individuals: A Nationwide Cohort Study. *Front Med (Lausanne)*, 10, pp.1190885.
- Ganji, V., Al-Obahi, A., Yusuf, S., Dookhy, Z., & Shi, Z., 2020. Serum Vitamin D is Associated with Improved Lung Function Markers but not with Prevalence of Asthma, Emphysema, and Chronic Bronchitis. *Sci Rep.*, 10(1), pp.11542.
- Ghosh, A.J., Moll, M., Hayden, L.P., Bon, J., Regan, E., & Hersh, C.P., 2020. Vitamin D Deficiency is Associated with Respiratory Symptoms and Airway Wall Thickening in Smokers with and without COPD: A Prospective Cohort Study. *BMC Pulm Med.*, 20(1), pp.123.
- Ham, J., Kim, J., Ko, Y.G., & Kim, H.Y., 2022. The Dynamic Contribution of Neutrophils in the Chronic Respiratory Diseases. *Allergy Asthma Immunol Res.*, 14(4), pp.361–78.
- Hamhoum, A.S., & Aljefree, N.M., 2022. Knowledge and Attitudes towards Vitamin D among Health Educators in Public Schools in Jeddah, Saudi Arabia: A Cross-Sectional Study. *Healthcare (Basel)*, 10(12), pp.2358.
- Herath, P., Wimalasekera, S., Amarasekara, T., Fernando, M., & Turale, S., 2022. Effect of Cigarette Smoking on Smoking Biomarkers, Blood Pressure and Blood Lipid Levels Among Sri Lankan Male Smokers. *Postgrad Med J.*, 98(1165), pp.848–54.
- Huish, S.A., Jenkinson, C., Dunn, J.A., Meredith, D.J., Bland, R., & Hewison, M., 2021. Low Serum 1,25(OH)2D3 in End-Stage Renal Disease: is Reduced 1α-Hydroxylase the Only Problem?. *Endocr Connect*, 10(10), pp.1291–8.
- Karnati, S., Seimetz, M., Kleefeldt, F., Sonawane, A., Madhusudhan, T., Bachhuka, A., Kosanovic, D., Weissmann, N., Krüger, K., & Ergün, S., 2021. Chronic Obstructive Pulmonary Disease and the Cardiovascular System: Vascular Repair and Regeneration as a Therapeutic Target. *Front Cardiovasc Med.*, 8, pp.649512.
- Kauser, H., Palakeel, J.J., Ali, M., Chaduvula, P., Chhabra, S., Lamsal, L.S., Ramesh, V., Opara, C.O., Khan, F.Y., Kabiraj, G., & Mohammed, L., 2022. Factors Showing the Growing

- Relation Between Vitamin D, Metabolic Syndrome, and Obesity in the Adult Population: A Systematic Review. *Cureus*, 14(7), pp.e27335.
- Lorensia, A., Muntu, C.M., Suryadinata, R.V., & Septiani, R., 2021. Effect of Lung Function Disorders and Physical Activity on Smoking and Non-Smoking Students. *J Prev Med Hyg.*, 62(1), pp.E89–E96.
- Lorensia, A., Suryadinata, R.V., & Inu, I.A., 2022. Comparison of Vitamin D Status And Physical Activity Related With Obesity in Student. *Journal of Applied Pharmaceutical Science*, 12(4), pp.108-18.
- Lorensia, A., Suryadinata, R.V., Rahmawati, R.K., & Septiani, R., 2024. The Effect of Smoking Habit on Vitamin D Status in Adults in Indonesia. *KEMAS*, 19(3), pp.410-421.
- Masa, J.F., Pépin, J.L., Borel, J.C., Mokhlesi, B., Murphy, P.B., & Sánchez-Quiroga, M.Á., 2019. Obesity Hypoventilation Syndrome. *Eur Respir Rev.*, 28(151), pp.180097.
- Mirza, I., Mohamed, A., Deen, H., Balaji, S., Elsabbahi, D., Munasser, A., Naquiallah, D., Abdulbaseer, U., Hassan, C., Masrur, M., Bianco, F.M., Ali, M.M., & Mahmoud, A.M., 2022. Obesity-Associated Vitamin D Deficiency Correlates with Adipose Tissue DNA Hypomethylation, Inflammation, and Vascular Dysfunction. *Int J Mol Sci.*, 23(22), pp.14377.
- Palma, G., Sorice, G.P., Genchi, V.A., Giordano, F., Caccioppoli, C., D'Oria, R., Marrano, N., Biondi, G., Giorgino, F., & Perrini, S., 2022. Adipose Tissue Inflammation and Pulmonary Dysfunction in Obesity. *Int J Mol Sci.*, 23(13), pp.7349.
- Song, K., Park, G., Choi, Y., Oh, J.S., Choi, H.S., Suh, J., Kwon, A., Kim, H.S., & Chae, H.W., 2020. Association of Vitamin D Status and Physical Activity with Lipid Profile in Korean Children and Adolescents: A Population-Based Study. *Children (Basel)*, 7(11), pp.241.
- Stanojevic, S., Kaminsky, D.A., Miller, M.R., Thompson, B., Aliverti, A., Barjaktarevic, I., Cooper, B.G., Culver, B., Derom, E., Hall, G.L., Hallstrand, T.S., Leuppi, J.D., MacIntyre, N., McCormack, M., Rosenfeld, M., Swenson, E.R., 2022. ERS/ATS Technical Standard on Interpretive Strategies for Routine Lung Function Tests. *Eur Respir J.*, 60(1), pp.2101499.
- Svartengren, M., Cai, G.H., Malinovschi, A., Theorell-Haglöw, J., Janson, C., Elmståhl, S., Lind, L., Lampa, E., & Lindberg, E., 2020. The Impact of Body Mass Index, Central Obesity and Physical Activity on Lung Function: Results of the EpiHealth Study. *ERJ Open Res.*, 6(4), pp.00214–2020.
- Thomas, E.T., Guppy, M., Straus, S.E., Bell, K.J.L., & Glasziou, P., 2019. Rate of Normal Lung Function Decline in Ageing Adults: A Systematic Review of Prospective Cohort Studies. *BMJ Open*, 9(6), pp.e028150.
- Tian, T., Jiang, X., Qin, R., Ding, Y., Yu, C., Xu, X., & Song, C., 2023. Effect of Smoking on Lung Function Decline in a Retrospective Study of a Health Examination Population in Chinese Males. *Front Med (Lausanne)*, 9, pp.843162.
- Tuckey, R.C., Cheng, C.Y.S., & Slominski, A.T., 2019. The Serum Vitamin D Metabolome: What we Know and what is Still to Discover. *J Steroid Biochem Mol Biol.*, 186, pp.4–21.
- Turner, M.E., Rowsell, T.S., White, C.A., Kaufmann, M., Norman, P.A., Neville, K., Petkovich, M., Jones, G., Adams, M.A., & Holden, R.M., 2022. The Metabolism of 1,25(OH)<sub>2</sub>D<sub>3</sub> in Clinical and Experimental Kidney Disease. *Sci Rep.*, 12(1), pp.10925.
- Wannamethee, S.G., Welsh, P., Papacosta, O., Lennon, L., & Whincup, P., 2021. Vitamin D Deficiency, Impaired Lung Function and Total and Respiratory Mortality in a Cohort of Older Men: Cross-Sectional and Prospective Findings from The British Regional Heart Study. *BMJ Open*, 11(12), pp.e051560.
- Weir, C.B., & Jan, A., 2024. BMI Classification Percentile And Cut Off Points. StatPearls. Treasure Island (FL): StatPearls Publishing.
- Yang, L., Zhao, H., Liu, K., Wang, Y., Liu, Q., Sun, T., Chen, S., & Ren, L., 2021. Smoking Behavior and Circulating Vitamin D Levels in Adults: A Meta-Analysis. *Food Sci Nutr.* 9(10), pp.5820–32.

# Demographic Factors and BMI on Declined Lung Function and Vitamin D Levels in Active Smokers

**Pandu Laksono**

Faculty of Pharmacy, University of Surabaya

Author

**Amelia Lorensia**

Faculty of Pharmacy, University of Surabaya

Author

<https://orcid.org/0000-0002-9746-0606>

**Heru Wijono**

Faculty of Medicine, University of Surabaya

Author

**DOI:** <https://doi.org/10.15294/kemas.v21i1.4547>

**Keywords:** vitamin D, lung function, BMI, smoking, demographics

## Abstract

Cigarettes are one of the causes of health problems in the world. Smoking has been known to have a direct impact on reducing lung function. Smoking increases the risk of vitamin D deficiency. Vitamin D is a vital nutrient as a lung immunomodulator. Deficient levels will cause problems with lung health, especially in active smokers. This cross-sectional research using multivariate path analysis and the SEM-PLS method has three objectives. The direct influence of three independent variables, including obesity, smoking, and demographics. Regarding vitamin D status and lung function, analyzing the direct effect of vitamin D on lung function, and thirdly, analyzing the indirect one of the three independent variables on lung function through vitamin D levels. This research took time from October 2023 to January 2024 and involved 47 active smoker respondents whose vitamin D level status and lung function were measured. There was a significant direct effect of obesity level on vitamin D levels in the active smoker population ( $p < 0.05$ ;  $f\text{-square} = 2.889$ ). While demographic factors ( $p > 0.05$ ;  $f\text{-square} = 0.030$ ) and smoking frequency ( $p > 0.05$ ;  $f\text{-square} = 0.003$ ) did not have a direct significant effect. Demographic factors, obesity, and smoking frequency don't have significant direct effect on lung function in active smokers ( $p > 0.05$ ).

 PDF

Published  
2025-07-30

Article ID  
4547

Issue  
[Vol. 21 No. 1 \(2025\)](#)

Section  
Articles

Share



## Similar Articles

- Adriel Wiemputra Wangsa, Wahyu Pamungkasih, [Household Wood Fuel Usage and Lung Cancer Predictor Symptoms in Primary Care: A Retrospective Cross-Sectional Study for Lung Cancer Early Detection](#), *Jurnal Kesehatan Masyarakat*: Vol. 21 No. 1 (2025).



- Indra Pratama Dana, Kanti Yunika, [Relation between Nose Scale and Sleep Disorder Breathing Among Spice Factory Workers in Semarang.](#), *Jurnal Kesehatan Masyarakat: Vol. 20 No. 4 (2025)*.
- Harlyanti Muthma'innah Mashar, Lamia Diang Mahalia, Sukmawati Ahmad Damiti, [Mineral Content and Antioxidant Capacity of Cookies Formulated with Spinach and Pangas Catfish.](#), *Jurnal Kesehatan Masyarakat: Vol. 21 No. 1 (2025)*.
- Ake Royke Calvin Langingi, Grace Irene Viodyta Watung, Sudirman Sudirman, Mareyke Yolanda Lusia Sepang, Siska Sibua, Ignatia Yohana Rembet, [Multifactorial Risk Factors of Hypertension in Patients Aged 45-55 Years in Kota Kotamobagu: A Cross-Sectional Study.](#), *Jurnal Kesehatan Masyarakat: Vol. 21 No. 1 (2025)*.
- Fenti Dewi Pertiwi, Evy Damayanthi, Rimbawan Rimbawan, [Sun Exposure on the Incidence of Allergies in Adult Women.](#), *Jurnal Kesehatan Masyarakat: Vol. 20 No. 4 (2025)*.
- Indah Tri Susilowati, Noviana Dewi, Dewi Saroh, [The Neurotoxic Impact Of Lead On The Appearance Of Antisocial Behavior In Batik Dye Workers.](#), *Jurnal Kesehatan Masyarakat: Vol. 20 No. 3 (2025)*.
- Machrumnizar Machrumnizar, Adang Bachtiar, Rina K. Kusumaratna, [The Double Burden: A Bibliometric Analysis on Tuberculosis with Diabetes Mellitus Comorbidity.](#), *Jurnal Kesehatan Masyarakat: Vol. 21 No. 1 (2025)*.
- Sri Nurlaily Z, Rahma Dewi Agustini, Nurhidayah, [Stunting Among Children Aged 6-59 Months in Gorontalo, Indonesia.](#), *Jurnal Kesehatan Masyarakat: Vol. 20 No. 3 (2025)*.
- Indra Elisabet Lalangpuling, Nurmila Sunati, Michael V.L. Tumbol, Kevin G. Pascoal, Jasman Jasman, [Intestinal Parasitic Infections and their Relationship with Healthy Living Behavior and Nutritional Status in Children.](#), *Jurnal Kesehatan Masyarakat: Vol. 20 No. 3 (2025)*.
- Alfitri Alfitri, Neviyarni Neviyarni, Firman Firman, Netrawati Netrawati, Rihaliza Rihaliza, Mahalul Azam, Elsa Yuniarti, [Exhaustive Analysis of the Study of Antiretroviral Adherence Factor From HIV/AIDS Patients.](#), *Jurnal Kesehatan Masyarakat: Vol. 20 No. 3 (2025)*.

1-10 of 15 [Next](#) →

You may also [start an advanced similarity search](#) for this article.

## MENU

---

SUBMISSION

DOWNLOAD

 ABOUT JOURNAL

CONTACT


 PRIVACY STATEMENT

 REGISTER

LOGIN

## PARAMETER INDEXING

---



#### ACCREDITATION RANKING



Scopus®



Journal & Publication Development Center  
Universitas Negeri Semarang

# KEMAS : Jurnal Kesehatan Masyarakat

Published by Universitas Negeri Semarang, Indonesia

✉ [kemas@mail.unnes.ac.id](mailto:kemas@mail.unnes.ac.id)



## About the Journal

Jurnal Kesehatan Masyarakat starting in 2024 migrates from OJS 2 to OJS 3 to better secure from various unwanted things, including journal hacking and so on. To submit, the author please visit the new website page of our journal at the link <https://journal.unnes.ac.id/journals/kemas>

**MIGRATION OFFICIAL STATEMENT [HERE](#)**

KEMAS: Jurnal Kesehatan Masyarakat [P-ISSN 1858-1196 | E-ISSN 2355-3596] published by Universitas Negeri Semarang in collaboration with [Ikatan Ahli Kesehatan Masyarakat Indonesia](#) (IAKMI Tingkat Pusat) and [Jejaring Nasional Pendidikan Kesehatan](#) (JNPK). KEMAS publishes the article based on research or equivalent to research results in public health or other disciplines related to public health that has not been loaded/published by other media. The journal contains articles about epidemiology and biostatistics, health policy and administration, public health nutrition, environmental health, occupational health and safety, health promotion, reproductive health, maternal and child health, and other related articles in public health. The journal can be used by health practitioners, health caregivers, teachers, medical students, and people who are interested in public health issues. The journal was first published in July 2005. Since 2022, the journal were published quarterly a year in July, October, January, and April. Starting October 2022, articles in KEMAS Journal have been accepted for indexing to [Scopus](#).

## Abstracting & Indexing

[GARUDA](#) | [SINTA](#) | [DOAJ](#) | [DIMENSION](#) | [SCOPUS](#)

## Current Issue

Vol. 21 No. 1 (2025)

DOI: <https://doi.org/10.15294/kemas.v21i1>

Published: 2025-07-30

## Articles

**In Silico and In Vitro Approach of Preeclampsia Prophylaxis from Water of Kalianda Kopyor**

DOI: <https://doi.org/10.15294/kemas.v21i1.3468>

Fitriana Fitriana, Soetrisno Soetrisno, Sri Sulistyowati, Dono Indarto (Author)

Article ID 3468

858-868

**Demographic Factors and BMI on Declined Lung Function and Vitamin D Levels in Active Smokers**

**DOI:** <https://doi.org/10.15294/kemas.v21i1.4547>

Pandu Laksono, Amelia Lorensia, Heru Wijono (Author)  
**Article ID** 4547

869-880



**Household Wood Fuel Usage and Lung Cancer Predictor Symptoms in Primary Care: A Retrospective Cross-Sectional Study for Lung Cancer Early Detection**

**DOI:** <https://doi.org/10.15294/0c6ksf18>

Adriel Wiemputra Wangsa, Wahyu Pamungkasih (Author)  
**Article ID** 5184

881-891



**The Double Burden: A Bibliometric Analysis on Tuberculosis with Diabetes Mellitus Comorbidity**

**DOI:** <https://doi.org/10.15294/kemas.v21i1.10742>

Machrumnizar Machrumnizar, Adang Bachtiar, Rina K. Kusumaratna (Author)  
**Article ID** 10742

892-902



**Mineral Content and Antioxidant Capacity of Cookies Formulated with Spinach and Pangas Catfish**

**DOI:** <https://doi.org/10.15294/qh9vdp63>

Harlyanti Muthma'innah Mashar, Lamia Diang Mahalia, Sukmawati Ahmad Damiti (Author)  
**Article ID** 13408

903-911



**Infertilization of dengue vector (*Aedes aegypti*) with Bromelin Solution and Extract From Ananas comosus**

**DOI:** <https://doi.org/10.15294/kemas.v21i1.15261>

Isnawati Isnawati, Muhammad Ir'fai, Ferry Kriswandana, Wanti Wanti , Muhammad Pahrudin, Muhammad Rasyid Ridha (Author)  
**Article ID** 15261

912-919



**Identification and Treatment for Depressive Disorder: Descriptive Study from Indonesia**

**DOI:** <https://doi.org/10.15294/kemas.v21i1.14298>

Dharmady Agus, Nicholas Hardi, Fransisca Theresia, Aila Johanna, Rima Sisca Fanuela, Ika Suswanti (Author)  
**Article ID** 14298

920-929



**Multifactorial Risk Factors of Hypertension in Patients Aged 45-55 Years in Kota Kotamobagu: A Cross-Sectional Study**

**DOI:** <https://doi.org/10.15294/kemas.v21i1.20561>

Ake Royke Calvin Langingi, Grace Irene Viodyta Watung, Sudirman Sudirman, Mareyke Yolanda Lusia Sepang, Siska Sibua, Ignatia Yohana Rembet (Author)  
**Article ID** 20561

938-947



**Strategies to Improve the Performance of Medical Record Officers Through Structural Equation Model Analysis**

**DOI:** <https://doi.org/10.15294/kemas.v21i1.21183>

MR Tawil, Anita Anita, Muarif Leo, Kuswinton Kuswinton, Muhammad Tasjidin Teheni, Wilda Fatmala (Author); Niska Salsiani Sinta, Yohanis Rongre (Translator)  
**Article ID** 21183

948-958



DOI: <https://doi.org/10.15294/kemas.v21i1.13796>

Jemimah Kezia Lee, Ferdi Antonio (Author)

959-972

Article ID 13796



Benzene, Toluene, Xylene Levels and Subjective Complaints in Shoe Workshops

DOI: <https://doi.org/10.15294/kemas.v21i1.14960>

Taufik Ashar, Devi Nuraini Santi, Evi Naria (Author)

973-982

Article ID 14960



Anemia in Pregnancy Based on Rural and Urban Areas

DOI: <https://doi.org/10.15294/kemas.v21i1.22585>

Catur Wulandari, Pratiwi Hariyani Putri, Anugrah Linda Mutiarani, Farah Nuriannisa, Farda Jamalia Hisbullah, Farah Nur Laily Mabruroh, Jauharotul Lu'luah, Miftakhul Solekha (Author)

983-992

Article ID 22585



[View All Issues](#) >

MENU

SUBMISSION

DOWNLOAD

ABOUT JOURNAL

CONTACT

PRIVACY STATEMENT

REGISTER

LOGIN

PARAMETER INDEXING







Can't write to file  
"/home/sinta/public\_html/public/tmp/776.png". Check  
that the process running PHP has enough permission.

#### ACCREDITATION RANKING

---



# Scopus®



Journal & Publication Development Center  
Universitas Negeri Semarang

[Home](#) / Editorial Team

## Editorial Team

### Editor-in-Chief

- Prof. Dr. dr. Oktia Woro Kasmini Handayani, M.Kes, {SCOPUS ID : 57192428885} {Public Health Nutrition} Universitas Negeri Semarang, Indonesia

### Editorial Advisory Regional America

- Dr. Gina Samaan, {SCOPUS ID : 6602382950} {Epidemiology} US. Centers for Disease Control and Prevention, United States

### Editorial Advisory Regional Asia

- Prof. Kathirvelu Baskar, {SCOPUS ID : 55092286200} {Entomology} Loyola College India, Entomology Research Institute, India
- Dr. Khalid M. Al Aboud, {SCOPUS ID : 7003345190} {Dermatology} King Faisal Specialist Hospital and Research Centre, Saudi Arabia
- Prof. Dato'. Syed Mohamed Aljunid, {SCOPUS ID : 6504304159} {Health Economics, Policy and Management} Universiti Kebangsaan Malaysia, Malaysia
- Dr. Dina Nur Anggraini Ningrum, {SCOPUS ID : 57195329470} {Health Information System} Taipei Medical University, Taiwan, Province of China
- Dr. Mahalul Azam, {SCOPUS ID: 57194196255} {Medical} Universitas Negeri Semarang, Indonesia
- Dr. Songpol Tornee, {SCOPUS ID : 6506180249} {Health Education} Shrinakharinwirot University, Thailand

### Editorial Advisory Regional Australia

- Prof. Doune Macdonald, PhD, {SCOPUS ID : 7401463393} {Health Education} University of Queensland, Australia

### Editorial Advisory Regional Africa

- Assoc. Prof. Dr. Henry Odhianoson Imhonde, {SCOPUS ID : 36069265600} {Psychology} Ambrose Alli University, Nigeria

### Editorial Board

- Muhammad Azinar, S.K.M, M.Kes, {SCOPUS ID : 57194193079} {Health Promotion} Universitas Negeri Semarang, Indonesia
- Nur Siyam, S.K.M, M.PH, {SCOPUS ID : 57222668801} {Maternal and Child Health} Universitas Negeri Semarang, Indonesia
- Efa Nugroho, S.K.M, M.Kes, {SCOPUS ID : 57192436111} {Reproduction Health} Universitas Negeri Semarang, Indonesia

### Administration

- Satria Adi Rachim, Universitas Negeri Semarang, Indonesia
- Widiyanto Widiyanto, Universitas Terbuka, Indonesia

## MENU

✓ DOWNLOAD

📘 ABOUT JOURNAL

👤 CONTACT


🔒 PRIVACY STATEMENT

👤 REGISTER

➡ LOGIN

## PARAMETER INDEXING

---



## ACCREDITATION RANKING

---



# Scopus<sup>®</sup>



Journal & Publication Development Center  
Universitas Negeri Semarang

[Home](#) / [About the Journal](#)

## About the Journal

### Focus and Scope

KEMAS: Jurnal Kesehatan Masyarakat is an international, peer-reviewed journal. It publishes original papers, reviews and short reports on all aspects of the science, philosophy, and practice of public health.

It is aimed at all public health practitioners and researchers and those who manage and deliver public health services and systems. It will also be of interest to anyone involved in provision of public health programmes, the care of populations or communities and those who contribute to public health systems in any way.

Published twelve times a year, KEMAS: Jurnal Kesehatan Masyarakat considers submissions on any aspect of public health across age groups and settings.

These include:

- Public health practice and impact
- Epidemiology and Biostatistic
- Applied Epidemiology
- Need or impact assessments
- Health service effectiveness, management and re-design
- Health Protection including control of communicable diseases
- Health promotion and disease prevention
- Evaluation of public health programmes or interventions
- Public health governance, audit and quality
- Public health law and ethics
- Health policy and administration
- Capacity in public health systems and workforce
- Public health nutrition
- Environmental health
- Occupational health and safety
- Reproductive health
- Maternal and child health

This is not an exhaustive list and the Editors will consider articles on any issue relating to public health.

KEMAS: Jurnal Kesehatan Masyarakat also publishes invited articles, reviews and supplements from leading experts on topical issues.

### Peer Review Process

All manuscripts submitted to this journal must follow focus and scope, and author guidelines of this journal. The submitted manuscripts must address scientific merit or novelty appropriate to the focus and scope. All manuscripts must be free from plagiarism contents. All authors are suggested to use plagiarism detection software to do the similarity checking. Editors check the plagiarism detection of articles in this journal by using a Turnitin software.

The research article submitted to this journal will be double blind reviewed at least 2 {two} or more expert reviewers. The reviewers give scientific valuable comments improving the contents of the manuscript.

Final decision of articles acceptance will be made by Editors according to reviewers comments. Publication of accepted articles including the sequence of published articles will be made by Editor in Chief by considering sequence of accepted date and geographical distribution of authors as well as thematic issue.

### Open Access Policy



This journal provides immediate open access to its content on the principle that making research freely available to the public supports a greater global exchange of knowledge.

This journal is open access journal which means that all content is freely available without charge to users or / institution. Users are allowed to read, download, copy, distribute, print, search, or link to full text articles in this journal without asking prior permission from the publisher or author. This is in accordance with Budapest Open Access Initiative.

## **Publication Ethics and Malpractice Statement**

Our Publication Ethics and Publication Malpractice Statement are based on COPEs Best Practice Guidelines for Journal Editors. As such, this journal follows the COPE [Code of Conduct and Best Practice Guidelines for Journal Editors](#) and the [Code of Conduct for Journal Publishers](#).

A selection of key points is included below, but you should always refer to the three documents listed above for full details.

### **Duties of Editors**

#### *Fair play and editorial independence*

Editors evaluate submitted manuscripts exclusively on the basis of their academic merit {importance, originality, study's validity, clarity} and its relevance to the journal's scope, without regard to the authors' race, gender, sexual orientation, ethnic origin, citizenship, religious belief, political philosophy or institutional affiliation. Decisions to edit and publish are not determined by the policies of governments or any other agencies outside of the journal itself. The Editor-in-Chief has full authority over the entire editorial content of the journal and the timing of publication of that content.

#### *Confidentiality*

Editors and editorial staff will not disclose any information about a submitted manuscript to anyone other than the corresponding author, reviewers, potential reviewers, other editorial advisers, and the publisher, as appropriate.

#### *Disclosure and conflicts of interest*

Editors and editorial board members will not use unpublished information disclosed in a submitted manuscript for their own research purposes without the authors' explicit written consent. Privileged information or ideas obtained by editors as a result of handling the manuscript will be kept confidential and not used for their personal advantage. Editors will recuse themselves from considering manuscripts in which they have conflicts of interest resulting from competitive, collaborative, or other relationships/connections with any of the authors, companies or institutions connected to the papers; instead, they will ask another member of the editorial board to handle the manuscript.

#### *Publication decisions*

The editors ensure that all submitted manuscripts being considered for publication undergo peer-review by at least two reviewers who are expert in the field. The Editor-in-Chief is responsible for deciding which of the manuscripts submitted to the journal will be published, based on the validation of the work in question, its importance to researchers and readers, the reviewers' comments, and such legal requirements as are currently in force regarding libel, copyright infringement and plagiarism. The Editor-in-Chief may confer with other editors or reviewers in making this decision.

#### *Involvement and cooperation in investigations*

Editors {in conjunction with the publisher and/or society} will take responsive measures when ethical concerns are raised with regard to a submitted manuscript or published paper. Every reported act of unethical publishing behaviour will be looked into, even if it is discovered years after publication. AP-SMART editors follow the COPE *Flowcharts* when dealing with cases of suspected misconduct. If, on investigation, the ethical concern is well-founded, a correction, retraction, expression of concern or other note as may be relevant, will be published in the journal.

### **Duties of Reviewers**

#### *Contribution to editorial decisions*

Peer review assists editors in making editorial decisions and, through editorial communications with authors, may assist authors in improving their manuscripts. Peer review is an essential component of formal scholarly communication and lies at the heart of scientific endeavour. AP-SMART shares the view of many that all scholars who wish to contribute to the scientific process have an obligation to do a fair share of reviewing.

#### *Promptness*

Any invited referee who feels unqualified to review the research reported in a manuscript or knows that its prompt review will be impossible should immediately notify the editors and decline the invitation to review so that alternative reviewers can be contacted.

#### *Confidentiality*

Any manuscripts received for review are confidential documents and must be treated as such; they must not be shown to or discussed with others except if authorized by the Editor-in-Chief {who would only do so under exceptional and specific circumstances}. This applies also to invited reviewers who decline the review invitation.

#### *Standards of objectivity*

Reviews should be conducted objectively and observations formulated clearly with supporting arguments so that authors can use them for improving the manuscript. Personal criticism of the authors is inappropriate.

#### *Acknowledgement of sources*

Reviewers should identify relevant published work that has not been cited by the authors. Any statement that is an observation, derivation or argument that has been reported in previous publications should be accompanied by the relevant citation. A reviewer should also notify the editors of any substantial similarity or overlap between the manuscript under consideration and any other manuscript {published or unpublished} of which they have personal knowledge.

#### *Disclosure and conflicts of interest*

Any invited referee who has conflicts of interest resulting from competitive, collaborative, or other relationships or connections with any of the authors, companies or institutions connected to the manuscript and the work described therein should immediately notify the editors to declare their conflicts of interest and decline the invitation to review so that alternative reviewers can be contacted.

Unpublished material disclosed in a submitted manuscript must not be used in a reviewers own research without the express written consent of the authors. Privileged information or ideas obtained through peer review must be kept confidential and not used for the reviewers personal advantage. This applies also to invited reviewers who decline the review invitation.

### **Duties of Authors**

#### *Reporting standards*

Authors of original research should present an accurate account of the work performed and the results, followed by an objective discussion of the significance of the work. The manuscript should contain sufficient detail and references to permit others to replicate the work. Review articles should be accurate, objective and comprehensive, while editorial 'opinion' or perspective pieces should be clearly identified as such. Fraudulent or knowingly inaccurate statements constitute unethical behaviour and are unacceptable.

#### *Data access and retention*

Authors may be asked to provide the raw data of their study together with the manuscript for editorial review and should be prepared to make the data publicly available if practicable. In any event, authors should ensure accessibility of such data to other competent professionals for at least 10 years after publication {preferably via an institutional or subject-based data repository or other data centre}, provided that the confidentiality of the participants can be protected and legal rights concerning proprietary data do not preclude their release.

#### *Originality and plagiarism*

Authors should ensure that they have written and submit only entirely original works, and if they have used the work and/or words of others, that this has been appropriately cited. Publications that have been influential in determining the nature of the work reported in the manuscript should also be cited. Plagiarism takes many forms, from "passing off" another's paper as the author's own, to copying or paraphrasing substantial parts of another's paper {without attribution}, to claiming results from research conducted by others. Plagiarism in all its forms constitutes unethical publishing behaviour and is unacceptable.

#### *Multiple, duplicate, redundant or concurrent submission/publication*

Papers describing essentially the same research should not be published in more than one journal or primary publication. Hence, authors should not submit for consideration a manuscript that has already been published in another journal. Submission of a manuscript concurrently to more than one journal is unethical publishing behaviour and unacceptable.

The publication of some kinds of articles {such as clinical guidelines, translations} in more than one journal is sometimes justifiable, provided that certain conditions are met. The authors and editors of the journals concerned must agree to the secondary publication, which must reflect the same data and interpretation of the primary document. The primary reference must be cited in the secondary publication.

#### *Authorship of the manuscript*

Only persons who meet these authorship criteria should be listed as authors in the manuscript as they must be able to take public responsibility for the content: {i} made significant contributions to the conception, design, execution, data acquisition, or analysis/interpretation of the study; and {ii} drafted the manuscript or revised it critically for important intellectual content; and {iii} have seen and approved the final version of the paper and agreed to its submission for publication. All persons who made substantial contributions to the work reported in the manuscript {such as technical help, writing and editing assistance, general support} but who do not meet the criteria for authorship must not be listed as an author, but should be acknowledged in the "Acknowledgements" section after their written permission to be named as been obtained. The corresponding author should ensure that all appropriate coauthors {according to the above definition} and no inappropriate coauthors are included in the author list and verify that all coauthors have seen and approved the final version of the manuscript and agreed to its submission for publication.

#### *Disclosure and conflicts of interest*

Authors should "at the earliest stage possible {generally by submitting a disclosure form at the time of submission and including a statement in the manuscript} disclose any conflicts of interest that might be construed to influence the results or their interpretation in the manuscript. Examples of potential conflicts of interest that should be disclosed include financial ones such as honoraria, educational grants or other funding, participation in speakers bureaus, membership, employment, consultancies, stock ownership, or other equity interest, and paid expert testimony or patent-licensing arrangements, as well as non-financial ones such as personal or professional relationships, affiliations, knowledge or beliefs in the subject matter or materials discussed in the manuscript. All sources of financial support for the work should be disclosed {including the grant number or other reference number if any}.

#### *Acknowledgement of sources*

Authors should ensure that they have properly acknowledged the work of others, and should also cite publications that have been influential in determining the nature of the reported work. Information obtained privately {from conversation, correspondence or discussion with third parties} must not be used or reported without explicit, written permission from the source. Authors should not use information obtained in the course of providing confidential services, such as refereeing manuscripts or grant applications, unless they have obtained the explicit written permission of the author{s} of the work involved in these services.

#### *Hazards and human or animal subjects*

If the work involves chemicals, procedures or equipment that have any unusual hazards inherent in their use, the authors must clearly identify these in the manuscript. If the work involves the use of animals or human participants, the authors should ensure that all procedures were performed in compliance with relevant laws and institutional guidelines and that the appropriate institutional committee{s} has approved them; the manuscript should contain a statement to this effect. Authors should also include a

statement in the manuscript that informed consent was obtained for experimentation with human participants. The privacy rights of human participants must always be observed.

#### *Peer review*

Authors are obliged to participate in the peer review process and cooperate fully by responding promptly to editors requests for raw data, clarifications, and proof of ethics approval, patient consents and copyright permissions. In the case of a first decision of "revisions necessary", authors should respond to the reviewers comments systematically, point by point, and in a timely manner, revising and re-submitting their manuscript to the journal by the deadline given.

#### *Fundamental errors in published works*

When authors discover significant errors or inaccuracies in their own published work, it is their obligation to promptly notify the journals editors or publisher and cooperate with them to either correct the paper in the form of an erratum or to retract the paper. If the editors or publisher learns from a third party that a published work contains a significant error or inaccuracy, then it is the authors obligation to promptly correct or retract the paper or provide evidence to the journal editors of the correctness of the paper.

### **Duties of the Publisher**

#### *Handling of unethical publishing behaviour*

In cases of alleged or proven scientific misconduct, fraudulent publication or plagiarism, the publisher, in close collaboration with the editors, will take all appropriate measures to clarify the situation and to amend the article in question. This includes the prompt publication of an erratum, clarification or, in the most severe case, the retraction of the affected work. The publisher, together with the editors, shall take reasonable steps to identify and prevent the publication of papers where research misconduct has occurred, and under no circumstances encourage such misconduct or knowingly allow such misconduct to take place.

#### *Access to journal content*

The publisher is committed to the permanent availability and preservation of scholarly research and ensures accessibility by partnering with organizations and maintaining our own digital archive.

### **Section A: Publication and authorship**

1. All submitted papers are subject to strict peer-review process by at least two international reviewers that are experts in the area of the particular paper.
2. Review process are blind peer review.
3. The factors that are taken into account in review are relevance, soundness, significance, originality, readability and language.
4. The possible decisions include acceptance, acceptance with revisions, or rejection.
5. If authors are encouraged to revise and resubmit a submission, there is no guarantee that the revised submission will be accepted.
6. Rejected articles will not be re-reviewed.
7. The paper acceptance is constrained by such legal requirements as shall then be in force regarding libel, copyright infringement and plagiarism.
8. No research can be included in more than one publication.

### **Section B: Authors' responsibilities**

1. Authors must certify that their manuscripts are their original work.
2. Authors must certify that the manuscript has not previously been published elsewhere.
3. Authors must certify that the manuscript is not currently being considered for publication elsewhere.
4. Authors must participate in the peer review process.
5. Authors are obliged to provide retractions or corrections of mistakes.
6. All Authors mentioned in the paper must have significantly contributed to the research.
7. Authors must state that all data in the paper are real and authentic.
8. Authors must notify the Editors of any conflicts of interest.
9. Authors must identify all sources used in the creation of their manuscript.
10. Authors must report any errors they discover in their published paper to the Editors.

### **Section C: Reviewers' responsibilities**

1. Reviewers should keep all information regarding papers confidential and treat them as privileged information.
2. Reviews should be conducted objectively, with no personal criticism of the author

3. Reviewers should express their views clearly with supporting arguments
4. Reviewers should identify relevant published work that has not been cited by the authors.
5. Reviewers should also call to the Editor in Chief's attention any substantial similarity or overlap between the manuscript under consideration and any other published paper of which they have personal knowledge.
6. Reviewers should not review manuscripts in which they have conflicts of interest resulting from competitive, collaborative, or other relationships or connections with any of the authors, companies, or institutions connected to the papers.

#### **Section D: Editors' responsibilities**

1. Editors have complete responsibility and authority to reject/accept an article.
2. Editors are responsible for the contents and overall quality of the publication.
3. Editors should always consider the needs of the authors and the readers when attempting to improve the publication.
4. Editors should guarantee the quality of the papers and the integrity of the academic record.
5. Editors should publish errata pages or make corrections when needed.
6. Editors should have a clear picture of a research's funding sources.
7. Editors should base their decisions solely on the papers' importance, originality, clarity and relevance to publication's scope.
8. Editors should not reverse their decisions nor overturn the ones of previous editors without serious reason.
9. Editors should preserve the anonymity of reviewers.
10. Editors should ensure that all research material they publish conforms to internationally accepted ethical guidelines.
11. Editors should only accept a paper when reasonably certain.
12. Editors should act if they suspect misconduct, whether a paper is published or unpublished, and make all reasonable attempts to persist in obtaining a resolution to the problem.
13. Editors should not reject papers based on suspicions, they should have proof of misconduct.
14. Editors should not allow any conflicts of interest between staff, authors, reviewers and board members.

Sources:

- [ELSEVIER: Elsevier publishing ethics resource kit](#)
- [COPE: Responsible research publication: international standards for authors](#)
- [COPE: Cope's new code of conduct](#)
- [COPE: Responsible research publication: International standards for editors](#)
- [COPE: Cope short guide to ethical editing for new editors](#)
- [COPE: Cope ethical guidelines for peer reviewers](#)
- [COPE: The editorial board follows the guidelines for retracting articles issued by COPE](#)
- [COPE: Code of conduct for journal publishers](#)
- [COPE: Cope retraction guidelines](#)

#### **Policy of Screening for Plagiarism**

All manuscripts must be free from plagiarism contents. All authors are suggested to use plagiarism detection software to do the similarity checking. Editors check the plagiarism detection of articles in this journal by using a Turnitin software.

#### **Author Guidelines**

The manuscript texts are written in English or Indonesia. Manuscripts will be first reviewed by editorial boards. The main text of a manuscript must be submitted as a Word document {.doc} or Rich Text Format {.rtf} file. The manuscript consists of 5000 words {minimum}, well-typed in single column on A4 size paper, use 12 pt of Times New Roman. The manuscript contains an original work and have potentially contribute to the highly scientific advancement.

The manuscript should contain the following section in this order:

a. Title

Title of articles in English should describe the main content of manuscripts, be informative, concise, and not too wordy {12-15 words only}, and does not contain formulas.

b. The authors name

Full name without academic degrees and titles, written in capital letters. Manuscript written by groups needs to supplemented by complete contact details.

c. Name of affiliation for each author

The author name should be accompanied by complete affiliation address, postal code number, telephone number and email address.

d. Abstract

Written briefly in English in one paragraph of 150-200 words, containing background, research



objectives, methodology, results, conclusion of the study and your research contributions to science.

e. Keywords

Written in English 3-5 words or groups of words, written alphabetically.

f. Introduction

Explaining the background, problems, importance of research, brief literature review that relates directly to research or previous findings that need to be developed, and ended with a paragraph of research purposes. A balance must be kept between the pure and applied aspects of the subject. The introduction is presented in the form of paragraphs of approximately 1000 words.

g. Methods

Make sure that work can be repeated according to the details provided. It contains technical information of the study presented clearly. Therefore, readers can conduct research based on the techniques presented. Materials and equipment specifications are necessary. Approaches or procedures of study together with data analysis methods must be presented.

h. Results and Discussion

Well-prepared tables and or figures must be of significant feature of this section, because they convey the major observations to readers. Any information provided in tables and figures should no longer be repeated in the text, but the text should focus on the importance of the principal findings of the study. In general, journal papers will contain three-seven figures and tables. Same data can not be presented in the form of tables and figures. The results of the study are discussed to address the problem formulated, objectives and research hypotheses. It is highly suggested that discussion be focused on the why and how of the research findings can happen and to extend to which the research findings can be applied to other relevant problems.

i. Conclusion

Conclusion should be withdrawn on the basis of research findings, formulated concerns and research purposes. Conclusion is presented in one paragraph without numerical form of expression. Explain your research contributions to science.

j. Acknowledgement

Contributors who are not mentioned as authors should be acknowledged, and their particular contribution should be described. All sources of funding for the work must be acknowledged, both the research funder and the grant number {if applicable} should be given for each source of funds

k. References

Manuscripts are written by using standard citation application {Mendeley/Endnote/Zotero}. APA {American Psychological Association} reference style is required.

## Copyright Notice

Authors retain copyright and grant the journal right of first publication with the work simultaneously licensed under a [Creative Commons Attribution License](#) that allows others to share the work with an acknowledgement of the works authorship and initial publication in this journal.

Authors are able to enter into separate, additional contractual arrangements for the non-exclusive distribution of the journal's published version of the work {e.g., post it to an institutional repository or publish it in a book}, with an acknowledgement of its initial publication in this journal.

Authors are permitted and encouraged to post their work online {e.g., in institutional repositories or on their website} prior to and during the submission process, as it can lead to productive exchanges, as well as earlier and greater citation of published work {See [The Effect of Open Access](#)}.

## Author Fees

This journal charges the following author fees.

Article Submission: 0.00 {IDR}

Article Publication: 3.300.000 {IDR}

## MENU

✓ DOWNLOAD

📘 ABOUT JOURNAL

👤 CONTACT


🔒 PRIVACY STATEMENT

👤 REGISTER

➡ LOGIN

## PARAMETER INDEXING

---



## ACCREDITATION RANKING

---



# Scopus<sup>®</sup>



Journal & Publication Development Center  
Universitas Negeri Semarang

# Scopus Indexed. IF: 2.2

Rapid Publication, Reliable Peer-review, Low APCs. Indexed in Scopus.

Curr. Mole

## Kemas



COUNTRY	SUBJECT AREA AND CATEGORY	PUBLISHER	SJR 2024
Indonesia	Medicine Health Policy Public Health, Environmental and Occupational Health	Universitas Negeri Semarang	0.141 Q4
<div><div></div><div>Universities and research institutions in Indonesia</div></div> <div><div></div><div>Media Ranking in Indonesia</div></div>			H-INDEX 4
PUBLICATION TYPE	ISSN	COVERAGE	INFORMATION
Journals	18581196, 23553596	2022-2024	<a href="#">Homepage</a> <a href="#">How to publish in this journal</a> <a href="mailto:oktia_woro@yahoo.co.id">oktia_woro@yahoo.co.id</a>

### SCOPE


KEMAS: Jurnal Kesehatan Masyarakat is an international, peer-reviewed journal. It publishes original papers, reviews and short reports on all aspects of the science, philosophy, and practice of public health. It is aimed at all public health practitioners and researchers and those who manage and deliver public health services and systems. It will also be of interest to anyone involved in provision of public health programmes, the care of populations or communities and those who contribute to public health systems in any way. Published twelve times a year, KEMAS: Jurnal Kesehatan Masyarakat considers submissions on any aspect of public health across age groups and settings.

Join the conversation about this journal

Buy vitamins and supplements

**SCImago Graphica**

Explore, visually communicate and make sense of data with our [new data visualization tool](#).





Loading comments...

Developed by:



Powered by:



Follow us on @ScimagoJR

Scimago Lab, Copyright 2007-2025. Data Source: Scopus®

EST MODUS IN REBUS  
Horatio (Satire 1.1.100)

[Legal Notice](#)

[Privacy Policy](#)





# Source details

Kemas

Open Access ⓘ

Years currently covered by Scopus: from 2019 to 2025

Publisher: Universitas Negeri Semarang


ISSN: 1858-1196 E-ISSN: 2355-3596

Subject area: Medicine: Health Policy Medicine: Public Health, Environmental and Occupational Health

Source type: Journal

[View all documents >](#)

[Set document alert](#)

 [Save to source list](#)

CiteScore 2024  
**0.7** ⓘ

SJR 2024  
**0.141** ⓘ

SNIP 2024  
**0.197** ⓘ

CiteScore CiteScore rank & trend Scopus content coverage

---

## About Scopus

[What is Scopus](#)

[Content coverage](#)

[Scopus blog](#)

[Scopus API](#)

[Privacy matters](#)

## Language

[日本語版を表示する](#)

[查看简体中文版本](#)

[查看繁體中文版本](#)

[Просмотр версии на русском языке](#)

## Customer Service

[Help](#)

[Tutorials](#)

[Contact us](#)

---

## ELSEVIER

[Terms and conditions ↗](#) [Privacy policy ↗](#) [Cookies settings](#)

All content on this site: Copyright © 2025 Elsevier B.V. ↗, its licensors, and contributors. All rights are reserved, including those for text and data mining, AI training, and similar technologies. For all open access content, the relevant licensing terms apply. We use cookies to help provide and enhance our service and tailor content. By continuing, you agree to the use of cookies ↗.



# Demographic Factors and BMI on Declined Lung Function and Vitamin D Levels in Active Smokers

*by* Amelia Lorensia

---

**Submission date:** 19-Aug-2025 10:06AM (UTC+0700)

**Submission ID:** 2731708937

**File name:** Demographic\_Factors\_and\_BMI.pdf (545.18K)

**Word count:** 7694

**Character count:** 39235

# Demographic Factors and BMI on Declined Lung Function and Vitamin D Levels in Active Smokers

## ORIGINALITY REPORT

10%

SIMILARITY INDEX

9%

INTERNET SOURCES

10%

PUBLICATIONS

2%

STUDENT PAPERS

## PRIMARY SOURCES

1	<a href="http://perpustakaan.poltekkes-malang.ac.id">perpustakaan.poltekkes-malang.ac.id</a> Internet Source	2%
2	<a href="http://ijbmi.org">ijbmi.org</a> Internet Source	1%
3	"Handbook of vitamin D in human health", Brill, 2013 Publication	1%
4	<a href="http://www.frontiersin.org">www.frontiersin.org</a> Internet Source	1%
5	Birgitta Annabel Luissa, So Yohanes Jimmy, David Tjahjana. "Investigation of The Antecedents of The Digital Mindset and Its Influence on Employee Intention to Stay in Generation Z: Empirical Study in The Indonesian Creative Industry", The South East Asian Journal of Management, 2024 Publication	1%
6	Lu Yang, Hang Zhao, Ke Liu, Yichao Wang, Qianqian Liu, Tiantian Sun, Shuchun Chen, Luping Ren. "Smoking behavior and circulating vitamin D levels in adults: A meta-analysis", Food Science & Nutrition, 2021 Publication	1%
7	<a href="http://enrichment.iocspublisher.org">enrichment.iocspublisher.org</a> Internet Source	1%

8

vitamindwiki.com  
Internet Source

1 %

9

www.nature.com  
Internet Source

1 %

10

dinastipub.org  
Internet Source

1 %

11

Submitted to Surabaya University  
Student Paper

1 %

12

pubmed.ncbi.nlm.nih.gov  
Internet Source

1 %

Exclude quotes      On  
Exclude bibliography      On

Exclude matches      < 1 %



## Demographic Factors and BMI on Declined Lung Function and Vitamin D Levels in Active Smokers

Pandu Laksono<sup>1</sup>, Amelia Lorensia<sup>1✉</sup>, Heru Wijono<sup>2</sup>

<sup>1</sup>Faculty of Pharmacy, University of Surabaya (Universitas Surabaya (UBAYA)) – Indonesia

<sup>2</sup>Faculty of Medicine, University of Surabaya (Universitas Surabaya (UBAYA)) – Indonesia

### Article Info

**Article History:**  
Submitted December 20XX  
Accepted September 20XX  
Published January 20XX

**Keywords:**  
vitamin D; lung  
function; BMI;  
smoking; demographics

**DOI**  
[https://doi.org/xxxxxxx/  
kemas.vxxxx.xxxx](https://doi.org/xxxxxxx/kemas.vxxxx.xxxx)

### Abstract

Cigarettes are one of the causes of health problems in the world. Smoking has been known to have a direct impact on reducing lung function. Smoking increases the risk of vitamin D deficiency. Vitamin D is a vital nutrient as a lung immunomodulator. Deficient levels will cause problems with lung health, especially in active smokers. This cross-sectional research using multivariate path analysis and the SEM-PLS method has three objectives. The direct influence of three independent variables, including obesity, smoking, and demographics. Regarding vitamin D status and lung function, analyzing the direct effect of vitamin D on lung function, and thirdly, analyzing the indirect one of the three independent variables on lung function through vitamin D levels. This research took time from October 2023 to January 2024 and involved 47 active smoker respondents whose vitamin D level status and lung function were measured. There was a significant direct effect of obesity level on vitamin D levels in the active smoker population ( $p < 0.05$ ;  $f\text{-square} = 2.889$ ). While demographic factors ( $p > 0.05$ ;  $f\text{-square} = 0.030$ ) and smoking frequency ( $p > 0.05$ ;  $f\text{-square} = 0.003$ ) did not have a direct significant effect. Demographic factors, obesity, and smoking frequency don't have significant direct effect on lung function in active smokers ( $p > 0.05$ ).

### Introduction

Indonesia is one of the countries with the highest number of smokers in the world. Based on data from the 2021 Global Adult Tobacco Survey (GATS) released by the Ministry of Health of the Republic of Indonesia (KEMENKES RI), the smoking prevalence of the adult population in Indonesia reached 33.5% in 2021 with an addition of 8 million people over the last 10 years (CDC, 2021). Exposure to cigarette smoke can cause inflammation of the airways and accumulation of mucus in the lungs, resulting in symptoms of shortness of breath and accelerating the decline in lung function. It is often associated with the emergence of lung diseases such as lung cancer, chronic obstructive pulmonary disease (COPD), asthma, and tuberculosis (Chung *et*

*al.*, 2023). Smokers experience decreased lung function compared to non-smokers, which can be measured using spirometry through a decrease in forced expiratory volume in 1 second (FEV1) values reaching >50 mL per year (Lorensia *et al.*, 2021). The higher the intensity of smoking, the lower the rate of decline in FEV1 and FVC. Greater than non-smokers. Further reduction in lung function due to smoking will have an impact on various lung health problems, such as COPD, including emphysema, chronic bronchitis, and asthma (Tian *et al.*, 2023).

Another mechanism that also plays a role in causing damage or decreased lung function is the involvement of levels of a vitamin in the body, namely vitamin D. Vitamin D has a protective mechanism for lung function

✉ Correspondence Address:  
Faculty of Pharmacy, University of Surabaya (Universitas Surabaya (UBAYA)) –  
Indonesia  
Email: amelia.lorensia@gmail.com

through increasing the secretion of the antimicrobial peptide cathelicidin, decreasing chemokine production, inhibiting dendritic cell activation and changing cell activation. T. This cellular mechanism is vital for the response of the lung organ to the threat of infection and the development of allergic lung diseases such as asthma (Bishop *et al.*, 2020). A decrease in the production of the inactive form of vitamin D (25(OH)D) in lung epithelial cells is thought to be caused by exposure to smoke due to smoking activities (Lorensia *et al.*, 2024). A meta-analysis result by Yang *et al.* (2021) of 24 studies with 11,340 participants showed that levels of vitamin D in the inactive form 25(OH)D were lower in smokers than non-smokers (Yang *et al.*, 2021). In addition, the expression level of vitamin D receptors can also be influenced by exposure to cigarette smoke (Ahn *et al.*, 2021). A study by Ghosh *et al.* (2020) also stated that vitamin D deficiency plays a role in changes in lung structure and decreased lung function.

There are other influencing factors that can be part of the impact of decreased lung function besides smoking, namely, obesity. Individuals who are obese show reduced lung volume and capacity when compared to people of normal weight. Larger fat deposits in the abdominal area produce greater resistance to diaphragm contraction, thereby inhibiting respiratory ventilation mechanisms (Cao *et al.*, 2022). Based on the explanation above, several previous studies have been conducted that examined the effect of vitamin D on lung function. However, there have been no studies that have examined this by involving factors such as obesity levels, smoking, and demographics in Indonesia, which is one of the countries with the highest prevalence of smoking, accompanied by consequences in the form of increased death rates and lung disease sufferers due to smoking. Therefore, research will be carried out that will examine the influence of demographic factors, BMI, and smoking intensity, on the decline in lung function and vitamin D levels in active smokers. There is a study by Abi-Ayad *et al.* (2023) who measured smokers' vitamin D levels using blood plasma samples, found that vitamin D deficiency was associated with lower lung function conditions, lung function in this case FEV1, FVC, and FEV1/FVC measured

using a spirometer experienced faster decrease in smoking subjects. It shows that adequate serum vitamin D levels are associated with a protective effect against the detrimental effects of smoking on lung function.

## 1 Method

This study used an observational clinical trial with a cross-sectional design where the data collection stage was carried out once at a time. The independent variables in this study were demographic factors, BMI, and smoking intensity. The dependent variable in this study was the lung function value (percentage of FEV1/FVC ratio). The mediating variable in this study was vitamin D levels. Demographic factors consist of age and education level. Age is a measure of the patient's length of life, which is calculated based on the patient's date of birth until they become a research respondent. Educational level is a measure of the respondent's level of education as evidenced by possession of the latest educational certificate. Body Mass Index, or BMI, is a value obtained from the mass and height of each sample individual. BMI can be calculated using a formula by dividing the individual's body weight in kilograms (kg) by the square of body height in meters squared ( $m^2$ ) (Weir & Jan, 2023). Smoking intensity is the habit of smoking tobacco cigarettes. Smoker classification can be calculated using the Brinkman Index (IB)=number of cigarettes smoked per day x length of smoking (years) (Herath *et al.*, 2022). Lung function measurements using a handheld spirometer. The level of lung function impairment based on the percentage of the FEV1/FVC ratio is divided into 4 categories, namely normal-mild obstruction, moderate obstruction, severe obstruction, and very severe obstruction (Stanojevic, 2021).

The vitamin D levels that will be measured are the most abundant metabolite form in serum, namely 25(OH)D. 25(OH) levels reflect skin production of vitamin D3 and vitamin D (D2 and D3) from food, 25(OH) D has a half-life (The long  $t_{1/2}$ ) in the blood circulation is 3-4 weeks compared to the active vitamin D metabolite 1,25(OH)2D which only has a short  $t_{1/2}$  of around 4-6 hours (Tuckey *et al.*, 2019). Methods for observing vitamin



D levels (25(OH)D), which are the ELFA (Enzyme Linked Fluorescent Assay) method used is the VIDAS tool from bioMerieux. The type of specimen that can be used in testing can be serum or blood plasma. In this study, serum 25(OH)D levels were classified as deficient (<20 ng/mL), insufficiency (20-29 ng/mL), and normal (30-100 ng/mL) (Ahn *et al.*, 2021; Amrein *et al.*, 2020).

The population is active smokers located in the Mejoyo area, Rungkut District, Surabaya City. The accessible population is active smoking respondents in the Mejoyo 2 RT 6 Surabaya area who have filled out the questionnaire, can be found, and are not included in the exclusion criteria. The sample in this study was part of the affordable population who met the criterias: (1) aged 18-60 years; (2) didn't use vape; (3) no history of COVID-19 infection; (4) no history of diseases or conditions that can affect serum vitamin D levels; (5) not taking supplements containing vitamin D during the last 1 month before checking 25(OH)D levels; and (6) no history of diseases or conditions that can affect lung function. The minimum number of sample was calculated using the Slovin method. So the number of samples required in this research was 45 people. The sampling technique was purposive sampling.

The first meeting involved a recording of demographic data for the entire sample through a direct interview process with research respondents, and then continued with measuring height and weight. The researcher measured the height of the sample using a height

measuring device with the Onemed brand type HT701 Wireless. Body weight measurements were carried out by researchers using a SPEEDS brand digital weight scale, type LX040-8 USB, which has a sensitivity of one digit after the comma. Lung function measurements were carried out by researchers who had received special training from clinicians using a Contec SP10 handheld spirometer to obtain FEV1 and FVC values. The lung function measurement procedure for each respondent was repeated 3 times. The results in milliliters (mL) are recorded to calculate the percentage of the FEV1/FVC ratio. Measurement of Vitamin D or 25(OH)D levels using the VIDAS® 25 OH Vitamin D TOTAL (VITD) tool with the ELFA technique carried out by a standardized laboratory in Surabaya.

The aim of carrying out multivariate analysis is to determine the magnitude of the influence between variables determined based on the  $P_{value}$  and  $t_{statistic}$  value using the Structural Equation Modeling with Partial Least Squares (SEM-PLS) method with the help of the SmartPLS application. SEM-PLS is a powerful analysis method and is often referred to as soft modeling, because it eliminates the assumptions of Ordinary Least Square (OLS) regression such as data must be normally distributed in a multivariate manner and there is no problem of multicollinearity between independent variables, SEM-PLS can be used to testing weak theories and weak data (small samples and data normality problems). The analysis of the magnitude




Figure 1. Path Analysis

#### Information :

- i. Direct influence of obesity level on lung function
- ii. Direct influence of obesity level on vitamin D levels
- iii. Direct effect of smoking on vitamin D levels
- iv. Direct effect of smoking on lung function
- v. Direct influence of demographics on vitamin D levels
- vi. Direct influence of demographics on lung function
- vii. Direct influence of vitamin D levels on lung function

of the effect in this research will estimate the value of the  $P_{value}$  and also the value of the  $t_{statistic}$ . If the test results show a  $P_{value} < 0.05$  and a  $t_{statistic} \geq t_{table}$  value, then the influence of the two variables is said to be significant. It comes along with the conclusion that there is an influence of the independent variable on the dependent variable of active smoking. However, if the  $P_{value} > 0.05$  and the  $t_{statistic} < t_{table}$  value, then the influence of the two variables is said to be insignificant, with the conclusion that there is no influence of the independent variable on the dependent variable in active smokers. An overview of the path analysis model used in the research can be seen in **Figure 1**.

**Result and Discussion**

This research was conducted from October 2023 to March 2024, located in the Mejoyo 2 RT 6 Surabaya, Kalirungkut, East Surabaya. This research has received an ethical certificate number 232/KE/IX/2023 from the University of Surabaya. Based on the preliminary study, the affordable population was 54 smokers in the area, and 47 people were willing and met the research subject criteria. Most of the respondents were the largest early elderly, 23 respondents (48.93%). Based on BMI measurements, the normal BMI group was the largest, namely 29 respondents (61.70%).

Based on the measurement of education level, the Secondary Education group was the largest, namely 20 respondents (42.55%). Based on smoking frequency, the moderate smoker group was the largest, namely 27 respondents (57.44%). Data on the distribution of respondents' characteristics can be seen in Table 1.

In this study, vitamin D (25(OH)D) levels were measured in 47 respondents using the ELFA method with the VIDAS® tool from Biomerieux. Based on the classification of vitamin D levels of the total respondents measured, the insufficiency group was the largest group with 23 respondents (48.93%). The results of the percentage value of the FEV1/FVC ratio, of the total respondents who were measured, the moderate obstruction group had the largest, namely 21 respondents (44.68%) (Table 2).

The implementation of multivariate statistical analysis in this research was carried out using a quantitative technical approach, where there were two analyzes used, namely descriptive analysis and hypothesis testing or statistical analysis using the Partial Least Square (PLS) method using the SmartPLS program. There are 5 (five) variables involved in this research, which include obesity levels, smoking, demographics, vitamin D, and lung function.

Table 1. Demographic Profile of Respondents

		Frequency (n=47)	Percentage (%)	$\bar{x} \pm SD$
Age (years)	Late Adulthood (36-45)	10	21.27	51.20±6.43
	Early Seniors (46-55)	23	48.93	
	Late Seniors (56-60)	14	29.78	
	Basic Education (elementary school-junior high school)	19	40.42	
	Secondary Education (senior high school) higher education	20 8	42.55 17.02	
BMI (kg/m²)	Normal (18.5-25.0)	29	61.70	
	Fat (Overweight) (25.1-27.0)	14	29.78	
	Obesity (>27)	4	8.51	
B r i n k m a n Index (BI)	Light (<200)	14	29.78	
	Moderate (200-599)	27	57.44	
	High (>600)	6	12.76	

$\bar{x}$  = average; SD= standard deviation  
Source: Primary Data, 2024

Table 2. Distribution of Vitamin D and Lung Function Examination

		Frequency (n=47)	Percentage (%)	$\bar{x} \pm SD$
Vitamin D Classification -25(OH) D Content (ng/mL)	Deficiency (<20)	15	31.91	23.52±8.75
	Insufficiency (20-29)	23	48.93	
	Normal (30-100)	9	19.14	
Lung Function Classification (FEV1/ FVC ratio) (%)	Normal-Mild Obstruction (≥ 80)	7	14.89	95.43±5.72
	Moderate Obstruction (50- 79)	21	44.68	
	Severe Obstruction (30-49)	10	21.27	
	Very Severe Obstruction (<30)	9	19.14	

$\bar{x}$  = average; SD= standard deviation

Source: Primary Data, 2024

The research model evaluation stage in PLS consists of measuring model evaluation and structural model evaluation. The measurement model in this research consists of a reflective measurement model for the variables Levels of Obesity, Smoking, Demography, Vitamin D, and Lung Function by examining loading factor values  $\geq 0.50$ , composite reliability  $\geq 0.60$ , and average variance extracted (AVE)  $\geq 0.50$ . The level of obesity is measured by BMI (X1.2), where the outer loading value is between 0.688-0.936, which shows that BMI is strongly correlated in explaining the obesity level variable. Every change in BMI on obesity levels has a more significant effect. The smoking variable is measured from the Brinkman index (IB) of smoking frequency (X2.1), where the outer loading value is between 0.243-0.996, which shows that IB is correlated in explaining the smoking variable. The level of reliability of the motivation variable is well accepted, with a composite reliability value (0.618)  $> 0.60$ . Any change in smoking frequency on smoking activity has a more significant effect.

Demographic variables are measured by 2 (two) indicators, namely age (X3.1) and education level (X3.2), where the outer loading value is between 0.243-0.996, which shows that these two indicators correlate in explaining the obesity level variable. The level of reliability of the motivation variable is acceptable, even though the composite reliability value (0.406) is  $< 0.60$ , but has an AVE value close to 5.0.

Between the two measurement indicators, the validity of demographic variables appears to be more strongly reflected by the education level indicator (X3.3). It means that every change in age demographics has a more significant effect. The vitamin D variable is measured directly with 1 (one) indicator, namely the 25(OH)D level, where the outer loading value is 1,000, which shows that this indicator has a strong correlation in explaining the vitamin D variable. The level of reliability of the vitamin D variable is acceptable, with the composite reliability value (1,000)  $> 0.60$ . The validity measurement of the vitamin D variable looks strong, as reflected by the 25(OH)D level indicator (Y). The lung function variable is measured directly with 1 (one) indicator, namely the P<sub>value</sub> of the percentage of the FEV1/FVC ratio, where the outer loading value is 1,000, which shows that this indicator has a strong correlation in explaining the lung function variable. The level of reliability of the vitamin D variable is acceptable, with a composite reliability value (1,000)  $> 0.60$ . The validity measurement of lung function variables appears to be strong, as reflected by the percentage indicator of the FEV1/FVC ratio (Z).

Structural model evaluation is a form of evaluation to determine the influence of various endogenous variables on exogenous ones by observing the R-square (R<sup>2</sup>) value (coefficient determination), if the R<sup>2</sup> value = 0.19 then the exogenous influence on the endogenous is weak,

Table 3. Evaluation Results of Reflective Measurement Model, Inner VIF Measurement

Variables	Measurement Item Code	Indicators	Outer Loading	Composite Reliability	AVE	VIF
Obesity Rate	X1.2	BMI	0.936	0.802	0.675	1.178
Cigarette Frequency	X2.1	Smoking Frequency	0.996	0.618	0.525	1.024
Demographics	X3.1	Age	0.137	0.406	0.485	1.007
	X3.2	Level of education	0.975			1.007
Vitamin D	Y	25(OH)D levels	1.000	1.000	1.000	1.000
Lung Function	Z	FEV1/FVC ratio	1.000	1.000	1.000	1.000

AVE=average variance extracted; BMI=Body Mass Index; FEV<sub>1</sub>=Forced Expiratory Volume in 1 second; VIF = Variance Inflated Factor

Source: Primary Data, 2024

if the R<sup>2</sup>value=0.33 the exogenous influence on the endogenous is moderate, if the R<sup>2</sup>value=0.67 the exogenous influence on the endogenous is strong. Next, hypothesis testing is carried out to find out the significance of the influence of the observed research variables by observing the path coefficient value. The influence of the level of obesity, smoking, and demographics on vitamin D, with an R<sup>2</sup>value of 0.782, means that the endogenous variable vitamin D is influenced by 78.2% by the exogenous variables the level of obesity, smoking, and demographics, while 21.8% is influenced by other factors outside the variable. It can be concluded that the influence of exogenous variables on endogenous variables is strong. The influence of the level of obesity, smoking, demographics, vitamin D on lung function with an R<sup>2</sup>value of 0.190, meaning that the endogenous variable lung function is influenced by 19% by the exogenous variable the level of obesity, smoking, demographics, and vitamin D, while 81% is influenced by other factors in outside the variables studied. It can be concluded that the influence of exogenous variables on endogenous variables is weak.

Next, the evaluation of the structural model with path coefficients was carried out in three stages: The first stage, namely checking the absence of multicollinearity between variables and the inner VIF (Variance Inflated Factor). If the estimation results show an inner VIF value <5, then the level of multicollinearity between variables is low. All indicators have an inner VI value <5, so the estimates of all variables and indicators in SEM-PLS are not robust (not biased). The second stage, hypothesis testing, is carried out between variables by looking at the

t<sub>statistic</sub> value and P<sub>value</sub>. If the t<sub>statistic</sub> = calculation result is greater than the t<sub>table</sub> (2.0166) and the P<sub>value</sub> of the test results is <0.05, then there is a significant influence between the variables. The third stage, analysis of the results of the f-square value is carried out, namely the influence of variables at the structural level with the criteria f-square 0.02 ≤ no effect, 0.02 ≤ f-square ≤ 0.14 small effect, 0.15 ≤ f-square ≤ 0.35 has a medium effect, and >0.35 has a high effect. The path coefficient assesses the magnitude of the direct influence of exogenous variables on endogenous variables; the magnitude of the influence ranges from -1 (negative influence) to +1 (positive influence).


Hypothesis testing is not only direct observation between exogenous variables and endogenous variables, but also tested indirectly through mediating variables (vitamin D). The SEM-PLS model can also be useful as a predictor in developing strategies to improve health promotion related to the influence of obesity, smoking, and demographic factors on lung function and vitamin D levels directly, as well as on lung function indirectly through vitamin D levels in active smokers. Education provided in terms of increasing vitamin D (25(OH)D) levels is by reducing body weight through BMI indicators and reducing the frequency of smoking. It also applies to implementing education to improve lung function by improving or increasing vitamin D levels (Figure 2).

Body mass index (BMI) is the ratio of body weight to height squared. Most of the respondents had normal BMI values, namely 29 respondents (61.70%) and safe limit abdominal

Table 4. Structural Model Evaluation Results, consisting of Research Hypothesis Testing and Effect Size (f-square)

	Hypothesis	Path coefficient	t <sub>statistic</sub>	P <sub>value</sub>	f-square	Interpretations
Direct Hypothesis	Is there a direct effect of obesity level on vitamin D in active smokers?	0.908	9.497	0.000	2.889	There is a direct influence of the level of obesity on vitamin D levels in active smokers. The significant influence of the level of obesity on vitamin D has a high influence value.
	Is there a direct effect of the level of obesity on lung function in active smokers?	-0.090	1.076	0.283	0.030	There is no direct effect of the level of obesity on lung function in active smokers. The obesity level variable on vitamin D has a moderate influence value.
	Is there a direct effect of smoking on vitamin D in active smokers?	0.030	0.265	0.791	0.003	There is no direct effect of smoking on vitamin D in active smokers. The cigarette variable on vitamin D has a small influence value.
	Is there an effect of smoking on lung function in active smokers?	0.815	0.512	0.609	0.161	There is no direct effect of smoking on lung function in active smokers. The smoking variable on lung function has a small influence value.
	Is there a direct influence of demographics on vitamin D in active smokers?	-0.319	1.229	0.220	0.098	There is no direct influence of demographics on vitamin D in active smokers. Demographic variables on vitamin D have no influence.
	Is there a direct influence of demographics on lung function in active smokers?	0.122	0.654	0.513	0.016	There is no direct influence of demographics on lung function in active smokers. Demographic variables on lung function have no influence value.
	Is there a direct effect of vitamin D on lung function in active smokers?	-0.791	0.500	0.617	0.168	There is no direct effect of vitamin D on lung function in active smokers. The vitamin D variable on lung function has a moderate influence value.
Indirect Hypothesis	Is there an effect of obesity level on lung function through vitamin D levels in active smokers?	-	0.256	0.798	-	There is no influence of obesity level on lung function through vitamin D levels in active smokers. Vitamin D does not mediate the indirect relationship between obesity level and lung function.
	Is there an effect of smoking on lung function through vitamin D levels in active smokers?	-	1.087	0.277	-	There is no effect of smoking on lung function through vitamin D levels in active smokers. Vitamin D does not mediate the indirect relationship between smoking and lung function.
	Is there a demographic influence on lung function through vitamin D levels in active smokers?	-	0.454	0.650	-	There is no demographic influence on lung function through vitamin D levels in active smokers. Vitamin D did not mediate the indirect relationship between demographic variables and lung function.

Source: Primary Data, 2024



Source: Primary Data, 2024

Figure 2. SEM-PLS Model After Bootstrapping with  $P_{\text{value}}$  = between Variables Based on SmartPLS analysis

circumference of 38 respondents (80.85%). It is supported by the results of research in Indonesia which analyzed the BMI picture and found that the majority of respondents had a normal BMI of 45%. 75 Another study in Indonesia which aimed to find out the factors related to the incidence of central obesity in adults found that there was a prevalence Central obesity with excessive abdominal circumference based on age 25-34 years (22.9%) and 35-44 years (33.5%), indicates that there are more respondents with abdominal circumference within safe limits. The results of this study found 4 respondents (8.51%) in the peripheral obesity category ( $\text{BMI} > 27 \text{ kg/m}^2$ ) and 9 people (19.14%) with central obesity (abdominal circumference  $> 90 \text{ cm}$ ). Based on the evaluation of the structural model (inner model) using SmartPLS version 3, there was a significant direct influence between the obesity level variable on vitamin D levels ( $P_{\text{value}} < 0.05$  and  $t_{\text{statistic}} > t_{\text{table}}$ ) in active smokers. It is supported by a theory that states the relationship between obesity mechanisms in causing a decrease in vitamin D levels, including three mechanisms that can explain the relationship between deficiency in vitamin D levels in obese individuals. Individuals who are obese experience decreased exposure to sunlight compared to non-obese individuals (Mirza *et al.*, 2022). The release of adiponectin from fatty tissues was inversely correlated with

body weight and BMI suggesting a link between vitamin D deficiency and insulin resistance (Kausar *et al.*, 2022).

The evaluation of the control model (inner model) found no direct significant influence between the obesity level variable on lung function ( $P_{\text{value}} > 0.05$  and  $t_{\text{statistic}} < t_{\text{table}}$ ) in active smokers. These results have conclusions that are the opposite of several theories which state that obesity harms the lung organs, one of which is the development of OHS which is defined as a combination of obesity ( $\text{BMI} \geq 30 \text{ kg/m}^2$ ), hypercapnia (arterial  $\text{CO}_2 \geq 45 \text{ mmHg}$ ) and the presence of breathing disorders during sleep which causes alveolar hypoventilation (Masa *et al.*, 2019; Palma *et al.*, 2022). Research results that are not in accordance with theory or previous research can be caused by only a small portion of respondents being classified as obese, whereas based on lung function examinations also the majority of respondents (44.68%) included in the moderate obstruction category. Individuals who are obese show a decrease in lung volume and capacity when compared to people with a normal body weight, this is because in people with a normal BMI the diaphragm contracts to push the contents of the stomach down and forward without any obstacles from excess fat deposits, when Likewise, contraction of the external intercostal muscles (muscles that play a role in



the inspiration process) can push the ribs up and forward without resistance (Svartengren *et al.*, 2020). Therefore, in this study, the majority of respondents had a normal BMI and the majority were in the obstruction category, moderate (27.65%), because the majority of respondents had breathing patterns that were not influenced by obesity.

Most of the smokers were moderate smokers (57.44%). Evaluation of the structural model (inner model) found that there was no direct significant influence between smoking variables on vitamin D levels ( $P_{value} > 0.05$  and  $t_{statistic} < t_{table}$ ) in active smokers. Smoking, ultraviolet radiation, and age are considered important factors that contribute to the skin aging process in humans by increasing wrinkles on the skin surface. In addition, tobacco smoke can affect the expression level of vitamin D receptors. If the expression of vitamin D receptors increases, the physiological effects of vitamin D will be better, one of which is the effect on lung function (Yang *et al.*, 2021). Most of the respondents fell into the moderate level with IB. Based on the evaluation of the inner control model using SmartPLS, it was found that there was no direct significant influence between smoking variables on lung function ( $P_{value} > 0.05$  and  $t_{statistic} < t_{table}$ ) in active smokers. These results have conclusions that are the opposite of several theories, which state that cigarettes contain various dangerous substances and their pathophysiological mechanisms, which can disrupt lung function. Nicotine was found to be chemotactic for human neutrophils, neutrophils being the first cells recruited in the process of lung inflammation due to cigarette smoke, which ultimately causes airway obstruction (Ham *et al.*, 2022). Cigarettes cause airway inflammation, which occurs more precisely in the bronchioles, causing lung remodeling. - The lungs lose their elasticity during the air exchange process, which then results in chronic obstructive airway limitations (Karnati *et al.*, 2021). Inappropriate research results can be caused by other factors or variables that are not observed but can influence, among others, genetics, physical activity, food intake, knowledge, attitudes, and behavior, sun exposure, skin color, and air pollution.

Patient demographics, which include age and level of education, are one of the factors that have been widely studied and influence the reduction in vitamin D levels. Based on the evaluation of the structural model (inner model), the results found that there is no direct significant influence between demographic variables on vitamin D levels ( $P_{value} > 0.05$  and  $t_{statistic} < t_{table}$ ) in active smokers. These results are aligned with the theory regarding demographic relationships, whether studied in terms of age or education level, which can influence vitamin D levels. In terms of age, physiologically an individual can experience a decrease in kidney function, resulting in reducing the production of the active metabolite 1,25(OH)D with increasing age, due to a decline in the activity of the kidney enzyme 1 $\alpha$ -hydroxylase which converts 25(OH)D to 1,25(OH)D. Serum 1,25(OH)D levels are inversely proportional to serum creatinine levels and proportional to glomerular function rate (GFR). Aging is not only associated with a decrease in kidney function, but is also associated with a decrease in vitamin D production in the skin, in the form of a decline in the concentration of 7-dehydrocholesterol in the epidermis and a reduced response to UVB light exposure in elderly individuals compared to young adults, resulting in a decrease in the formation of previtamin D by 50% (Turner *et al.*, 2022; Huish *et al.*, 2021). The higher the level of education, the greater the knowledge and awareness of the importance of adequate vitamin D nutrition (Hamhoum & Aljefree, 2022). Therefore, this is a limitation of this research because there are several factors that were not observed but tend to influence vitamin D levels, including the influence of physical activity, knowledge and attitudes towards vitamin D, and genetics.

Age and education level are also considered factors that influence lung function. Based on the evaluation of the inner control model using SmartPLS, there was no direct significant influence between demographic variables on lung function ( $P_{value} > 0.05$  and  $t_{statistic} < t_{table}$ ) in active smokers. These results have conclusions that are the opposite of several theories, which state that demographics, both in terms of age and level of education, can influence lung function, especially in active



smokers. Other research results show that the older a person is, the lower the FEV1 value will be. In particular, the smaller the FVC value due to advanced age, the more significant the decrease in FEV1/FVC (Thomas *et al.*, 2019). It is different from the results of this study, where there was no influence of demographics on lung function. This result can be caused by factors that influence these two variables (demography and lung function), each of these two variables can be influenced by smoking factors and the level of obesity studied in this study to factors that were not examined in this study but can be contribute to changes in lung function such as physical activity, knowledge, attitudes and behavior, etc. as previously explained.

The largest number of respondents belonged to the vitamin D insufficiency and moderate obstruction group (25.53%). Based on the evaluation of the inner control model, it was found that there was no direct significant influence of the vitamin D variable on lung function ( $P_{\text{value}} > 0.05$  and  $t_{\text{statistic}} < t_{\text{table}}$ ) in active smokers. These results have conclusions that are the opposite of several theories, which state that vitamin D can affect lung function, especially in active smokers. It is not aligned with previous theory and research, which states that there is an inverse relationship between vitamin D levels and lung function. Previous research by Ganji *et al.* (2020) involved 11,983 respondents aged  $\geq 20$  years. This study combines three NHANES data from 2007-2008, 2009-2010, and 2011-2012. The results of the study concluded that serum 25(OH)D levels were directly related to FVC and FEV1; in other words, serum 25(OH)D levels were associated with improved lung function values in healthy people, but not with the prevalence of asthma, emphysema, and chronic bronchitis. Other research also supports previous research, such as Wannamethee *et al.* (2021), who used a prospective cohort design involving 3575 male respondents (60-79 years), concluded that male respondents with COPD tend to experience vitamin D deficiency when compared to male respondents with normal lung function. In contrast to the results of this study, there was no effect of vitamin D on lung function. This result can be caused by factors that influence these two variables (vitamin D and lung function),

each of these two variables can be influenced by obesity, smoking and demographic factors studied in this study to factors that were not studied such as genetics, physical activity, food intake, knowledge, attitudes and behavior, sun exposure, skin color, air pollution.

The results of the evaluation of the structural model (inner model) found that there was no influence of the level of obesity on lung function through vitamin D levels in active smokers. Therefore, vitamin D does not mediate the indirect relationship between obesity levels and lung function. This result could be caused by factors that influence these two variables (vitamin D and lung function), each of these three variables can be influenced by smoking and demographic factors studied in this study to factors not studied such as genetics, physical activity, food intake, knowledge, attitudes and behavior, sun exposure, skin color, air pollution and so on, as previously explained.

This research still has several limitations, including other factors that cannot be controlled such as genetic factors, physical activity, knowledge, attitudes and behavior, intensity of exposure to sunlight, use of sunscreen, air pollution, food intake cannot be controlled in this research so that it can influence the research results. Current knowledge finds that a lack of physical activity is an important risk factor for vitamin D deficiency. Various observational studies show that maintaining vitamin D nutritional status is related to physical activity/exercise habits, where the level of physical activity is significantly positively correlated with 25(OH)D levels (Song *et al.*, 2020; Loreja *et al.*, 2022). Variations in sun exposure can cause vitamin D levels to fluctuate. Indoor activity may be a factor in these fluctuations and may cause a decrease in vitamin D levels. A systematic review and meta-analysis aimed at identifying whether physical activity or exercise indoors compared to outdoors has a significant effect on vitamin D levels through subgroup analysis and multivariate meta-regression (Bársan *et al.*, 2023).

### Conclusion

Obesity level factors (BMI) have a significant direct effect on vitamin D levels in active smokers. The demographic factors and smoking frequency do

not have a direct significant effect on vitamin D in active smokers. Demographic factors, obesity level, and smoking frequency do not have a significant direct effect on lung function in active smokers. Vitamin D has no direct significant effect on lung function in active smokers. Demographic factors, obesity level, and smoking frequency do not have a significant effect on lung function via vitamin D in active smokers. Therefore, it is recommended to maintain body weight and lose weight, especially in obese populations, to reduce the incidence of vitamin D deficiency. So it is necessary to carry out further research on the influence of other factors on vitamin D levels and lung function, such as genetic factors, physical activity, knowledge and attitudes, and behavior, intensity of exposure to sunlight, use of sunscreen, air pollution, and food intake that were not observed in this study to develop theories about the influence of broader factors.

#### Acknowledgment

This research was funded by the Research and Community Service Institute (LPPM) of Universitas Surabaya.

#### References

- Abi-Ayad, M., Nedjar, I., & Chabni, N., 2023. Association between 25-Hydroxy Vitamin D and Lung Function (FEV1, FVC, FEV1/FVC) in Children and Adults with Asthma: A Systematic Review. *Lung India*, 40(5), pp.449–56.
- Ahn, K.M., Kim, S.S., Lee, S.Y., Lee, S.H., & Park, H.W., 2021. Vitamin D Deficiency and Lung Function Decline in Healthy Individuals: A Large Longitudinal Observation Study. *Respir Med*, 182, pp.106395.
- Amrein, K., Scherkl, M., Hoffmann, M., Neuwersch-Sommeregger, S., Köstenberger, M., Tmava, B.A., Martucci, G., Pilz, S., & Malle, O., 2020. Vitamin D Deficiency 2.0: An Update on the Current Status Worldwide. *Eur J Clin Nutr*, 74(11), pp.1498–513.
- Bărsan, M., Chelaru, V.E., Răsnoveanu, A.G., Popa, Ș.L., Socaciu, A.I., & Bădulescu, A.V., 2023. Difference in Levels of Vitamin D between Indoor and Outdoor Athletes: A Systematic Review and Meta-Analysis. *Int J Mol Sci*, 24(8), pp.7584.
- Bishop, E.L., Ismailova, A., Dimeloe, S., Hewison, M., & White, J.H., 2020. Vitamin D and Immune Regulation: Antibacterial, Antiviral, Anti-Inflammatory. *JBM Plus*, 5(1), pp.e10405.
- Cao, Y., Li, P., Wang, Y., Liu, X., & Wu, W., 2022. Diaphragm Dysfunction and Rehabilitation Strategy in Patients with Chronic Obstructive Pulmonary Disease. *Front Physiol*, 13, pp.872277.
- CDC., 2021. GATS (Global Adult Tobacco Survey) Comparison Fact Sheet Indonesia 2011 & 2021. *Global Adult Tobacco Survey*, pp.1–2.
- Chung, C., Lee, K.N., Han, K., Shin, D.W., & Lee, S.W., 2023. Effect of Smoking on the Development of Chronic Obstructive Pulmonary Disease in Young Individuals: A Nationwide Cohort Study. *Front Med (Lausanne)*, 10, pp.1190885.
- Ganji, V., Al-Obahi, A., Yusuf, S., Dookhy, Z., & Shi, Z., 2020. Serum Vitamin D is Associated with Improved Lung Function Markers but not with Prevalence of Asthma, Emphysema, and Chronic Bronchitis. *Sci Rep*, 10(1), pp.11542.
- Ghosh, A.J., Moll, M., Hayden, L.P., Bon, J., Regan, E., & Hersh, C.P., 2020. Vitamin D Deficiency is Associated with Respiratory Symptoms and Airway Wall Thickening in Smokers with and without COPD: A Prospective Cohort Study. *BMC Pulm Med*, 20(1), pp.123.
- Ham, J., Kim, J., Ko, Y.G., & Kim, H.Y., 2022. The Dynamic Contribution of Neutrophils in the Chronic Respiratory Diseases. *Allergy Asthma Immunol Res*, 14(4), pp.361–78.
- Hamhoum, A.S., & Aljefree, N.M., 2022. Knowledge and Attitudes towards Vitamin D among Health Educators in Public Schools in Jeddah, Saudi Arabia: A Cross-Sectional Study. *Healthcare (Basel)*, 10(12), pp.2358.
- Herath, P., Wimalasekera, S., Amarasekara, T., Fernando, M., & Turale, S., 2022. Effect of Cigarette Smoking on Smoking Biomarkers, Blood Pressure and Blood Lipid Levels Among Sri Lankan Male Smokers. *Postgrad Med J*, 98(1165), pp.848–54.
- Huish, S.A., Jenkinson, C., Dunn, J.A., Meredith, D.J., Bland, R., & Hewison, M., 2021. Low Serum 1,25(OH)2D3 in End-Stage Renal Disease: is Reduced 1 $\alpha$ -Hydroxylase the Only Problem?. *Endocr Connect*, 10(10), pp.1291–8.
- Karnati, S., Seimetz, M., Kleefeldt, F., Sonawane, A., Madhusudhan, T., Bachhuka, A., Kosanovic, D., Weissmann, N., Krüger, K., & Ergün, S., 2021. Chronic Obstructive Pulmonary Disease and the Cardiovascular System: Vascular Repair and Regeneration as a Therapeutic Target. *Front Cardiovasc Med*, 8, pp.649512.
- Kauser, H., Palakeel, J.J., Ali, M., Chaduvula, P., Chhabra, S., Lamsal, L.S., Ramesh, V., Opara, C.O., Khan, F.Y., Kabiraj, G., & Mohammed, L., 2022. Factors Showing the Growing

- Relation Between Vitamin D, Metabolic Syndrome, and Obesity in the Adult Population: A Systematic Review. *Cureus*, 14(7), pp.e27335.
- Lorensia, A., Muntu, C.M., Suryadinata, R.V., & Septiani, R., 2021. Effect of Lung Function Disorders and Physical Activity on Smoking and Non-Smoking Students. *J Prev Med Hyg*, 62(1), pp.E89–E96.
- Lorensia, A., Suryadinata, R.V., & Inu, I.A., 2022. Comparison of Vitamin D Status And Physical Activity Related With Obesity in Student. *Journal of Applied Pharmaceutical Science*, 12(4), pp.108-18.
- Lorensia, A., Suryadinata, R.V., Rahmawati, R.K., & Septiani, R., 2024. The Effect of Smoking Habit on Vitamin D Status in Adults in Indonesia. *KEMAS*, 19(3), pp.410-421.
- Masa, J.F., Pépin, J.L., Borel, J.C., Mokhlesi, B., Murphy, P.B., & Sánchez-Quiroga, M.Á., 2019. Obesity Hypoventilation Syndrome. *Eur Respir Rev*, 28(151), pp.180097.
- Mirza, I., Mohamed, A., Deen, H., Balaji, S., Elabbahi, D., Munasser, A., Naquiallah, D., Abdulbaseer, U., Hassan, C., Masrur, M., Bianco, F.M., Ali, M.M., & Mahmoud, A.M., 2022. Obesity-Associated Vitamin D Deficiency Correlates with Adipose Tissue DNA Hypomethylation, Inflammation, and Vascular Dysfunction. *Int J Mol Sci*, 23(22), pp.14377.
- Palma, G., Soric, G.P., Genchi, V.A., Giordano, F., Caccioppoli, C., D'Oria, R., Marrano, N., Biondi, G., Giorgino, F., & Perrini, S., 2022. Adipose Tissue Inflammation and Pulmonary Dysfunction in Obesity. *Int J Mol Sci*, 23(13), pp.7349.
- Song, K., Park, G., Choi, Y., Oh, J.S., Choi, H.S., Suh, J., Kwon, A., Kim, H.S., & Chae, H.W., 2020. Association of Vitamin D Status and Physical Activity with Lipid Profile in Korean Children and Adolescents: A Population-Based Study. *Children (Basel)*, 7(11), pp.241.
- Stanojevic, S., Kaminsky, D.A., Miller, M.R., Thompson, B., Aliverti, A., Barjaktarevic, I., Cooper, B.G., Culver, B., Derom, E., Hall, G.L., Hallstrand, T.S., Leuppi, J.D., MacIntyre, N., McCormack, M., Rosenfeld, M., Swenson, E.R., 2022. ERS/ATS Technical Standard on Interpretive Strategies for Routine Lung Function Tests. *Eur Respir J*, 60(1), pp.2101499.
- Svartengren, M., Cai, G.H., Malinowski, A., Theorell-Haglöw, J., Janson, C., Elmstahl, S., Lind, L., Lampa, E., & Lindberg, E., 2020. The Impact of Body Mass Index, Central Obesity and Physical Activity on Lung Function: Results of the EpiHealth Study. *ERJ Open Res*, 6(4), pp.00214–2020.
- Thomas, E.T., Guppy, M., Straus, S.E., Bell, K.J.L., & Glasziou, P., 2019. Rate of Normal Lung Function Decline in Ageing Adults: A Systematic Review of Prospective Cohort Studies. *BMJ Open*, 9(6), pp.e028150.
- Tian, T., Jiang, X., Qin, R., Ding, Y., Yu, C., Xu, X., & Song, C., 2023. Effect of Smoking on Lung Function Decline in a Retrospective Study of a Health Examination Population in Chinese Males. *Front Med (Lausanne)*, 9, pp.843162.
- Tuckey, R.C., Cheng, C.Y.S., & Slominski, A.T., 2019. The Serum Vitamin D Metabolome: What we Know and what is Still to Discover. *J Steroid Biochem Mol Biol*, 186, pp.4–21.
- Turner, M.E., Rowsell, T.S., White, C.A., Kaufmann, M., Norman, P.A., Neville, K., Petkovich, M., Jones, G., Adams, M.A., & Holden, R.M., 2022. The Metabolism of 1,25(OH)<sub>2</sub>D<sub>3</sub> in Clinical and Experimental Kidney Disease. *Sci Rep*, 12(1), pp.10925.
- Wannamethee, S.G., Welsh, P., Papacosta, O., Lennon, L., & Whincup, P., 2021. Vitamin D Deficiency, Impaired Lung Function and Total and Respiratory Mortality in a Cohort of Older Men: Cross-Sectional and Prospective Findings from The British Regional Heart Study. *BMJ Open*, 11(12), pp.e051560.
- Weir, C.B., & Jan, A., 2024. BMI Classification Percentile And Cut Off Points. StatPearls. Treasure Island (FL): StatPearls Publishing.
- Yang, L., Zhao, H., Liu, K., Wang, Y., Liu, Q., Sun, T., Chen, S., & Ren, L., 2021. Smoking Behavior and Circulating Vitamin D Levels in Adults: A Meta-Analysis. *Food Sci Nutr*, 9(10), pp.5820–32.