



Research Article

Influence of historical use of antibiotics toward antibiotic resistance

Fawandi Fuad Alkindi^{[1]*}, Rika Yulia^[1], Fauna Herawati^[1], Abdul Kadir Jaelani^[2]

¹ Department of Pharmacy, Faculty of Pharmacy, University of Surabaya, Surabaya, Indonesia

² Regional General Hospital Bangil, Pasuruan, Indonesia

* Email: alkindifawandi@staff.ubaya.ac.id Telp: (+62) 896 6446 2224

ARTICLE INFO

Article History

Received March 20, 2019

Revised July 1, 2019

Accepted July 10, 2019

Keywords

Resistance

Antimicrobial

Infection

Prior antibiotic use

Bangil

Doi

10.22219/farmasains.v4i1.7901

ABSTRACT

A WHO's Global Surveillance of Antibiotic Resistance showed there is an increase of antimicrobial resistance in Asia between 2013 until 2014. Many studies showed that there is a correlation between prior antibiotic use with antimicrobial resistance case. The primary objective of this study is to analyze the relationship between prior antibiotic use with antimicrobial resistance. It was a retrospective and descriptive study conducted at Bangil Regional General Hospital. The data collected from the medical record and microbiological test from the patient at the internal ward. Chi-square analysis used for the statistic. This study showed that prior antibiotic use increased 0,399 bigger for antimicrobial resistance rate ($p=0,001$).

How to cite: Alkindi, F. F., Yulia, R., Herawati, F., & Jaelani, A. K. (2019). Influence of historical use of antibiotics toward antibiotic resistance. *Farmasains : Jurnal Farmasi dan Ilmu Kesehatan*, 4(1), 9-12. Doi: <https://doi.org/10.22219/farmasains.v4i1.7901>.

INTRODUCTION

Global Surveillance of Antibiotic Resistance conducted by the World Health Organization (WHO) shows an increase in the incidence of antibiotic resistance from 2013 to 2014 on the Asian continent. Southeast Asia is one region that has a high prevalence of antibiotic resistance and occurs in almost all classes of antibiotics (World Health Organization, 2013). Antibiotic resistance is the ability of a microorganism to fight or resist the effects of one or more antibiotics (World Health Organization, 2015). Resistance can occur in cases of different infectious diseases or bacteria.

Research conducted in Indonesia in the period 1998-1999 showed for *Salmonella spp.* experience resistance to several antibiotics such as ampicillin (7%), chloramphenicol (15%), tetracycline (20%), cotrimoxazole (10%), ceftriaxone (2%), kanamycin (10%), norfloxacin (1%), cephalothin (7%), neomycin (3%) (Lestari, Severin & Verbrugh, 2012). Several factors are thought to be the cause of resistance, including bacterial gene mutations, horizontal gene transfer between bacterial species, and inappropriate use of antibiotics.

In addition to the things mentioned above, several studies show that there is a relationship between

antibiotic resistance and antibiotic use over a certain period (prior antibiotic use) before the patient diagnosed with resistance. Research conducted by Kuster et al. (2014) showed that prior antibiotic use has a relationship with the occurrence of resistance in a patient at three months after the last time using the antibiotic in question.

Research on this subject is very little, so the authors are interested in observing this because knowing the presence or absence of the influence of prior antibiotic use on the occurrence of resistance can help the clinician in determining which therapy is more appropriate for the patient. The use of 3 months as a reference from the time of last antibiotic use was due to a study conducted by Kuster et al. (2014), which showed that the possibility of resistance from an antibiotic to return to baseline level was 90 days after the last dose gave to patients.

RESEARCH METHODS

This study is retrospective and descriptive. Data obtained from medical records and bacterial culture results of all internal medicine patients in Bangil Regional General Hospital in July to December 2016.

Based on medical records obtained and then traced to patients diagnosed with an infection and using antibiotics so that patients who had infections but did not get antibiotics and outpatients included in the exclusion criteria in this study. Patients who are known to have an infection and get antibiotics will then be traced to a history of antibiotic use for three months before the patient is admitted to the hospital and observed the occurrence of resistance in these patients based on the results of the culture.

The analytical method for proving the relationship between antibiotic use after patients hospitalized with resistance was tested statistically using the chi-square method. The results of processing the next data will show in tables and diagrams.

RESULT AND DISCUSSIONS

Patients who were hospitalized in the internal hospital ward Bangil in July to December 2016 reached 4303 people with the highest number of patients, namely 846 people in November and the lowest as many as 600 people in July. The total patients through BPJS funding were 2348 patients (54.57%), private insurance as many as 4 patients (0.093%), GAKIN APBD 90 patients (2.1%), Jamkesmas 2 patients (0.046%), SPM 1247 patients (28.98%), and general as many as 612 patients (14.22%).

The value of 54.57% of patients whose funding

Table 1. Antibiotic sensitivity profile in the group of patients with a history of antibiotic use over the past three months.

Culture Diagnosis	Total
The complication of procedures, not elsewhere classified	33
Infection of obstetric surgical wound	30
Pneumonia	23
Osteomyelitis	12
Cutaneous abscess	9
Type 2 diabetic with gangrene	7
Bacteremia	2
Vulvovaginitis	2
Rheumatic heart disease	2
Abses submandibular	1
Acute pharyngitis	1
Burn of an unspecified body region	1
Corneal ulcer	1
Diare and gastroenteritis presume infectious origin	1
Fracture	2
skin ulcer	1
Urine tract infection, site not specified	1
Staphylococcal arthritis elbow	1
Sepsis due to E.coli	1
Cellulitis	2
Nontoxic goiter	1
Meningitis	1
Laryngeal diphtheria	1
Intracranial injury, unspecified	1
Impetigo bullous	1
Generalized skin eruption due to drugs and medicaments taken internally	1
Total	140

through BPJS resembles the coverage of BPJS in East Java Province which amounted to 59.16% in 2016 (Kementrian Kesehatan Republik Indonesia [Kemenkes RI], 2017). The number of patients whose funding through BPJS on inpatient wards also resembled the coverage of BPJS participation in 2015 in East Java Province which amounted to 55.95% (Kemenkes RI, 2017).

The most common diagnosis of the cause of culture is Complication of procedures, not elsewhere classified as many as 33 patients followed by infection in the surgical wound which also includes cesarean surgery as many as 30 patients then pneumonia as many as 19 patients.

Table 2. Antibiotic sensitivity profiles in the group of patients with a history of antibiotic use over the past three months.

Antibiotics	Sensitivity	
	Sensitive	Resistent
Amikacin	1	-
Amoxicillin	-	-
Ampisillin	-	-
Azythromycine	-	-
Benzathine Penicillin	-	-
Metronidazole	-	-
Cefadroxil	-	-
Cefazolin	1	-
Cefixime	-	-
Ceftazidime	6	3
Ceftriaxone	2	-
Cefuroxime	1	-
Ciprofloxacin	1	1
Doxyciline	-	1
Gentamicin	3	-
Total	15	5

Table 3. Antibiotic sensitivity profile in the patient group without a history of antibiotic use in the last three months.

Antibiotic Class	Sensitivity	
	Sensitive	Resistent
Penicillin	203	352
Cephalosporin	146	336
Aminoglycoside	76	117
Macrolide	34	40
Amphenicol	20	18
Carbapenem	59	37
Lincosamide	25	20
Streptomycin	-	-
Imidazole	-	-
Anti TBC	18	6
Quinolone	155	257
Others	170	118
Total	906	1301

Observations were made regarding the history of antibiotic use by patients over the past three months. Of the 140 samples, there were 78 samples where the patient had a history of antibiotic use in the last three months before the culture was carried out and the remaining 62 patients had no history of

antibiotic use. Patients who have a history of antibiotic use in the last three months have observed sensitivity. From the results of these observations for ceftazidime, antibiotics had the highest number of sensitive results (6 samples) while for antibiotics that had the highest resistance level was cefazolin (14 samples). Cefadroxil is the most commonly found antibiotic in the history of antibiotic use with as many as 35 samples, but there is no sensitivity data from the antibiotic.

From patients who did not have a history of antibiotic use, meropenem became an antibiotic with the highest number of sensitive samples as many as 59 samples then followed by piperacillin + tazobactam and norfloxacin (52 samples) and cotrimoxazole (44 samples)

Based on observations in groups with a history of use, antibiotics that showed sensitive results found in 15 samples and 54 samples showed resistance. The antibiotics that have the most number of sensitive samples are ceftazidime with six samples while for antibiotics, which show that the most resistant amount is cefazolin with 14 samples. Sensitivity for groups that do not have history is as many as 906 antibiotic samples tested showing 1301 sensitive and resistant results. The antibiotics that showed the most sensitive results were meropenem with 59 samples and Ampicillin, showing the most resistant results as many as 87 samples. Antibiotic sensitivity from history groups with no history showed significantly different results where the likelihood of antibiotic resistance was higher in the history group compared to the group without history. It indicated by the calculation of OR from the sensitivity of the two groups where the possibility of antibiotic resistance in patients who have a history of antibiotic use is 0.399 times greater than patients without a history of antibiotics ($p = 0.001$).

It supported by research conducted by Kuster et al. (2014), which showed that antibiotic resistance in patients was caused more by patients having a history of antibiotic use before being hospitalized. Penicillin, cephalosporins, and quinolones decreased their resistance faster than with macrolide. Antibiotic resistance other than the macrolide can reach the baseline level after 90 days after the last antibiotic administration. Cephalosporin antibiotics mentioned in the study took 17 to 65 days after the last administration to return to sensitivity in pneumonia patients, whereas macrolide such as azithromycin require a longer time of 49-78 days to reach back to painful conditions. It needs between

each antibiotic to reach the half-life of the plasma (Bolon, 2009; Ferrero et al., 1990; Foulds, Shepard & Johnson, 1990; Humbert, Spyker, Fillastre & Leroy, 1979; Maholtra et al., 2007) influences the baseline.

Other studies that support this idea include the results of a study conducted by Chung et al., (2007), wherein the bacterium *Haemophilus influenza*, the resistance level of new amoxicillin will approach the baseline after > 12 weeks. In other research, the old, low dose and the last time used for amoxicillin in urinary tract infections can affect the level of resistance, which will only return at baseline after 1-3 months (Hillier et al., 2007). Other studies say that considering the last time given antibiotics themselves with what antibiotics will give next referring to antibiotics requires time to replenish the mucous membrane of the bacteria so that the amount will exceed the number of resistant strains of the bacteria (Levin & Rozen, 2006).

CONCLUSION

This study does have limitations related to the lack of data in the medical record related to the history of antibiotic use in hospitalized patients but based on calculations comparing the antibiotic profile in patients with and without a history of antibiotic use for the last 3 months shows that patients who have a history of antibiotic use over the past 3 months has been shown to increase the likelihood of higher resistance compared to patients without a history of antibiotic use over the past 3 months

REFERENCES

- Bolon, M. K. (2011). The newer fluoroquinolones. *The Medical clinics of North America*, 95(4), 793-817.
- Chung, A., Perera, R., Brueggemann, A. B., Elamin, A. E., Harnden, A., Mayon-White, R., ... & Mant, D. (2007). Effect of antibiotic prescribing on antibiotic resistance in individual children in primary care: prospective cohort study. *BMJ*, 335(7617), 429.
- Ferrero, J. L., Bopp, B. A., Marsh, K. C., Quigley, S. C., Johnson, M. J., Anderson, D. J., ... & Cavanaugh, J. H. (1990). Metabolism and disposition of clarithromycin in man. *Drug Metabolism and Disposition*, 18(4), 441-446.
- Foulds, G., Shepard, R. M., & Johnson, R. B. (1990). The pharmacokinetics of azithromycin in human serum and tissues. *Journal of Antimicrobial Chemotherapy*, 25(suppl_A), 73-82.
- Hillier, S., Roberts, Z., Dunstan, F., Butler, C., Howard, A., & Palmer, S. (2007). Prior antibiotics and risk of antibiotic-resistant community-acquired urinary tract infection: a case-control study. *Journal of antimicrobial chemotherapy*, 60(1), 92-99.
- Humbert, G., Spyker, D. A., Fillastre, J. P., & Leroy, A. (1979). Pharmacokinetics of amoxicillin: dosage nomogram for patients with impaired renal function. *Antimicrobial agents and chemotherapy*, 15(1), 28-33.
- Kementerian Kesehatan RI. 2017. Data dan Informasi Profil Kesehatan Indonesia 2016. Jakarta, Indonesia: Author.
- Kuster, S. P., Rudnick, W., Shigayeva, A., Green, K., Baqi, M., Gold, W. L., ... & Simor, A. E. (2014). Previous antibiotic exposure and antimicrobial resistance in invasive pneumococcal disease: results from prospective surveillance. *Clinical Infectious Diseases*, 59(7), 944-952.
- Lestari, E. S., Severin, J. A., & Verbrugh, H. A. (2012). Antimicrobial resistance among pathogenic bacteria in Southeast Asia. *Southeast Asian Journal of Tropical Medicine & Public Health*, 43(2), 385-422.
- Levin, B. R., & Rozen, D. E. (2006). Non-inherited antibiotic resistance. *Nature Reviews Microbiology*, 4(7), 556.
- Malhotra-Kumar, S., Lammens, C., Coenen, S., Van Herck, K., & Goossens, H. (2007). Effect of azithromycin and clarithromycin therapy on pharyngeal carriage of macrolide-resistant streptococci in healthy volunteers: a randomised, double-blind, placebo-controlled study. *The Lancet*, 369(9560), 482-490.
- World Health Organization. (2013). The world health report 2013: research for universal health coverage. Geneva, Switzerland: Authors.
- World Health Organization. (2015). Global Antimicrobial Resistance Surveillance System: Manual for Early Implementation. Geneva, Switzerland: Authors.

Farmasains : Jurnal Farmasi dan Ilmu Kesehatan

[HOME](#)[ABOUT](#)[LOGIN](#)[REGISTER](#)[SEARCH](#)[CURRENT](#)[ARCHIVES](#)[ANNOUNCEMENTS](#)

[Home](#) > [Farmasains : Jurnal Farmasi dan Ilmu Kesehatan](#)

Farmasains : Jurnal Farmasi dan Ilmu Kesehatan

Journal Title : Farmasains : Jurnal Farmasi dan Ilmu Kesehatan
Initials : Farmasains
Abbreviation : Farmasains. J. Farm. Il. Kes
Frequency : 2 issues per year (April and November)
DOI : 10.22219
Print ISSN : 2086-3373
Online ISSN : 2620-987X
Editor-in-Chief : Siti Rofida
Managing Editor : Amalyah Dina Anggraeni and M. Artabah Muchlisin
Publisher : Universitas Muhammadiyah Malang
Citation Analysis : [Google scholar](#)

Farmasains is a scientific publication for widespread research and criticism topics in pharmacy science studies. Farmasains is published twice a year in April and November since June 2010 by Department of Pharmacy University of Muhammadiyah Malang. One volume of Farmasains is published in two year calendar.

Visitors

	ID 1,909
	US 365
	KR 11
	IN 8
	MY 6
	ZA 3
	UA 3
	RU 3
	SG 3
	HK 2

5,957

 FLAG counter



Editorial Team

Editor in Chief

Siti Rofida, University of Muhammadiyah Malang, Indonesia

Managing Editors

Amaliyah Dina Anggraeni, Department of Pharmacy, Faculty of Health Science, University of Muhammadiyah Malang, Malang, East Java, Indonesia

M. Artabah Muchlisin, M.Farm, Department of Pharmacy, Faculty of Health Science, University of Muhammadiyah Malang, Malang, East Java, Indonesia

Reviewers

Dr. Susi Ari Kristina, S.Farm., M.Kes., Apt., Department of Pharmaceutics, Faculty of Pharmacy, Universitas Gadjah Mada, Yogyakarta, Indonesia, Indonesia

Dr. Roihatul Mutiah, SF., M. Kes., Apt., Department of Pharmacy, Faculty of Science and Technology, Islamic State Maulana Malik Ibrahim University, Malang, East Java, Indonesia, Indonesia

Anita Sukmawati, M.Si., Apt., Ph.D., Universitas Muhammadiyah Surakarta, Solo, Centre of Java, Indonesia

Helmy Yusuf, S.Si., M.Sc., Apt., Ph.D., Department of Pharmaceutics, Faculty of Pharmacy, University of Airlangga, Surabaya, East Java, Indonesia

Drs. Didik Hasmono, MS., Apt., Department of Clinical Pharmacy, Faculty of Pharmacy, University of Airlangga, Surabaya, East Java, Indonesia

Author Guidelines

Abstract is a brief description of the entire writing, containing the main issues discussed, the reasons for the research, the objectives, methods, and results of the study. Abstract is written not more than 250 words in which there are no tables / images and libraries (Cambria size 11, space 1)

KEYWORDS: Keyword-1, Keyword-2, Keyword-3, Keyword-4, Keyword-5.

Use semicolon (;) as a separator

INTRODUCTION (11 pt)

Article titles are written in size 18 and use the Cambria font type. The beginning of a word starts with a capital letter except for conjunctions beginning with a lowercase letter. Writing the Latin names of plants in the title with a capital letter at the beginning of the first word (genus), lowercase letters in the second word (Species) and tilted (italic).

In this section, the author must contain the background, issue / problem / urgency / rationalization, and explain the phenomenon of the actual problem under study, supported by references and the results of previous related studies that have been done by themselves or others, and explain the existence of research in that context . It is better not to contain detailed literature searches or summary results, not a review of the subject area, but a clear statement must be written about the question being discussed. The introduction must be designed in a rational sentence and the reader can know the importance of research conducted by the author.

The scientific text is written in 5-10 pages. All sources referenced or quoted must be written in the bibliography. Introduction without using the section title and written using the special first line 1 cm paragraph format. Do not give paragraph spacing between adjacent paragraphs.

(Cambria size 11, space 1)

II. RESEARCH METHOD (11 pt)

The method contains an explanation of the type or nature of research, data sources, data, data collection techniques, data collection instruments, data collection procedures, and data analysis methods

The research method must be well explained, so that readers can repeat the experiment that has been done by the author. All tools, instruments, chemicals, etc. must include the name of the product and the manufacturer. Genes, strains, age, and average body weight of the test animals must be clearly stated in this section. Plant samples must be written from the origin and authenticity.

(Cambria size 11, space 1)

RESULTS AND DISCUSSION (11 pt)

The results and discussion must answer the problems and objectives of the study. The results are written concisely and clearly. Can be presented in the sub-section, without using numbers. The title of the section is written in capital letters at the beginning of the word and bold. If in the sub-section there are sub-sections, the writing of the sub-subab title is written in capital letters at the beginning of the word tilted (italic). Writing paragraphs in each sub-section uses the 1cm special first line paragraph format.

Tables, Graphs and Images.

Tables, graphs, pictures, and / or photos (if any) do not occur overlapping in their compilation, given numbers, titles, and complete information and are quoted in the text. Tables, graphs, images, and / or photos are numbered in the order they appear. Tables and images must be clearly legible and can be printed properly because the text will be printed in a black and white format (for writers who want proof of print). Images, tables, graphics, photos must be placed as close as possible to the related text. Referral, quotation, or the inclusion of images, tables, and so on using numbering, not with the words "as follows", "as below", and so on.

Picture

Writing the title of the image is under the picture in left. Diagrams are classified in the form of images, therefore the rules for giving a description of a diagram are adjusted to the rules of writing.

Table

Inclusion of tables / data that are too long (more than one page) should be avoided. Tables only use horizontal lines or minimize the use of vertical lines. Each table column must be given a heading.

Interpretation of results must be written in the discussion of the research article or in some circumstances it may be desirable to combine the results and discussion in one section. Writing paragraphs on the results and discussion is written using a special first line paragraph format 1 cm.
(Cambria size 11, space 1)

IV. CONCLUSION (11 pt)

Conclusions must answer the problems and objectives of the study. Conclusions are not summaries nor are they rewritten from the discussion. Writing paragraphs in conclusions using the special first line 0.8 cm paragraph format. (Times New Roman size 12, space 1).

REFERENCES (11 pt)

The references referred to are at least 10 libraries, with a composition of 70% being primary sources and should come from the results of research, ideas, theories or concepts that have been published in journals, both printed and electronic. The 30% reference referred to from the book is the result of the publication of the last 10 years, with the exception of classical references which are used as material for historical studies.

Writing bibliography and quotations using the APA or American Psychological Association style. In addition, for writing bibliography and citations it is recommended to use the Mendeley or Zootero application.

Library taken from journal publications

Writer's name. (year). Title of article in journal publication. Publisher Name italicized, Volume (issue), page.

Example:

Hasson, S. S. A. A., Al-Busaidi, J. K. Z., & Sallam, T. A. (2015). The past, current and future trends in DNA vaccine immunisations. *Asian Pacific Journal of Tropical Biomedicine*, 5 (5), 344-353. [https://doi.org/10.1016/S2221-1691\(15\)30366-X](https://doi.org/10.1016/S2221-1691(15)30366-X)

Library taken from books

Author's name. (year). The title of the book is italicized (edition). City: Name of Book Publisher.

Example:

Poedjiadi, A. (1994). *Basics of Biochemistry (xi)*. Jakarta: University of Indonesia Publisher.

Library taken from the Website Page

MOH, R. (2014). Eliminate the Myth About Cancer. Retrieved from <http://www.depkes.go.id/article/print/201407070001/hulir-mitos-tentang-kanker.html> (Accessed August 21, 2018)

[Home](#) > [Archives](#) > **Vol 4, No 1 (2019)**

Vol 4, No 1 (2019)

DOI: <https://doi.org/10.22219/farmasains.v4i1>

Sedang Dlaam Proses

Table of Contents

Medical Plant Inventory of Dayak Benuaq Tribe Society Intu Lingau Village, Nyuatan Subdistrict, Kutai Barat, Kalimantan Timur

Indah Lestari, Lailiyatus Syafah

Abstract views : 8



History Effect of Antibiotic Use on Resistance

Fawandi Fuad Alkindi, Rika Yulia, Fauna Herawati, Abdul Kadir Jaelani

Abstract views : 7



Clinical Study of The Efficacy and Safety of Herbal Medicine for Hyperuricemia

Agus Triyono

Abstract views : 19



Method Validation of Sodium Benzoat Levels in Fruit Juice Drinks Circulating in Malang

Engrid Juni Astuti, Roha Fakhri Naufal Ilham, Januar Rahman

Abstract views : 14



Quantitative Struture-Cytotoxic Activity Relationship of Phenylthiourea derivatives from ChemBL database on Sirtuin-1 receptor by in silico

Anindi Lupita Nasyanka

Abstract views : 10



Study of In Silico of COX-2 on Indomethacin and Diclofenac as Nonsteroidal Antiinflammatory Drugs (NSAID)

Janatun Na'imah

Abstract views : 8



Gel Formulations Containing Sumbawa's Horse Milik With Carbomer as Gelling Agent

Dian Ermawati, Uswatun Chasanah, Herika Juli Andani, Khaldi Nisrina

Abstract views : 15



Business Strategy PT. XYZ in Marketing CITric Acid For Food and Beverage Ingredients Industry

Agus Hartanto

Abstract views : 7



History Effect of Antibiotic Use on Resistance

Fawandi Fawad Aikindi, Rika Yulia, Fauna Herawati, Abdul Kadir Jaelani



Abstract

A WHO's *Global Surveillance of Antibiotic Resistance* showed there is an increase of antimicrobial resistance in Asia between 2013 until 2014. Many studies showed that there is a correlation between prior antibiotic use with antimicrobial resistance case. The primary objective of this study is to analyze the relationship between prior antibiotic use with antimicrobial resistance. It was a retrospective and descriptive study which conducted at Bangil regional general hospital. The data collected from the medical record and microbiological test from the patient at the internal ward. Chi-square analysis used for the statistic. This study showed that prior antibiotic use increased 0,399 bigger for antimicrobial resistance rate ($p=0,001$)

Keywords

Resintention, Antibiotic, infection, Bangil

References

- World Health Organization. 2013. The world health report 2013: research for universal health coverage. Geneva, Switzerland: World Health Organization.
- World Health Organization. Global Antimicrobial Resistance Surveillance System: Manual for Early Implementation. 2015: WHO Library Cataloguing-in-Publication Data
- Lestari ES., Severin AJ., Verbrugh AH. Antimicrobial Resistance Among Pathogenic Bacteria In Southeast Asia. *Southeast Asian J Trop Med Public Health*. Vol 43 No. 2 March 2012
- Kuster SP, Rudnick W, Shigayeva A, Green K, Baqi M, Gold WL, et al. Previous antibiotic exposure and antimicrobial resistance in invasive pneumococcal disease: Results from prospective surveillance. *Clin Infect Dis*. 2014;59(7):944-52.
- Kementerian Kesehatan RI. Data dan Informasi Profil Kesehatan Indonesia 2016. Kementerian Kesehatan RI [Internet]. 2017;100. Available from: [http://www.depkes.go.id/resources/download/pusdatin/lain-lain/Data dan Informasi Kesehatan Profil Kesehatan Indonesia 2016 - smaller size - web.pdf](http://www.depkes.go.id/resources/download/pusdatin/lain-lain/Data%20dan%20Informasi%20Kesehatan%20Profil%20Kesehatan%20Indonesia%202016%20-%20smaller%20size%20-%20web.pdf)
- Malhotra-Kumar S, Lammens C, Coenen S, Van Herck K, Goossens H. Effect of azithromycin and clarithromycin therapy on pharyngeal carriage of macrolide-resistant streptococci in healthy volunteers: a 515 randomised, double-blind, placebo-controlled study. *Lancet* 2007; 369:482-90.
- Ferrero JL, Bopp BA, Marsh KC, et al. Metabolism and disposition of clarithromycin in man. *Drug Metab Dispos* 1990; 18:441-6.
- Foulds G, Shepard RM, Johnson RB. The pharmacokinetics of azithro- 540 mycin in human serum and tissues. *J Antimicrob Chemother* 1990; 25(suppl A): 73-82.
- Bolan MK. The newer fluoroquinolones. *Infect Dis Clin North Am* 2009; 23:1027-51.
- Humbert G, Spyker DA, Fillastre JP, Leroy A. Pharmacokinetics of 545 amoxicillin: dosage nomogram for patients with impaired renal function. *Antimicrob Agents Chemother* 1979; 15:28-33.
- Chung A, Perera R, Brueggemann AB, et al. Effect of antibiotic prescribing on antibiotic resistance in individual children in primary care: pro- spective cohort study. *BMJ* 2007; 335:429.
- Hillier S, Roberts Z, Dunstan F, Butler C, Howard A, Palmer S. Prior antibiotics and risk of antibiotic-resistant community-acquired urinary 535 tract infection: a case-control study. *J Antimicrob Chemother* 2007; 60:92-9.
- Levin BR, Rozen DE. Non-inherited antibiotic resistance. *Nat Rev Microbiol* 2006; 4:556-62.

DOI: <https://doi.org/10.22219/farmessains.v4i1.7901>