

Analysing Factors Contributing to Length of Stay (LoS) Among Patients with COVID-19 in Indonesian Hospital Setting

Sonny Edbert^{1,4}, Yosi Irawati Wibowo^{2,5*}, Adji Prayitno Setiadi^{3,6}

¹Pharmacy Department, Faculty of Mathematics and Natural Sciences, Universitas Lambung Mangkurat, Banjarbaru, South Kalimantan, Indonesia

²Pharmacist Professional Education Department, Faculty of Mathematics and Natural Sciences, Universitas Lambung Mangkurat, Banjarbaru, Kalimantan Selatan, Indonesia

³Pharmacy Department, Faculty of Pharmacy, Universitas Muhammadiyah Kalimantan Timur, Samarinda, East Kalimantan, Indonesia

⁴Faculty of Pharmacy, Universitas Surabaya, Surabaya, East Java, Indonesia

⁵Centre for Medicines Information and Pharmaceutical Care (CMIPC), Faculty of Pharmacy, Universitas Surabaya, Surabaya, East Java, Indonesia

⁶Department of Clinical and Community Pharmacy, Faculty of Pharmacy, Universitas Surabaya, Surabaya, East Java, Indonesia

Email: yosi_wibowo@staff.ubaya.ac.id

ABSTRAK

Penyakit COVID-19 masih menjadi ancaman konstan bagi kesehatan masyarakat global. Penggunaan obat dan lama rawat inap (LoS) merupakan indikator utama kualitas pelayanan di rumah sakit. Penelitian ini bertujuan untuk mengamati LoS dan faktor yang mempengaruhinya (termasuk karakteristik pasien dan penggunaan obat) pada pasien rawat inap COVID-19 di Rumah Sakit X di Surabaya, Indonesia. Penelitian observasional dilakukan dengan menggunakan rekam medis pasien rawat inap COVID-19 di Rumah Sakit X selama tahun 2022. Data karakteristik pasien dan profil obat dianalisis secara deskriptif. Kesesuaian terapi obat ditentukan berdasarkan pedoman terapi. Uji korelasi dilakukan untuk hubungannya dengan LoS. Total terdapat 203 pasien rawat inap COVID-19, meliputi: neonatus/anak/remaja (12,32%), dewasa (66,5%), dan lanjut usia (21,2%). Sebagian besar pasien laki-laki (54,2%) dengan keparahan sedang (80,8%), dan 36,4% memiliki komorbiditas. Kesesuaian terapi pada pasien neonatus/anak/remaja, dewasa, dan lanjut usia adalah: 12% (3/25), 76% (103/135), 60% (26/43), secara berurutan. Rata-rata LoS adalah 7,0±3,11 hari, dimana LoS terendah pada neonatus/anak/remaja (6,1±2,43 hari) dan tertinggi pada lansia (7,4±3,02 hari). Faktor yang secara signifikan mempengaruhi LoS adalah status vaksinasi ($p=0,000$) dan tingkat keparahan COVID-19 ($p=0,002$), namun tidak dipengaruhi oleh kesesuaian terapi ($p=0,864$). Hasil penelitian merupakan data awal untuk mengembangkan strategi perbaikan kualitas layanan untuk pasien COVID-19 rawat inap.

Kata Kunci: COVID-19, Faktor Risiko, Penggunaan Obat, Lama Rawat Inap, Rumah Sakit

ABSTRACT

COVID-19 disease remains a constant threat to global public health. Drug use and Length of Stay (LoS) are among main indicators of service quality in hospitals. Object of this research were understanding of LoS and the contributing factors (including patient characteristics and drug use) among COVID-19 inpatients at Hospital X Surabaya, Indonesia. Observational research was conducted using medical records of COVID-19 inpatients at Hospital X in 2022. Data on patient characteristics and drug profiles were analyzed descriptively. Drug therapy conformity was determined based on guidelines. Correlation tests were used to determine factors associated to LoS. The results were 203 COVID-19 inpatients, including: neonate/child/adolescent (12.32%), adults (66.5%) and elderly (21.2%). Most patients were males (54.2%) with moderate severity (80.8%), and 36.4% had comorbidities. Therapy conformity in neonate/child/adolescent, adult and elderly patients were: 12% (3/25), 76% (103/135), 60% (26/43), respectively. The average LOS was 7.0 ± 3.1 days, where neonate/child/adolescent was the lowest (6.1 ± 2.43 days), and elderly was the highest (7.4 ± 3.02 days). Significant contributing factors to LoS included vaccination status ($p = 0.000$) and COVID-19 severity ($p = 0.002$), but not with therapy conformity ($p = 0.864$). The study findings provide a preliminary data to guide the development of strategies to improve service quality for hospitalised patients with COVID-19.

Keywords: COVID-19, Risk Factors, Medication Use, Length of Stay, Hospital

I. INTRODUCTION

Coronavirus disease 2019 (COVID-19) is an infectious disease caused by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), and has been declared as a pandemic in March 2020 by the World Health Organization (WHO) (WHO, 2022a). WHO has categorized the SARS-CoV-2 virus into two groups, namely Variant of Interest (VOI) and Variant of Concern (VOC). In May 2021, WHO designated the Alpha (B.1.1.7), Beta (B.1.351), Gamma (P1) and Delta (B.1.671.2) variants as VOC variants; this was followed by the Omicron (B.1.1. 529) in November 2021 (Amalia, 2021; WHO,

2022). WHO designated Omicron as a VOC because of its high transmissibility and ability to avoid antibodies induced by vaccination or natural infection (Burhan & Rachmadi, 2022). As per 5 October 2022, the number of positive COVID-19 patients in Indonesia reached 6,439,292 cases with 158,165 deaths; while globally, there was more than 615,777,700 confirmed cases and 6,527,192 deaths (MoH-RI, 2021). Nowadays, even though COVID-19 is considered as manageable, the virus is capable of evolving and is going to remain a constant threat.

Hospitals are health facilities that have an important role in caring for

COVID-19 patients, especially in moderate to critical conditions (Burhan et al, 2022). In a pandemic situation, hospital burdens and health costs have increased significantly; but on the other hand, there are limited resources available in hospitals. Therefore, efficiency of hospital management warrants special attention. Length of Stay (LoS) is number of days in the hospital which is counted from the day of admission until discharge; and LoS is considered an important indicator for assessing the efficiency of hospital management and the quality of services provided to patients (quality patient care). The shorter the patient's length of stay, the more effective and efficient the service at the hospital is (Tai et al, 2020). In addition, shorter hospital stays reduce the burden of medical costs and increase bed turnover, and thereby increase hospital profit margins while lowering overall social costs (Guo et al, 2021). Therefore, it is important to map factors or characteristics that can shorten or lengthen patient LoS in hospitals.

To date, research evidence regarding factors influencing LoS in COVID-19 patients has been limited. In July 2022, the Centre for Disease Control and Prevention (CDC) released data on risk factors that influence the length of stay in hospitals for COVID-19 patients. Age has been considered as the strongest risk factor for worsening COVID-19. As of 23 May

2022 data, the proportion of deaths in the age group over 65 years was 42 times that of those aged 30-39 years. In addition, people from racial and ethnic minority groups have been at greater risk of hospitalization, ICU admission, and death from COVID-19 at younger ages. Comorbid groups that have had a high risk of worsening COVID-19 are asthma, cancer, cerebrovascular disease, chronic kidney or liver diseases, diabetes mellitus types 1 and 2, cardiovascular disorders, obesity, pregnancy and smoking (CDC, 2022). Apart from patient factors, research evidence shows that the quality of treatment and the type of drug chosen can also influence LoS in hospitals (Khosravizadeh et al, 2016), although this has not been specifically observed in COVID-19 patients.

Hospital X is a health facility with special ward for airborne infectious diseases, such as COVID-19, in Surabaya, East Java, Indonesia. While limited data is available on LoS and the contributing factors (i.e., patient characteristics and drug use) among COVID-19 patients in Indonesia, this research is conducted to provide a preliminary data to guide stakeholders and professional health workers in allocating health resources more effectively and efficiently, as well as in developing strategies to optimize the quality of services for hospitalised patients

with COVID-19, which remains a constant public health issue in Indonesia.

II. METHOD

A. Study design and Setting

This is an observational study conducted in Hospital X in Surabaya, East Java, Indonesia. Hospital X is a Type C private hospital which have special wards for airborne infectious disease, such as COVID-19. This study obtained approval from the Research Ethics Committee of Hospital X (No. 001/S/KET/KOMETIK/IV/2023).

B. Sample

This study included patients diagnosed with COVID-19 who was hospitalised in Hospital X during January-December 2022. COVID-19 diagnosis was confirmed with positive results on RT-PCR swab examination upon admission as recorded in medical records. Data was obtained from medical records which was stored at the Hospital Information System (SIMRS). Patients who were forced to return home or returned at their own request or referred to another hospital before considered recovered were excluded from this study. Patient data was followed from the day of admission until discharge from hospital (which was considered as recovered).

C. Data Collection

Data regarding patient characteristics (including age, gender, comorbidities, vaccine status, and degree of severity) and the drug therapy were obtained from patient medical record, and recorded in the data collection sheet per patient. The drug profiles were observed based on the 4th edition COVID-19 Management Guideline and the Hospital X guideline (Table 1 and 2). LoS was defined as number of days from admission to discharge from hospital as recorded in the medical record.

D. Data Analysis

Data on patient characteristics and drug therapy profiles were analysed descriptively. Drug therapy conformity was determined based on the 4th edition COVID-19 Management Guideline and Hospital X Guideline (Table I and Table II); it was considered 'conform' when the patient received all drugs listed in the guidelines, and 'not conform' if there was any drug which is not in accordance with the guidelines. Bivariate analysis was conducted to determine the associations between patient characteristic and therapy conformity (independent variable) with LoS (dependent variable).

Independent sample t-test was used to test the association between LoS and independent variables in the form of

categorical data if the data is normally distributed (Mann-Whitney U-Test if the data is not normally distributed), while Spearman's correlation was used for

independent variables in the form of nominal/ordinal data. IBM SPSS Statistics version 25 was used to assist with the analysis.

Tabel I. Pharmacological Therapy Guidelines for Children with COVID-19 (Burhan et al, 2022; RS X, 2022)

Level of Severity	COVID-19 Therapy	Other Therapy
Mild	-	• Vitamin C • Zinc • Other supplements • Symptomatic treatment
Moderate	• Antivirus: Remdesivir or Favipiravir or Molnupiravir • Antibiotic*	• Vitamin C • Zinc • Corticosteroid* • Symptomatic treatment
Severe-Critical	• Antivirus: Remdesivir or Favipiravir or Molnupiravir • Antibiotic*	• Vitamin C • Zinc • Corticosteroid or other inflammatory agents* • Anticoagulant* • Symptomatic treatment

*given with careful considerations after discussing with COVID-19 team at hospitals

Tabel II. Pharmacological Therapy Guidelines for Adults with COVID-19 (Burhan et al, 2022; RS X, 2022)

Level of Severity	COVID-19 Therapy	Other Therapy
Mild	• Antivirus: Favipiravir or Molnupiravir	• Vitamin C • Vitamin D • Symptomatic treatment
Moderate	• Antivirus: Remdesivir or Favipiravir or Molnupiravir	• Vitamin C • Vitamin D • Anticoagulant* • Symptomatic treatment
Severe-Critical	• Antivirus - Remdesivir - Favipiravir or Molnupiravir • Antibiotic (infection/sepsis)*	• Vitamin C • Vitamin D • Vitamin B1 • Corticosteroid* • Anti IL-6 (Tocilizumab)* • Anticoagulant* • Additional therapy: IvIG (Regdanvimab)* • Symptomatic treatment

Abbreviations: IVIG, intravenous immunoglobulin; IL, interleukin

*given with careful considerations after discussing with COVID-19 team at hospitals

III.RESULTS AND DISCUSSION

A total of 203 COVID-19 patients were hospitalised at Hospital X in 2022; 15 patients were excluded as they were discharged on their own requests. Of 203 patients included in this study, the majority were adults (66.5%). The distribution of patient age groups can be seen in Table III.

Table III. The distribution of age among COVID-19 patients (N=203)

Age groups	n	%
Neonate/Child/Adolescent (N=25)		
Neonates (0-1 year)	5	20.0
Children (2-11 years)	8	32.0
Adolescent (12-18 years)	12	48.0
Adult (N=135)		
19-30 years	43	31.9
31-42 years	52	38.5
43-54 years	20	14.8
55-60 years	20	14.8
Elderly (N=43)		
61-69 years	17	39.5
70-78 years	18	41.9
79-87 years	6	14.0
88-98 years	2	4.6

It was reported by Statsenko et al. (2022) that adult and elderly patients with COVID-19 tended to experience hypercoagulation and clot formation because of higher levels of APTT, D-Dimer, and fibrinogen compared to the younger group ($p < 0.001$). In addition, D-dimer levels deviated from clinical reference values ($> 0.5 \text{ ug/L}$) were seen only in older adults with COVID-19.

In all age groups, a higher percentage of male patients compared to females (Table IV). Statsenko et al. (2022) reported that male patients had a longer disease duration, thus required a more intensive treatment ($p < 0.01$). Likewise, based on the C-reactive protein (CRP) value, which is a marker of the severity of COVID-19, a higher value was found in males than females ($p < 0.05$). Apart from that, hormonal differences between males and females can also influence the COVID-19 patients' conditions. Wray & Arrowsmith (2021) stated that oestrogen can potentiate vitamin D' effects on immunity; on the other hand, male sex hormones might cause males more susceptible to COVID-19 and worsen the prognosis. Testosterone allegedly promotes viral entry by increasing the activity of the ACE2 receptor which is known to be the entry point for SARS-CoV-2. Testosterone also exerts immunosuppressive effects and can reduce the antibody response (Wray & Arrowsmith, 2021).

In this study, the majority of patients had received vaccination, at least the first dose (74.4%), and 36.4% had comorbidities (Table IV). Several studies show that vaccination can reduce morbidity due to COVID-19. Research by Li et al. (2022) involving COVID-19 patients aged ≥ 60 years reported that the risk of developing severe COVID-19 was greater

in the unvaccinated group compared to the vaccinated group. Furthermore, previous research shows that comorbidities, such as obesity, chronic kidney disease, diabetes, hypertension, kidney and cardiovascular disorders can potentially increase the risk of worsening COVID-19 (Russell et al, 2023). Use of medications among COVID-19 patients based on the age groups can be seen in **Table V**, while its conformity with the guidelines can be seen in **Table VI**.

Conformity of COVID-19 therapy with the guidelines for all age groups was 65% (118/203). In the neonate/child/adolescent group, most therapies were not according to guidelines (88%, 22/25); while in the adult and elderly groups, most therapy were appropriate (76%, 103/135; and 60%, 26/43; respectively). Inappropriate therapy has been mainly dominated by not giving supplementation and vitamins, namely vitamin C, vitamin D and zinc.

Table IV. Characteristics of COVID-19 Patients based on the Age Groups

Characteristics	Neonate/ Child/Adolescent (N=25)		Adult (N=135)		Elderly (N=43)		Total (N=203)	
	n	(%)	n	(%)	n	(%)	n	(%)
Gender								
Male	15	60.0	73	54.1	22	51.2	110	54.2
Female	10	40.0	62	45.9	21	48.8	93	45.8
Vaccination								
No data	13	52.0	30	22.2	9	20.9	52	25.6
1x vaccination	2	8.0	0	0.0	0	0.0	2	1.0
2x vaccination	8	32.0	61	45.2	17	39.5	86	42.4
3x vaccination	2	8.0	44	32.6	16	37.2	62	30.5
4x vaccination	0	0.0	0	0.0	1	2.3	1	0.5
Level of disease severity								
Mild	6	24.0	20	14.8	0	0.0	26	12.8
Moderate	17	68.0	110	81.5	37	86.0	164	80.8
Severe-Critical	2	8.0	5	3.7	6	14.0	13	6.4
Comorbidity*								
No comorbidity	24	96.0	86	63.7	19	44.2	129	63.5
With comorbidity (n=74):	1	4.0	49	36.3	24	55.8	74	36.4
Diabetes Mellitus	0	0.0	13	21.7	9	31.0	22	29.7
Hypertension	0	0.0	20	33.3	12	41.4	32	43.2
Respiratory diseases	0	0.0	11	18.3	1	3.4	12	16.2
Cardiovascular diseases	0	0.0	10	16.7	4	13.8	14	18.1
Renal diseases	0	0.0	2	3.3	0	0.0	2	2.7
Others (e.g. Parkinson)	1	0.0	4	6.7	3	10.3	7	9.4

* Each patient can have more than 1 comorbidity

The majority of neonate/child/adolescent patients were at moderate and severe levels of severity (19/25, 76%), where based on guidelines it was necessary to consider giving antivirals and antibiotics (Table I). Antivirals were given to 11 patients (44.0%), of which the majority were given Remdesivir (n=9). The use of Remdesivir for children under 12 years warrants special attention.

On April 25, 2022, the FDA released a statement that Remdesivir can be used in paediatric patients aged 28 days or older at least 3 kilograms with positive results of direct SARS-CoV-2 viral testing who are hospitalized or non-hospitalized, and have mild-to -moderate COVID-19 and are at high risk for progression to severe COVID-19, including hospitalization or death (Samuel *et al*, 2023).

Tabel V. Medications use among COVID-19 patients based on the age groups

Medications	Neonate/child/ adolescent (N=25)		Adult (N=135)		Elderly (N=43)	
	n	%	n	%	n	%
Vitamin C	17	68.0	128	94.8	38	88.4
Vitamin D	0	0.0	124	91.9	35	81.4
Vitamin B1	0	0.0	118	87.4	36	83.7
Zinc	8	32.0	0	0.0	0	0.0
Antivirus						
Favipiravir	2	18.2	21	17.5	2	5.0
Remdesivir	9	81.8	73	60.8	33	82.5
Molnupiravir	0	0.0	15	12.5	1	2.5
Remdesivir → Molnupiravir	0	0.0	5	4.2	1	2.5
Remdesivir → Favipiravir	0	0.0	2	1.7	2	5.0
Favipiravir → Remdesivir	0	0.0	2	1.7	0	0.0
Molnupiravir → Remdesivir	0	0.0	2	1.7	1	2.5
Total	11	44.0	120	88.9	40	93.0
Antibiotic						
Meropenem	1	6.7	23	28.4	14	50.0
Azithromycin	6	40.0	9	11.1	2	7.1
Moxifloxacin	1	6.7	7	8.6	4	25.0
Cefixime	1	6.7	2	2.5	3	10.7
Levofloxacin	1	6.7	35	43.2	11	39.3
Ceftriaxone	9	60.0	15	18.5	4	14.3
Cefadroxil	0	0.0	1	1.2	0	0.0
Cefoperazone-Sulbactam	0	0.0	1	1.2	0	0.0
Metronidazole	0	0.0	1	1.2	0	0.0
Total	15	60.0	81	60.0	28	65.1
Corticosteroid						
Dexamethasone	7	87.5	19	57.6	10	66.7
Methylprednisolone	1	12.5	14	42.4	5	33.3
Total	8	32.0	33	24.4	15	34.9
Anticoagulant						
Edoxaban	1	50.0	7	50.0	8	38.1

Medications	Neonate/child/ adolescent (N=25)		Adult (N=135)		Elderly (N=43)	
	n	%	n	%	n	%
Rivaroxaban	1	50.0	2	14.3	2	9.5
Fondaparinux	0	0.0	3	21.4	1	4.8
Heparin	0	0.0	0	0.0	1	4.8
Enoxaparin/Heparin	0	0.0	1	7.1	1	4.8
Enoxaparin	0	0.0	1	7.1	3	14.3
Fondaparinux/Rivaroxaban	0	0.0	0	0.0	2	9.5
Fondaparinux/Enoxaparin	0	0.0	0	0.0	1	4.8
Enoxaparin/Rivaroxaban	0	0.0	0	0.0	1	4.8
Rivaroxaban/Edoxaban/Heparin	0	0.0	0	0.0	1	4.8
Total	2	8.0	14	10.4	21	48.8
Anti IL-6						
Tocilizumab	0	0.0	2	1.5	2	4.7
Monoclonal antibody						
Regdanvimab	0	0.0	2	1.5	0	0
Symptomatic treatment*						
Antiulcer	10	40.0	110	81.5	42	97.9
Antitusive	15	60.0	65	48.1	20	46.5
Antirhinitis	12	48.0	47	34.8	9	20.9
Laxative	5	20.0	7	5.2	2	4.7
Sedative	1	4.0	10	7.4	5	11.6
Analgesic-Antipyretic	15	60.0	54	40.0	23	53.5
Antiemetic	9	36.0	40	29.6	19	44.2
Antihistamine	1	4.0	9	6.7	4	9.3
Analgesic	5	20.0	34	25.2	6	14.0
Antidyspnea	11	44.0	13	9.6	4	9.3
Antipharyngitis	3	12.0	22	16.3	5	11.6
Antimotility	3	12.0	4	3.0	1	2.3
	25	100.0	135	100.0	43	100.0

*Each patient can receive more than 1 type of symptomatic treatment

Table VI. Medication conformity with the guideline based on age groups and severity levels

Medication conformity	Levels of COVID-19 severity			Total N (%)
	Mild	Moderate	Severe	
Neonate/child/adolescent (n=25)				
Conform	3	0	0	3 (12.0)
Not conform	3	17	2	22 (88.0)
Adult (n=135)				
Conform	6	95	2	103 (76.0)
Not conform	14	15	3	32 (24.0)
Elderly (n=43)				
Conform	0	23	3	26 (60.0)
Not conform	0	14	3	17 (40.0)
All age groups (N=203)				
Conform	9	118	5	132 (65.0)
Not conform	17	46	8	71 (35.0)

In the adult patient group, the majority were at moderate and severe levels of severity (115/130, 85.1%); most patients (n=120) were received an antiviral, particularly Remdesivir (n=73). Based on the national guideline, Remdesivir is one of the antivirals of choice for adult patients with moderate or severe levels of severity (Table II) (Burhan et al., 2022; Awdisma et al., 2021). In this present study, antibiotics were administered to 60% of adult (mainly Levofloxacin) and paediatric (mainly Azithromycin) patients; administration of antibiotics should only be considered if there are indications of bacterial infection. Azithromycin is the antibiotic of choice if atypical pneumonia is suspected, however other types of antibiotics can be chosen according to the patient's clinical condition and the suspected cause of infection (Burhan et al., 2022).

Tabel VII. LoS of COVID-19 patients based on age groups

Length of Stay (LoS)	Mean $\pm$ SD	Median (range, min-max)
LoS neonate/child/adolescent (n=25), in days	6.1 $\pm$ 2.43	6 (3-15)
LoS adult (n=135), in days	7.1 $\pm$ 3.24	6 (3-25)
LoS elderly (n=43), in days	7.4 $\pm$ 3.02	6 (4-13)
LoS all age groups (N=203), in days	7.0 $\pm$ 3.11	6 (3-25)

The average LoS of all COVID-19 patients at hospital X was 7.0 ± 3.11 days; the highest average LoS was in the elderly group (7.4 ± 3.02 days) (Table VII).

Several previous studies showed a longer LoS of patients with COVID-19. Research by Oksuz et al. (2021) in Turkey involving 1,056 patients showed an average LoS of 8.0 days for hospitalized patients versus 14.8 days for patients in the Intensive Care Unit (ICU). Alimohamadi et al. (2022) reported that the >60 year age group had an average LoS of 16.60 days (95% CI 12.94-20.25), while the 40 year age group had the lowest LoS (10.15, 95% CI 4.90-15.39). Research by Baihaqi and Henny (2022) in Indonesia involving 48 COVID-19 patients hospitalized at Serui Hospital, Papua Province, reported that 26 subjects (54.2%) had a length of stay of >11 days. This present study might have lower LoS due to the better herd immunity formed with the increasing vaccination coverage.

Analysis of risk factors contributing to LOS can be seen in Table VIII. Significant associations were reported for vaccination status ($p= 0.000$) and level of COVID-19 severity ($p= 0.002$). Previous studies have shown that factors associated with LOS vary widely. A systematic review in 2022 reported that patient age, gender, ethnicity, and number of comorbidities significantly influence the LOS of COVID-19 patients (Alimohamadi et al., 2022).

Further, research by Oksuz et al. (2021) in Turkey (n=1,056 patients) showed that level of COVID-19 severity significantly influenced LOS. Baihaqi and Henny (2022) in Papua, Indonesia, showed that the factor associated with the length of stay for COVID-19 patients was the patient's clinical condition, namely fever (p=0.003; OR 10.59; 95% CI 2.22-50.49) and Neutrophil-Lymphocyte Ratio - NLR (p=0.034; OR 4.55; 95% CI 1.12 -18.49); while age and gender did not significantly influence LOS. In parallel, Ramatillah et al. (2021) in Jakarta, Indonesia, reported that gender (p=0.450) and age (p=0.226) did not have a significant correlation with LoS or clinical outcomes for COVID-19 patients. Moreover, Triamyanti et al. (2022) in Bangil, Indonesia, found that the use of antibiotics and antivirals had no effect on the recovery rate of COVID-19 patients; while comorbidity showed a significant correlation with duration of treatment (p=0.002) and clinical condition (p= 0.014) of COVID-19 patients (20). An Indonesian multi-centres study with multifactorial

approaches would be required to confirm these findings.

Study limitation. This study used a population approach to evaluate the medication use conformity where patients' drug therapy were generally compared with the drug of choice in the 4th edition COVID-19 Management Guideline, which was considered a benchmark in the management of COVID-19 patients with the dominant Omicron variant in 2022. Thus, the use of medications have not yet been evaluated for each individual patient, such as drug interactions, polypharmacy and Drug Related Problems (DRP). Further research would be required to evaluate the quality of drug use at the individual level. Apart from that, this research was conducted retrospectively using data from patient medical records, hence some missing data might be an issue. Hospital X, however, have used electronic medical records stored at SIMRS, thus minimum missing data are expected (this is particularly regarding patient data before entering the hospital).

Tabel VIII. Associations between patient characteristics and drug therapy conformity with LOS

Risk factors	Median LOS, in days (range min-max)	<i>p-value</i>
Gender		0.726 ^a
Male (n=110)	6 (3-15)	
Female (n=93)	6 (4-25)	
Age groups		0.122 ^b
Neonate/Child/Adolescent (n=25)	6 (3-15)	
Adult (n=135)	6 (3-25)	

Risk factors	Median LOS, in days (range min-max)	<i>p-value</i>
Elderly (n=43)	6 (4-13)	
Vaccination		0.000 ^b
1-2x vaccination (n=88)	7 (4-15)	
3-4x vaccination (n=63)	6 (3-16)	
Level of severity		0.002 ^b
Mild (n=26)	5 (3-13)	
Moderate (n=164)	6 (3-25)	
Severe-Critical (n=13)	9 (4-15)	
Comorbidity*		0.201 ^a
Yes (n=55)	6 (3-19)	
No (n=148)	6 (3-25)	
Therapy conformity		0.864 ^a
Conform	7 (4-25)	
Not conform	6 (3-15)	

*Comorbidities included: diabetes mellitus, hypertension, respiratory/cardiovascular/renal disorders

^ausing *Mann-Whitney U Test*

^busing *Spearman's correlation*

IV. CONCLUSION

This study provided insights on the patients' characteristics and drug use contributing to LoS among COVID-19 patients in an Indonesian setting. Patients history of vaccination status and level of severity warrants considerations when treating patients hospitalised with COVID-19. In addition, optimising drug utilisation according to the guidelines should be encouraged. Findings from this study should provide a basis in improving services for hospitalised patients with COVID-19, which is still evolving and remains a constant threat.

CONFLICT OF INTEREST

All authors declare that there is no conflict of interest in this research.

ACKNOWLEDGEMENT

We would like to thank the Director as well as all the staff of Hospital X in Surabaya for their technical assistances during the data collection.

REFERENCES

- Alimohamadi, Y., Yekta, E.M., Sepandi, M., Sharafoddin, M., Arshadi, M., & Hesari, E. (2022). Hospital length of stay for COVID-19 patients: a systematic review and meta-analysis. *Multidiscip Respir Med*, 17(1), 856. <https://doi.org/10.4081/mrm.2022.856>
- Amalia H. (2021). Omicron penyebab COVID-19 sebagai variant of concern. *J Biomedika dan Kesehatan*, 4(4), 139-141. <https://doi.org/10.18051/jbiomedkes.2021.v4.139-141>
- Awdisma, W.M., Nosa, U.S., Hasmono, D., Nurmainah, N. (2021). Kajian literatur: tinjauan remdesivir sebagai pilihan terapi pada COVID-19. *J*

- Pharmascience*, 8(2), 121-132.
<http://dx.doi.org/10.20527/jps.v8i2.11650>
- Baihaqi, F.A., & Rumaropen, H. (2022). Faktor-faktor yang berhubungan dengan lama rawat inap pasien COVID-19 di RSUD Serui Provinsi Papua: Studi potong lintang. *J Penyakit Dalam Indones*, 8(4), 187.
<https://doi.org/10.7454/jpdi.v8i4.627>
- Burhan, E., & Rachmadi, R.A. (2022). Omicron surge and the future of COVID-19 vaccinations. *Med J Indones*, 31(1), 80-84.
<https://doi.org/10.13181/mji.bc.226066>
- Burhan, E., Susanto, A.D., Nasution, S.A., et al. (2022). *Pedoman Tatalaksana COVID-19 Edisi 4*. Jakarta: PDPI, PERKI, PAPDI, PERDATIN, IDAI
- Centers for Disease Control and Prevention (CDC). (2022). *Underlying medical conditions associated with higher risk for severe COVID-19: information for healthcare professionals*. Available at: <https://www.cdc.gov/coronavirus/2019-ncov/hcp/clinical-care/underlyingconditions.html> (Accessed June 21, 2022).
- Di Gennaro, F., Pizzol, D., Marotta, C., et al. (2020). Coronavirus diseases (COVID-19) current status and future perspectives : A narrative review. *J Environ Res Public Health*, 17(8), 2690.
<https://doi.org/10.3390/ijerph17082690>
- Guo, A., Lu, J., Tan, H., Kuang, Z., Luo, Y., Yang, T., Xu, J., Yu, J., Wen, C., & Shen, A. (2021). Risk factors on admission associated with hospital length of stay in patients with COVID-19: A retrospective cohort study. *Sci Rep*, 11, 7310.
<https://doi.org/10.1038/s41598-021-86853-4>
- Khosravizadeh, O., Vatankhah, S., Bastani, P., Kalhor, R., Alirezaei, S., & Doosty, F. (2016). Factors affecting length of stay in teaching hospitals of a middle-income country. *Elec Physician*, 8(10), 3042-3047.
<https://doi.org/10.19082/3042>
- Li, M., Liu, Q., Wu, D., Tang, L., Wang, X., Yan, T., An, Z., Wang, Z., Gao, G.F., Wang, F., & Zheng H. (2022). Association of COVID-19 vaccination and clinical severity of patients infected with Delta or Omicron variants — China , May 21 , 2021 – February 28 , 2022. *China CDC Wkly*, 4(14), 293-297.
<https://doi.org/10.46234/ccdcw2022.074>
- Ministry of Health Republic of Indonesia (MoH-RI). (2021). *Infeksi Emerging Kementerian Kesehatan*. Available at: <https://covid19.kemkes.go.id/> (Accessed March 9, 2022).
- Oksuz, E., Malhan, S., Gonen, M.S., Kutlubay, Z., Keskindemirci, Y., & Tabak, F. (2021). COVID-19 healthcare cost and length of hospital stay in Turkey: retrospective analysis from the first peak of the pandemic. *Health Econ Rev*, 11(1), 1-12.
<https://doi.org/10.1186/s13561-021-00338-8>
- Ramatillah, D.L., & Isnaini, S. (2021). Treatment profiles and clinical outcomes of COVID-19 patients at private hospital in Jakarta. *PLoS One*, 16, 1-11.
<https://doi.org/10.1371/journal.pone.0250147>
- Russell, C.D., Lone, N.I., & Baillie, J.K. (2023). Comorbidities, multimorbidity and stages of COVID-19 pathogenesis. *Nature Medicine*, 29, 334-343.
<https://doi.org/10.1038/s41591-022-02156-9>
- Samuel, A.M., Hacker, L.L., Zebracki, J., et al. (2023). Remdesivir Use in pediatric patients for SARS-CoV-2 treatment: single academic center study. *Pediatr Infect Dis J*, 42(4), 310-314.
<https://doi.org/10.1097/INF.00000000>

- [000003814](#)
 Statsenko, Y., Al Zahmi, F., Habuza, T., et al. (2022). Impact of age and sex on COVID-19 severity assessed from radiologic and clinical findings. *Front Cell Infect Microbiol*, 11, 777070. <https://doi.org/10.3389/fcimb.2021.777070>
- Tai, P.Q., Toan, D.T.T., Son, D.T., et al. (2020). Factors associated with the duration of hospitalization among COVID-19 patients in Vietnam: A survival analysis. *Epidemiol Infect*, 148, e114. <https://doi.org/10.1017/S0950268820001259>
- Triamyanti, V.Y., Yulia, R., Herawati, F., & Jaelani, A.K. (2022). Evaluasi persentase kesesuaian dengan Pedoman 3 Penggunaan Antibiotik dan Antivirus pada Pasien Covid-19 non- ventilator di RSUD Bangil. *MEDFARM J Farm dan Kesehat*, 11(2), 127-141. <https://doi.org/10.48191/medfarm.v11i2.94>
- World Health Organization (WHO). (2020). Coronavirus disease 2019 situation report 51 - 11th March 2020. *WHO Bulletin*, 2633.
- World Health Organisation (WHO). (2022). *Tracking SARS-COV-2 Variants*. Available at: <https://www.who.int/activities/tracking-SARS-CoV-2-variants> (Accessed July 1, 2022).
- Wray S., & Arrowsmith, S. (2021). The physiological mechanisms of the sex-based difference in outcomes of COVID-19 Infection. *Front Physiol*. 12, 627260. <https://doi.org/10.3389/fphys.2021.627260>

[Home](#) > [Archives](#) > **Vol 12, No 2 (2025)**

VOL 12, NO 2 (2025)

JURNAL PHARMASCIENCE

DOI: <http://dx.doi.org/10.20527/jps.v12i2>

TABLE OF CONTENTS

JURNAL PHARMASCIENCE

Potensi Ekstrak Etanol Rimpang Temulawak (Curcuma Xanthorrhiza Roxb.) sebagai Antibakteri: Literature Review

Abstract view : 235 times

DOI: [10.20527/jps.v12i2.19262](https://doi.org/10.20527/jps.v12i2.19262)

Sri Rahayu

PDF (BAHASA INDONESIA)
268-278

Uji Aktivitas Antioksidan, Fenolik Total, dan Flavonoid Total dari Kombinasi Ekstrak Etanol Daun Pandan (Pandanus Amaryllifolius) dan Daun Jeruk Purut (Citrus Hystrix)

Abstract view : 202 times

DOI: [10.20527/jps.v12i2.20426](https://doi.org/10.20527/jps.v12i2.20426)

Ika Ayu Mentari, Jundy Eko Wahyudi

PDF (BAHASA INDONESIA)
279-287

Standarisasi dan Uji Aktivitas Penangkapan Radikal Bebas Fraksi Daun Ketapang (Terminalia catappa L.)

Abstract view : 103 times

DOI: [10.20527/jps.v12i2.20860](https://doi.org/10.20527/jps.v12i2.20860)

Sri Wahdaningsih, Robby Najini

PDF (BAHASA INDONESIA)
288-298

Tinjauan Sistematis terhadap Penerapan HOT-Fit sebagai Framework Evaluasi Keberhasilan Sistem Informasi Rumah Sakit

Abstract view : 70 times

DOI: [10.20527/jps.v12i2.21542](https://doi.org/10.20527/jps.v12i2.21542)

Fathur Rahman, Lutfan Lazuardi, Anna Wahyuni Widayanti

PDF (BAHASA INDONESIA)
299-313

Integrated LC-HRMS Analysis, Network Pharmacology, and Molecular Docking to Predict the Potential of Syzygium polyanthum in Alleviating Rheumatoid Arthritis

Abstract view : 131 times

DOI: [10.20527/jps.v12i2.23762](https://doi.org/10.20527/jps.v12i2.23762)

Nurlely Nurlely, Arief Nurrochmad, Nanang Fakhrudin, Sitarina Widiyarni

PDF
314-334

Comparison Of HbA1c Levels in Type II Diabetes Mellitus Patients Receiving A Combination of Oral Hypoglycemic Drugs and Insulin Degludec-Aspart Versus Insulin Degludec-Aspart Monotherapy

Abstract view : 67 times

DOI: [10.20527/jps.v12i2.22242](https://doi.org/10.20527/jps.v12i2.22242)

Helmina Wati, Syahrizal Ramadhani, Dita Ayulia Dwi Sandi, Esty Restiana Rusida, Muhammad Raffyqurrahman

PDF
335-345

Formulation and Optimization of Marigold Flower (Tagetes erecta L.) Extract Emulgel using Response Surface Methodology

Abstract view : 179 times

DOI: [10.20527/jps.v12i2.21173](https://doi.org/10.20527/jps.v12i2.21173)

Dinda Rahmanda Priatna, Yenni Puspita Tanjung, Tubagus Akmal

PDF
346-366

Effectiveness of Clozapine and Its Potential Drug-Drug Interactions in Patients with Schizophrenia: A Study in Referral Psychiatric Hospital

Abstract view : 46 times

DOI: [10.20527/jps.v12i2.23111](https://doi.org/10.20527/jps.v12i2.23111)

Salma Fajar Puspita, Rifa Salsabila, Wahyu Ratna Dewi, Vitarani Dwi Ananda Ningrum

PDF
367-379

Review: Pengantaran Nanopartikel Berbasis Lipid Dalam Pemanfaatan Senyawa Bahan Alam Yang Mengandung Antioksidan Untuk Dermatologi

Abstract view : 95 times

DOI: [10.20527/jps.v12i2.20868](https://doi.org/10.20527/jps.v12i2.20868)

Panji Bachrudin, Garnadi Jafar, Yani Mulyani

PDF (BAHASA INDONESIA)
380-400

Tinjauan : Antielastase dari Berbagai Tumbuhan untuk Kosmetik

Abstract view : 36 times

DOI: [10.20527/jps.v12i2.21963](https://doi.org/10.20527/jps.v12i2.21963)

Retty Handayani, Delin Arlina, Syifa Nurisma Ekaputri, Siti Nurajijah, Siti Nurmulyanti Sopiana, Farisha Rahma Fadilla, Aghna Urfa

PDF (BAHASA INDONESIA)
401-423

Bibliometric Analysis of Medicinal Plants Research for Diabetes Mellitus in Indonesia

Abstract view : 44 times

DOI: [10.20527/jps.v12i2.23703](https://doi.org/10.20527/jps.v12i2.23703)

Khoerul Anwar, Nanang Fakhrudin, Retno Murwanti, Agung Endro Nugroho

PDF
424-440

Analysing Factors Contributing to Length of Stay (LoS) Among Patients with COVID-19 in Indonesian Hospital Setting

Abstract view : 26 times

DOI: [10.20527/jps.v12i2.22557](https://doi.org/10.20527/jps.v12i2.22557)

Sonny Edbert, Yosi Irawati Wibowo, AdjI Prayitno Setiadi

PDF
441-454

Potensi Anti-inflamasi Ekstrak Euphorbia heterophylla Desf.: Penelitian In Vitro dan In Vivo

Abstract view : 75 times

DOI: [10.20527/jps.v12i2.23814](https://doi.org/10.20527/jps.v12i2.23814)

Elis Susilawati, Intan Pertiwi, Aulia Nurfazri Istiqomah, Yasminwar Susilawati, Jutti Levita, Sri Adi Sumiwi

PDF (BAHASA INDONESIA)
455-467

Molecular Docking Analysis Of Phenolic and Flavonoid Compounds from Eichhornia Crassipes for Antidiabetic Activity Through Interaction with PPAR- γ (5Y2O) and A-Glucosidase (3TOP)

Abstract view : 44 times

DOI: [10.20527/jps.v12i2.23583](https://doi.org/10.20527/jps.v12i2.23583)

Khoirunnisa Muslimawati, Taufik Muhammad Fakh, Nabila Hadiah Akbar, Aditya Maulana Perdana Putra, Nazhipah Isnani

PDF
468-481

Innovative Gel Mask A. jiringa (Jack) Nielsen: Effects of HPMC and Palm Stem Starch on Physical Characteristics and Antibacterial Activity

Abstract view : 34 times

DOI: [10.20527/jps.v12i2.22397](https://doi.org/10.20527/jps.v12i2.22397)

Nani Kartinah, Arnia Ulpah, Pratika Viogenta, Amalia Khairunnisa

PDF
482-492

GC-MS Profiling and Inhibitory Effect on UV-C Exposure Induced Erythrocyte Damage of Sungkai Leaf: A Potential Novel Antioxidant Agent

Abstract view : 24 times

DOI: [10.20527/jps.v12i2.23081](https://doi.org/10.20527/jps.v12i2.23081)

Mashuri Mashuri, Eko Suhartono, Noor Muthmainah, Lisda Hayatie, Nadia Salma Nazwa Assyfa, Hanna Dwi Aprilia Husna, Helma Febrianti

PDF
492-501

Acne Fighter Microemulsion Gel mengandung Basil Oil (Ocimum sanctum)

Abstract view : 50 times

DOI: [10.20527/jps.v12i2.23773](https://doi.org/10.20527/jps.v12i2.23773)

Noridafi Noridafi, Alya Zainah, Nahdiya Nahdiya, Salsabila Azzahra Rusda, Syifa Nur Cahyani, Mia Fitriana

PDF (BAHASA INDONESIA)
502-511

Aktivitas Antioksidan Ekstrak dan Antihyperpigmentasi Serum Biji Alpukat (Persea americana Mill.) sebagai Inhibitor Enzim Tirosinase

Abstract view : 77 times

DOI: [10.20527/jps.v12i2.21458](https://doi.org/10.20527/jps.v12i2.21458)

Fatimah Zahra, Arini Khaerunnisa, Dhyneu Dwi Jayantie, Eva Vauziah, Irfan Iryana

PDF (BAHASA INDONESIA)
512-524

Cytotoxicity and Pro-Apoptotic Effect of Dichloromethane Fraction of Nauclea subdita Stem Bark on HepG2 Liver Cancer Cell Modeling

Abstract view : 99 times

DOI: [10.20527/jps.v12i2.22330](https://doi.org/10.20527/jps.v12i2.22330)

Chesya Nabila, Masriani Masriani, Rini Muharini

PDF
525-541

Efek Kombinasi Ekstrak Etanol Rimpang Kunyit (Curcuma longa L.) dan Daun Pegagan (Centella asiatica (L.) Urb) terhadap Ekspresi Gen eNOS pada Jaringan Hati Tikus Obes

Abstract view : 48 times

DOI: [10.20527/jps.v12i2.23813](https://doi.org/10.20527/jps.v12i2.23813)

Agus Sulaeman, Salim Agustian, Elis Susilawati, Soni Muhsinin, Marita Kaniawati, Marita Feliana

PDF (BAHASA INDONESIA)
542-557

Formulasi Lip Tint Alami Ekstrak Stroberi: Peran Tween 80 dan Propilen Glikol Dalam Optimasi Stabilitas dan Aktivitas Antioksidan dengan Menggunakan Design Expert Software

Abstract view : 65 times

DOI: [10.20527/jps.v12i2.23810](https://doi.org/10.20527/jps.v12i2.23810)

Benni Iskandar, Revina Tasya Azkiyah, Wira Noviana Suhery, Rudi Hendra

PDF (BAHASA INDONESIA)
558-581

Jurnal Pharmascience Published by:

Program Studi Farmasi Universitas Lambung Mangkurat

Banjarmaru, Indonesia

Jurnal Pharmascience is indexed by:



This is an open acces article under a Creative Commons Attribution-NonCommercial 4.0 International License.

SINTA 4



SINTA 2

KERJA SAMA



Google Scholar Citation
: JPS in google scholar

	All	Since 2021
Citations	3585	3005
h-index	32	30
i10-index	101	88

ADDITIONAL MENU

[ONLINE SUBMISSION](#)

[KEBIJAKAN OPEN ACCES](#)

[AUTHOR GUIDELINES](#)

[EDITORIAL TEAM](#)

[REVIEWER](#)

[KEBIJAKAN PLAGIARISME](#)

[AUTHOR\(S\) FEE](#)

[JOURNAL HISTORY](#)

[CONTACT US](#)

USER

Username

Password

☐ Remember me

Journal Template



SUPPORT TOOLS



Visitors

ID 434.788	JP 410
US 5.540	PH 372
CN 5.269	IN 366
SG 2.428	CA 203
MY 457	RU 191

Pageviews: 960.017



00478161

[View My Stats](#)

NOTIFICATIONS

[View](#)

[Subscribe](#)

LANGUAGE

Select Language

JOURNAL CONTENT

Search

Search Scope

Browse

[By Issue](#)

[By Author](#)

[By Title](#)

[By Sections](#)

[By Identify Types](#)

[Other Journals](#)

INFORMATION

[For Readers](#)

[For Authors](#)

[For Librarians](#)

JURNAL PHARMASCIENCE

Publikasi Resmi Penelitian Bidang Kefarmasian dan Kesehatan

ISSN-Print : 2355-5386; ISSN-Online: 2460-9560

https://ppjp.ulm.ac.id/journal/index.php/pharmascience
jps@ulm.ac.id

HOME ABOUT LOGIN REGISTER SEARCH CURRENT ARCHIVES ANNOUNCEMENTS FOCUS&SCOPE ETIKA PUBLIKASI

Home > About the Journal > Editorial Team

EDITORIAL TEAM

EDITOR-IN-CHIEF

Prof. Dr. apt. Sutomo Sutomo, Program Studi Farmasi Universitas Lambung Mangkurat, Indonesia

EDITORIAL TEAM

Dita Ayulia Dwi Sandi, Prodi Farmasi, FMIPA Universitas Lambung Mangkurat, Indonesia
apt. Normaidah Normaidah, Program Studi Pendidikan Profesi Apoteker Universitas Lambung Mangkurat, Indonesia
Nur Mahdi, Universitas Lambung Mangkurat
Nurul Mardiaty, Program Studi Farmasi Universitas Lambung Mangkurat
Dhania Novitasari, Fakultas Farmasi, Universitas Padjadjaran, Indonesia
Fauziah Fauziah, Universitas Harapan Bangsa, Indonesia

Jurnal Pharmascience Published by:

Program Studi Farmasi Universitas Lambung Mangkurat

Banjarbaru, Indonesia

Jurnal Pharmascience is indexed by:



This is an open acces article under a Creative Commons Attribution-NonCommercial 4.0 International License.

SINTA 4



SINTA 2

KERJA SAMA



Google Scholar Citation
: JPS in google scholar

	All	Since 2021
Citations	3585	3005
h-index	32	30
i10-index	101	88

ADDITIONAL MENU

ONLINE SUBMISSION

KEBIJAKAN OPEN ACCES

AUTHOR GUIDELINES

EDITORIAL TEAM

REVIEWER

KEBIJAKAN PLAGIARISME

AUTHOR(S) FEE

JOURNAL HISTORY

CONTACT US

USER

Username
Password
☐ Remember me

Journal Template



SUPPORT TOOLS



Visitors

ID 434.788	JP 410
US 5.540	PH 372
CN 5.269	IN 366
SG 2.428	CA 203
MY 457	RU 191

Pageviews: 960.017



00478163

View My Stats

NOTIFICATIONS

View
Subscribe

LANGUAGE

Select Language

English

Submit

JOURNAL CONTENT

Search
Search Scope

All

Search

Browse
By Issue
By Author
By Title
By Sections
By Identify Types
Other Journals

INFORMATION

For Readers
For Authors
For Librarians



JURNAL PHARMASCIENCE

 PROGRAM STUDI FARMASI UNIVERSITAS LAMBUNG MANGKURAT

 P-ISSN : 23555386 <> E-ISSN : 24609560  Subject Area : Health

4.25
Impact

2843
Google Citations

Sinta 2
Current Accreditation

 Google Scholar

 Garuda

 Website

 Editor URL

History Accreditation

2017

2018

2019

2020

2021

2022

2023

2024

2025

2026

2027

2028

Garuda

Google Scholar

Tingkat Kepuasan Pasien Terhadap Pelayanan Resep Di Apotek Kabupaten Karanganyar

Program Studi Farmasi FMIPA Universitas Lambung Mangkurat Jurnal Pharmascience Vol 12, No 1 (2025): Jurnal Pharmascience 132-142

 2025  DOI: 10.20527/jps.v12i1.20293  Accred : Sinta 2

Pengaruh Suhu Pengeringan terhadap Aktivitas Antioksidan Ekstrak Daun Ungu (Graptophyllum pictum L. Griff)

Program Studi Farmasi FMIPA Universitas Lambung Mangkurat Jurnal Pharmascience Vol 12, No 1 (2025): Jurnal Pharmascience 157-166

 2025  DOI: 10.20527/jps.v12i1.21010  Accred : Sinta 2

Nanopartikel Sebagai Sistem Penghantaran Oral untuk Low Molecular Weight Heparin (LMWH) dalam Pengobatan Kolitis Ulseratif

Program Studi Farmasi FMIPA Universitas Lambung Mangkurat Jurnal Pharmascience Vol 12, No 1 (2025): Jurnal Pharmascience 84-102

 2025  DOI: 10.20527/jps.v12i1.20144  Accred : Sinta 2

Potensi Ekstrak Etanol Daun Enceng Gondok (Eichhornia crassipes) sebagai Terapi Penyembuh Luka Bakar pada Tikus Putih Jantan

Program Studi Farmasi FMIPA Universitas Lambung Mangkurat Jurnal Pharmascience Vol 12, No 1 (2025): Jurnal Pharmascience 233-242

 2025  DOI: 10.20527/jps.v12i1.20326  Accred : Sinta 2

Pemisahan Senyawa Flavonoid pada Tanaman Keji Beling (Strobhilantes crispa)

Program Studi Farmasi FMIPA Universitas Lambung Mangkurat Jurnal Pharmascience Vol 12, No 1 (2025): Jurnal Pharmascience 1-8

 2025  DOI: 10.20527/jps.v12i1.16886  Accred : Sinta 2

Uji Aktivitas Antidiabetes Ekstrak Etanol 96% Daun Bungur (Lagerstroemia speciosa) Secara In Vivo

Program Studi Farmasi FMIPA Universitas Lambung Mangkurat Jurnal Pharmascience Vol 12, No 1 (2025): Jurnal Pharmascience 181-189

 2025  DOI: 10.20527/jps.v12i1.21737  Accred : Sinta 2

Analisis Kualitatif dan Kuantitatif Kandungan Senyawa Flavonoid dan Fenolik pada Ekstrak Etanol Daun Mangga Kasturi (Mangifera Casturi Kosterm.)

Program Studi Farmasi FMIPA Universitas Lambung Mangkurat Jurnal Pharmascience Vol 12, No 1 (2025): Jurnal Pharmascience 167-180

 2025  DOI: 10.20527/jps.v12i1.21350  Accred : Sinta 2

Analisis Kuantitatif dan Kualitatif Senyawa Obat : Inovasi Dalam Preparasi Sampel Biologis [Review]

Program Studi Farmasi FMIPA Universitas Lambung Mangkurat Jurnal Pharmascience Vol 12, No 1 (2025): Jurnal Pharmascience 31-49

 2025  DOI: 10.20527/jps.v12i1.19569  Accred : Sinta 2

Kemampuan Daya Hambat Beberapa Produk Sampo Antiketombe Dengan Kandungan Bahan Aktif Berbeda Terhadap Jamur Candida albicans

Program Studi Farmasi FMIPA Universitas Lambung Mangkurat Jurnal Pharmascience Vol 12, No 1 (2025): Jurnal Pharmascience 9-18

 2025  DOI: 10.20527/jps.v12i1.18749  Accred : Sinta 2


Skrining Fitokimia dan Aktivitas Antibakteri Ekstrak Metanol Daun Kalangkala (Litsea garciae Vidal)

Program Studi Farmasi FMIPA Universitas Lambung Mangkurat Jurnal Pharmascience Vol 12, No 1 (2025): Jurnal Pharmascience 190-198

 2025  DOI: 10.20527/jps.v12i1.21785  Accred : Sinta 2

View more ...

Citation Per Year By Google Scholar



Year	Citation
2018	100
2019	150
2020	230
2021	400
2022	550
2023	750
2024	550
2025	0
2026	0

Journal By Google Scholar

	All	Since 2021
Citation	2843	2638
h-index	28	27
i10-index	88	82

View more ...