



Bacterial lipoxygenases for fatty acid oxyfunctionalization: advances and future directions

Ruth Chrisnasari^{1,2,3} · Tom A. Ewing² · Willem J. H. van Berkel¹ · Jean-Paul Vincken¹ · Marie Hennebelle¹

Received: 12 December 2025 / Revised: 13 April 2026 / Accepted: 14 April 2026
© The Author(s) 2026

Abstract

The shift to a biobased economy aims to reduce dependence on fossil fuels by converting biological materials, such as plant-derived and algae-derived fatty acids, into valuable compounds. The enzymatic oxyfunctionalization of fatty acids is of special interest because it enables the eco-friendly synthesis of a wide range of products with unique properties. Lipoxygenases (LOXs) are key oxyfunctionalization enzymes featuring broad substrate specificities and distinct regioselectivities. Among them, bacterial LOXs remain underexplored compared to their eukaryotic counterparts, representing an untapped resource with unique structural and functional traits. This review highlights recent advances in bacterial LOXs, emphasizing their versatile oxidative capabilities via dioxygenase and hydroperoxide isomerase activities. Key strategies, including protein engineering and oxygen level modulation, are discussed for modifying substrate specificity, regioselectivity, and preferred catalytic pathway. Multi-enzyme cascade reactions further expand the scope of LOX-derived products, unlocking new applications in biotechnological contexts. We address challenges in harnessing bacterial LOXs' biocatalytic potential and propose solutions to optimize their performance. This work enhances understanding of bacterial LOXs and highlights their potential for transformative biotechnological applications in fatty acid derivatization.

Keywords Dioxygenase · Hydroperoxide isomerase · Lipoxygenase · Oxyfunctionalization · Oleochemicals · Oxidoreductase · Oxylipins

Introduction

The growing interest in a biobased economy reflects the need to reduce reliance on fossil resources by sustainably converting renewable biomass into value-added chemicals and materials. In this context, fatty acids derived from plant and algal oils are abundant and widely available feedstocks that serve as important starting materials for the production of biobased chemicals (de Espinosa And Meier 2011). Functionalizing fatty acids through oxyfunctionalization—a

process that introduces peroxy (OOH), hydroxyl (OH), epoxy, carbonyl (C=O), and carboxyl (COOH) groups—is crucial for enhancing their reactivity and enabling the synthesis of diverse valuable products. In addition, depending on the position of oxyfunctionalization within the fatty acid, derivative products of varying chain lengths can be obtained, which in turn affect their properties and functionalities (Revol-Cavalier et al. 2025).

One promising approach to achieve fatty acid oxyfunctionalization is through enzymatic conversion, which can be carried out by enzymes such as fatty acid hydratases (FAHs), cytochrome P450 monooxygenases (CYPs), peroxygenases (PXGs), and lipoxygenases (LOXs) (Feussner And Wastermack 1998; Konkel And Schunck 2011; Aranda et al. 2018; Bornscheuer 2018; Hagedoorn et al. 2021). Among these enzymes, lipoxygenases (LOXs) offer distinct advantages that make them highly attractive for industrial applications. Unlike CYPs, LOXs do not require costly cofactors or complex redox partner systems, significantly reducing operational costs and simplifying reaction setups, which is beneficial for large-scale processes. In addition, LOXs exhibit


✉ Marie Hennebelle
marie.hennebelle@wur.nl

¹ Laboratory of Food Chemistry, Wageningen University & Research, Bornse Weiland 9, Wageningen 6708 WG, The Netherlands

² Wageningen Food & Biobased Research, Wageningen University & Research, Bornse Weiland 9, Wageningen 6708 WG, The Netherlands

³ Faculty of Biotechnology, University of Surabaya (UBAYA), Surabaya 60293, Indonesia

A Dioxygenase



B Hydroperoxide isomerase (HPI)




Fig. 1 Lipoxygenase-catalyzed reactions. **A** The enantio- and regioselective dioxygenation activity of LOX converts polyunsaturated fatty acids (PUFAs) containing a (1Z,4Z)-pentadiene structural unit into fatty acid hydroperoxides (FAHPs) containing a conjugated diene pattern. The dioxygenation enantioselectivity of LOX is either *R*-selective or *S*-selective. The regioselectivity shown in the figure at C5 of the pentadiene moiety is an example; alternative regioselectivity at C1 is also possible. **B** The hydroperoxide isomerase (HPI) activity of LOX converts FAHPs into epoxy alcohols or ketones

tive or *S*-selective. The regioselectivity shown in the figure at C5 of the pentadiene moiety is an example; alternative regioselectivity at C1 is also possible. **B** The hydroperoxide isomerase (HPI) activity of LOX converts FAHPs into epoxy alcohols or ketones

a broader substrate scope than FAHs and offer more diverse regioselectivities than FAHs, CYPs, and PXGs, resulting in the efficient production of various oxyfunctionalized fatty acids.

LOXs (EC 1.13.11.x) are non-heme iron, or sometimes manganese-dependent enzymes, that catalyze the regio- and enantioselective dioxygenation of polyunsaturated fatty acids (PUFAs) leading to the formation of fatty acid hydroperoxides (FAHPs) (Fig. 1A). FAHPs are key intermediates, serving as precursors for a range of valuable chemical compounds, including flavors and fragrances (Johannsen et al. 2020, 2021; Ribeaucourt et al. 2022), oleochemicals (Song et al. 2013; Seo et al. 2015; Yu et al. 2018; Kim And Park 2019), and lipid mediators (An et al. 2021). In addition to dioxygenation activity, a few LOXs from soybean, mammals, fungi and bacteria have been reported to have hydroperoxide isomerase (HPI) activity, which transforms FAHPs into epoxy alcohols or ketones (Fig. 1B) (Garssen et al. 1976; Yu et al. 2003; Cristea And Oliw 2006; Zheng and Brash 2010). The presence of HPI activity in LOX is of interest because hydroxy, epoxy or keto groups typically impart special characteristics to fatty acids, such as increased viscosity and reactivity, that can be

used in a wide range of applications (Hou 2009). Moreover, unsaturated epoxy alcohols serve as precursors for the synthesis of biologically active molecules (Riera And Moreno 2010). The presence of HPI activity in LOXs enables further diversification of reaction products, making them uniquely capable of generating a broader range of fatty acid derivatives. Their unique combination of versatility in transforming diverse PUFA substrates and ability to generate regioselective products highlights LOXs as promising biocatalysts for biotechnological applications.

LOXs have been studied across diverse biological kingdoms, including mammals, plants, fungi, and bacteria. Most early research and reviews focused on eukaryotic LOXs, especially their biochemical characteristics (Brash 1999; Coffa et al. 2005; Andreou And Feussner 2009) and applications (Nanda And Yadav 2003; Joo and Oh 2012; Heshof et al. 2016; Hayward et al. 2017; An et al. 2021). Unlike their well-characterized eukaryotic counterparts, bacterial LOXs remained largely unexplored until the past decade, when phylogenetic analyses first proposed them as a separate subfamily with unique features (Hansen et al. 2013). Since then, increasing efforts to explore and characterize novel bacterial LOXs have

been reported (Newie et al. 2016; An et al. 2018a; Goloshchapova et al. 2018; Ellamar et al. 2019; Qi et al. 2020; Lee et al. 2022; Oh et al. 2022; Chrisnasari et al. 2024b, 2025a). These enzymes exhibit broad substrate specificity toward polyunsaturated fatty acids (PUFAs, C18–C22) and often display distinct and desirable selectivity (An et al. 2021; Chrisnasari et al. 2022).

The rapid expansion of available bacterial LOX sequences in databases (>4700 sequences in 2022) combined with the fact that less than 0.5% have been biochemically characterized, highlights both the diversity and the significant knowledge gaps within this enzyme family. A comprehensive phylogenetic analysis of bacterial LOXs, including sequence clustering, structural prediction, and identification of residues influencing regioselectivity, has been reported in our previous review (Chrisnasari et al. 2022), in which bacterial LOXs were grouped into nine distinct phylogenetic clusters predicted to possess unique structural and functional characteristics. In addition, another review summarized the use of bacterial LOXs for the biocatalytic synthesis of lipid mediators (An et al. 2021).

Building on this foundation, this review provides a critical update on bacterial LOXs, offering deeper insights into factors influencing their catalytic reactions and innovative strategies for modulating their activity. We also explore how multi-enzyme cascade reactions expand the scope of LOX-derived products and their applications. Finally, we discuss the key challenges in harnessing the biocatalytic potential of these enzymes and propose solutions to advance their use in industrial and biotechnological contexts.

LOX-catalyzed reactions

Dioxygenase activity: factors influencing regioselectivity and substrate preference

LOX catalyzes the regioselective dioxygenation of fatty acids through five distinct steps (Fig. 2): (1) The ferric iron (Fe^{3+}) cofactor initiates the reaction by abstracting a hydrogen atom at the center of the pentadiene structure of the fatty acid substrate (Lehnert and Solomon 2003) the unpaired electron is transferred to the ferric iron and the proton is transferred to the hydroxide ligand coordinated to the iron, yielding ferrous iron (Fe^{2+}), water, and a lipid alkyl radical; (2) The unpaired electron in the lipid alkyl radical undergoes rearrangement to either the [+2] or the [−2] position relative to the abstracted hydrogen; (3) A dioxygen molecule is introduced leading to the formation of a fatty acid peroxy radical ($\text{ROO}\bullet$); (4) The fatty acid peroxy radical is then reduced by an electron from the ferrous iron (Fe^{2+}) and protonated, resulting in the formation of FAHP and the iron is re-oxidized to its ferric form

(Fe^{3+}); (5) The FAHP is then released and the enzyme is ready to start the next catalytic cycle (Hamberg And Samuelsson 1967; Egmond et al. 1972; Hamberg et al. 1998; Lehnert And Solomon 2003).

The regioselectivity of LOX dioxygenation is determined by specific structural characteristics of the enzyme, such as the position and size of the oxygen migration channel (Knapp et al. 2001; Coffa and Brash 2004; Newcomer and Brash 2015; Collazo And Klinman 2016) and the depth of the substrate-binding pocket (Vogel et al. 2010). A key residue in the oxygen channel, known as the Coffa site, was found to determine the regio- and enantioselectivity of LOXs by directing the oxygen molecule through a glycine/alanine switch (Coffa and Brash 2004; Coffa et al. 2005). With respect to residues influencing the depth of the substrate-binding pocket, studies on mammalian LOXs have identified three key amino acid determinants (triad determinants): Sloane determinant (SD) (Sloane et al. 1991), Borngräber 1 determinant (BD1) (Borngräber et al. 1996), and Borngräber 2 determinant (BD2) (Borngräber et al. 1999). The role of the Coffa site and triad determinants in determining the regio- and enantioselectivity of bacterial LOXs has been summarized in our previous review (Chrisnasari et al. 2022).

The corresponding residues of triad determinants in other LOXs are typically identified through amino acid sequence alignment. However, this method is not always accurate for bacterial LOXs due to their structural heterogeneity. For instance, the crystal structures of LOX from *Pseudomonas aeruginosa* (Pa-LOX) revealed that Pa-LOX exhibits a large U-shaped substrate-binding pocket, similar to the binding pockets commonly found in LOXs from mammals, plants, and corals (Fig. 3A) (Newcomer and Brash 2015). Multiple sequence alignments identified amino acid residues corresponding to the triad determinants in Pa-LOX, i.e., E369 (BD1), M434 and F435 (SD), and I608 (BD2) (Chrisnasari et al. 2022). However, the position of BD1 was relatively distant from the bottom of the pocket, while E373 was positioned closer to it, suggesting a larger contribution of this residue in determining the depth of the binding pocket (Fig. 3A). A similar observation was made for the AlphaFold model of *Burkholderia thailandensis* LOX (Bt-LOX), which belongs to the same bacterial LOX cluster as Pa-LOX (Chrisnasari et al. 2024b).

Compared to Pa-LOX, LOX from *Cyanobacter* sp. (Cya-LOX) exhibits a smaller, straight substrate-binding pocket, which is structurally very different from that of Pa-LOX and other eukaryotic LOXs (Fig. 3B). As a consequence, the residues forming the bottom of the substrate-binding pocket in Cya-LOX (Fig. 3C), i.e., E355, Y360, V365, F413, and V406, do not correspond to any of the triad determinants. A similar phenomenon is observed in the AlphaFold model




Fig. 2 Reaction mechanism of lipooxygenase (LOX). LOX catalyzes the oxygenation of fatty acids through five reaction steps: hydrogen abstraction, radical rearrangement, oxygen insertion, radical reduc-

tion-protonation, and product release. The figure is adapted from (Chrisnasari et al. 2024a) with some modification


of *Microcystic aeruginosa* LOX (Ma-LOX), which clusters with Cya-LOX within the bacterial LOX phylogeny (Chrisnasari et al. 2025a). The residues forming the bottom of the substrate-binding pocket in Cya-LOX are also positioned at the bottom of the Ma-LOX pocket, whereas the residues corresponding to the triad determinants are not located near the bottom of the binding pocket. These findings collectively indicate that identifying triad determinants in bacterial LOXs solely through amino acid sequence alignment with LOXs from other kingdoms may not be reliable. However, as demonstrated for Bt-LOX (Chrisnasari et al. 2024b), sequence alignment combined with structural modeling is useful for predicting triad determinants within the same bacterial LOX cluster.

Over the past three decades, extensive efforts have been made to modify residues at the bottom of the substrate-binding pocket to alter the dioxygenation regioselectivity of LOXs (Table 1). In mammalian LOXs, introducing larger

residues typically constrains the pocket, limiting substrate entry and shifting regioselectivity toward the methyl end of the substrate (Borngräber et al. 1996). Conversely, smaller residues deepen the pocket, allowing further substrate penetration and shifting oxygenation closer to the carboxyl end (Sloane et al. 1991; Borngräber et al. 1996, 1999).

Compared to mammalian LOXs, bacterial LOXs exhibit more varied outcomes when trying to direct the regioselectivity of dioxygenation through protein engineering. For instance, introducing smaller residues at the bottom of the substrate-binding pocket in bacterial LOXs results in a loss of regioselectivity, leading to the production of hydroperoxides at random positions (An et al. 2018a). In other cases, replacement with smaller residues, as expected, can deepen the substrate-binding cavity, allowing deeper substrate penetration and shifting oxygenation closer to the carboxyl end of the fatty acid, as demonstrated for *Pseudomonas aeruginosa* LOX engineered to produce 8S-hydroperoxy and

Fig. 3 Comparison of substrate-binding pocket architectures in representative bacterial and mammalian lipoxygenases. (A) Comparison of the substrate-binding pockets of *Pseudomonas aeruginosa* LOX (Pa-LOX, PDB 5IR5) and human 15B-LOX (PDB 4NRE). Both enzymes exhibit a relatively large U-shaped substrate-binding pocket typical of many lipoxygenases. Residues forming the bottom triad determinants (BD1, SD, BD2), which are associated with control of regioselectivity, are indicated. Despite the similar overall pocket architecture, a different relative position of BD1 can be observed. (B) Superimposition of the substrate-binding pockets of Pa-LOX and *Cyanothece* sp. LOX (Cya-LOX, PDB 5EK8). Pa-LOX displays a larger U-shaped channel (cyan), whereas Cya-LOX exhibits a narrower and more linear substrate channel (grey), illustrating structural diversity among bacterial LOXs that may influence substrate positioning and oxygenation patterns. (C) Residues located at the bottom of the Cya-LOX substrate-binding pocket, including the corresponding triad determinant positions (BD1, SD, BD2). The iron cofactor is shown as an orange sphere



11S-hydroperoxy eicosatetraenoic acid (Seo et al. 2025). However, such modifications do not universally alter regioselectivity; in some instances, mutations have little or no effect on regioselectivity or may even lead to enzyme inactivation (Banthiya et al. 2016; Goloshchapova et al. 2018; Chrisnasari et al. 2024b). An unexpected shift in regioselectivity toward the methyl end has been observed for the L445A variant of Bt-LOX. Especially with ω -3 PUFAs, this SD variant demonstrated a shift in the preferred dioxygenation position from the ω -5 to the ω -2 carbon, a regioselectivity that has not been reported for any other LOX before (Chrisnasari et al. 2024b). Structural modeling and docking experiments suggested that the L445A substitution caused a smaller and reshaped substrate-binding pocket, promoting a shallower penetration for ω -3 PUFAs. With several ω -6 PUFAs, the L445A variant of Bt-LOX facilitated dioxygenation closer to the carboxylic acid terminus, indicating a shift in substrate orientation (Chrisnasari et al. 2024b), consistent

with observations reported for *Myxococcus xanthus* LOX (An et al. 2018a).

Given the complexity of substrate-binding pocket dynamics, the effects of bottom-pocket residue substitutions cannot be universally generalized across all LOXs. Structural changes introduced by these substitutions can alter the pocket's shape, size, and residue positioning, resulting in diverse regioselectivity outcomes. Therefore, detailed product profiling, structural analysis, and computational modeling are crucial for understanding and predicting the effects of such modifications in individual LOXs.

Besides affecting regioselectivity, the shape and size of the substrate-binding pocket likely explain the distinct substrate specificities observed among bacterial LOXs. Pa-LOX and Bt-LOX, which have spacious U-shaped substrate-binding pockets, prefer ω -6 and longer PUFAs (Banthiya et al. 2016; Chrisnasari et al. 2024b). In contrast, Cya-LOX, with its smaller and straighter pocket, is active toward linoleic

Table 1 Effects of bottom pocket residue substitutions on LOX regioselectivity

Enzyme	Enzyme variant*	Regioselectivity (%) **						Ref
		15-	12-	11-	9-	8-	5-	
<i>Mammalian LOXs</i>								
Human 15-LOX	Wild-type	90	10	-	-	-	-	(Sloane et al. 1991)
	M 418 V (SD)	60	40	-	-	-	-	
	M 418V/ I 417A/ Q 416 K (SD)	5	95	-	-	-	-	
Rabbit reticulocyte 15-LOX	Wild-type	97	3	-	-	-	-	(Borngräber et al. 1996)
	F 353L (BD1)	28	72	-	-	-	-	
	F 353L/ I 593 Y (BD1/BD2)	21	79	-	-	-	-	
Porcine leukocyte 12-LOX	Wild-type	3	97	-	-	-	-	(Borngräber et al. 1996)
	L 353 F (BD1)	94	6	-	-	-	-	
Human 12/15-LOX	Wild-type	86	14	-	-	-	-	(Vogel et al. 2010)
	F 353L (BD1)	12	88	-	-	-	-	
	I 418 A (SD)	6	94	-	-	-	-	
	M 419 V (SD)	42	58	-	-	-	-	
	I 593 A (BD2)	65	35	-	-	-	-	
	F 353L/ I 418 F (BD1/SD)	64	36	-	-	-	-	
	F 35L/ I 418F/ I 593 A (BD1/SD/BD2)	70	30	-	-	-	-	
<i>P. pygmaeus</i> 12/15-LOX	Wild-type	86	14	-	-	-	-	(Vogel et al. 2010)
	F 353 V (BD1)	6	94	-	-	-	-	
	I 418 A (SD)	18	82	-	-	-	-	
	M 419 V (SD)	60	40	-	-	-	-	
	I 593 A (BD2)	60	40	-	-	-	-	
<i>Bacterial LOXs</i>								
<i>Myxococcus xanthus</i> LOX	Wild-type	10	90	-	-	-	-	(An et al. 2018a)
	T 397 Y (BD1)	94	6	-	-	-	-	
	T 397 A (BD1)	28	11	-	-	26	35	
	A 461 I (SD)	33	18	-	-	13	36	
	A 461 F (SD)	41	40	-	-	3	16	
	I 664 Y (BD2)	35	65	-	-	-	-	
<i>Myxococcus fulvus</i> LOX	Wild-type	5	95	-	-	-	-	(Goloshchapova et al. 2018)
	G 359 F (BD1)	-	-	-	-	-	-	
	F 424 I / I 425 M (SD)	5	95	-	-	-	-	
<i>Pseudomonas aeruginosa</i> LOX	Wild-type	100	-	-	-	-	-	(Banthiya et al. 2016)
	E 369 A (BD1)	100	-	-	-	-	-	
	M 434 V (SD)	100	-	-	-	-	-	
	F 435L (SD)	100	-	-	-	-	-	
	L 612V	100	-	-	-	-	-	
<i>Pseudomonas aeruginosa</i> LOX	Wild-type	97	-	-	-	-	3	(Seo et al. 2025)
	M 434G (SD)	6	-	61	-	20	13	
	Y 609G	2	-	-	-	98	-	
	Y 609A	-	-	-	-	100	-	
	Y 609V	6	-	-	-	88	6	
<i>Burkholderia thailandensis</i> LOX	Wild-type	98	2	-	-	-	-	(Chrisnasari et al. 2024b)
	I 618 A (BD1)	-	-	-	-	-	-	
	L 445 A (SD)	46	3	-	2	6	43	
	F 446 V (SD)	96	4	-	-	-	-	

*Larger residue is indicated in bold. Corresponding residues to the triad determinants are indicated in parentheses

**Reported regioselectivity when using arachidonic acid (AA, C20:4 ω -6) as substrate. Regioselectivity (%): percentage of product formed at a specific oxygenation position relative to the sum of all regioisomers formed

acid (LA, C18:2 ω -6), α -linolenic acid (ALA, C18:3 ω -3), arachidonic acid (AA, C20:4 ω -6), and eicosapentaenoic acid (EPA, C20:5 ω -3), although its precise substrate preference remains unreported (Newie et al. 2016). Ma-LOX has a similarly small pocket and shows a preference for LA and ALA over longer PUFAs (Chrisnasari et al. 2025a). These findings highlight the critical role of pocket shape and size in determining the substrate specificity of LOX.

Bacterial LOXs generally exhibit distinct preferences for specific PUFA substrates. For instance, LOXs from *Myxococcus fulvus* and *Pseudomonas aeruginosa* PAO1 preferentially act on longer-chain PUFAs (Banthiya et al. 2016; Goloshchapova et al. 2018). In contrast, LOX from *Microcystic aeruginosa* demonstrates a strong preference for shorter-chain PUFAs (Chrisnasari et al. 2025a). In addition, LOXs from *Burkholderia thailandensis*, *Pseudomonas* sp. 42A2, *Calothrix* sp. HK-06, *Sphingopyxis macrogoltabida*, and *Myxococcus xanthus* demonstrate high activity towards LA, but reduced activity towards ALA (Busquets et al. 2004; An et al. 2015, 2018a; Qi et al. 2020; Kim et al. 2022; Chrisnasari et al. 2024b).

Given the inherent substrate preferences of LOXs, the ability to engineer these enzymes to modify their substrate specificity has great potential for broadening their applications. A striking example involves the L445A substitution in Bt-LOX. As explained above, this substitution caused a positional shift in nearby residues, reducing the size of the substrate-binding pocket and changing its shape. These changes altered the enzyme's substrate penetration depth and resulted in a shift in substrate preference from ω -6 PUFAs to ω -3 PUFAs (Chrisnasari et al. 2024b).

Double dioxygenase activity and its possible mechanisms

In addition to catalyzing single dioxygenation reactions, some LOXs also demonstrate a double dioxygenation reaction. Double dioxygenation reactions are of particular interest due to the potential anti-inflammatory effects of the derived dihydroxy fatty acids (Basil And Levy 2015; Ishihara et al. 2019). Recently, several bacterial LOXs have been reported to demonstrate a pronounced double dioxygenation activity (Kim et al. 2021, 2022; Lee et al. 2022; Oh et al. 2022). Despite these intriguing findings, the precise mechanisms underlying double dioxygenation in LOXs remain poorly understood. The initial explanation for double dioxygenation reactions focused on variations in substrate orientation within the enzyme's binding pocket, as reported for soybean LOX-1, which catalyzes the formation of 5,15-dihydroperoxy fatty acids from arachidonic acid (AA, C20:4 ω -6) (Van Os et al. 1981). In this scenario, the initially formed 15-hydroperoxy fatty acid re-enters the

enzyme in a head-first orientation and undergoes a second oxygenation at the C5 position to generate the double dioxygenation product (Fig. 4A). While this scenario provides an interesting perspective, it fails to account for the distinct dioxygenation patterns observed in other LOXs, including the two bacterial LOXs discussed below.

An alternative explanation suggests that variations in substrate penetration depth play a key role in double dioxygenation, as reported for *Sphingopyxis macrogoltabida* LOX (Sm-LOX), which generates 9S,15S-dihydroperoxy and 11S,17S-dihydroperoxy fatty acids from C20 and C22 PUFAs, respectively (Kim et al. 2022). In this case, when using AA as a substrate, the enzyme first oxygenates at the C9 position to produce a 9-hydroperoxy fatty acid. This product then re-enters the binding pocket, where a second




Fig. 4 Proposed mechanisms for double dioxygenation activity in LOXs. **A** Initially, double dioxygenation was ascribed to variations in substrate orientation within the binding pocket (carboxyl end-first or methyl end-first). **B** Another potential mechanism involves differences in substrate penetration depth, caused by a blocking residue (shown in grey semicircle). **C** Recently, a novel mechanism for double dioxygenation in LOXs has been proposed. A shallow substrate-binding pocket may enable oxygen insertion via two distinct pathways: through the substrate entrance and a putative oxygen channel. The direction of oxygen insertion is indicated by blue arrows, and the iron cofactor is shown as an orange sphere. Newly added hydroperoxide groups are highlighted in blue

oxygenation occurs at the C15 position. The hydroperoxide group at C9 hinders further substrate penetration, in part due to the presence of a blocking residue in the pocket, preventing oxygenation at other positions (Kim et al. 2022) (Fig. 4B).

Another mechanism for the LOX-mediated double dioxygenation reaction has been proposed for Ma-LOX (Chrisnasari et al. 2025a). The AlphaFold model of Ma-LOX revealed a shallow substrate-binding pocket, which suggests that oxygen insertion may occur via two distinct pathways, through the substrate entrance and a putative oxygen channel. This mechanism leads to the formation of 9,16-dihydroperoxy fatty acids from α -linolenic acid (ALA; C18:3 ω -3). Upon the first oxygenation at the C16 position producing a 16-hydroperoxy fatty acid, the hydroperoxide group at this position prevents further oxygen insertion through the oxygen channel. Consequently, oxygenation at the C13 position is blocked, and instead, oxygen insertion occurs at the C9 position through the substrate entrance (Fig. 4C). The same principle applies when the first dioxygenation occurs at the C9 position; the hydroperoxide group at this position hinders the insertion of oxygen from the substrate entrance, thus the dioxygenation at C12 is not possible. However, oxygen insertion from the putative oxygen channel to the C16 position is still possible. This deviating mechanism underscores the complexity of double dioxygenation activity in LOX catalysis.

The oxygen diffusion pathway in LOX catalysis has been previously studied through molecular dynamics simulations. In soybean LOX-1, eight possible channels for oxygen diffusion have been identified. However, computational and mutagenesis studies strongly support only one dominant channel that directs molecular oxygen to a specific region of the active site, ensuring the regio- and enantio-specificity of the product (Collazo And Klinman 2016). Similar to soybean LOX, studies on mammalian LOXs have identified four oxygen channels. However, only one channel—localized opposite to the nonheme iron—exhibits high oxygen affinity at the active site (Saam et al. 2007). The critical role of an oxygen channel opposite to the nonheme iron in determining LOX catalytic specificity has also been proposed in some reviews (Kuhn et al. 2005; Newcomer and Brash 2015).

A recent study reported the presence of two primary oxygen channels that direct oxygen to the substrate-binding pocket, based on conserved motifs and helical structures found across the LOX family, including plants, mammals, and bacteria (Manivarma et al. 2025). However, these two primary oxygen channels predominantly lead to the formation of a single oxidized product, indicating that only one channel is primarily responsible for LOX catalysis, while the other is located relatively far from the non-heme iron. Interestingly, the two conserved motifs and structures observed

in other LOXs (Manivarma et al. 2025) were also identified in Ma-LOX, aligning with the proposed bidirectional oxygen flow (Chrisnasari et al. 2025a). This observation supports the hypothesis that dioxygen molecules can enter the enzyme's active site from two different directions—both through the putative oxygen channel and the substrate entrance. In addition, it highlights the role of Ma-LOX's shallower substrate-binding pocket in facilitating its double dioxygenation activity, a feature not commonly observed in other LOXs.

The variability of proposed double-dioxygenation mechanisms across different LOXs highlights the complexity of this catalytic activity. Different enzymes appear to adopt a distinct strategy to achieve double dioxygenation, suggesting that no single model can be universally applied. Therefore, unraveling the mechanisms driving double dioxygenation requires a comprehensive approach, combining detailed product profiling, structural analysis, and computational modeling to predict enzyme-substrate interactions. Only through this integrated approach can we begin to fully understand the unique catalytic capabilities of individual LOXs and their potential applications in biocatalysis.

Factors affecting hydroperoxide isomerase activity (HPI)

In addition to their primary dioxygenase activity, a few LOXs from soybean, mammals, fungi, and bacteria have been reported to have hydroperoxide isomerase (HPI) activity (Garssen et al. 1976; Yu et al. 2003; Cristea And Oliw 2006; Zheng and Brash 2010; Chrisnasari et al. 2025a, 2025b). These two reactions are catalyzed by LOX at different oxidation states of the non-heme iron: dioxygenase activity requires the ferric (Fe^{3+}) state, while HPI activity requires the ferrous (Fe^{2+}) state. Newly isolated enzymes are typically in the ferrous state, which is inactive toward PUFAs. They require a priming reaction with FAHPs to convert the iron to its ferric state (Schilstra et al. 1992, 1993), thereby, activating the enzyme to enter the dioxygenase catalytic cycle (Schilstra et al. 1994) (Fig. 5). An alkoxyl radical or epoxy-allylic radical is postulated as an intermediate in the LOX activation process (Yu et al. 2003; Zheng and Brash 2010). The alkoxyl radical or epoxy-allylic radical subsequently dissociates through an oxygen-dependent dissociation process, leading to the formation of epoxy-allylic hydroperoxides, which are then further transformed into epoxy-allylic ketones (Zheng and Brash 2010). LOX activation is a single-turnover event, meaning that once the enzyme is oxidized to its ferric form, it will remain in this form and no longer react with FAHPs (de Groot et al. 1975). An exception applies to LOXs that exhibit HPI activity, as the ferrous form of these enzymes can convert FAHPs into




Fig. 5 Modulating dioxigenase and hydroperoxide isomerase (HPI) activity of LOX. Activation of the non-heme iron cofactor of LOX by FAHP (typically generated through auto-oxidation) leads to the formation of an alkoxy-allylic or epoxy-allylic radical intermediate. In the presence of oxygen, these radical intermediates dissociate via an oxygen-dependent pathway to form an epoxy-allylic hydroperoxide, which is subsequently transformed into an epoxy-allylic ketone. The free ferric enzyme then catalyzes a dioxigenase reaction, converting PUFA into FAHP. LOX activation is a single-turnover event, meaning that once the enzyme is oxidized to its ferric form, it will remain in this form and no longer react with FAHPs. Under O₂-saturated con-

ditions, both the oxygen-dependent dissociation of the radical intermediates and the dioxigenase reaction become more pronounced. Conversely, under O₂-limited conditions, the radical intermediate can be transformed into an epoxy alcohol or ketone by HPI activity, with the iron in LOX being regenerated to its ferrous state upon product release. When continuous regeneration of the ferrous enzyme occurs due to HPI activity, the FAHP produced by dioxigenase activity can enter the HPI cycle and be further transformed into either an epoxy alcohol or ketone (indicated by a dashed arrow). This figure is adopted from Chrisnasari et al. (2025b)

epoxy alcohols or ketones, reverting to their ferrous state after product release (Yu et al. 2003; Zheng and Brash 2010).

HPI activity in those LOXs typically occurs only under specific conditions. For instance, soybean LOX-1 exhibits HPI activity in the presence of excess FAHP and under anaerobic conditions (Garssen et al. 1976). HPI activity in the manganese LOX from *Gaeumannomyces avenae* is induced by the G316A mutation, which appears to reduce the size of the oxygen channel, limiting oxygen access and promoting HPI activity (Cristea And Oliw 2006). Human epidermal LOX3 (eLOX3), despite being an isoform of human 15-LOX2 with its high dioxigenase activity (Kobe et al. 2014), is atypical as it either entirely lacks dioxigenase activity (Yu et al. 2003) or requires a prolonged lag phase to initiate it (Zheng and Brash 2010). The absence of an oxygen channel in this enzyme may explain its high HPI activity and lack of dioxigenase activity (Fig. S1). Bacterial LOXs from *Burkholderia thailandensis* and *Microcystic aeruginosa* exhibit strong HPI activity at high enzyme concentrations (Chrisnasari et al. 2025a, 2025b). From this, it was concluded that elevated enzyme concentrations increase

oxygen consumption, thereby creating oxygen-limited conditions that favor HPI activity.

The dual functionality of LOXs, which exhibit both dioxigenation and HPI activity, expands the catalytic repertoire of these enzymes but also poses challenges in product purification. The products of dioxigenase activity (FAHPs) and the products of HPI activity (epoxy alcohols or ketones) have different potential applications which will be discussed in the next section. Harnessing the full potential of LOXs requires a deeper understanding of how to control their activity, steering the enzyme toward either the dioxigenase or HPI pathway, depending on the desired outcome. A study on Bt-LOX revealed that by controlling oxygen concentration in the reaction environment, HPI and dioxigenase activity can be modulated (Chrisnasari et al. 2025b) (Fig. 5). Under O₂-saturated conditions, HPI activity was suppressed, while dioxigenase activity was enhanced (Faillace et al. 2025; Chrisnasari et al. 2025b). Conversely, under O₂-limited conditions, dioxigenase activity persisted, and the resulting fatty acid hydroperoxides were efficiently converted into epoxy alcohols and

ketones. It appears that the low residual oxygen levels initially present support the conversion of PUFAs into their corresponding FAHPs, which are subsequently transformed into epoxy alcohols and ketones due to pronounced HPI activity in an oxygen-limited environment. These findings highlight an effective way to modulate LOX activity and direct the enzyme toward a specific catalytic pathway. Modulating LOX activity represents a major advancement in PUFA derivatization, broadening both the product scope and the applicability of LOXs.

Potential applications of LOXs

LOX in the production of flavors and fragrances

Flavors and fragrances (F&F) are important components in consumer goods such as food, cosmetics, and household products. While chemical synthesis dominates F&F production due to its efficiency, it comes with drawbacks such as

high energy consumption, toxic catalysts, and reliance on fossil-based materials (Johannsen et al. 2020, 2021; Ribeaucourt et al. 2022). This has led to increasing demand for natural alternatives in F&F production. Natural extraction methods, such as distillation, often yield extremely low amounts, driving the demand for sustainable biotechnological alternatives. Biotransformation using enzymes is an emerging and highly desirable approach to meet this demand.

Among F and F compounds, fatty aldehydes and alcohols constitute a prominent class of molecules widely used in food and perfume industries, most notably those in the C3–C12 range (Johannsen et al. 2020, 2021; Ribeaucourt et al. 2022). Fatty aldehydes of different chain lengths exhibit distinct scents (Ribeaucourt et al. 2022). Various of these aldehydes are potentially accessible from fatty acids, but their synthesis requires regioselective biocatalysts capable of generating a specific aldehyde. The diverse regioselectivity and broad substrate preferences observed in bacterial LOXs (Chrisnasari et al. 2022) enable the enzymatic production of fatty aldehydes with varying chain lengths (Table 2).

Table 2 Potential application of bacterial LOXs for fatty aldehydes production

Bacterial LOX source	Ref.	Regio-selectivity	PUFA Substrate*	Fatty aldehydes	Structure	Scent
L445A variant <i>B. thailandensis</i>	(Chrisnasari et al. 2024b)	ω -2 LOX	ALA, EPA	Propanal, Propionaldehyde		floral and fruity
<i>P. aeruginosa</i>	(Banthiya et al. 2016; Deschamps et al. 2016)	ω -5 LOX	LA, ALA, GLA, AA	Hexanal, Caproaldehyde		Green, grassy fruity
<i>B. thailandensis</i>	(An et al. 2015; Sim et al. 2015; Chrisnasari et al. 2024b)					
<i>M. xanthus</i>	(An et al. 2018a)					
<i>Rivularia sp.</i>	(Qi et al. 2020)					
<i>Calothrix sp.</i>						
<i>Tolypothrix bouillei</i>						
<i>Microcystis aeruginosa</i>	(Chrisnasari et al. 2025a)					
<i>Cyanothece sp.</i>	(Newie et al. 2016)	ω -6 LOX	ALA			
<i>A. marina</i>	(Gao et al. 2010)					
<i>M. xanthus</i>	(An et al. 2018a)	ω -8 LOX	AA, EPA, DHA	Nonanal, Pelargonal-dehyde		Soapy, citrus-like, rose-like, floral
<i>M. fulvus</i>	(Goloshchapova et al. 2018)					
<i>Nostoc sp.</i>	(Andreou et al. 2008)	ω -9R LOX	LA			
<i>Cyanothece sp.</i>	(Newie et al. 2016)					
<i>A. marina</i>	(Gao et al. 2010)					
<i>S. macrogoltabida</i>	(Kim et al. 2022)	ω -9S LOX	LA, ALA			
<i>S. macrogoltabida</i>	(Kim et al. 2022)	ω -11 LOX	AA, EPA, DPA, DHA	Dodecanal, Lustraldehyde		Waxy, fatty, soapy, reminiscent of violets

*PUFA substrates: linoleic acid (LA, C18:2 ω -6), α -linolenic acid (ALA, C18:3 ω -3), γ -linolenic acid (GLA, C18:3 ω -6), arachidonic acid (AA, C20:4 ω -6), eicosapentaenoic acid (EPA, C20:5 ω -3), docosapentaenoic acid (DPA, C22:5 ω -3), docosahexaenoic acid (DHA, C22:6 ω -3)

FAHPs generated by LOXs can be enzymatically cleaved into aldehydes and oxo-fatty acids by hydroperoxide lyases (Brühlmann et al. 2013; Brühlmann And Bosijokovic 2016) (Fig. 6). The aldehydes may be further converted into the corresponding alcohols by alcohol dehydrogenases (ADHs), which can provide desirable sensory properties or improved stability for certain applications. Such enzymatic strategies may be particularly attractive for food applications, where high selectivity and mild reaction conditions are desirable. Alternatively, hydrogenation, i.e., chemical reduction using metal catalysts, enables both saturation of remaining carbon-carbon double bonds and reduction of aldehydes to alcohols, thereby offering an efficient route for converting unsaturated aldehydes into saturated alcohols (Milone et al. 2007; Ide et al. 2012), which is especially relevant for non-food applications. The remaining oxo-fatty acids can be further transformed into oleochemicals such as ω -hydroxy fatty acids, α,ω -dicarboxylic acids, and ω -amino carboxylic acids, as discussed in the next section. By leveraging their diverse regioselectivities and broad substrate preferences, bacterial LOXs emerge as highly promising biocatalysts for sustainable F&F production.

LOX in the production of oleochemicals

The FAHPs generated by LOXs could potentially serve as key intermediates in producing a wide range of oleochemicals, including medium-chain-length to long-chain-length ω -hydroxy fatty acids, ω -amino-carboxylic acids, and α,ω -dicarboxylic acids (Yadav and Srinivas 1997; Köckritz And Martin 2011; Song et al. 2013; Seo et al. 2015; Yu et al. 2018; Kim And Park 2019). These oleochemicals are essential for manufacturing diverse chemical products and intermediates, such as nylons and other polyamides, polyesters, resins, hot-melt adhesives, powder coatings, corrosion inhibitors, lubricants, plasticizers, and greases. Importantly, the chain length of these oleochemicals, which affects their properties and functionalities, depends on the position of the hydroperoxide group on the fatty acid. By selecting specific regioselective LOXs and fatty acid substrates, products of varying lengths can be achieved.

FAHPs generated by LOXs can be cleaved into their oxo-fatty acid derivatives using hydroperoxide lyases, which can then be further converted biocatalytically into ω -hydroxy fatty acid; α,ω -dicarboxylic acid, and ω -amino carboxylic




Fig. 6 Schematic multistep enzymatic synthesis of PUFA-derived aldehydes and alcohols involving distinct regioselective lipxygenases, hydroperoxide lyases, and alcohol dehydrogenases. Hydrogenation efficiently converts unsaturated aldehydes into saturated alcohols by facilitating both the saturation of unsaturated bonds and

the reduction of aldehydes to alcohols. Enzymes and newly introduced functional groups are highlighted in blue. Some of these reaction cascades are hypothetical in the sense that the regioselectivity of enzymes catalyzing the follow-up steps, i.e., hydroperoxide lyases, may have to be identified or developed

acid using alcohol dehydrogenase, aldehyde dehydrogenase, and ω -amino transferase, respectively (Song et al. 2013; Seo et al. 2015) (Fig. 7). To saturate the remaining unsaturated bonds in the resulting products, chemical hydrogenation can be performed (Žula et al. 2022; Quaranta et al. 2022; Spiekermann And Seidensticker 2024).

A significant innovation has been achieved through the L445A variant of Bt-LOX due to its unique regioselectivity at the $\omega - 2$ position, enabling the production of $\omega - 2$ FAHPs from $\omega - 3$ PUFAs (Chrisnasari et al. 2024b). This novel regioselectivity expands the range of fatty acid transformations by facilitating the synthesis of longer-chain oxo-fatty acids (Fig. 6). These oxo-fatty acids can subsequently be converted into ω -hydroxycarboxylic acids, α,ω -dicarboxylic acids, and ω -amino carboxylic acid derivatives (Björkhem 1972; Grechkin and Hamberg 2004; Grechkin et al. 2006), which were previously unattainable with natural LOX regioselectivities. Such extended-chain compounds hold considerable promise as precursors for biobased polymers and specialty chemicals. By demonstrating both natural and novel regioselectivities, bacterial LOXs serve as promising biocatalysts for sustainable oleochemical production, thus developing eco-friendly alternatives for the chemical industry.

LOX for the production of PUFAs-derived oxylipins

PUFA-derived oxylipins are structurally diverse and biologically active oxygenated molecules found across various organisms, including animals, plants, and microorganisms. These compounds play a critical role as signaling molecules in physiological processes such as homeostasis, cell communication, and responses to inflammation and infections (Blée 2002; Serhan 2014; Zhang et al. 2025; De Bartolo et al. 2025). In mammals, oxylipins are often referred to as lipid mediators, and include leukotrienes, lipoxins, hepoxilins, trioxilins, resolvins, protectins, and maresins. These LOX-derived oxylipins are composed of dihydroxy or trihydroxy fatty acids, and fatty acid epoxy alcohols (Fig. 8) (An et al. 2021; Lee et al. 2025). The specific functionality of these oxylipins depends on the chain-length and the degree of unsaturation of the PUFA substrate as well as the type, position, and chirality of the attached functional group (An et al. 2021; Revol-Cavalier et al. 2025; Lee et al. 2025). Research suggests that most biologically active oxylipins exist in different chiral forms and their bioactivity appears to be influenced by their chirality (Oh et al. 2011; Isobe et al.





Fig. 7 Schematic representation of the multistep enzymatic synthesis of PUFA-derived ω -hydroxy fatty acids, α,ω -dicarboxylic acids, and ω -amino-carboxylic acids. The final hydrogenation step is typically carried out using chemical methods

Fig. 8 Typical chemical structures of oxylipins derived from lipoxygenase (LOX), including leukotriene (LT), lipoxin (LX), hepoxilin (Hx), trioxilin (TrX), resolvins (Rev), protectin D (PD), and maresin (MaR)



2013; Blum et al. 2019; Ambaw et al. 2020). As a result, there is significant interest in producing enantioselective oxylipins through enzymatic catalysis.


Bacterial LOXs are particularly promising for synthesizing diverse oxylipins due to their distinct substrate preferences, regio- and enantioselectivities (An et al. 2018b; Chrisnasari et al. 2022). Harnessing both the dioxygenase and HPI activities of LOX enables the generation of a broad spectrum of bioactive oxylipins (Fig. 9), especially if subsequent enzymatic or chemical derivatization processes can preserve the chirality of the compounds.

The regioselective dioxygenation of PUFAs by LOX produces specific FAHPs, which can undergo a second dioxygenation step to form dihydroperoxy fatty acids. These

intermediates are then reduced to dihydroxy fatty acids either enzymatically or through chemical reduction. Some bacterial LOXs have been shown to exhibit double dioxygenation activity (Kim et al. 2021, 2022; Lee et al. 2022; Oh et al. 2022), which represents a promising green route for synthesizing dihydroxy fatty acids. The growing interest in these compounds, particularly due to their potential anti-inflammatory properties (Basil And Levy 2015; Ishihara et al. 2019), highlights the significance of bacterial LOXs as powerful tools in oxylipin biosynthesis.

An alternative route for oxylipin production involves the epoxidation of FAHPs via the HPI activity of LOX, resulting in fatty acid epoxy alcohols. Depending on the positions of the epoxy and hydroxy groups, these epoxy

Fig. 9 Schematic multistep enzymatic synthesis of PUFA-derived oxylipins involving lipoxygenases and epoxide hydrolase. Reduction of fatty acid hydroperoxide to fatty acid hydroxide (indicated by dashed arrow) can be done enzymatically or chemically



alcohols can function as biologically active molecules known as hepoxilins (An et al. 2018b). Further hydrolysis of the epoxide group by epoxide hydrolases produces trihydroxy fatty acids, another class of biologically significant molecules called trioxilins (Riera And Moreno 2010; An et al. 2018b). Both hepoxilins and trioxilins have garnered significant attention in recent years due to their anti-inflammatory, anti-infective, antibacterial, antiviral, anti-apoptotic, neuroprotective, and tissue-healing properties (An et al. 2018b). LOXs that exhibit dual functionality of dioxygenase and HPI activities, e.g., LOXs from soybean, *Gaeumannomyces avenae*, *Burkholderia thailandensis*, and *Microcystis aeruginosa* (Garssen et al. 1976; Cristea And Oliw 2006; Chrisnasari et al. 2025a, 2025b), facilitate the efficient synthesis of oxylipins by enabling the direct conversion of PUFAs into epoxy alcohols. The dual functionality of these LOXs underscores their versatility and establishes them as robust biocatalysts for producing structurally complex oxylipins.

Current challenges and outlook

Although bacterial LOXs hold significant potential due to their diverse dioxygenation and hydroperoxide isomerase activities, scaling up their use for large-scale applications remains challenging. Heterologous production of bacterial LOXs usually is performed in *Escherichia coli* (Andreou et al. 2008; Lu et al. 2013b; Neau et al. 2014; Sim et al. 2015; Newie et al. 2016; Qian et al. 2017). However, heterologous production in *E. coli* results in extensive formation of inclusion bodies (Vidal-Mas et al. 2005; Lu et al. 2013b) and production of enzymes with low iron loading (Andreou et al. 2010; Deschamps et al. 2016; Goloshchapova et al. 2018; Chrisnasari et al. 2024b). Various strategies, such as lowering temperature and inducer concentrations, have been employed to balance gene expression and protein folding efficiency (Andreou et al. 2008; An et al. 2015, 2018a; Pang et al. 2020; Chrisnasari et al. 2024b). Yet, these adjustments were not always successful.

To fully harness the potential of bacterial LOXs, a paradigm shift in expression systems and optimization strategies is essential. The recent successful overexpression of *Pseudomonas aeruginosa* LOX in the yeast *Komagataella phaffii* (formerly known as *Pichia pastoris*) offers a promising alternative (Hashem et al. 2020). This yeast system facilitates the secretion of LOX into the supernatant, significantly simplifying downstream processing compared to *E. coli*. However, key factors, such as the iron content of the expressed protein, remain unassessed. Therefore, further evaluation of the different expression systems is crucial to unlocking the full potential of bacterial LOXs for various applications.

In addition to production challenges, some bacterial LOXs have been reported to be thermally unstable (Busquets et al. 2004; Qian et al. 2017; Chrisnasari et al. 2024b). Similarly, low thermostability has also been reported for soybean mini-LOX, an enzyme that lacks the N-terminal beta-barrel domain found in native soybean LOX (Di Venere et al. 2003). The low stability of this mini-LOX is attributed to greater hydration compared to the native enzyme (Di Venere et al. 2003). As most bacterial LOXs have been predicted not to contain an N-terminal beta-barrel domain (Chrisnasari et al. 2022), it is possible that these enzymes are more hydrated and thus less stable. This finding highlights the urgent need for strategies to enhance the stability of bacterial LOXs. Rational protein engineering through mutagenesis, guided by computer-aided approaches such as FRESCO (Wijma et al. 2018) or computationally assisted structure-guided design (Chi et al. 2026), offers a viable solution. Techniques such as site-directed mutagenesis (Guo et al. 2014), deletion of the highly flexible region (Lu et al. 2014), and fusion with self-assembling amphipathic peptides (Lu et al. 2013a) have demonstrated success in improving the thermostability of bacterial LOXs.

Overcoming these challenges requires a multidisciplinary approach that integrates advanced expression systems, enhanced protein engineering, and novel stabilization techniques. Addressing these limitations will not only pave the way for the efficient large-scale production of bacterial LOXs, but also unlock their full potential in biotechnological applications. However, applying bacterial LOXs across various industries requires balancing the benefits of “green” production with the often higher costs of enzymatic biocatalysis. Therefore, LOXs are particularly well-suited for synthesizing high-value products, such as lipid mediators, where these advantages justify the investment.

Conclusions

Bacterial LOXs are promising biocatalysts for fatty acid oxygenation, utilizing regioselective dioxygenation and hydroperoxide isomerization activities. Strategies such as protein engineering and oxygen level modulation effectively

modify substrate specificity, regioselectivity, and the preferred catalytic pathway. Multi-enzyme cascades incorporating LOXs facilitate the production of diverse flavors, fragrances, oleochemicals, and oxylipins. This review also highlights efficient pathways for synthesizing oxylipins via LOX-mediated double dioxygenation and combined dioxygenase-hydroperoxide isomerase activities. To fully realize their potential, advancements in expression systems and enzyme thermostability are essential. This review not only enhances our understanding of bacterial LOX enzymology but also provides a foundation for their broader adoption in sustainable and innovative biotechnological applications.

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1007/s00253-026-13837-7>.

Authors contribution RC: Conceptualization; data analysis and visualization; writing—original draft, review, and editing. TE and MH: Conceptualization; writing—review and editing; supervision of RC. JPV and WB: Writing—review and editing; supervision of RC.

Funding This study was funded by Indonesian Endowment Fund for Education (LPDP) (grant number 0006605/BIO/D/BUDI-2019).

Data availability No datasets were generated or analysed during the current study.

Declarations

Ethical approval This article does not contain any studies with human participants or animals performed by any of the authors.

Competing interests The authors declare no competing interests.

Open Access This article is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License, which permits any non-commercial use, sharing, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if you modified the licensed material. You do not have permission under this licence to share adapted material derived from this article or parts of it. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by-nc-nd/4.0/>.

References

- Ambaw YA, Fuchs D, Raida M, Mazengia NT, Torta F, Wheelock CE, Wenk MR, Tong L (2020) Changes of tear lipid mediators after eyelid warming or thermopulsation treatment for meibomian gland dysfunction. *Prostaglandins Other Lipid Mediat*. <https://doi.org/10.1016/j.prostaglandins.2020.106474>
- An JU, Kim BJ, Hong SH, Oh DK (2015) Characterization of an omega-6 linoleate lipoxygenase from *Burkholderia thailandensis*

- and its application in the production of 13-hydroxyoctadecadienoic acid. *Appl Microbiol Biotechnol* 99:5487–5497. <https://doi.org/10.1007/s00253-014-6353-8>
- An JU, Hong SH, Oh DK (2018a) Regiospecificity of a novel bacterial lipoxygenase from *Myxococcus xanthus* for polyunsaturated fatty acids. *Biochim Biophys Acta Mol Cell Biol Lipids* 1863:823–833. <https://doi.org/10.1016/j.bbalip.2018a.04.014>
- An JU, Song YS, Kim KR, Ko YJ, Yoon DY, Oh DK (2018) Biotransformation of polyunsaturated fatty acids to bioactive heptoxilins and trioxilins by microbial enzymes. *Nat Commun* 9:1–10. <https://doi.org/10.1038/s41467-017-02543-8>
- An JU, Kim SE, Oh DK (2021) Molecular insights into lipoxygenases for biocatalytic synthesis of diverse lipid mediators. *Prog Lipid Res*. <https://doi.org/10.1016/j.plipres.2021.101110>
- Andreou A, Feussner I (2009) Lipoxygenases - structure and reaction mechanism. *Phytochemistry* 70:1504–1510. <https://doi.org/10.1016/j.phytochem.2009.05.008>
- Andreou A, Vanko M, Bezakova L, Feussner I (2008) Properties of a mini 9R-lipoxygenase from *Nostoc* sp. PCC 7120 and its mutant forms. *Phytochemistry* 69:1832–1837. <https://doi.org/10.1016/j.phytochem.2008.03.002>
- Andreou A, Göbel C, Hamberg M, Feussner I (2010) A bisallylic mini-lipoxygenase from *Cyanobacterium Cyanotheca* sp. that has an iron as cofactor. *J Biol Chem* 285:14178–14186. <https://doi.org/10.1074/jbc.M109.094771>
- Aranda C, Olmedo A, Kiebish K, del Río JC, Martínez AT, Gutiérrez A (2018) Selective epoxidation of fatty acids and fatty acid methyl esters by fungal peroxygenases. *ChemCatChem* 10:3964–3968. <https://doi.org/10.1002/cctc.201800849>
- Banthiya S, Kalmes J, Galemou Yoga E, Ivanov I, Carpena X, Hamberg M, Kuhn H, Scheerer P (2016) Structural and functional basis of phospholipid oxygenase activity of bacterial lipoxygenase from *Pseudomonas aeruginosa*. *Biochimica et Biophysica Acta (BBA) - Molecular and Cell Biology of Lipids* 1861:1681–1692. <https://doi.org/10.1016/j.bbalip.2016.08.002>
- Basil MC, Levy BD (2015) Specialized pro-resolving mediators: endogenous regulators of infection and inflammation. *Nat Rev Immunol* 16(1):51–67. <https://doi.org/10.1038/NRI.2015.4>
- Björkhem I (1972) On the role of alcohol dehydrogenase in ω -oxidation of fatty acids. *Eur J Biochem* 30:441–451. <https://doi.org/10.1111/j.1432-1033.1972.tb02116.x>
- Blée E (2002) Impact of phyto-oxylipins in plant defense. *Trends Plant Sci* 7:315–322. [https://doi.org/10.1016/S1360-1385\(02\)02290-2](https://doi.org/10.1016/S1360-1385(02)02290-2)
- Blum M, Dogan I, Karber M, Rothe M, Schunck WH (2019) Chiral lipidomics of monoepoxy and monohydroxy metabolites derived from long-chain polyunsaturated fatty acids. *J Lipid Res* 60:135–148. <https://doi.org/10.1194/jlr.M089755>
- Borngräber S, Kuban RJ, Anton M, Kühn H (1996) Phenylalanine 353 is a primary determinant for the positional specificity of mammalian 15-lipoxygenases. *J Mol Biol* 264:1145–1153. <https://doi.org/10.1006/jmbi.1996.0702>
- Borngräber S, Browner M, Gillmor S, Gerth C, Anton M, Fletterick R, Kuhn H (1999) Shape and specificity in mammalian 15-lipoxygenase active site. The functional interplay of sequence determinants for the reaction specificity. *J Biol Chem* 274:37345–37350. <https://doi.org/10.1074/JBC.274.52.37345>
- Bornscheuer U (2018) Lipid modification by enzymes and engineered microbes. Academic Press and AOCS Press, London
- Brash AR (1999) Lipoxygenases: occurrence, functions, catalysis, and acquisition of substrate. *J Biol Chem* 274:23679–23682. <https://doi.org/10.1074/jbc.274.34.23679>
- Brühlmann F, Bosijokovic B (2016) Efficient biochemical cascade for accessing green leaf alcohols. *Org Process Res Dev* 20:1974–1978. <https://doi.org/10.1021/ACS.OPRD.6B00303>
- Brühlmann F, Bosijokovic B, Ullmann C, Auffray P, Fourage L, Wahler D (2013) Directed evolution of a 13-hydroperoxide lyase (CYP74B) for improved process performance. *J Biotechnol* 163:339–345. <https://doi.org/10.1016/j.jbiotec.2012.11.005>
- Busquets M, Deroncelé V, Vidal-Mas J, Rodríguez E, Guerrero A, Manresa A (2004) Isolation and characterization of a lipoxygenase from *Pseudomonas* 42A2 responsible for the biotransformation of oleic acid into (S)-(E)-10-hydroxy-8-octadecenoic acid. *Antonie van Leeuwenhoek* 85:129–139. <https://doi.org/10.1023/B:ANTO.0000020152.15440.65>
- Chi H, Xia B, Shen J, Jiang Q, Lu Z, Lu F, Zhu P, Zhu X (2026) Overcoming the stability–activity trade-off in lipoxygenase by integrated computational-assisted structure-guided design. *J Agric Food Chem* 74:8834–8850. <https://doi.org/10.1021/ACS.JAFC.5C16524>
- Chrisnasari R, Hennebelle M, Vincken J-P, van Berkel WJH, Ewing TA (2022) Bacterial lipoxygenases : biochemical characteristics, molecular structure and potential applications. *Biotechnol Adv*. <https://doi.org/10.1016/j.biotechadv.2022.108046>
- Chrisnasari R, Ewing TA, Hilgers R, van Berkel WJH, Vincken J-P, Hennebelle M (2024) Versatile ferrous oxidation–xylene orange assay for high-throughput screening of lipoxygenase activity. *Appl Microbiol Biotechnol*. <https://doi.org/10.1007/s00253-024-13095-5>
- Chrisnasari R, Hennebelle M, Nguyen KA, Vincken J-P, van Berkel WJH, Ewing TA (2024b) Engineering the substrate specificity and regioselectivity of *Burkholderia thailandensis* lipoxygenase. *N Biotechnol* 84:64–76. <https://doi.org/10.1016/j.nbt.2024b.09.007>
- Chrisnasari R, Chen S, Hilgers R, van Vliet D, van Berkel WJH, Vincken J-P, Ewing TA, Hennebelle M (2025a) Catalytic versatility of lipoxygenase from *Microcystis aeruginosa*. *N Biotechnol* 91:36–51. <https://doi.org/10.1016/j.nbt.2025a.11.005>
- Chrisnasari R, Hilgers R, Li G, Vincken J-P, van Berkel WJH, Hennebelle M, Ewing TA (2025b) Modulating dioxygenase and hydroperoxide isomerase activities in *Burkholderia thailandensis* lipoxygenase. *Enzyme Microb Technol* 191. <https://doi.org/10.1016/J.ENZMICTEC.2025b.110709>
- Coffa G, Brash AR (2004) A single active site residue directs oxygenation stereospecificity in lipoxygenases: stereocontrol is linked to the position of oxygenation. *Proc Natl Acad Sci U S A* 101:15579–15584. <https://doi.org/10.1073/pnas.0406727101>
- Coffa G, Schneider C, Brash AR (2005) A comprehensive model of positional and stereo control in lipoxygenases. *Biochem Biophys Res Commun* 338:87–92. <https://doi.org/10.1016/j.bbrc.2005.07.185>
- Collazo L, Klimman JP (2016) Control of the position of oxygen delivery in soybean lipoxygenase-1 by amino acid side chains within a gas migration channel. *J Biol Chem* 291:9052–9059. <https://doi.org/10.1074/jbc.M115.709154>
- Cristea M, Oliu EH (2006) A G316A mutation of manganese lipoxygenase augments hydroperoxide isomerase activity: mechanism of biosynthesis of epoxyalcohols. *J Biol Chem* 281:17612–17623. <https://doi.org/10.1074/jbc.M510311200>
- De Bartolo A, Romeo N, Angelone T, Rocca C (2025) Specialized pro-resolving mediators as emerging players in cardioprotection: from inflammation resolution to therapeutic potential. *Acta Physiol*. <https://doi.org/10.1111/apha.70062>
- de Espinosa LM, Meier MAR (2011) Plant oils: the perfect renewable resource for polymer science?! *Eur Polym J* 47:837–852. <https://doi.org/10.1016/j.eurpolymj.2010.11.020>
- de Groot JJMC, Garssen GJ, Veldink GA, Vliegthart JFG, Boldingh J, Egmond MR (1975) On the interaction of soybean lipoxygenase-1 and 13-L-hydroperoxylinoleic acid, involving yellow and purple coloured enzyme species. *FEBS Lett* 56:50–54. [https://doi.org/10.1016/0014-5793\(75\)80109-8](https://doi.org/10.1016/0014-5793(75)80109-8)

- Deschamps JD, Oguniola AF, Li JBJ, Yasgar A, Flitter BA, Freedman CJ, Melvin JA, Nguyen JVMH, David J, Jadhav A, Simeonov A, Bomberger JM, Theodore R (2016) Biochemical/cellular characterization and inhibitor discovery of *Pseudomonas aeruginosa* 15-lipoxygenase. *Biochemistry* 55:3329–3340. <https://doi.org/10.1021/acs.biochem.6b00338>
- Di Venere A, Salucci ML, Van Zadelhoff G, Veldink G, Mei G, Rosato N, Finazzi-Agrò A, Maccarrone M (2003) Structure-to-function relationship of mini-lipoxygenase, a 60-kDa fragment of soybean lipoxygenase-1 with lower stability but higher enzymatic activity. *J Biol Chem* 278:18281–18288. <https://doi.org/10.1074/jbc.M212122200>
- Egmond MR, Vliegthart JFG, Boldingh J (1972) Stereospecificity of the hydrogen abstraction at carbon atom n-8 in the oxygenation of linoleic acid by lipoxygenases from corn germs and soya beans. *Biochem Biophys Res Commun* 48:1055–1060. [https://doi.org/10.1016/0006-291X\(72\)90815-7](https://doi.org/10.1016/0006-291X(72)90815-7)
- Ellamar JB, Kim IH, Hou CT, Park HS, Kim HR (2019) Isolation and identification of a variant strain of *Pseudomonas aeruginosa* PR3 with enhanced production of 7,10-dihydroxy-8(E)-octadecenoic acid. *Biocatal Agric Biotechnol*. <https://doi.org/10.1016/j.bcab.2019.101068>
- Faillace E, Brunini-Bronzini de Caraffa V, Gambotti C, Berti L, Maury J, Vincenti S (2025) Optimizing the second step of the biocatalytic process for green leaf volatiles production: synthesis by soybean 13-lipoxygenase of 13-hydroperoxides from hydrolyzed hempseed oil. *Biocatal Agric Biotechnol*. <https://doi.org/10.1016/j.bcab.2025.103548>
- Feussner I, Wasternack C (1998) Lipoxygenase catalyzed oxygenation of lipids. *Lipid - Fett* 100:146–152. [https://doi.org/10.1002/\(sici\)1521-4133\(1998\)100:4/5<146::aid-lipi146>3.0.co;2-d](https://doi.org/10.1002/(sici)1521-4133(1998)100:4/5<146::aid-lipi146>3.0.co;2-d)
- Gao B, Boeglin WE, Brash AR (2010) Omega-3 fatty acids are oxygenated at the n-7 carbon by the lipoxygenase domain of a fusion protein in the cyanobacterium *Acaryochloris marina*. *Biochim Biophys Acta Mol Cell Biol Lipids* 1801:58–63. <https://doi.org/10.1016/J.BBALIP.2009.09.004>
- Garssen GJ, Veldink GA, Vliegthart JFG, Boldingh J (1976) The formation of threo-11-hydroxy-*trans*-12: 13-epoxy-9-*cis*-octadecenoic acid by enzymic isomerisation of 13-*l*-hydroperoxy-9-*cis*, 11-*trans*-octadecadienoic acid by soybean lipoxygenase-1. *Eur J Biochem* 62:33–36. <https://doi.org/10.1111/j.1432-1033.1976.tb10094.x>
- Goloshchapova K, Stehling S, Heydeck D, Blum M, Kuhn H (2018) Functional characterization of a novel arachidonic acid 12S-lipoxygenase in the halotolerant bacterium *Myxococcus fulvus* exhibiting complex social living patterns. *MicrobiologyOpen* 8:1–17. <https://doi.org/10.1002/mbo3.775>
- Grechkin AN, Hamberg M (2004) The “heterolytic hydroperoxide lyase” is an isomerase producing a short-lived fatty acid hemiacetal. *Biochimica et Biophysica Acta (BBA) - Molecular and Cell Biology of Lipids* 1636:47–58. <https://doi.org/10.1016/j.bbalip.2003.12.003>
- Grechkin AN, Brühlmann F, Mukhtarova LS, Gogolev YV, Hamberg M (2006) Hydroperoxide lyases (CYP74C and CYP74B) catalyze the homolytic isomerization of fatty acid hydroperoxides into hemiacetals. *Biochimica et Biophysica Acta (BBA) - Molecular and Cell Biology of Lipids* 1761:1419–1428. <https://doi.org/10.1016/j.bbalip.2006.09.002>
- Guo F, Zhang C, Bie X, Zhao H, Diao H, Lu F, Lu Z (2014) Improving the thermostability and activity of lipoxygenase from *Anabaena* sp. PCC 7120 by directed evolution and site-directed mutagenesis. *J Mol Catal B Enzym* 107:23–30. <https://doi.org/10.1016/j.molcatb.2014.05.016>
- Hagedoorn PL, Hollmann F, Hanefeld U (2021) Novel oleate hydratases and potential biotechnological applications. *Appl Microbiol Biotechnol* 105:6159–6172. <https://doi.org/10.1007/s00253-021-11465-x>
- Hamberg M, Samuelsson B (1967) On the specificity of the oxygenation of unsaturated fatty acids catalyzed by soybean lipoxidase. *J Biol Chem* 242:5329–5335. [https://doi.org/10.1016/S0021-9258\(18\)99432-9](https://doi.org/10.1016/S0021-9258(18)99432-9)
- Hamberg M, Su C, Oliw E (1998) Manganese lipoxygenase. Discovery of a bis-allylic hydroperoxide as product and intermediate in a lipoxygenase reaction. *J Biol Chem* 273:13080–13088. <https://doi.org/10.1074/JBC.273.21.13080>
- Hansen J, Garreta A, Benincasa M, Fusté MC, Busquets M, Manresa A (2013) Bacterial lipoxygenases, a new subfamily of enzymes? A phylogenetic approach. *Appl Microbiol Biotechnol* 97:4737–4747. <https://doi.org/10.1007/s00253-013-4887-9>
- Hashem C, Stollerfoht H, Rinnofner C, Steinberger S, Winkler M, Pichler H (2020) Secretion *Pseudomonas aeruginosa* lipoxygenase by *Pichia pastoris* upon glycerol feed. *Biotechnol J* 15. <https://doi.org/10.1002/biot.202000089>
- Hayward S, Cilliers T, Swart P (2017) Lipoxygenases: from isolation to application. *Compr Rev Food Sci Food Saf* 16:199–211. <https://doi.org/10.1111/1541-4337.12239>
- Heshof R, De Graaff LH, Villaverde JJ, Silvestre AJD, Haarmann T, Dalsgaard TK, Buchert J (2016) Industrial potential of lipoxygenases. *Crit Rev Biotechnol* 36:665–674. <https://doi.org/10.3109/07388551.2015.1004520>
- Hou CT (2009) Biotechnology for fats and oils: new oxygenated fatty acids. *N Biotechnol* 26:2–10. <https://doi.org/10.1016/j.nbt.2009.05.001>
- Ide MS, Hao B, Neurock M, Davis RJ (2012) Mechanistic insights on the hydrogenation of α,β -unsaturated ketones and aldehydes to unsaturated alcohols over metal catalysts. *ACS Catal* 2:671–683. <https://doi.org/10.1021/cs200567z>
- Ishihara T, Yoshida M, Arita M (2019) Omega-3 fatty acid-derived mediators that control inflammation and tissue homeostasis. *Int Immunol* 31:559–567. <https://doi.org/10.1093/INTIMM/DXZ001>
- Isobe Y, Arita M, Iwamoto R, Urabe D, Todoroki H, Masuda K, Inoue M, Arai H (2013) Stereochemical assignment and anti-inflammatory properties of the omega-3 lipid mediator resolvin E3. *J Biochem* 153:355–360. <https://doi.org/10.1093/jb/mvs151>
- Johannsen J, Meyer F, Fieg G, Waluga T (2020) Process for the preparation of fatty aldehydes from renewable resources. *Chem Ing Tech* 92:777–781. <https://doi.org/10.1002/cite.201900111>
- Johannsen J, Baek G, Fieg G, Waluga T (2021) An innovative approach for fatty acid reduction to fatty aldehydes. *Green Chem Lett Rev* 14:454–460. <https://doi.org/10.1080/17518253.2021.1943006>
- Joo YC, Oh DK (2012) Lipoxygenases: potential starting biocatalysts for the synthesis of signaling compounds. *Biotechnol Adv* 30:1524–1532. <https://doi.org/10.1016/j.biotechadv.2012.04.004>
- Kim SK, Park YC (2019) Biosynthesis of ω -hydroxy fatty acids and related chemicals from natural fatty acids by recombinant *Escherichia coli*. *Appl Microbiol Biotechnol* 103:191–199
- Kim TH, Lee J, Kim SE, Oh DK (2021) Biocatalytic synthesis of dihydroxy fatty acids as lipid mediators from polyunsaturated fatty acids by double dioxygenation of the microbial 12S-lipoxygenase. *Biotechnol Bioeng* 118:3094–3104. <https://doi.org/10.1002/bit.27820>
- Kim SE, Lee J, An JU, Kim TH, Oh CW, Ko YJ, Krishnan M, Choi J, Yoon DY, Kim Y, Oh DK (2022) Regioselectivity of an arachidonate 9S-lipoxygenase from *Sphingopyxis macrogoltabida* that biosynthesizes 9S,15S- and 11S,17S-dihydroxy fatty acids from C20 and C22 polyunsaturated fatty acids. *Biochim Biophys Acta Mol Cell Biol Lipids*. <https://doi.org/10.1016/J.BBALIP.2021.159091>
- Knapp MJ, Seebeck FP, Klinman JP (2001) Steric control of oxygenation regiochemistry in soybean lipoxygenase-1. *J Am Chem Soc* 123:2931–2932. <https://doi.org/10.1021/ja003855k>

- Kobe MJ, Neau DB, Mitchell CE, Bartlett SG, Newcomer ME (2014) The structure of human 15-lipoxygenase-2 with a substrate mimic. *J Biol Chem* 289:8562–8569. <https://doi.org/10.1074/jbc.M113.543777>
- Köckritz A, Martin A (2011) Synthesis of azelaic acid from vegetable oil-based feedstocks. *Eur J Lipid Sci Technol* 113:83–91. <https://doi.org/10.1002/ejlt.201000117>
- Konkel A, Schunck WH (2011) Role of cytochrome P450 enzymes in the bioactivation of polyunsaturated fatty acids. *Biochim Biophys Acta Proteins Proteom* 1814:210–222. <https://doi.org/10.1016/j.bbapap.2010.09.009>
- Kuhn H, Saam J, Eibach S, Holzhütter HG, Ivanov I, Walther M (2005) Structural biology of mammalian lipoxygenases: enzymatic consequences of targeted alterations of the protein structure. *Biochem Biophys Res Commun* 338:93–101
- Lee J, An J-U, Kim T-H, Ko Y-J, Park J-B, Oh D-K (2022) Discovery and engineering of a microbial double-oxygenating lipoxygenase for synthesis of dihydroxy fatty acids as specialized pro-resolving mediators. *ACS Sustain Chem Eng* 8:16172–16183. <https://doi.org/10.1021/acssuschemeng.0c04793>
- Lee TE, Shin KC, Park JB, Oh DK (2025) Specialized pro-resolving mediators: biosynthetic pathways, biocatalytic synthesis, and applications. *Biotechnol Adv*. <https://doi.org/10.1016/j.biotechadv.2025.108682>
- Lehnert N, Solomon EI (2003) Density-functional investigation on the mechanism of H-atom abstraction by lipoxygenase. *J Biol Inorg Chem* 8:294–305. <https://doi.org/10.1007/S00775-002-0415-6>
- Lu X, Liu S, Zhang D, Zhou X, Wang M, Liu Y, Wu J, Du G, Chen J (2013) Enhanced thermal stability and specific activity of *Pseudomonas aeruginosa* lipoxygenase by fusing with self-assembling amphipathic peptides. *Appl Microbiol Biotechnol* 97:9419–9427. <https://doi.org/10.1007/s00253-013-4751-y>
- Lu X, Zhang J, Liu S, Zhang D, Xu Z, Wu J, Li J, Du G, Chen J (2013) Overproduction, purification, and characterization of extracellular lipoxygenase of *Pseudomonas aeruginosa* in *Escherichia coli*. *Appl Microbiol Biotechnol* 97:5793–5800. <https://doi.org/10.1007/s00253-012-4457-6>
- Lu X, Liu S, Feng Y, Rao S, Zhou X, Wang M, Du G, Chen J (2014) Enhanced thermal stability of *Pseudomonas aeruginosa* lipoxygenase through modification of two highly flexible regions. *Appl Microbiol Biotechnol* 98:1663–1669. <https://doi.org/10.1007/s00253-013-5039-y>
- Manivarma T, Nowak W, Tyurina YY, Tyurin VA, Bayir H, Kagan VE, Mikulska-Ruminska K (2025) The presence of substrate warrants oxygen access tunnels toward the catalytic site of lipoxygenases. *Redox Biol*. <https://doi.org/10.1016/j.redox.2025.103636>
- Milone C, Crisafulli C, Ingoglia R, Schipilliti L, Galvagno S (2007) A comparative study on the selective hydrogenation of α,β unsaturated aldehyde and ketone to unsaturated alcohols on Au supported catalysts. *Catal Today* 122:341–351. <https://doi.org/10.1016/j.cattod.2007.01.011>
- Nanda S, Yadav JS (2003) Lipoxygenase biocatalysis: a survey of asymmetric oxygenation. *J Mol Catal B Enzym* 26:3–28. [https://doi.org/10.1016/S1381-1177\(03\)00146-2](https://doi.org/10.1016/S1381-1177(03)00146-2)
- Neau DB, Bender G, Boeglin WE, Bartlett SG, Brash AR, Newcomer ME (2014) Crystal structure of a lipoxygenase in complex with substrate: the arachidonic acid-binding site of 8R-lipoxygenase. *J Biol Chem* 289:31905–31913. <https://doi.org/10.1074/jbc.M114.599662>
- Newcomer ME, Brash AR (2015) The structural basis for specificity in lipoxygenase catalysis. *Protein Sci* 24:298–309. <https://doi.org/10.1002/pro.2626>
- Newie J, Andreou A, Neumann P, Einsle O, Feussner I, Ficner R (2016) Crystal structure of a lipoxygenase from *Cyanotheca* sp. may reveal novel features for substrate acquisition. *J Lipid Res* 57:276–286. <https://doi.org/10.1194/JLR.M064980>
- Oh SF, Vickery TW, Serhan CN (2011) Chiral lipidomics of E-series resolvins: aspirin and the biosynthesis of novel mediators. *Biochim Biophys Acta Mol Cell Biol Lipids* 1811:737–747. <https://doi.org/10.1016/j.bbailip.2011.06.007>
- Oh CW, Kim SE, Lee J, Oh DK (2022) Bioconversion of C20- and C22-polyunsaturated fatty acids into 9S,15S- and 11S,17S-dihydroxy fatty acids by *Escherichia coli* expressing double-oxygenating 9S-lipoxygenase from *Sphingopyxis macrogoltabida*. *J Biosci Bioeng* 134:14–20. <https://doi.org/10.1016/J.JBIOOSC.2022.04.001>
- Pang C, Liu S, Zhang G, Zhou J, Du G, Li J (2020) Combinatorial strategy towards the efficient expression of lipoxygenase in *Escherichia coli* at elevated temperatures. *Appl Microbiol Biotechnol* 104:10047–10057. <https://doi.org/10.1007/s00253-020-10941-0>
- Qi Y-K, Zheng Y-C, Zhang Z-J, Xu J-H (2020) Efficient transformation of linoleic acid into 13(S)-hydroxy-9,11-(Z,E)-octadecadienoic acid using putative lipoxygenases from *Cyanobacteria*. *ACS Sustain Chem Eng* 8:5558–5565. <https://doi.org/10.1021/acssuschemeng.9b07457>
- Qian H, Xia B, He Y, Lu Z, Bie X, Zhao H, Zhang C, Lu F (2017) Expression, purification, and characterization of a novel acidic lipoxygenase from *Myxococcus xanthus*. *Protein Expr Purif* 138:13–17. <https://doi.org/10.1016/j.pep.2017.05.006>
- Quaranta E, Dibenedetto A, Colucci A, Cornacchia D (2022) Partial hydrogenation of FAMES with high content of C18:2 dienes. Selective hydrogenation of tobacco seed oil-derived biodiesel. *Fuel*. <https://doi.org/10.1016/j.fuel.2022.125030>
- Revol-Cavalier J, Quaranta A, Newman JW, Brash AR, Hamberg M, Wheelock CE (2025) The octadecanoids: synthesis and bioactivity of 18-carbon oxygenated fatty acids in mammals, bacteria, and fungi. *Chem Rev* 125:1–90. <https://doi.org/10.1021/acs.chemrev.3c00520>
- Ribeaucourt D, Bissaro B, Lambert F, Lafond M, Berrin JG (2022) Biocatalytic oxidation of fatty alcohols into aldehydes for the flavors and fragrances industry. *Biotechnol Adv*. <https://doi.org/10.1016/j.biotechadv.2021.107787>
- Riera A, Moreno M (2010) Synthetic applications of chiral unsaturated epoxy alcohols prepared by Sharpless asymmetric epoxidation. *Molecules* 15:1041–1073. <https://doi.org/10.3390/molecules15021041>
- Saam J, Ivanov I, Walther M, Holzhü H-G, Kuhn H, Klinman JP (2007) Molecular dioxygen enters the active site of 12/15-lipoxygenase via dynamic oxygen access channels. *PNAS* 104:13319–13324
- Schilstra MJ, Veldink GA, Verhagen J, Vliegthart JFG (1992) Effect of lipid hydroperoxide on lipoxygenase kinetics. *Biochemistry* 31:7692–7699. <https://doi.org/10.1021/bi00148a033>
- Schilstra MJ, Veldink GA, Vliegthart JFG (1993) Kinetic analysis of the induction period in lipoxygenase catalysis. *Biochemistry* 32:7686–7691. <https://doi.org/10.1021/bi00081a012>
- Schilstra MJ, Veldink GA, Vliegthart JFG (1994) The dioxygenation rate in lipoxygenase catalysis is determined by the amount of Iron(III) lipoxygenase in solution. *Biochemistry* 33:3974–3979. <https://doi.org/10.1021/bi00179a025>
- Seo JH, Lee SM, Lee J, Park JB (2015) Adding value to plant oils and fatty acids: biological transformation of fatty acids into ω -hydroxycarboxylic, α,ω -dicarboxylic, and ω -aminocarboxylic acids. *J Biotechnol* 216:158–166. <https://doi.org/10.1016/j.jbiotec.2015.10.024>
- Seo EJ, Lee HR, Hwang SY, Oh DK, Kwon YU, Köchl K, Nestl B, Park JB, Bornscheuer UT (2025) Substrate-binding cavity engineering of the lipoxygenase from *Pseudomonas aeruginosa* to produce 8S- and 11S-hydroxyeicosatetraenoic acids.

- ACS Sustain Chem Eng 13:80–90. <https://doi.org/10.1021/acssuschemeng.4c05400>
- Serhan CN (2014) Pro-resolving lipid mediators are leads for resolution physiology. *Nature* 510:92–101. <https://doi.org/10.1038/nature13479>
- Sim DH, Shin KC, Oh DK (2015) 13-hydroxy-9Z,11E-octadecadienoic acid production by recombinant cells expressing *Burkholderia thailandensis* 13-lipoxygenase. *J Am Oil Chem Soc* 92:1259–1266. <https://doi.org/10.1007/s11746-015-2694-4>
- Sloane DL, Leung R, Cralk CS, Sigal E (1991) A primary determinant for lipoxygenase positional specificity. *Nature* 354:149. <https://doi.org/10.1038/354149a0>
- Song JW, Jeon EY, Song DH, Jang HY, Bornscheuer UT, Oh DK, Park JB (2013) Multistep enzymatic synthesis of long-chain α,ω -dicarboxylic and ω -hydroxycarboxylic acids from renewable fatty acids and plant oils. *Angew Chem Int Ed* 52:2534–2537. <https://doi.org/10.1002/anie.201209187>
- Spiekermann ML, Seidensticker T (2024) Catalytic processes for the selective hydrogenation of fats and oils: reevaluating a mature technology for feedstock diversification. *Catal Sci Technol* 14:4390–4419. <https://doi.org/10.1039/d4cy00488d>
- Van Os CPA, Rijke-Schilder GPM, Van Halbeek H, Verhagen J, Vliegthart JFG (1981) Double dioxygenation of arachidonic acid by soybean lipoxygenase-1 kinetics and regio-stereo specificities of the reaction steps. *Biochim Biophys Acta* 663:177–193. [https://doi.org/10.1016/0005-2760\(81\)90204-6](https://doi.org/10.1016/0005-2760(81)90204-6)
- Vidal-Mas J, Busquets M, Manresa A (2005) Cloning and expression of a lipoxygenase from *Pseudomonas aeruginosa* 42A2. *Antonie Van Leeuwenhoek* 87:245–251. <https://doi.org/10.1007/s10482-004-4021-1>
- Vogel R, Jansen C, Roffeis J, Reddanna P, Forsell P, Claesson HE, Kuhn H, Walther M (2010) Applicability of the triad concept for the positional specificity of mammalian lipoxygenases. *J Biol Chem* 285:5369–5376. <https://doi.org/10.1074/JBC.M109.057802>
- Wijma HJ, Fürst MJLJ, Janssen DB (2018) A computational library design protocol for rapid improvement of protein stability: FRESKO. *Methods Mol Biol* 1685:69–85. https://doi.org/10.1007/978-1-4939-7366-8_5
- Yadav JS, Srinivas Dale (1997) Titanocene induced deoxygenation of di-epoxides - a convenient regioselective preparation of 1,3-diols. *Chem Lett CL-970345:905–906*. <https://doi.org/10.1246/cl.1997.905>
- Yu Z, Schneider C, Boeglin WE, Marnett LJ, Brash AR (2003) The lipoxygenase gene ALOXE3 implicated in skin differentiation encodes a hydroperoxide isomerase. *PNAS* 100:9162–9167. <https://doi.org/10.1073/pnas.1633612100>
- Yu JM, Liu YY, Zheng YC, Li H, Zhang XY, Zheng GW, Li CX, Bai YP, Xu JH (2018) Direct access to medium-chain α,ω -dicarboxylic acids by using a Baeyer-Villiger monooxygenase of abnormal regioselectivity. *Chembiochem* 19:2049–2054. <https://doi.org/10.1002/cbic.201800318>
- Zhang Q, Wang Y, Zhu J, Zou M, Zhang Y, Wu H, Jin T (2025) Specialized pro-resolving lipid mediators: a key player in resolving inflammation in autoimmune diseases. *Sci Bull (Beijing)* 70:778–794. <https://doi.org/10.1016/j.scib.2024.07.049>
- Zheng Y, Brash AR (2010) On the role of molecular oxygen in lipoxygenase activation: comparison and contrast of epidermal lipoxygenase-3 with soybean lipoxygenase-1. *J Biol Chem* 285:39876–39887. <https://doi.org/10.1074/jbc.M110.180794>
- Žula M, Grilc M, Likozar B (2022) Hydrocracking, hydrogenation and hydro-deoxygenation of fatty acids, esters and glycerides: mechanisms, kinetics and transport phenomena. *Chem Eng J*. <https://doi.org/10.1016/j.cej.2022.136564>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Applied and Microbiology Biotechnology

Volume 109 Number 1 December 2025

- | | | | |
|---|---|----|--|
| 1 | Targeted knockdown of ATM, ATR, and PDEδ increases Gag HIV-1 VLP production in HEK293 cells
A. Díaz-Maneh · P. Pérez-Rubio · C.R. Granes · L. Bosch-Molist · J. Lavado-García · F. Gòdia · L. Cervera | 6 | Exploring <i>Tetraselmis chui</i> microbiomes—functional metagenomics for novel catalases and superoxide dismutases
J.F.H. Macdonald · Y. Han · Y. Astafyeva · L. Bergmann · M. Gurschke · P. Dirksen · P. Blümke · Y.K.H. Schneider · M. Alawi · S. Lippemeier · J.H. Andersen · I. Krohn |
| 2 | Precision tumor treatment utilizing bacteria: principles and future perspectives
Z. Liu · L. Wang · P. Wu · L. Yuan | 7 | Characterization and design of dipeptide media formulation for scalable therapeutic production
J. Song · S.-Y. Park · D.-Y. Lee |
| 3 | Anxiety of microbially synthesized Fe₃O₄-SPIONs on embryonic/larval ontogeny in red tilapia (<i>Oreochromis</i> sp.)
S.S. Abouelkheir · M.M. Mourad | 8 | A Lambda-evo (λ_{evo}) phage platform for Zika virus E_{DIII} protein display
H. Negrete-Méndez · G. Valencia-Toxqui · E. Martínez-Peñañiel · O. Medina-Contreras · F. Fernández-Ramírez · E. Morales-Ríos · L.J. Navarro-González · J.M. Torres-Flores · L. Kameyama |
| 4 | Identification and monitoring of cell heterogeneity from plasmid recombination during limonene production
L. Gelain · J.W. Yeoh · G.S. Hossain · S. Alfenore · S. Guillouet · H. Ling · C.L. Poh · N. Gorret · J.L. Foo | 9 | Microaerobic biodegradation of aromatic hydrocarbon mixtures: strategies for efficient nitrate and oxygen dosage
D.C. Aydin · A. Aldas-Vargas · T. Grotenhuis · H. Rijnaarts |
| 5 | Evaluation of MALDI-TOF for identification of <i>Vibrio cholerae</i> and <i>Vibrio parahaemolyticus</i> from growth on agar media
S. Banerjee · A. Flint · M.B. Brosseau · K. Weedmark · B. Shutinoski | 10 | Effects of gut microbiota and metabolites on the host defense peptide expression
Y. Jin · T. Gong · X. Lu · Y. Wang · Y. Cheng |

(Continued on inside front cover)



Applied Microbiology and Biotechnology

Publishing model: **Open access**

[Explore open access funding](#) [Select institution](#)

▼ **Journal menu**

Editorial board

Editor-In-Chief

Hermann J. Heipieper PhD

Helmholtz Centre for Environmental Research, Leipzig, Germany

Managing Editor

Dorothea Kessler, Dr. rer. nat.

Independent Researcher, Heidelberg, Germany

Associate Editors

Carla C. C. R. de Carvalho PhD

Department of Bioengineering, Instituto Superior Técnico, Lisbon, Portugal;
University of Lisbon, Lisbon, Portugal

Łukasz Chrzanowski PhD

Poznań University of Technology, Poznan, Poland

Ursula Kües, Dr. rer. nat.

Büsgeninstitut, University of Göttingen, Göttingen, Germany

Bing Li PhD

Institute of Environment and Ecology, Tsinghua Shenzhen International
Graduate School, Tsinghua University, Beijing, China

Thomas Mansell PhD

Chemical & Biological Engineering, Iowa State University, Ames, United States

Zhilong Wang PhD

Shanghai Jiao Tong University, Shanghai, China

Kunihiko Watanabe PhD

Editors

Hal Alper PhD

McKetta Department of Engineering, Cockrell School of Engineering, The University of Texas at Austin, Austin, United States

George N. Bennett PhD

Department of Biochemistry & Cell Biology, Rice University, Houston, United States

Przemyslaw Bernat PhD

University of Łódź, Lodz, Poland

Uwe T. Bornscheuer PhD

Department of Biotechnology & Enzyme Catalysis, Institute of Biochemistry, Universität Greifswald, Greifswald, Germany

Axel A. Brakhage, Dr. rer. nat.

Department of Molecular and Applied Microbiology, Leibniz-Institut für
Naturstoff-Forschung und Infektionsbiologie e. V. – Hans-Knöll-Institut (HKI),
Jena, Germany

Gerhard H. Braus, Dr. sc. nat.

Department of Microbiology and Genetics, Institute of Molecular Microbiology
and Genetics, University of Göttingen, Göttingen, Germany

Guo-qiang (George) Chen PhD

Department of Biology, Tsinghua University, Beijing, China

Douglas J. Cork PhD

Department of Biological and Chemical Sciences, Illinois Institute of Technology,
Chicago, United States

Uwe Deppenmeier PhD

Institut für Mikrobiologie und Biotechnologie, University of Bonn, Bonn,
Germany

Gerrit Eggink PhD

Food and Biobased Research and AlgaePARC, Wageningen University & Research,
Wageningen, The Netherlands

Shinya Fushinobu PhD

Department of Biotechnology, The University of Tokyo, Tokyo, Japan

Navdeep Grover PhD

Heparin Applied Research Center, Rensselaer Polytechnic Institute, Troy, United States

Urszula Guzik PhD

University of Silesia in Katowice, Katowice, Poland

Bruce K. Hamilton PhD

U.S. National Science Foundation, Alexandria, United States

Yoshiteru Hashimoto MD

Graduate School of Life and Environmental Sciences, University of Tsukuba, Tsukuba, Japan

Michael Hecker PhD

Microbial Physiology and Molecular Biology, Universität Greifswald, Greifswald, Germany

Soon-Kwang Hong PhD

Division of Bioscience and Bioinformatics, Myongji University, Seoul, South Korea

Masayuki Inui PhD

Research Institute of Innovative Technology for the Earth, Kyoto, Japan

Atsuhiko Ishida PhD

Hiroshima University, Hiroshima, Japan

Nemat O. Keyhani PhD

Department of Biological Sciences, University of Illinois Urbana-Champaign, Urbana, United States

Tasuo Kurihara PhD

Kyoto University Institute for Chemical Research, Kyoto, Japan

Sang Yup Lee PhD

Department of Chemical and Biomolecular Engineering, Korea Advanced Institute of Science and Technology, Daejeon, South Korea

Xiangyang Liu PhD

Department of Biological Sciences, University of Wisconsin–Milwaukee,
Milwaukee, United States

Z. Lewis Liu PhD

National Center for Agricultural Utilization Research, Peoria, United States

Mark C.M. van Loosdrecht PhD

Department of Biotechnology, Delft University of Technology, Delft, The
Netherlands

José M. Luengo PhD

Department of Molecular Biology, Universidad de León, León, Spain

Yanping Mao PhD

Shenzhen University, Shenzhen, China

Diethard Mattanovich, Dr. rer. nat. techn.

Department of Biotechnology, BOKU University, Vienna, Austria

Ian Menz PhD

School of Life Sciences, University of Technology Sydney, Sydney, Australia

Hisaaki Mihara PhD
Ritsumeikan University, Kyoto, Japan

Jörg Nickelsen PhD
Ludwig-Maximilians-Universität München, Munich, Germany

Jens Nielsen PhD
Chalmers University of Technology, Gothenburg, Sweden

Kevin O'Connor PhD
School of Biomolecular and Biomedical Science, Science Centre, University
College Dublin, Dublin, Ireland

Masayuki Okuyama PhD
Division of Applied Bioscience, Research Faculty of Agriculture, Hokkaido
University, Sapporo, Japan

E. Terry Papoutsakis PhD
Delaware Biotechnology Institute, University of Delaware, Newark, United States

Prashant Phale PhD

Indian Institute of Technology Bombay, Mumbai, India

Bernd H.A. Rehm PhD

Centre for Cell Factories and Biopolymers, Griffith Institute for Drug Discovery,
Nathan;Queensland, Australia

Badal C. Saha PhD

Agricultural Research Service, United States Department of Agriculture,
Washington, United States

Carmen Sanchez PhD

Department of Biotechnology, Autonomous University of Tlaxcala, Tlaxcala City,
Mexico

Thomas H. Scheper, Dr. rer. nat.

Institute of Technical Chemistry, Leibniz University Hannover, Hanover,
Germany

Frank-Rainer Schmidt PhD

Prozessentwicklung Chemie, Industriepark Höchst, Sanofi-Aventis Deutschland,
Frankfurt, Germany

Helmut Schwab PhD

Department of Molecular Biotechnology, Graz University of Technology, Graz,
Austria

Yasushi Shigeri PhD

Department of Chemistry, Wakayama Medical University, Wakayama, Japan

Daniel K.Y. Solaiman PhD

Eastern Regional Research Center, Agricultural Research Service, United States
Department of Agriculture, Washington, United States

A.J.J. Straathof PhD

Department of Biotechnology, Delft University of Technology, Delft, The
Netherlands

Andras Tancsics PhD

Hungarian University of Agriculture and Life Sciences, Godollo, Hungary

Dirk Tischler PhD

Ruhr University Bochum, Bochum, Germany

Kenji Ueda PhD

Life Science Research Center, College of Bioresource Sciences, Nihon University,
Tokyo, Japan

Roland Ulber, Dr. rer. nat.

Department of Mechanical and Process Engineering, Institute of Bioprocess
Engineering, Rheinland–Pfälzische Technische Universität Kaiserslautern–
Landau, Landau, Germany

Paul J. Weimer PhD

Department of Bacteriology, University of Wisconsin–Madison, Madison, United
States

Volker F. Wendisch PhD

Genetics of Prokaryotes, Faculty of Biology & CeBiTec, Bielefeld University,
Bielefeld, Germany

Josef Winter, Dr. rer. nat.

Institute of Bioengineering and Biotechnology of Wastewater, Karlsruhe
Institute of Technology, Karlsruhe, Germany

Marta Wozniak–Karczewska PhD

Poznań University of Technology, Poznan, Poland

Yu Xia PhD

Southern University of Science and Technology, Shenzhen, China

Shotaro Yamaguchi PhD

Medical Enzymes Division, Amano Enzyme (Japan), Nagoya, Japan

Yeo Joon Yoon PhD

College of Pharmacy, Seoul National University, Seoul, South Korea

Takeshi Zendo PhD

Department of Bioscience and Biotechnology, Faculty of Agriculture, Kyushu University, Fukuoka, Japan

Tong Zhang PhD

Environmental Microbiome Engineering and Biotechnology Lab, Department of Civil Engineering, University of Hong Kong, Pok Fu Lam, Hong Kong

Past Editor-In-Chief

Alexander Steinbüchel

University of Münster, Münster, Germany

Editorial Board Members

Tanja Narancic PhD

School of Biomolecular and Biomedical Science, University College Dublin,
Dublin, Ireland



Applied Microbiology and Biotechnology

Publishing model: Open access

[Explore open access funding](#) [Select institution](#)

▼ Journal menu

[Search all Applied Microbiology and Biotechnology articles](#) →




Volume 110, Issue 1

December 2026


261 articles in this issue

Heterologous production of a plant biostimulant in *Streptomyces albidoflavus*




Research | Open access | 20 June 2026 | Article: 261

Construction of a novel reporter strain indicating inhibition of cell wall biosynthesis




Research | Open access | 28 August 2026 | Article: 260

Biocontrol of mushroom crop mycoparasites by novel *Bacillus velezensis* strains




Efficient biotechnological production of *N*-(Hydroxycinnamoyl)tyramines by utilizing plant-derived enzymes in a bench-scale bioreactor with in situ product precipitation



Research | Open access | 26 August 2026 | Article: 258

Oxygen-based soft-sensing of glucose demand in Chinese hamster ovary fed-batch cultures across scales




Research | Open access | 17 June 2026 | Article: 257



Part of 1 collection:


Single Use Technologies in Bioprocessing

Fusion with immunoglobulin Fc fragment enhances the immunogenicity of the African swine fever virus p30/p54 fusion protein




Research | Open access | 15 June 2026 | Article: 256

Activation of a gene cluster in *Streptomyces lavendulae* subsp. *lavendulae* CCM 3239 induced strevertene production




Research | Open access | 15 June 2026 | Article: 255

Gut microbiome composition and cellulolytic bacteria associated with the carpenter bee *Xylocopa frontalis*




Research | Open access | 12 June 2026 | Article: 254

Streamlined cell-free protein synthesis using wheat germ extract




Research | Open access | 11 June 2026 | Article: 253

A constitutive promoter toolbox for tunable gene expression in *Acinetobacter baumannii*



Research | Open access | 11 June 2026 | Article: 252

Biomass and coccolith productions in *Chrysotila* sp. in high density cultivation system




Research | Open access | 10 June 2026 | Article: 251



Part of 1 collection:

Cultivation of Microorganisms: New Media and Techniques

Biogenic synthesis and characterization of selenium nanoparticles by halotolerant *Bacillus sonorensis* 2MNHR



Research | Open access | 15 August 2026 | Article: 250



Part of 1 collection:


Emerging Applications of Nanoparticles in Biotechnology and Medicine

Biotransformation-guided production of quinalizarin as an antibacterial agent against *Acinetobacter baumannii*




Research | Open access | 14 August 2026 | Article: 249

Correction: Bacterial communities in domestic washing machines associated with user-perceived odours



Correction | Open access | 14 August 2026 | Article: 248

Innovative epitope-based ELISA assay for detecting Group B *Streptococcus* carriage in pregnant women




Research | Open access | 09 June 2026 | Article: 247

Retraction Note: Biotechnology for propagation and secondary metabolite production in *Bacopa monnieri*


Retraction Note | Open access | 11 August 2026 | Article: 246

Comparative genomics reveals immunomodulatory and anti-obesity traits of *Lactococcus lactis* subsp. *lactis*




Research | Open access | 06 June 2026 | Article: 245

Evaluating yeasts as biocontrol agents against pathogenic and spoilage microorganisms in agri-food: action of volatile organic compounds (VOCs)




Research | Open access | 06 June 2026 | Article: 244

First experimentally characterised *Proteus* phage endolysins have distinct metal cofactor dependencies



Research | Open access | 06 June 2026 | Article: 243

Chitosan nanoparticles as adjuvants to enhance the biodegradation of PET by *Ideonella sakaiensis*



Research | Open access | 06 June 2026 | Article: 242




Part of 1 collection:

Emerging Applications of Nanoparticles in Biotechnology and Medicine

Retraction Note: Successive use of microorganisms to remove chromium from wastewater

Retraction Note | Open access | 10 August 2026 | Article: 241

Perfusion development and its potential for cell therapy manufacturing with adherent cells




Mini Review | Open access | 03 July 2026 | Article: 240



Part of 1 collection:

Single Use Technologies in Bioprocessing

Rational development of an mRNA antibody cocktail with extended therapeutic window against ricin intoxication




Research | Open access | 04 June 2026 | Article: 239



Part of 1 collection:


Emerging Applications of Nanoparticles in Biotechnology and Medicine

Molecular mechanism underlying biofloc formation of *Acidovorax soli* Q11 and its application potential




Research | Open access | 04 June 2026 | Article: 238

Fungal signatures in a cadaver and a separated upper limb: insights from a forensic case study




Research | Open access | 04 June 2026 | Article: 237

A cumate-inducible promoter system for tuned gene expression and magnetosome overproduction in magnetotactic bacteria



Research | Open access | 08 August 2026 | Article: 236

***Escherichia coli* DH5 α -loaded fenugreek extract: an effective anticancer combination against human breast cancer cell lines**




Research | Open access | 08 August 2026 | Article: 235

Extrachromosomally co-encoded toxin-immunity pairs in yeast and their application potential




Mini Review | Open access | 26 May 2026 | Article: 234

SphB1 export system for antigen presentation to the respiratory tract by a live pertussis vaccine




Research | Open access | 29 May 2026 | Article: 233

Inhalation of cadmium oxide nanoparticles alters intestinal and pulmonary microbiomes in mice



Research | Open access | 28 July 2026 | Article: 232

Bacterial communities in domestic washing machines associated with user-perceived odours




Research | Open access | 28 July 2026 | Article: 231

Retraction Note: Biotechnological interventions and genetic diversity assessment in *Swertia* sp.: a myriad source of valuable secondary metabolites


Retraction Note | Open access | 28 July 2026 | Article: 230

Cultivar and tree age jointly shape rhizosphere bacterial diversity and community assembly in *Macadamia integrifolia*




Research | Open access | 26 June 2026 | Article: 229

Evaluation of biobased carriers derived from agri-food waste for biostimulants delivery in horticulture



Research | Open access | 28 May 2026 | Article: 228


High-yield production of α -glucosylglycerol using *Xanthomonas campestris* WU-9701 derived XgtA



Research | Open access | 28 May 2026 | Article: 227


Restoration of urban agriculture soil by autochthonous fungal biodiversity

Research | Open access | 28 May 2026 | Article: 226




Effect of enzymatic saccharification on edible seaweed (*Ulva* sp.) fermentation

Research | Open access | 28 May 2026 | Article: 225




Identification of targets for high penicillin G production in an industrial strain of *Penicillium rubens*

Research | Open access | 28 May 2026 | Article: 224




FliW regulates biofilm formation in *Geobacter sulfurreducens* through interaction with CsrA

Research | Open access | 28 May 2026 | Article: 223




Secretion systems shaping host–pathogen interactions in *Burkholderia* species

Mini Review | Open access | 22 May 2026 | Article: 222




Calcium enhances hydrolytic and transglucosylation activities in α -L-fucosidase from *Thermotoga maritima* through catalytic loop stabilization: an MD simulation study

Research | Open access | 23 May 2026 | Article: 221



A comparative analysis of monascins and ankaflavins biosynthesis in *Monascus purpureus* and its albino variant

Research | Open access | 23 May 2026 | Article: 220




Neisseria subflava Type 6 Secretion System competition with bacterial and fungal species




Research | Open access | 23 May 2026 | Article: 219

Impact of direct cell–cell contact in a three–species wine yeast consortium




Research | Open access | 23 May 2026 | Article: 218

Analytic attributes of the 16S rRNA gene sequencing methodology for human gut microbiota characterization




Research | Open access | 22 May 2026 | Article: 217

Neuroprotective effects of cell-free supernatant from *Pediococcus pentosaceus* TAP041 against glycation- and inflammation-associated stress responses




Research | Open access | 22 May 2026 | Article: 216

Improving autotrophic D–lactate production by heterologous expression of a pyruvate formate lyase in an LDHD–expressing *Acetobacterium woodii* strain




Research | Open access | 10 July 2026 | Article: 215

Quantitative proteomics identifies GntR as a novel potential DIVA antigen for bovine brucellosis




Research | Open access | 21 May 2026 | Article: 214

Divergent metallothionein strategies underlie copper tolerance in *Saccharomyces* species relevant to winemaking



Research | Open access | 21 May 2026 | Article: 213

Adapting prime editing with split prime editors in *Escherichia coli* and its application to *Staphylococcus aureus* genome editing



Research | Open access | 04 June 2026 | Article: 212



Applied Microbiology and Biotechnology

Publishing model: Open access

[Explore open access funding](#) [Select institution](#)

▼ Journal menu

[Search all Applied Microbiology and Biotechnology articles](#) →



Volume 110, Issue 1

December 2026


261 articles in this issue

LysR-type transcriptional regulator TtdR regulates both ethylene glycol and polyhydroxyalkanoate (PHA) metabolism in *Pseudomonas umsongensis* GO16




Research | Open access | 19 May 2026 | Article: 211

Engineering a robust cell-surface display platform in the multi-stress-tolerant yeast *Issatchenkia orientalis*




Research | Open access | 19 May 2026 | Article: 210

Endophytic *Aspergillus glaucus* as a novel epothilone B producer: Antiproliferative activity and in silico analysis



Evaluation of vector system in *Saccharopolyspora erythraea* and construction of new replicative vector




Research | Open access | 14 May 2026 | Article: 208

Relief of bHD1 autoinhibition and modular coupling support *MAT-A* pathway activation in *Flammulina filiformis*




Research | Open access | 15 May 2026 | Article: 207

Synergistic attenuation of *Aeromonas dhakensis* virulence by dual deletion of PhoBR and KdpE two-component systems and its protective immunity in zebrafish




Research | Open access | 15 May 2026 | Article: 206

Computationally guided enzyme engineering for regioselective synthesis of fucosylated human milk oligosaccharides




Research | Open access | 14 May 2026 | Article: 205

Human anti- α -hemolysin scFvs mitigate toxin-mediated damage in a mouse model of MRSA pneumonia



Research | Open access | 14 May 2026 | Article: 204

Removal and analysis of foodborne-associated pathogens via peptide functionalized nanoparticles




Research | Open access | 06 July 2026 | Article: 203



Part of 1 collection:


Emerging Applications of Nanoparticles in Biotechnology and Medicine

Expression and identification of a novel high-activity recombinant humanized type I collagen SynthCol1 in *Pichia pastoris* GS115




Research | Open access | 11 May 2026 | Article: 202

H₂-dependent reduction of rubredoxin by FrhAGB hydrogenase in *Thermococcus onnurineus* NA1




Research | Open access | 11 May 2026 | Article: 201

Immobilization of TtAlaDH on magnetic nanoparticles: effects on stability and reuse




Research | Open access | 09 May 2026 | Article: 200

Characterization of three glycerol transporters with broad sugar specificities from *Aspergillus niger*




Research | Open access | 08 May 2026 | Article: 199

Engineering natural *Saccharomyces cerevisiae* isolates for enhanced one-step cellulosic ethanol production




Research | Open access | 08 May 2026 | Article: 198

Bacterial lipoxygenases for fatty acid oxyfunctionalization: advances and future directions




Mini Review | Open access | 07 May 2026 | Article: 197

Real-time NMR monitoring of HMF biotransformation by *Ectopseudomonas oleovorans* CECT 5344 R1D




Research | Open access | 07 May 2026 | Article: 196

Characterization of a novel bifunctional endo- β -1,4-glucanase/mannanase from *Aspergillus oryzae*



Research | Open access | 07 May 2026 | Article: 195

Enhancing reproducibility in mixing time determination of stirred tank reactors via automated analysis and standardized inter-laboratory trials




Research | Open access | 30 June 2026 | Article: 194



Part of 1 collection:


Single Use Technologies in Bioprocessing

Molecular engineering of *Komagataella phaffii* for venom toxin production



Mini Review | Open access | 07 May 2026 | Article: 193

Improving the reproducibility of mixing-time experiments in stirred single-use bioreactors




Research | Open access | 29 June 2026 | Article: 192



Part of 1 collection:

Single Use Technologies in Bioprocessing

Comparative production of recombinant capripoxvirus in Vero cells using microcarrier- and macrocarrier-based dynamic cell culture systems under serum-free conditions




Research | Open access | 29 June 2026 | Article: 191




Part of 1 collection:

Single Use Technologies in Bioprocessing

A new Kayvirus vB_SauM-MUHD-1 combats Methicillin-resistant *Staphylococcus aureus* wound infections



Comparative genomic analysis of RNA-binding proteins across some drug resistant and sensitive *Staphylococcus aureus*




Research | Open access | 24 June 2026 | Article: 189

Development of a *Ganoderma lucidum* bioreactor for dichlorodiphenyltrichloroethane contaminated soil treatment




Research | Open access | 15 June 2026 | Article: 188

Intracellular β -glucosidase regulates cellulase expression and development in *Aspergillus nidulans*




Research | Open access | 04 May 2026 | Article: 187

Fecal microbiota transplantation from different pig breeds alters fat deposition and gut microbiota in mice



Research | Open access | 04 May 2026 | Article: 186

B-vitamin supplementation unlocks DHA production of *Aurantiochytrium limacinum* on molasses-based fermentation residue




Research | Open access | 01 May 2026 | Article: 185



Part of 1 collection:


Cultivation of Microorganisms: New Media and Techniques

2-Keto-3-deoxy-D-xylonate and 2-oxo-4-hydroxybutyrate as natural and artificial effectors of transcription factors regulating D-xylonate operons in *E. coli*




Research | Open access | 01 May 2026 | Article: 184

Correction: The PurR family transcriptional regulator promotes butenyl-spinosyn production in *Saccharopolyspora pogona*



Correction | Open access | 18 June 2026 | Article: 183

Isolation and genomic characterization of a lytic bacteriophage against multidrug-resistant *Proteus mirabilis*



Research | Open access | 29 April 2026 | Article: 182

Evaluation of sterilizable 3D-printed cell culture platforms using multiple FDM materials and their effects on cell behavior




Research | Open access | 29 April 2026 | Article: 181



Part of 1 collection:


Single Use Technologies in Bioprocessing

Optimization of biological sulfuric acid generation by *Acidithiobacillus thiooxidans* in a stirred-tank reactor system




Research | Open access | 14 May 2026 | Article: 180

Mitigating protocatechuic acid toxicity in *Vibrio natriegens* enables poly-3-hydroxybutyrate production from an aromatic carbon source




Research | Open access | 27 April 2026 | Article: 179

Bacteriophage survival and ARG dissemination from wastewater treatment plants to the environment



Mini Review | Open access | 27 April 2026 | Article: 178

Understanding time-growth rate relationship for slow-growing microbial cells in retentostat culturing systems




Research | Open access | 25 April 2026 | Article: 177



Part of 1 collection:

Cultivation of Microorganisms: New Media and Techniques

Piezoelectric cold atmospheric plasma inactivates dairy cattle bacteria and potentiates antibiotics in vitro



Research | Open access | 24 April 2026 | Article: 176

Leaf mold extract agar facilitates recovery of soil bacterial diversity beyond *Bacillus* dominance




Research | Open access | 22 April 2026 | Article: 175



Part of 1 collection:


Cultivation of Microorganisms: New Media and Techniques

Vibrio proteolyticus DCF12.2 postbiotic modulates intestinal metabolic and immune pathways in zebrafish




Research | Open access | 22 April 2026 | Article: 174

Developing an oncolytic Newcastle disease virus production process with EB66 cells



Research | Open access | 04 June 2026 | Article: 173

Engineering of ω -transaminases for advanced applications




Mini Review | Open access | 04 June 2026 | Article: 172

Metagenomic insights into urolithin formation from rambutan rind extract by rat faecal-derived microbiome



Research | Open access | 27 April 2026 | Article: 171

Polycationic nanopeptide-fused endolysins for the control of *Mannheimia haemolytica*




Research | Open access | 08 May 2026 | Article: 170

Correction: The combination of mupirocin and *Kayvirus* broadens the decolonization effect against *Staphylococcus aureus*


Correction | Open access | 04 June 2026 | Article: 169

Bacillus velezensis EU07 suppresses *Fusarium graminearum* via transcriptomic reprogramming




Research | Open access | 08 May 2026 | Article: 168

A hydrolase from *Serratia liquefaciens* IMD717 with esterase, dehalogenase and *N*-deformylase activities




Research | Open access | 20 April 2026 | Article: 167

Characterization of a membrane-bound PQQ-dependent D-lactate dehydrogenase in *Gluconobacter japonicus*




Research | Open access | 20 April 2026 | Article: 166

Effects of fermented mulberry leaves on rumen fermentation and microbiota in rumen fluid of heat-stressed Hu sheep



Research | Open access | 20 April 2026 | Article: 165

Glycerol-mediated improvement of heterologous aurachin D production in *E. coli*




Research | Open access | 30 May 2026 | Article: 164



Part of 1 collection:


Cultivation of Microorganisms: New Media and Techniques

Development of a gut microbiota model for the analysis of bacterial modifications of xenobiotics



Research | Open access | 30 May 2026 | Article: 163

The p40 fusion domain as a scaffold for producing functional inclusion bodies



Research | Open access | 18 April 2026 | Article: 162



Part of 1 collection:

Emerging Applications of Nanoparticles in Biotechnology and Medicine

< [Previous](#) [1](#) 2 [3](#) ... [6](#) [Next](#) >



Applied Microbiology and Biotechnology

Publishing model: Open access

[Explore open access funding](#) [Select institution](#)

▼ Journal menu

[Search all Applied Microbiology and Biotechnology articles](#) →




Volume 110, Issue 1

December 2026


261 articles in this issue

Beyond multilocus barcoding: harnessing phylogenomics and genome mining for next-generation *Trichoderma*-based bioproducts




Mini Review | [Open access](#) | 28 May 2026 | Article: 161

Ribosomal protein RPSA is a promising target for the prevention of *Streptococcus suis* serotype 2



Research | [Open access](#) | 17 April 2026 | Article: 160

Redox and physiological responses of wheat to wheat streak mosaic virus under liposomal bionanocomposite seed treatment




Research | Open access | 17 April 2026 | Article: 159



Part of 1 collection:


Emerging Applications of Nanoparticles in Biotechnology and Medicine

An engineered *Pseudomonas stutzeri* strain overexpressing the sigma factor AlgU promotes maize growth under salt stress




Research | Open access | 16 April 2026 | Article: 158

The combination of mupirocin and *Kayvirus* broadens the decolonization effect against *Staphylococcus aureus*




Research | Open access | 16 April 2026 | Article: 157

Development of a double-antibody sandwich ELISA for detecting Peste des petits ruminants virus infection



Research | Open access | 15 April 2026 | Article: 156

Characterization and evaluation of a novel polylactic acid-made bioreactor for large-scale adherent cell expansion




Research | Open access | 22 May 2026 | Article: 155



Part of 1 collection:


Single Use Technologies in Bioprocessing

Engineering *Penicillium expansum* antifungal proteins unveils new clues about their mode of action




Research | Open access | 20 April 2026 | Article: 154

Motility and growth *Burkholderia* two-component: a new regulatory system of *Burkholderia seminalis*




Research | Open access | 07 May 2026 | Article: 153

Isolation, characterization and biocontrol of a novel bacteriophage against *Ralstonia pseudosolanacearum*




Research | Open access | 15 April 2026 | Article: 152

Structural and mechanistic insights into the divergence of pterin deaminase and sepiapterin deaminase




Mini Review | Open access | 16 April 2026 | Article: 151

Development of a double-monoclonal antibody sandwich ELISA for the detection of Citrus tristeza virus




Research | Open access | 23 March 2026 | Article: 150

Optimized expression and antimicrobial activity of rainbow trout defensin 3 in *Komagataella phaffii*




Research | Open access | 16 April 2026 | Article: 149

One-step in vivo assembly of a 63 kb-long biosynthetic gene cluster via multiple recombination in *Aspergillus oryzae*



Research | Open access | 07 May 2026 | Article: 148

Biocalorimetry for the biotechnological use of natural and synthetic macromolecules




Mini Review | Open access | 21 May 2026 | Article: 147

Application of clearance strategies for extractables and leachables and advances in process modeling




Mini Review | Open access | 20 May 2026 | Article: 146

A novel *Azospirillum vistecanum* strain from methane digestate promotes plant growth via indole-3-acetic acid biosynthesis




Research | Open access | 11 April 2026 | Article: 145

A cold-adapted and detergent-stable α -amylase from marine bacterium *Photobacterium gaetbulicola* Gung47^T




Research | Open access | 11 April 2026 | Article: 144

Carbohydrate concentration and type drive product selectivity to a mixture of volatile fatty acids or lactic acid in thermophilic mixed-culture fermentation




Research | Open access | 11 April 2026 | Article: 143

Recent advances in pathway engineering for the biosynthesis of 1,4-butanediol




Mini Review | Open access | 10 April 2026 | Article: 142

Exploiting the ability of *Bacillus subtilis* to synthesize surfactin under oxygen limitation




Research | Open access | 14 May 2026 | Article: 141

Understanding pellet population heterogeneity of *Aspergillus niger* in stirred tank and rocking motion bioreactors



Research | Open access | 08 May 2026 | Article: 140

Evolutionary engineering of *Geobacillus thermoleovorans* for growth on adipic acid and 1,4-butanediol




Single-use solutions for PUPSIT: requirements, challenges, and solutions

Mini Review | Open access | 01 May 2026 | Article: 138




Part of 1 collection:

Single Use Technologies in Bioprocessing




Correction: Ornithine lipids and other acyloxyacyl amino lipids: the coming-of-age story of a group of non-canonical membrane lipids

Correction | Open access | 01 May 2026 | Article: 137




A *katE katG* double-knockout *E. coli* strain eliminates the risk of catalase contamination in recombinant proteins

Research | Open access | 22 April 2026 | Article: 136




Characterization of antibodies against the replication protein (Rep) encoded by bovine meat and milk factors (BMMFs)

Research | Open access | 16 April 2026 | Article: 135



Investigation of regioselectivity and thermostability of free and immobilized *Pleurotus citrinopileatus* lipase

Research | Open access | 07 April 2026 | Article: 134




Foot-and-mouth disease virus-like particle mRNA vaccine induces strong humoral and cellular immunity

Research | Open access | 06 April 2026 | Article: 133




Part of 1 collection:

Emerging Applications of Nanoparticles in Biotechnology and Medicine




Enhanced biomass and desulfurization by recombinant *Rhodococcus qingshengii* IGTS8 in bioreactor cultures



Research | Open access | 06 April 2026 | Article: 132

Identification and characterization of prostaglandin F_{2α} 9-dehydrogenase from *Rhodotorula kratochvilovae*




Research | Open access | 06 April 2026 | Article: 131

Research progress on nucleic acid amplification–based detection technologies for phytopathogenic fungi




Mini Review | Open access | 02 April 2026 | Article: 130

Physiological characterization of anaerobic cometabolic transformation of sulfamethoxazole by *Nitratidesulfovibrio vulgaris* Hildenborough



Research | Open access | 31 March 2026 | Article: 129

Co-expression, purification, and characterization of an acidophilic and n-hexane-tolerant lipase with its foldase from *Burkholderia gladioli* Bsp-1



Research | Open access | 28 March 2026 | Article: 128

Lab Scale Closed-Loop Recycling of Polycarbonate Bioreactors for Sustainable Process Development




Research | Open access | 27 March 2026 | Article: 127




Part of 1 collection:

Single Use Technologies in Bioprocessing

Reusability of immobilised lipase in the production of omega-3 oils from squid viscera



A fast workflow to explore active enzymes from environmental samples through functional metagenomics




Research | Open access | 25 March 2026 | Article: 125

Retraction Note: Expression of liver-targeting peptide modified recombinant human endostatin and preliminary study of its biological activities


Retraction Note | Open access | 25 March 2026 | Article: 124

Enzyme-constrained genome-scale model of *Yarrowia lipolytica* predicts growth-phase specific metabolic engineering targets




Research | Open access | 24 March 2026 | Article: 123

Testing the biodegradability of difficult compounds: a future challenge for the OECD/ISO standardization




Mini Review | Open access | 24 March 2026 | Article: 122

Environmental and chemical modulation of *Staphylococcus aureus* Newman biofilm formation




Research | Open access | 24 March 2026 | Article: 121

Development of canine parvovirus-neutralizing monoclonal antibodies from natural host and their germline gene usage



Research | Open access | 24 March 2026 | Article: 120

Optimized green zinc oxide/chitosan/amoxicillin nanocomposite against *Pseudomonas aeruginosa*




Research | Open access | 24 March 2026 | Article: 119



Part of 1 collection:


Emerging Applications of Nanoparticles in Biotechnology and Medicine

Revolutionizing sweetness: the multifaceted health benefits of fermented stevia




Mini Review | Open access | 24 March 2026 | Article: 118

Traditional plant urologicals curb early uropathogenic *E. coli* infection and strengthen host innate defense



Research | Open access | 23 March 2026 | Article: 117

Hanseniaspora uvarum and *Hanseniaspora opuntiae* differ in their growth and in the production of volatile organic compounds in wine




Research | Open access | 23 March 2026 | Article: 116

Evaluation of the biological activities of *Bacillus cereus*-synthesized titanium dioxide nanoparticles



Insights into the antimicrobial and antibiofilm activities of ionic liquids and their mechanisms of action




Mini Review | Open access | 20 March 2026 | Article: 112



▼ Journal menu


December 2026

Transcriptomic insights into fruiting body malformations in *Lentinula edodes*




Research | Open access | 17 March 2026 | Article: 108

The potential of plant-derived triterpenoids as biological nitrification inhibitors




Research | Open access | 17 March 2026 | Article: 107

Eucalyptus lignin production coupled with Cellic® CTec3 HS enzymatic hydrolysis to monosaccharides



Research | Open access | 16 March 2026 | Article: 106

Development of a high-throughput label-free method for enrichment of antimetabolite-resistant yeast strains using pico-litre agarose beads




Research | Open access | 16 March 2026 | Article: 105

Effects of sialylated oligosaccharides on the N- and O-glycosylation of etanercept in recombinant CHO cells




Research | Open access | 13 March 2026 | Article: 104

Optimizing cryopreservation protocols of *Saccharomyces eubayanus* using heat transfer modeling



Research | Open access | 13 March 2026 | Article: 103

Cellular responses to prolonged non-thermal plasma exposure in *Schizosaccharomyces pombe*




Natural food-derived antimicrobials against beer spoilage microorganisms




Research | Open access | 11 March 2026 | Article: 101

Plant and soil transcriptomics reveal the consequences of bioaugmentation of co-contaminated soil with *Pseudomonas qingdaonensis* ZCR6 during bacteria-assisted phytoremediation




Research | Open access | 11 March 2026 | Article: 100

Biocontrol potential of *Aspergillus* section *Nigri* strains against *Aspergillus carbonarius*



Research | Open access | 09 March 2026 | Article: 99

Growth and protein content of *Cupriavidus necator* on organic acids derived from fermented grey starch




Research | Open access | 09 March 2026 | Article: 98



Part of 1 collection:


Cultivation of Microorganisms: New Media and Techniques

Ornithine lipids and other acyloxyacyl amino lipids: the coming-of-age story of a group of non-canonical membrane lipids




Mini Review | Open access | 08 March 2026 | Article: 97

Enhanced fructooligosaccharides synthesis by engineered *Trichoderma atroviride* β -fructofuranosidase




Research | Open access | 08 March 2026 | Article: 96

Probiotic characterization and spore production optimization of *Lysinibacillus* sp. MK212927




Research | Open access | 08 March 2026 | Article: 95

Abiotic stress effects on polyunsaturated fatty acid biosynthesis in *Chlorococcum oleofaciens* and *Leptolyngbya* sp.




Research | Open access | 08 March 2026 | Article: 94

Microbial-induced calcite precipitation by indigenous alkaliphilic bacteria: a dual-enzyme strategy for crack-healing in cementitious materials




Research | Open access | 06 March 2026 | Article: 93

Rerouting reductant flux via protein tethering enhances biohydrogen production in *Thermococcus kodakarensis*




Research | Open access | 06 March 2026 | Article: 92

Unveiling a catalytically promiscuous feruloyl esterase from *Clostridium acetobutylicum*




Research | Open access | 04 March 2026 | Article: 91

Biofilm formation directly correlates with cell viability in *Candida tropicalis* on polypropylene



Research | Open access | 04 March 2026 | Article: 90

Unique gut microbiota and metabolomic profiling as biomarker of post-transplant recovery in acute-on-chronic liver failure after liver transplantation




Research | Open access | 03 March 2026 | Article: 89

Editorial expression of concern: Role and fate of SP100 protein in response to Rep-dependent nonviral integration system


Editorial Expression of Concern | Open access | 04 March 2026 | Article: 88

Genome mining and screening for plastic-degrading potential in marine bacteria




Research | Open access | 03 March 2026 | Article: 87

S-equal producing bacteria: isolation and identification from Albino Wistar rat gut microbiota



Research | Open access | 02 March 2026 | Article: 86

Development of novel microbial synthetic consortia for the production of antimicrobial fermentates containing caproate




Research | Open access | 27 February 2026 | Article: 85

Fusarium enrichment alters the soil microbial community and network structure of Brassica crops



Research | Open access | 27 February 2026 | Article: 84

Nanoparticle-based approaches for bacterial detection and therapy




Mini Review | Open access | 26 February 2026 | Article: 83



Part of 1 collection:


Emerging Applications of Nanoparticles in Biotechnology and Medicine

Characterization, genomic analysis, and decontamination application of a novel *Vibrio* phage BUCT787 on salmon fillets



Research | Open access | 26 February 2026 | Article: 82

Evaluation of the delivery of an anti-*Listeria* endolysin via CRISPR-Cas9 engineered probiotic *Saccharomyces boulardii*




Research | Open access | 26 February 2026 | Article: 81

Correction: Virus-like particle (VLP)-based indirect ELISA (iELISA) for the detection of beak and feather disease virus (BFDV) antibodies


Correction | Open access | 26 February 2026 | Article: 80

CO₂-driven biosurfactant synthesis by bacteria within CCUS




Mini Review | Open access | 25 February 2026 | Article: 79

The visualizable and marker-free gene editing platform mediated by CRISPR/Cas9 in *Coleophoma empetri*



Research | Open access | 25 February 2026 | Article: 78

Robust *Pichia kudriavzevii* for open production of methylotrophic protein




Research | Open access | 24 February 2026 | Article: 77

Multistep metabolic engineering of *Pichia pastoris* for biosynthesis of N-acetylneuraminic acid



Research | Open access | 24 February 2026 | Article: 76

Blue LED-assisted bioreactor cultivation enhances carotenoid accumulation in *Rhodotorula mucilaginosa* LB-01




Research | Open access | 24 February 2026 | Article: 75

Correction: Valorisation of spent mushroom substrate by secondary microbial fermentation

Correction | Open access | 23 February 2026 | Article: 74

A new-engineered integrative tool to target the terminal compartment of the *Streptomyces* chromosome



Research | Open access | 21 February 2026 | Article: 73

Clove oil and selenium nanoparticles as eco-friendly antifungal agents for photographic preservation



Research | Open access | 21 February 2026 | Article: 72



Part of 1 collection:


Emerging Applications of Nanoparticles in Biotechnology and Medicine

Aloe emodin disrupts *Candida albicans* mitochondrial iron homeostasis against its hyphal development and oral candidiasis




Research | Open access | 20 February 2026 | Article: 71

Green synthesis of silver nanoparticles by isolated *Bacillus velezensis* BS1 and their applications



Research | Open access | 19 February 2026 | Article: 70

A monoclonal antibody-based electrochemical immunosensor for porcine epidemic diarrhea virus



Research | Open access | 19 February 2026 | Article: 69




Part of 1 collection:

Emerging Applications of Nanoparticles in Biotechnology and Medicine

Nootkatone antifungal activity and calcium signaling-mediated resistance in *Aspergillus fumigatus*

Research | Open access | 18 February 2026 | Article: 68




Establishment and validation of a rat sepsis model aligned with sepsis-3 via *Staphylococcus aureus* tail vein infection



Research | Open access | 18 February 2026 | Article: 67

Acetoclastic versus hydrogenotrophic methanogenesis: defining how pH and alkalinity shape acetate metabolism in a haloalkaliphilic methanogenic community for biomethane production




Research | Open access | 16 February 2026 | Article: 66

Identification of fecal microbiome signatures associated with longevity through 16S rRNA sequencing in different age groups in China




Research | Open access | 16 February 2026 | Article: 65

Molecular breeding alters mycelium material properties in the white-rot fungus *Pleurotus ostreatus*




Research | Open access | 13 February 2026 | Article: 64

Elucidating carbohydrate preference and engineering glucose transport in *Caldimonas thermodepolymerans* for enhanced polyhydroxyalkanoate production



Research | Open access | 13 February 2026 | Article: 63

Rapid and visual dual-color lateral flow immunoassay for african swine fever virus antibody detection



Research | Open access | 11 February 2026 | Article: 62



Applied Microbiology and Biotechnology

Publishing model: Open access

[Explore open access funding](#) [Select institution](#)

▼ Journal menu

[Search all Applied Microbiology and Biotechnology articles](#) →




Volume 110, Issue 1

December 2026


261 articles in this issue

One-pot recombinase polymerase amplification and CRISPR/Cas12a assay for *Cryptococcus neoformans*




Research | Open access | 09 February 2026 | Article: 61

Antimicrobial effects of combined piezoelectric cold plasma, organic acids, and nanoemulsions against *Salmonella* Typhimurium and *Listeria monocytogenes* on pork




Research | Open access | 09 February 2026 | Article: 60

Cold-adapted carboxylesterases from Alcanivoracaceae active with a wide range of synthetic polyesters




Multimodal molecular mechanisms of octanoic acid (OA) against recurrent mastitis causing pathogens



Research | Open access | 05 February 2026 | Article: 58

Nutrient content of grass influences microbial communities in the phyllosphere more than climate change



Research | Open access | 04 February 2026 | Article: 57

Making GFP count: a validated framework for absolute protein quantification in precision fermentation




Research | Open access | 04 February 2026 | Article: 56

Correction: A novel cholesterol-producing *Pichia pastoris* strain is an ideal host for functional expression of human Na,K-ATPase $\alpha 3\beta 1$ isoform


Correction | Open access | 04 February 2026 | Article: 55

A multiplex assay based on capillary electrophoresis to detect *Mycobacterium tuberculosis* complex: development and clinical validation



Research | Open access | 02 February 2026 | Article: 54

Post-defecation exposure alters gut microbiota of forest musk deer with implications for conservation metagenomics



Research | Open access | 02 February 2026 | Article: 53

Arsenic efflux mechanisms in ectomycorrhizal mushrooms *Hebeloma bulbiferum* and *Hebeloma sinapizans*




Retraction Note: High efficient expression of a functional humanized single-chain variable fragment (scFv) antibody against CD22 in *Pichia pastoris*

Retraction Note | Open access | 31 January 2026 | Article: 51


A genetically encoded L-rhamnose biosensor for monitoring marine polysaccharide depolymerization

Research | Open access | 29 January 2026 | Article: 50




Bipolar, not tetrapolar: mating system determination in *Inonotus hispidus* through genomic and phenotypic analysis

Research | Open access | 29 January 2026 | Article: 49




Highly enantioselective reduction of benzophenones by engineered *Geotrichum candidum* alcohol dehydrogenase

Research | Open access | 29 January 2026 | Article: 48




Enhanced bacterial cellulose production by *Komagataeibacter xylinus* using agro-derived flour nitrogen sources

Research | Open access | 29 January 2026 | Article: 47




An integrated cell and medium engineering approach for production of a nanobody fusion in *Saccharomyces cerevisiae*


Research | Open access | 29 January 2026 | Article: 46



Heterologous expression and functional characterization of novel microcin B17 congeners from environmental *Pseudomonas* isolates




Elucidating the roles of essential genes in autotrophic metabolism and cell morphology of *Clostridium ljungdahlii* by CRISPRi




Research | Open access | 27 January 2026 | Article: 44

Response and degradation of indole-3-acetic acid by the plant growth-promoting rhizobacteria *Variovorax boronicumulans*



Research | Open access | 27 January 2026 | Article: 43

An improved CRISPR-Cas9 protein-based method for knocking out insect Sf9 cell genes




Research | Open access | 26 January 2026 | Article: 42

Antibiofilm and antimetabolic effects of disinfectants on *Pseudomonas aeruginosa* strains isolated from cosmetic manufacturing environments




Research | Open access | 26 January 2026 | Article: 41

Photoprotective and cryoprotective effect of the carotenoid bacterioruberin on peripheral blood mononuclear cells




Research | Open access | 27 January 2026 | Article: 40

Oxygen-driven nebulization of *Clostridium butyricum* prevents drug-resistant bacterial pneumonia




Research | Open access | 24 January 2026 | Article: 39


Microbial transglutaminase in food biotechnology: from biochemical mechanisms to industrial applications



N-Glycosylation of Band 3 is essential for the binding of *Ganoderma lucidum* immunomodulatory protein rLZ-8




Valorisation of spent mushroom substrate by secondary microbial fermentation




Mini Review | Open access | 20 January 2026 | Article: 30

Production of uncommon carotenoids and lipids by red yeasts utilizing agri-food residues and waste cooking oil




Research | Open access | 20 January 2026 | Article: 29

Priestia megaterium reduces hydrogen sulfide level in swine manure potentially through an assimilatory mechanism and microbiota modulation




Research | Open access | 19 January 2026 | Article: 28

Geographical traceability of bacteria based on a systematic stable isotope analysis method




Research | Open access | 19 January 2026 | Article: 27

Enhancing the heat tolerance of homolactic *Escherichia coli* through reverse metabolic engineering



Research | Open access | 17 January 2026 | Article: 26

Low-temperature two-stage probiotic fermentation enhances nutrition and safety of pig liquid feed




Research | Open access | 24 November 2025 | Article: 25

Metabolomic and functional profiling of *Lactiplantibacillus plantarum* strains reveals distinct probiotic, immunomodulatory, and antimicrobial signatures



Characterization of the oral microbiome in the diverse South African population by PacBio HiFi sequencing




Research | Open access | 14 January 2026 | Article: 23

Investigating soil and ATCC bacterial strains for their ability to synthesize anisotropic gold nanoparticles




Research | Open access | 12 January 2026 | Article: 22

Enhancing stability and safety of chimeric peptidoglycan hydrolases by linker engineering




Research | Open access | 12 January 2026 | Article: 21

Bacillus subtilis T7 encodes [FeFe]-hydrogenase and lignin-targeting LPMOs for consolidated bioprocessing




Research | Open access | 12 January 2026 | Article: 20

Investigation on physical and physiological properties of extracellular vesicles derived from Enterococcus faecalis




Research | Open access | 12 January 2026 | Article: 19

Distinct oral and fecal microbiota composition in preschool children with overweight/obesity: a cross-sectional study




Research | Open access | 12 January 2026 | Article: 18

Microalgae as a sustainable alternative to palm oil: fatty acid profiles under photoautotrophic and heterotrophic growth




Alternative malonyl-CoA pathways for microbial production of diverse products




Mini Review | Open access | 12 January 2026 | Article: 16

Characterization and genetic diversity of pseudomonads population from highbush blueberry in western Canada




Research | Open access | 12 January 2026 | Article: 15

Harnessing silica-coated silver nanoparticles for combating multidrug-resistant *Pseudomonas aeruginosa*




Research | Open access | 10 January 2026 | Article: 14

Addition of lactoferrin increases efficacy of three Kayviruses and limits the inflammatory response in pulmonary epithelial cells



Research | Open access | 10 January 2026 | Article: 13

Unlocking the Zn-enriching potential of industrial yeast strains—an experimental journey from metal analysis to proteomics



Research | Open access | 10 January 2026 | Article: 12



Applied Microbiology and Biotechnology

Publishing model: Open access

[Explore open access funding](#) [Select institution](#)

▼ Journal menu

[Search all Applied Microbiology and Biotechnology articles](#) →




Volume 110, Issue 1

December 2026


261 articles in this issue

The phage shock protein response of *Listeria monocytogenes* influences tolerance to the multi-peptide bacteriocin garvicin KS




Research | Open access | 10 January 2026 | Article: 11

Enzymatic galactosylation of phenolic compounds by permeabilized cells of *Pantoea anthophila*




Research | Open access | 10 January 2026 | Article: 10

Sustainable recovery from pig slurry using ionic liquid microbial fuel cells and microalgae consortia




Toward food-grade production of the *Bacteroides helcogenes* protein-glutamine glutaminase with an optimized *Bacillus subtilis* strain



Research | Open access | 10 January 2026 | Article: 8

Chb and nag genes drive *N,N'*-diacetylchitobiose metabolism in probiotic *Lactacaseibacillus paracasei*



Research | Open access | 10 January 2026 | Article: 7

Genome sequence of *Halovibrio* sp. HP20-59 as a promising polyhydroxybutyrate producer




Research | Open access | 10 January 2026 | Article: 6

Precision hydrolysis: tailored yeast processing enzymes for yeast-based products




Mini Review | Open access | 06 January 2026 | Article: 5

A newly isolated *Streptomyces nigra* strain for the biotechnological production of melanin



Research | Open access | 06 January 2026 | Article: 4

Cosolvent-induced spontaneous refolding of lipase




Research | Open access | 06 January 2026 | Article: 3

Engineering galactoside acetyltransferase for enhanced hesperetin-7-O-glucoside bioavailability



Redefining media blending mathematically: a systematic approach for screening of medium components





Applied Microbiology and Biotechnology

Germany | Universities and research institutions | Media Ranking

Country

Germany



SCIMAGO
INSTITUTIONS
RANKINGS



SCImago
Media Rankings

Subject Area and Category

Biochemistry, Genetics
and Molecular Biology
Biotechnology

Immunology and
Microbiology

Applied Microbiology
and Biotechnology

Medicine

Medicine
(miscellaneous)

Publisher

Springer Science and
Business Media
Deutschland GmbH

SJR 2025

1.008

H-Index

Q1 296

Publication type

Journals

ISSN

01757598, 14320614

Coverage

1984-2026

Information

[Home](#)

[How to publish in this journal](#)


AMBoffice@gmx.de

Scope

Mini-Reviews are short reviews critically summarizing the current state of and providing perspectives on methods and equipment. I ology and biotechnology ed and in

pipe technologies of biotechnological processes using prokaryotic or eukaryotic cells or enzymes and other proteins for production, conversion, degradation and detection of substances including bioreactor design. Biotechnologically relevant enzymes and proteins Characterization of hitherto undescribed, biotechnologically relevant enzymes and proteins or of known proteins with novel properties and purification of proteins on a technical scale. Bioenergy and biofuels Biotechnological processes for production of fuels, combustible chemicals and other forms of energy. Applied genetics and molecular biotechnology New vectors and methods for genetic transformation and all aspects of strain improvement employing recombinant DNA technology including metabolic engineering and enzyme engineering. Genomics, transcriptomics, proteomics Characterization and analyses of genomes, transcriptomes and proteomes. Applied microbial and cell physiology Physiological studies on biosynthesis, catabolism and biotransformation of biotechnologically relevant compounds. Environmental biotechnology All aspects of environmental processes using microorganisms and biodegradation of hazardous chemicals. Methods and protocols Novel methods and techniques or significant modifications of already described methods or new combinations of methods yielding significant improvements. Their novelty and reliability must be concisely presented.

Quartiles



Tersedia di mana saja.
Di area pedesaan
dan daerah terpencil sekalipun.

DIRANCANG OLEH **SPACEX**
Kecepatan tidak dijamin dan didasarkan pada metrik P80 nasional di Indonesia. Ketersediaan dan harga di setiap lokasi mungkin berbeda.

STARLINK

Find similar journals

All quartiles

All countries

All subject categories

Clear filters

Download

1 - AMB Express

90%

similarity

2 - World Journal of Microbiology and Biotechnology

84%

similarity


3 - Biotechnology Letters

79%


similarity

< >

SJR




Total Documents






External Cites per Doc




Cites per Doc



% International Collaboration









Documents related to SDGs (UN)

Coming soon



← Show this widget in your own website

Just copy the code below and paste within your html code:

```
<a href="https://www.scimago
```

SCImago Graphica

Explore, visually communicate and make sense of data with our [new data visualization tool](#).



Metrics based on Scopus® data as of March 2026

Tersedia di mana saja.

Di area pedesaan
dan daerah terpencil sekalipun.

DIRANCANG OLEH **SPACEX**

Kecepatan tidak dijamin dan didasarkan pada metrik P80 nasional di Indonesia. Ketersediaan dan harga di setiap lokasi mungkin berbeda.

STARLINK

Discover more



Scopus Database Access



Journal Ranking Reports



SCImago Lab Services



V

Veronica

4 years ago

what it is the impact factor of this journal according to SJR ?

reply



Melanie Ortiz

4 years ago

Dear Veronica, thank you very much for your comment. SCImago Journal and Country Rank uses Scopus data, our impact indicator is the SJR. We suggest you consult the Journal Citation Report for other indicators (like Impact Factor) with a Web of Science data source. Best Regards, SCImago Team

A

A.E.Z. Hasan

5 years ago

I will submit articles to your journal. Hopefully, the article will finish soon. best regards...aez hasan (biochemistry, IPB University)

reply



Melanie Ortiz

5 years ago

Dear Hasan, thanks for your participation! Best Regards, SCImago Team

Hello; Good morning. We have done a study - the isolation purification and characterization of a phytochemical which has anti-Pseudomonas activity against Pseudomonas aeruginosa isolated from used industrial coolant- Will this be appropriate for your journal ?

We have also standardised the procedure for using the extract in coolant to extend the life of the coolant which is the basis for an Indian patent application. Do we have a chance of publishing this in your esteemed journal.

reply



Melanie Ortiz

5 years ago

Dear Vinitha,

thank you for contacting us.

We are sorry to tell you that SCImago Journal & Country Rank is not a journal. SJR is a portal with scientometric indicators of journals indexed in Elsevier/Scopus. Unfortunately, we cannot help you with your request, we suggest you visit the journal's homepage (See scope and submission/author guidelines) or contact the journal's editorial staff , so they could inform you more deeply.

Best Regards, SCImago Team

D

Dudha Raj Malla Thakuri

6 years ago

i want to published artical

reply



Melanie Ortiz

6 years ago

Dear Dudha, thank you very much for your comment, we suggest you look for author's instructions/submission guidelines in the journal's website. Best

Regards, SCImago Team

Email



Message



I'm not a robot

reCAPTCHA

Send

The users of Scimago Journal & Country Rank have the possibility to dialogue through comments linked to a specific journal. The purpose is to have a forum in which general doubts about the processes of publication in the journal, experiences and other issues derived from the publication of papers are resolved. For topics on particular articles, maintain the dialogue through the usual channels with your editor.



Services

Journal Rankings
Country Rankings
Journal Value

About us

Who we are
Contact us

SCImago's products

SCImago Journal Country & Rank
SCImago Institutions Ranking
SCImago Media Ranking
SCImago Iber
SCImago Arab
SCImago Graphica
Ediciones Profesionales de la Información

All the information shown in the SCImago Journal Country & Rank website can be used for non-commercial purposes as long as it is cited.

How to cite:

SCImago, (n.d.). SJR-SCImago Journal Country & Rank [Portal]. Retrieved (Date), from <https://www.scimagojr.com>

Powered by:
Scopus®